

# Chryso splenol-C Increases Contraction by Augmentation of Sarcoplasmic Reticulum $\text{Ca}^{2+}$ Loading and Release via Protein Kinase C in Rat Ventricular Myocytes<sup>S</sup>

J. Wang, T.N. Trinh, A. T. V. Vu, J.C. Kim, A.T.N. Hoang, C.J. Ohk, Y.H. Zhang, C.M. Nguyen, and S.H. Woo

*Pathophysiology Laboratory, College of Pharmacy, Chungnam National University, Yuseong-gu, Daejeon, South Korea (J.W., T.N.T., A.T.V.V., S.H.W.); NEXEL Co., Ltd. 8F, 55 Magokdong-ro, Gangseo-gu, Seoul, South Korea (J.C.K.); Institute of Natural Products Chemistry, VAST, Hanoi, Vietnam (A.T.N.H., C.M.N.); and Department of Physiology and Biomedical Sciences, College of Medicine, Seoul National University, Jongro-gu, Seoul, South Korea (C.J.O., Y.H.Z.)*

Received July 8, 2021; accepted October 11, 2021

## ABSTRACT

Naturally found chryso splenol-C (4',5,6-trihydroxy-3,3',7-trimethoxyflavone) increases the contractility of cardiac myocytes independent of  $\beta$ -adrenergic signaling. We investigated the cellular mechanism for chryso splenol-C-induced positive inotropy. Global and local  $\text{Ca}^{2+}$  signals, L-type  $\text{Ca}^{2+}$  current ( $I_{\text{Ca}}$ ), and contraction were measured from adult rat ventricular myocytes using two-dimensional confocal  $\text{Ca}^{2+}$  imaging, the whole-cell patch-clamp technique, and video-edge detection, respectively. Application of chryso splenol-C reversibly increased  $\text{Ca}^{2+}$  transient magnitude with a maximal increase of ~55% within 2- to 3-minute exposures ( $\text{EC}_{50} \cong 21 \mu\text{M}$ ). This chemical did not alter  $I_{\text{Ca}}$  and slightly increased diastolic  $\text{Ca}^{2+}$  level. The frequency and size of resting  $\text{Ca}^{2+}$  sparks were increased by chryso splenol-C. Chryso splenol-C significantly increased sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$  content but not fractional release. Pretreatment of protein kinase C (PKC) inhibitor but not  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II (CaMKII) inhibitor abolished the stimulatory effects of chryso splenol-C on  $\text{Ca}^{2+}$  transients and  $\text{Ca}^{2+}$  sparks. Chryso splenol-C-induced positive inotropy was removed by the

inhibition of PKC but not CaMKII or phospholipase C. Western blotting assessment revealed that PKC- $\delta$  protein level in the membrane fractions significantly increase within 2 minutes after chryso splenol-C exposure with a delayed (5-minute) increase in PKC- $\alpha$  levels in insoluble membrane. These results suggest that chryso splenol-C enhances contractility via PKC (most likely PKC- $\delta$ )-dependent enhancement of SR  $\text{Ca}^{2+}$  releases in ventricular myocytes.

## SIGNIFICANCE STATEMENT

Study shows that chryso splenol-C, a natural flavone showing a positive inotropic effect, increases SR  $\text{Ca}^{2+}$  releases on depolarizations and  $\text{Ca}^{2+}$  sparks with an increase of SR  $\text{Ca}^{2+}$  loading but not L-type  $\text{Ca}^{2+}$  current in ventricular myocytes. Chryso splenol-C-induced enhancement in contraction is eliminated by PKC inhibition, and it is associated with redistributions of PKC to the membrane. These indicate that chryso splenol-C enhances contraction via PKC-dependent augmentations of SR  $\text{Ca}^{2+}$  release and  $\text{Ca}^{2+}$  loading during action potentials.

## Introduction

Chryso splenol-C (4',5,6-trihydroxy-3,3',7-trimethoxyflavone) is a flavonoid compound contained in the medicinal plants *Pterocaulon sphacelatum* (Asteraceae) (Semple et al., 1999) and *Milium balansae* (Huong et al., 2004). It is known that *M. balansae* in particular has therapeutic effects for gastropathy and glomerulonephropathy. We have previously demonstrated that chryso splenol-C increases cell shortening in rat ventricular myocytes ( $\text{EC}_{50}$  of ~45  $\mu\text{M}$ ; Son et al., 2011). The positive

inotropic effect of chryso splenol-C was reversible and resistant to the inhibitors for  $\beta$ -adrenergic receptor and protein kinase A (PKA) (Son et al., 2011). The cellular mechanism for the chryso splenol-C-induced positive inotropic effect remains to be determined.

The contraction of mammalian cardiac myocytes is controlled by a sequence of events called excitation-contraction coupling, which includes the L-type  $\text{Ca}^{2+}$  current ( $I_{\text{Ca}}$ )-triggered gating of the  $\text{Ca}^{2+}$  release channels, ryanodine receptors (RyRs) on the sarcoplasmic reticulum (SR) membrane, and the release of  $\text{Ca}^{2+}$  from the SR (Beuckelmann and Wier, 1988; Näbauer et al., 1989; Niggli and Lederer, 1990). Confocal  $\text{Ca}^{2+}$  imaging of cardiac myocytes has shown that there are elementary SR  $\text{Ca}^{2+}$  release events via RyRs (“ $\text{Ca}^{2+}$  sparks”) underlying cardiac excitation-contraction coupling that are activated spontaneously or by  $I_{\text{Ca}}$  (Cheng et al., 1993; Cannell et al., 1994; Shacklock et al.,

This work was supported by National Research Foundation of Korea (NRF) grants funded by the Korean Government (MEST) (2017R1E1A1A01074504).

No author has an actual or perceived conflict of interest with the contents of this article.

[dx.doi.org/10.1124/molpharm.121.000365](https://doi.org/10.1124/molpharm.121.000365).

<sup>S</sup> This article has supplemental material available at [molpharm.aspetjournals.org](http://molpharm.aspetjournals.org).

**ABBREVIATIONS:**  $\tau$ , decay time constant; CaMKII,  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II; 2-D, two-dimensional; FWHM, full-width at half-maximal amplitude; GAPDH, glyceraldehyde-3-phosphate dehydrogenase;  $I_{\text{Ca}}$ , L-type  $\text{Ca}^{2+}$  current; PDBu, phorbol dibutyrate; PKA, protein kinase A; PKC, protein kinase C; PLC, phospholipase C; RyR, ryanodine receptor; SR, sarcoplasmic reticulum.

1995). After the contractile elements are activated, the released  $\text{Ca}^{2+}$  is removed from the cytosol mainly by the SR  $\text{Ca}^{2+}$  pump and the forward mode of the  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchanger (Negretti et al., 1993; Bassani et al., 1994).

In addition to PKA, the other key protein kinases  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II (CaMKII) and protein kinase C (PKC) are well known to regulate cardiac  $\text{Ca}^{2+}$  signaling and myocytes' contractility. These kinases are often involved in the mode of action for positive inotropic agents in cardiac myocytes (Guo et al., 2006; Zhou et al., 2009; Ogrodnik and Niggli, 2010; Kim et al., 2015; Bovo et al., 2017; Steinberg, 2018). Although no role of PKA signaling in chryso splenol-C-induced positive inotropy has been previously reported (Son et al., 2011), the possible roles of these kinases in the action of chryso splenol-C remain to be examined. It is thought that  $\text{Ca}^{2+}$ -dependent CaMKII activation can modulate  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release via phosphorylation in a number of excitation-contraction coupling proteins, including L-type  $\text{Ca}^{2+}$  channels and RyRs (Maier and Bers, 2007). The phosphorylation of RyRs by CaMKII can sensitize RyRs to  $\text{Ca}^{2+}$ , resulting in an increased SR  $\text{Ca}^{2+}$  leak (Guo et al., 2006; Pereira et al., 2007). Activation of PKC signaling is involved in positive inotropy and  $\text{Ca}^{2+}$  modulation under the stimulations of  $\alpha_1$ -adrenergic and endothelin receptors in cardiac myocytes, in which there are also significant controversies (Capogrossi et al., 1991; Asai et al., 1996; Woo and Lee, 1999a,b; Braz et al., 2004; Puglisi et al., 2011; Smyrnias et al., 2018).

In the present study, we investigated the cellular mechanisms underlying the positive inotropy of chryso splenol-C by examining global and local  $\text{Ca}^{2+}$  signals,  $I_{\text{Ca}}$ , and contraction in isolated rat ventricular myocytes using confocal  $\text{Ca}^{2+}$  imaging, the whole-cell patch-clamp technique, and video edge detection, respectively. In addition, the possible role of protein kinases in the effects of chryso splenol-C has been examined. We find that chryso splenol-C significantly increases SR  $\text{Ca}^{2+}$  release upon depolarization, resting  $\text{Ca}^{2+}$  sparks, and SR  $\text{Ca}^{2+}$  content and that the stimulatory effects of chryso splenol-C on  $\text{Ca}^{2+}$  release sites and contraction are abolished by PKC inhibition. Immunoblotting analysis showed chryso splenol-C-induced redistribution of PKC subtypes from soluble to the particulate compartment within 2 minutes, which is presumed to represent the activation of PKC. Our data indicate that chryso splenol-C elicits a positive inotropic effect via PKC (presumably  $\delta$  subtype)-mediated augmentation in  $\text{Ca}^{2+}$  signaling in ventricular myocytes.

## Materials and Methods

**Single Cell Isolation.** Ventricular myocytes were isolated from male Sprague-Dawley rats (200–300 g) as described previously (Kim et al., 2015). Rats were deeply anesthetized with sodium pentobarbital (150 mg/kg, i.p.), the chest cavity was opened, and hearts were excised. This surgical procedure was carried out according to the guiding principles for the care and use of experimental animals published by the Korean Food and Drug Administrations and Animal and Plant Quarantine Agency in South Korea and approved by Animal Care and Use Committees of the Chungnam National University (CNU-00368). The excised hearts were retrogradely perfused at 7 ml/min through the aorta (at 36.6°C), first for 3 minutes with  $\text{Ca}^{2+}$ -free Tyrode's solution composed of (in mM) 137 NaCl, 5.4 KCl, 10 HEPES, 1  $\text{MgCl}_2$ , and 10 glucose (pH 7.3); then with  $\text{Ca}^{2+}$ -free Tyrode's solution containing collagenase (1.4 mg/ml, type 1; Roche) and protease (0.14 mg/ml, type XIV; Sigma) for 12 minutes; and finally with Tyrode's solution containing 0.2 mM  $\text{CaCl}_2$  for 5 minutes. The ventricles of the digested heart

were then cut into several sections and subjected to gentle agitation to dissociate the cells. The freshly dissociated cells were stored at room temperature in Tyrode's solution containing 0.2 mM  $\text{CaCl}_2$ .

### Two-Dimensional Confocal $\text{Ca}^{2+}$ Imaging and Image Analysis.

Isolated myocytes were loaded with 3  $\mu\text{M}$  fluo-4 acetoxy-methyl ester (Invitrogen, USA) for 30 minutes. The dye-loaded cells were continuously superfused with 2 mM  $\text{Ca}^{2+}$ -containing normal Tyrode's solution (see above; pH 7.4). Intracellular  $\text{Ca}^{2+}$  fluorescence was imaged in 2-D using a laser scanning confocal imaging system (A1, Nikon, Japan) attached to an inverted microscope (Eclipse Ti, Nikon) fitted with a  $\times 60$  oil-immersion objective lens (Plan Apo, Numerical Aperture 1.4, Nikon). Dyes were excited at 488 nm using Ar ion laser (Ommichrome), and fluorescence emission at  $>510$  nm was detected. Images were recorded by NIS Elements AR software (v3.2, Nikon). To record global  $\text{Ca}^{2+}$  transients, cells were stimulated at 1 Hz using a pair of Pt electrodes connected to a stimulator (D-7806, Hugo Sachs Elektronik, Germany), and  $\text{Ca}^{2+}$  images were recorded at 60 or 120 Hz. To estimate the magnitudes of  $\text{Ca}^{2+}$  transients, the average resting fluorescence intensity ( $F_0$ ) was calculated from several frames immediately before electrical stimulation, and then tracings of global  $\text{Ca}^{2+}$  signals were shown as the average fluorescence of each area normalized relative to the  $F_0$  ( $F/F_0$ ).

To measure spark frequency,  $\text{Ca}^{2+}$  images were recorded at 30 Hz in 2-D, which allowed us to monitor the major part of the cell and compensate for the scarcity of resting  $\text{Ca}^{2+}$  sparks. Recording of spontaneous  $\text{Ca}^{2+}$  sparks was normally preceded by a train of electrical pulses at 1 Hz. Under this condition, the frequency of spontaneous sparks and SR  $\text{Ca}^{2+}$  content were stable during the experimental period.  $\text{Ca}^{2+}$  sparks were identified by a computerized algorithm in the "RealTimeMicroscopy" PC program as previously described (Kim et al., 2015). To calculate the frequency of  $\text{Ca}^{2+}$  sparks, the area of cell image was measured using the NIS Elements AR software (v3.2, Nikon). The focal  $\text{Ca}^{2+}$  releases were subjected to Gaussian approximations as previously described using the PC program "RealTimeMicroscopy" (Kim et al., 2015), which allowed routine measurements of the amplitude, width, and area of sparks (Fig. 4). Duration of focal  $\text{Ca}^{2+}$  releases was estimated as the time-to-peak area (Woo et al., 2003).

**Measurement of  $I_{\text{Ca}}$ .**  $I_{\text{Ca}}$  was recorded in the whole-cell configuration of the patch-clamp technique (Hamill et al., 1981) using an EPC7 amplifier (HEKA, Germany). The patch pipettes were made of glass capillaries (Kimble Glass Inc.) to have a resistance of 2–3 M $\Omega$  when filled with the internal solution containing (in mM) 110 CsCl, 20 TEA-Cl, 20 HEPES, 5 MgATP, and 15 EGTA, with the pH adjusted to 7.2 with CsOH. Outward  $\text{K}^+$  currents were suppressed by replacing internal  $\text{K}^+$  with Cs $^+$  and TEA $^+$ , and inward rectifier  $\text{K}^+$  current was suppressed by replacing external  $\text{K}^+$  with Cs $^+$ .  $\text{Na}^+$  current was inactivated by holding the membrane potential at  $-40$  mV. Trains of test pulses were to 0 mV for 120 milliseconds at 0.1 Hz. The measurement of  $I_{\text{Ca}}$  was carried out 7–8 minutes after rupture of the membrane with the patch pipette. Generation of voltage protocols and acquisition of data were carried out using pCLAMP (9.0, Molecular Devices) combined with an analog-to-digital converter (Digidata 1322, Molecular Devices). The series resistance was 1.5–5 times the pipette resistance and was electronically compensated through the amplifier. The current signals were digitized at 10 kHz. The percent suppression of  $I_{\text{Ca}}$  by interventions was evaluated after a gradual decrease in  $I_{\text{Ca}}$  by rundown was subtracted from the raw current. Peak detection was performed with Clampfit (9.0, Molecular Devices), and the time constant ( $\tau$ ) of inactivation of  $I_{\text{Ca}}$  was obtained with single exponential curve fitting using the equation:  $y = (A_i - A_f) \cdot \exp(-t/\tau) + A_f$ , wherein  $A_i$  and  $A_f$  are the initial ( $t = 0$ ) and final ( $t = \text{infinity}$ ) values of the parameter, and  $\tau$  is a time constant of exponential decay. Curve fitting was performed using OriginPro 8 SR0 software (OriginLab Corporation).

**Measurement of Cell Shortening.** Isolated myocytes were continuously superfused with normal Tyrode's solution containing 2 mM  $\text{Ca}^{2+}$ . Cells were field-stimulated with two paralleled Pt wires connected with an electrical stimulator (Stimulator I Hugo Sach

Elektronik, March-Hugstetten, Germany) at 1 Hz. Single cell shortening was detected with a video edge detector (Model VED-105; Crescent Electronics, Sandy, UT) connected with a CCD camera (LCL902C; Till Photonics, Graefelfing, Germany) and video monitor (Polychrome V system; Till Photonics) (Son et al., 2011). Signals from the edge detector were digitized by Digidata (1440A; Molecular Devices, Sunnyvale, CA) and then recorded with pClamp program (v10.3, Molecular Devices).

**Subcellular Fractionation and Western Blot Analysis.** To examine protein expression levels of PKC isoforms, rat ventricular myocytes were collected and treated with chrysosplenol-C (80  $\mu\text{M}$ ) for 2, 5, and 30 minutes. Cells were resuspended in lysis buffer without SDS (10 mM Tris-HCl, 1 mM phenylmethanesulfonyl fluoride, 1 mM  $\text{Na}_3\text{VO}_4$ , 0.5  $\mu\text{M}$  NaF, and protease inhibitors including 0.2  $\mu\text{g/ml}$  pepstatin, 1  $\mu\text{g/ml}$  aprotinin, 0.5  $\mu\text{g/ml}$  leupeptin, pH 7.4), sonicated five times by using a sonicator (Sonopuls, Berlin, Germany) for 2 seconds, and then centrifuged for 30 minutes at 100,000 g to obtain the cytosol. The resulting pellet was resuspended with lysis buffer containing 1% (w/v) SDS and 1% Triton-X 100 and incubated in ice for 30 minutes. Then, the sample was centrifuged at 100,000 g for 30 minutes. The supernatant was the Triton-soluble membrane fraction, and the pellet was considered to be the Triton-insoluble fraction. Protein concentrations were measured by BCA protein assay (Thermo Fisher Scientific, 23227). Approximately 25  $\mu\text{g}$  of proteins was run on 10% SDS-polyacrylamide gel. The proteins were transferred onto a nitrocellulose membrane, and the blots were sequentially probed with primary and secondary antibodies (anti-PKC- $\alpha$ , mouse monoclonal, 1:500, an epitope at the C terminus, sc-8393, Santa Cruz Biotechnology; anti-PKC- $\delta$ , mouse monoclonal, 1:500, an epitope at the C terminus, sc-8402, Santa Cruz Biotechnology; anti-PKC- $\epsilon$ , mouse monoclonal, 1:500, an epitope at the C terminus, sc-1681, Santa Cruz Biotechnology; anti- $\text{Na}^+\text{-K}^+$  ATPase  $\alpha 2$ , rabbit polyclonal, 1:500, an epitope at an N terminal, 07-674, Merck; anti-GAPDH, mouse monoclonal, 1:1000, ab8245, Abcam; secondary antibodies: mouse anti-rabbit IgG-horseradish peroxidase, 1:5000, sc-2357, Santa Cruz Biotechnology; goat anti-mouse IgG-horseradish peroxidase, 1:5000, sc-2005, Santa Cruz Biotechnology) using standard Western blot protocol. All blots were imaged using a ChemiDoc XRS densitometer (Bio-Rad) and quantified by Image J program.

**Solutions and Reagents.** Chrysosplenol-C was isolated from air-dried and ground leaves and branches of *M. balansae* as previously described (Son et al., 2011). The purity of chrysosplenol-C was  $98.1 \pm 0.99\%$  (three batches). Detailed physicochemical and structural information on chrysosplenol C has been reported previously (Son et al., 2011). Stock solutions of chrysosplenol-C (500 mM) were made in DMSO, and it was diluted in Tyrode solution for testing (DMSO  $\leq 0.08\%$  (v/v), e.g., 0.01% DMSO at 50  $\mu\text{M}$  chrysosplenol-C solutions). Same concentration of DMSO was added to Tyrode solutions without (control solutions and inhibitor-containing solutions) or with chrysosplenol-C. The drug solutions were applied to the cells by superfusion. The experiments were performed at room temperature (22–25°C).

Caffeine, chelerythrine, KN-92 (E)-N-(2-((3-(4-chlorophenyl)allyl)(methylamino) methyl)phenyl)-4-methoxybenzenesulfonamide), and KN-93 (2-[N-(2-hydroxyethyl)-N-(4-methoxybenzenesulfonyl)amino-N-(4-chlorocinnamyl)-N-methylbenzylamine) were purchased from Sigma-Aldrich (St. Louis, MO). GF109203X (bisindolylmaleimide I), U73122, and U73343 were supplied by Tocris Bioscience (Avonmouth, Bristol, BS11 9QD). Fluo-4 acetoxymethyl was from Thermo Fisher Scientific (Waltham MA).

**Statistics.** The numerical results are presented as mean  $\pm$  S.D. *n* indicates number of cells used. Paired or unpaired Student's *t* tests were used for statistical comparisons of most of the functional data depending on the experiments in single group. For comparison of unitary properties of single sparks, nonparametric Mann-Whitney test was performed (Fig. 4). For the Western blot data, statistical significance among the groups was determined using two-way ANOVA-related measures, with post hoc testing to control for multiple

comparisons. *p* values were adjusted for multiple comparisons using Bonferroni correction (see figure legends). The statistical significance was defined using a new *p* value threshold of 0.05/(number of tests) according to Bonferroni correction.


## Results

Chrysosplenol-C reversibly and stably enhances ventricular cell shortening (Son et al., 2011). To determine the cellular mechanism for the chrysosplenol-C-induced positive inotropic effect, we examined whether chrysosplenol-C affects  $\text{Ca}^{2+}$  releases from the SR during depolarizations. Figure 1, A and B shows that chrysosplenol-C (50  $\mu\text{M}$ ) reversibly increases cytosolic  $\text{Ca}^{2+}$  releases during depolarizations. The maximal effect of 50  $\mu\text{M}$  chrysosplenol-C was observed at about 2–3 minutes after exposure. Diastolic  $\text{Ca}^{2+}$  levels were also slightly but significantly increased by chrysosplenol-C (Fig. 1C). The magnitudes of  $\text{Ca}^{2+}$  transients were increased by chrysosplenol-C in a concentration-dependent manner with an  $\text{EC}_{50}$  of  $21 \pm 3.3 \mu\text{M}$  (Fig. 1D). Maximal effect on the  $\text{Ca}^{2+}$  transient magnitude (by  $\cong 58\%$  increase) by chrysosplenol-C was achieved at  $\sim 80 \mu\text{M}$ . Concentration dependence of the effects of chrysosplenol-C on  $\text{Ca}^{2+}$  transients was slightly lower than that of its positive inotropic effects ( $\cong 45 \mu\text{M}$ ) that were previously observed in rat ventricular myocytes (Son et al., 2011). The decay speed of  $\text{Ca}^{2+}$  transients was not altered by the application of chrysosplenol-C (Fig. 1E). These results suggest that chrysosplenol-C increases depolarization-induced SR  $\text{Ca}^{2+}$  release, thereby causing an increase in contraction.

$\text{Ca}^{2+}$  releases on depolarization are controlled by the  $\text{Ca}^{2+}$  influx through L-type  $\text{Ca}^{2+}$  channels (Beuckelmann and Wier, 1988; Näbauer et al., 1989; Niggli and Lederer, 1990). To understand the mechanism for the enhanced  $\text{Ca}^{2+}$  transients with the treatment of chrysosplenol-C, we next examined the effects of chrysosplenol-C on  $I_{\text{Ca}}$ . Figure 2 shows the effects of 50- and 80- $\mu\text{M}$  chrysosplenol-C on the whole-cell  $I_{\text{Ca}}$ . The  $I_{\text{Ca}}$  was not significantly altered by either concentration of chrysosplenol-C (Fig. 2A). We only observed a gradual rundown during the application of chrysosplenol-C, with no changes in its current-voltage relationship (Fig. 2B) or inactivation time constants (Fig. 2C). This result suggests that chrysosplenol-C-induced enhancement in  $\text{Ca}^{2+}$  transients may be mediated by  $I_{\text{Ca}}$ -independent mechanisms.

An increased SR  $\text{Ca}^{2+}$  release in the presence of chrysosplenol-C may be reflective of a higher propensity of RyR clusters to activate at a given  $\text{Ca}^{2+}$  concentration through changes in their  $\text{Ca}^{2+}$  sensitivity. The occurrence and size of  $\text{Ca}^{2+}$  sparks at rest represent the changes in  $\text{Ca}^{2+}$  sensitivity and altered properties of  $\text{Ca}^{2+}$  release through the  $\text{Ca}^{2+}$  release sites. Therefore, we assessed the effects of chrysosplenol-C on the spatiotemporal properties of spontaneous  $\text{Ca}^{2+}$  sparks. The treatment of chrysosplenol-C significantly increased the spark frequency (events/ $10^3 \mu\text{m}^2 \cdot \text{s}$ ) in a concentration-dependent manner ( $\text{EC}_{50} = 0.79 \pm 0.24 \mu\text{M}$ ; Fig. 3). At 50  $\mu\text{M}$ , it increased the spark frequency by approximately 2-fold (Fig. 3B).

In the next series of experiments, the unitary properties of  $\text{Ca}^{2+}$  sparks were measured before and after the application of chrysosplenol-C using 2-D  $\text{Ca}^{2+}$  imaging at 240 Hz. Representative images of spark growth and dissipation in the presence and absence of chrysosplenol-C (50  $\mu\text{M}$ ) are shown in Fig. 4. A Gaussian fitting to measure the time course of spark amplitude, width, and area (Fig. 4B; see Materials and Methods) showed




**Fig. 1.** Enhancement of  $\text{Ca}^{2+}$  transients by chryso splenol-C in rat ventricular myocytes. (A) 2-D confocal  $\text{Ca}^{2+}$  images recorded (120 Hz) from a representative myocyte at the time points marked by Roman numbers with arrowheads in the  $\text{Ca}^{2+}$  transient traces in (B). (B)  $\text{Ca}^{2+}$  transients measured from the series of confocal  $\text{Ca}^{2+}$  images before and after 2-minute chryso splenol-C (CP-C) (50  $\mu\text{M}$ ) exposure. (C) Summary of the effects of CP-C (50  $\mu\text{M}$ ) on the  $\text{Ca}^{2+}$  transient magnitude and diastolic  $\text{Ca}^{2+}$  level ( $n = 8$ ).  $p$  values obtained from paired  $t$  tests were written above the bar graphs. (D) Concentration-response curve for the stimulatory effect by CP-C on the  $\text{Ca}^{2+}$  transients (% increase) (1  $\mu\text{M}$ ,  $n = 5$ ; 10  $\mu\text{M}$ ,  $n = 5$ ; 50  $\mu\text{M}$ ,  $n = 23$ ; 80  $\mu\text{M}$ ,  $n = 8$ ; 200  $\mu\text{M}$ ,  $n = 5$ ). The curve was approximated by the Hill equation ( $y = y_{\text{start}} + (y_{\text{end}} - y_{\text{start}})(x^n/(k^n + x^n))$ ).  $\text{EC}_{50}(k) = 21 \pm 3.3 \mu\text{M}$ .  $\text{Max} = 57.9 \pm 8.59\%$ . (E) Upper: superimposed  $\text{Ca}^{2+}$  transient traces recorded in the absence and presence of 50  $\mu\text{M}$  CP-C after normalization to their peaks. Lower: summary of the effects of CP-C (50  $\mu\text{M}$ ) on the decay time constant ( $n = 8$ ). Considering the new  $p$  value after the Bonferroni correction,  $p$  value less than 0.017 (0.05/3  $t$  tests; equivalent to old 0.05) was considered to be significant. Con, control.

that  $\text{Ca}^{2+}$  spark amplitude slightly increased with no statistical significance after chryso splenol-C treatment (control: median = 1.64, 90% range = 1.00–2.71; chryso splenol-C: median = 1.76, 90% range = 0.88–2.15;  $P = 0.041$ ), whereas the spark width (FWHM, measured at peak area) was unaltered by chryso splenol-C (control: median = 1.64  $\mu\text{m}$ , 90% range = 1.01–2.23  $\mu\text{m}$ ; chryso splenol-C: median = 1.68  $\mu\text{m}$ , 90% range = 0.93–2.24  $\mu\text{m}$ ;  $P = 0.759$ ; Fig. 4, B and C). The peak area of  $\text{Ca}^{2+}$  sparks was significantly increased by chryso splenol-C (control: median = 3.98  $\mu\text{m}^2$ , 90% range = 1.47–8.56  $\mu\text{m}^2$ ; chryso splenol-C: median = 5.10  $\mu\text{m}^2$ , 90% range = 1.44–9.32  $\mu\text{m}^2$ ;  $P = 0.0056$ ). The release durations for individual sparks were not significantly increased by chryso splenol-C (control: median = 8 milliseconds, 90% range = 4–12 milliseconds; chryso splenol-C: median = 8 milliseconds, 90% range = 4–16 milliseconds;  $P = 0.049$ ) (Fig. 4, B and C). These results suggest that chryso splenol-C increases the amount of  $\text{Ca}^{2+}$  released from the release sites.

An increase of  $\text{Ca}^{2+}$  in the lumen of the SR can enhance not only the frequency of  $\text{Ca}^{2+}$  sparks but also the effectiveness of a given  $\text{Ca}^{2+}$  current to activate SR  $\text{Ca}^{2+}$  release (Han et al., 1994; Bassani et al., 1995; Janczewski et al., 1995; Satoh et al., 1997; Györke and Györke, 1998). Therefore, we further examined whether chryso splenol-C affects SR  $\text{Ca}^{2+}$  loading by

measuring caffeine (10 mM)-induced  $\text{Ca}^{2+}$  transients. The magnitudes of caffeine-induced  $\text{Ca}^{2+}$  transients were increased by 50  $\mu\text{M}$  chryso splenol-C (for 2–3 minutes) to ~135% with increases of  $\text{Ca}^{2+}$  transient magnitudes (Fig. 5, A and B). The fractional release representing the amount of  $\text{Ca}^{2+}$  release relative to SR  $\text{Ca}^{2+}$  content (Bassani et al., 1995) was further calculated as the ratio of the  $\text{Ca}^{2+}$  transient magnitude during electrical stimulation relative to the magnitude of caffeine-induced  $\text{Ca}^{2+}$  transients. The fractional release was not significantly changed by chryso splenol-C (Fig. 5B). The  $\tau$  of caffeine-induced  $\text{Ca}^{2+}$  transients were not significantly altered by chryso splenol-C (Fig. 5B, right). These results suggest that enhancements in  $\text{Ca}^{2+}$  spark occurrence and  $\text{Ca}^{2+}$  transients in the presence of chryso splenol-C may be caused by an increase in SR  $\text{Ca}^{2+}$  content.


Chryso splenol-C-induced positive inotropy has been demonstrated to be independent of  $\beta$ -adrenoceptor–PKA signaling (Son et al., 2011). We further examined whether other major kinases, including PKC and CaMKII, play a role in the enhancement of  $\text{Ca}^{2+}$  transients and  $\text{Ca}^{2+}$  sparks in the presence of chryso splenol-C. Interestingly, when the PKC was suppressed by the treatment of chelerythrine (2  $\mu\text{M}$ , 5–10 minute), the application of chryso splenol-C no longer increased




**Fig. 2.** No effect by chrysosplenol-C on  $I_{Ca}$ . (A) Left: superimposed  $I_{Ca}$  recorded in the absence and presence of 50 and 80  $\mu$ M chrysosplenol-C (CP-C) in a representative rat ventricular myocyte.  $I_{Ca}$  was recorded on depolarizing pulses from -40 to 0 mV at 0.1 Hz. Right: comparison of averaged  $I_{Ca}$  recorded under control condition and after the application of 50 and 80  $\mu$ M CR-C ( $n = 8$ ). (B) Averaged current-voltage relationships obtained in the absence and presence of 50 and 80  $\mu$ M CP-C ( $n = 6$ ). The currents were recorded during the voltage steps from -40 to +40 mV with 10-mV increment. Holding potential was -40 mV. (C) Averaged inactivation time constants of  $I_{Ca}$  measured under the control conditions and after the application of 50 and 80  $\mu$ M CP-C ( $n = 8$ ). The  $p$  values obtained from paired  $t$  tests were written above the bar graphs. con, control.

$Ca^{2+}$  transients (Fig. 6, A and B). Chelerythrine alone did not significantly alter the  $Ca^{2+}$  transients (Fig. 6, A and B) or SR  $Ca^{2+}$  content (Supplemental Fig. 1). Since PKC can activate CaMKII (Waxham and Aronowski, 1993), the stimulatory effects of chrysosplenol-C may be mediated by CaMKII, activated by PKC. To test this possibility, we examined the effects

of chrysosplenol-C on  $Ca^{2+}$  transient in the presence of KN-93, the CaMKII inhibitor. Previous reports have shown that KN-93 at a concentration of 1  $\mu$ M successfully eliminates CaMKII-dependent cardiac  $Ca^{2+}$  response (Lu et al., 2020). When this protocol to suppress CaMKII was used in rat ventricular myocytes, chrysosplenol-C (50  $\mu$ M) slightly increased



**Fig. 3.** Enhancement of  $Ca^{2+}$  spark occurrence by chrysosplenol-C in rat ventricular myocytes. (A) Upper: 2-D confocal  $Ca^{2+}$  images recorded for the period marked by the boxes in the lower panel. Lower: time course of  $Ca^{2+}$  spark occurrence per frame for 2-second recording period before and after treatment of chrysosplenol-C (CP-C; 50  $\mu$ M, 2 minutes). (B) Comparison of mean spark frequency measured in the absence and presence of CP-C (50  $\mu$ M).  $***P < 0.001$  vs. control,  $###P < 0.001$  vs. CP-C ( $n = 9$ ; paired  $t$  test). The  $p$  values obtained from paired  $t$  tests were written above the bar graphs. (C) Concentration-response (% increase) curve for the stimulatory effects by CP-C on the resting spark frequency (0.01  $\mu$ M, 0.1  $\mu$ M, and 1  $\mu$ M;  $n = 5$ ; 10, 50 and 80  $\mu$ M;  $n = 8$ ). The curve was approximated by the Hill equation ( $y = V_{max} \cdot x^n / (k^n + x^n)$ ).  $EC_{50}(k) = 0.79 \pm 0.24 \mu$ M. Hill coefficient ( $n$ ) =  $0.57 \pm 0.06$ . Con, control; Rec, recovery.




**Fig. 4.** Effects of chrysofenol-C on the unitary properties of  $\text{Ca}^{2+}$  sparks. (A) 2-D confocal  $\text{Ca}^{2+}$  images for representative  $\text{Ca}^{2+}$  sparks recorded under control condition and after 2-minute application of chrysofenol-C (CP-C) ( $50 \mu\text{M}$ ). Images were recorded at 240 Hz. (B) Time courses of amplitude, full-width at half-maximal amplitude (FWHM) and area of  $\text{Ca}^{2+}$  spark images shown in the (A) in the absence and presence of CP-C. (C) Distribution histograms for the peak amplitude, FWHM at the time showing peak area, and time-to-peak of area (release duration) of individual  $\text{Ca}^{2+}$  sparks in the absence and presence of CP-C ( $50 \mu\text{M}$ ). The Mann-Whitney test was used to evaluate the significant difference between control sparks ( $n = 81$ ) and sparks recorded in the presence of CP-C ( $n = 115$ ). Considering the new  $p$  value after the Bonferroni correction,  $p$  value less than 0.0125 (0.05/4  $t$  tests; equivalent to old 0.05) was considered to be significant.


$\text{Ca}^{2+}$  transients with no statistical difference ( $16 \pm 2.2\%$ ,  $n = 9$ , Fig. 6, C and D). Diastolic  $\text{Ca}^{2+}$  level was not altered by chrysofenol-C in the presence of chelerythrine or KN-93 (Table 1). There were no significant changes in the decay time constants of the  $\text{Ca}^{2+}$  transients recorded in the presence of these blockers with and without chrysofenol-C (Table 1).

We next tested whether chelerythrine or KN-93 alters the stimulatory effects of chrysofenol-C on the spark frequency.

Treatment with chelerythrine ( $2 \mu\text{M}$ ) alone did not significantly change the spark frequency (Fig. 6, E and F), whereas KN-93 ( $1 \mu\text{M}$ ) alone significantly decreased it in the resting ventricular myocytes (by  $56 \pm 8.8\%$ ,  $n = 8$ ,  $P < 0.01$ ; Fig. 6, G and H). Note that KN-93 alone at the concentrations used significantly reduced the caffeine-induced  $\text{Ca}^{2+}$  releases (Supplemental Fig. 1). In the myocytes preincubated with chelerythrine, chrysofenol-C failed to enhance spark occurrence



**Fig. 5.** Increase in SR  $\text{Ca}^{2+}$  loading with similar fractional release in the presence of chrysofenol-C (CP-C). (A)  $\text{Ca}^{2+}$  transients on depolarizations followed by 10 mM caffeine (caff)-induced  $\text{Ca}^{2+}$  transient measured in the same rat ventricular myocyte in the absence [control (Con)] and presence of CP-C ( $50 \mu\text{M}$ , 3 minutes). (B) Mean magnitude of caffeine-induced  $\text{Ca}^{2+}$  releases representing SR  $\text{Ca}^{2+}$  content (left), mean fractional release (middle), and mean  $\tau$  of caffeine-induced  $\text{Ca}^{2+}$  transients (right), measured before and after application of CP-C in rat ventricular myocytes ( $n = 8$ ; paired  $t$  test). Considering the new  $p$  value after the Bonferroni correction,  $p$  value less than 0.017 (0.05/3  $t$  tests; equivalent to old 0.05) was considered to be significant.



**Fig. 6.** Major role of PKC in the stimulatory effects by chrysosplenol-C (CP-C) on Ca<sup>2+</sup> transients and sparks. (A and C) Representative Ca<sup>2+</sup> transients sequentially recorded in a same rat ventricular myocyte exposed to the PKC inhibitor chelerythrine (2 μM) (A) or 1 μM KN-93, the CaMKII inhibitor (C) followed by an additional application of 50 μM CP-C for 3 minutes. (B and D) Summary of the effects of chelerythrine (Chel; *n* = 8, B) or KN-93 (D; *n* = 9) without and with CP-C on the magnitude of Ca<sup>2+</sup> transients. (E and G) Upper: representative sequential confocal Ca<sup>2+</sup> images selected during the period marked with boxes in the time courses below each series of images (lower panel) showing spontaneous Ca<sup>2+</sup> sparks in rat ventricular myocytes in the control solutions and after application of chelerythrine (E; 2 μM) or KN-93 (G; 1 μM) without and with CP-C (50 μM). Arrowheads indicate Ca<sup>2+</sup> sparks. Lower: time course of spark occurrence during 2-second-long imaging at 30 Hz in the cell shown above. (F and H) Summary of the effects of CP-C in the presence of 2 μM Chel (F; *n* = 8) or 1 μM KN-93 (H; *n* = 8). Considering the new *p* value after the Bonferroni correction, *p* value less than 0.017 (0.05/3 paired *t* tests; equivalent to old 0.05) was considered to be significant. Con, control.

(Fig. 6, E and F). However, in the presence of KN-93, chrysosplenol-C (50 μM) tended to increase the spark occurrence (by 32 ± 5.2%, *n* = 8; Fig. 6, G and H). It should be noted that the effect of chrysosplenol-C on the spark frequency was smaller in the presence of KN-93 compared with that under control conditions (compare with Fig. 3B; *P* < 0.01). These results suggest that PKC may play a key role in the enhancement of spark frequency and Ca<sup>2+</sup> transients in the presence of chrysosplenol-C and that CaMKII may also partly contribute to the enhancement of Ca<sup>2+</sup> transients and sparks under the control of PKC.

Next, we confirmed whether chrysosplenol-C-induced positive inotropic effect is mediated by PKC in rat ventricular myocytes. The application of chrysosplenol-C increased cell shortening by ~2-fold at the concentrations of 50 μM (Fig. 7A). This chrysosplenol-C-mediated positive inotropic effect was completely removed by preincubation of either chelerythrine or GF109203X, which are PKC inhibitors (Fig. 7, D and G). Consistent with the Ca<sup>2+</sup> data, the pretreatment of KN-93 (1 μM) tended to suppress the positive inotropic effect of chrysosplenol-C (Fig. 7, B and G), whereas its inactive analog KN-92 did not suppress the chrysosplenol-C-induced positive inotropic effect (Fig. 7, C and G).

TABLE 1

Effects of KN-93 or chelerythrine on diastolic  $\text{Ca}^{2+}$  level and transient decay time constant in the absence and presence of chryso splenol-C in rat ventricular myocytes

Data represent mean  $\pm$  S.D. The number in the parenthesis indicates number of cells. Paired *t* tests were done. Considering the new *p* value after the Bonferroni correction, *p* value less than 0.0125 (0.05/4) was considered to be significant.

	Diastolic $[\text{Ca}^{2+}]$ (F/F <sub>0</sub> )	Decay ( $\tau$ , ms)	n
Control	1	156 $\pm$ 52	12
KN-93	1.02 $\pm$ 0.0045	160 $\pm$ 51	12
KN-93 + CP-C	1.07 $\pm$ 0.026*	162 $\pm$ 76	12
Control	1	165 $\pm$ 34	8
Chel	1.06 $\pm$ 0.013	168 $\pm$ 44	8
Chel + CP-C	1.05 $\pm$ 0.018	169 $\pm$ 54	8

Chel, chelerythrine (2  $\mu\text{M}$ ); CP-C, chryso splenol-C.

KN-93 itself but not KN-92 slightly reduced cell shortening with no significance (Fig. 7, B and C), which is somewhat consistent with its effect on  $\text{Ca}^{2+}$  transients (Fig. 6C). The inhibition of phospholipase C (PLC), which generates the PKC substrate diacylglycerol and inositol 1,4,5-trisphosphate, using U73122 did not significantly attenuate chryso splenol-C-mediated positive inotropic effect (Fig. 7, E and G). Its inactive analog U73343 also showed no significant effect on the chryso splenol-C-induced positive inotropy (Fig. 7, F and G), although both chemicals tended to decrease contraction. These results support the major role of PKC in chryso splenol-C-induced positive inotropy.

The next experiments were designed to determine whether chryso splenol-C alters the activity of PKC in these myocytes. For this purpose, we examined the effects of chryso splenol-C on the subcellular distribution of individual PKC isoforms. Immunoblot analyses with antibodies that detect the three major PKC isoforms (PKC- $\alpha$ , PKC- $\delta$ , and PKC- $\epsilon$ ) expressed in rat ventricular myocytes (Goldberg et al., 1997; Simonis et al., 2002) were performed on cytosolic and membrane fractions of rat ventricular myocytes. The membrane fraction was further divided into Triton-soluble and Triton-insoluble membrane components because scaffolding proteins that play an important role in signal transduction are known to be generally Triton-insoluble (Yan et al., 1996; Anderson, 1998). We confirmed the subcellular fractions by detecting well known cytosolic and membrane markers, GAPDH and  $\text{Na}^+$ - $\text{K}^+$  ATPase, respectively, using immunoblotting (Fig. 8). Figure 8 demonstrates that PKC- $\alpha$  and PKC- $\delta$  preferentially partition to the insoluble membrane fractions from 2 minutes to 5 minutes after exposure to chryso splenol-C (80  $\mu\text{M}$ ). Note that the level of PKC- $\delta$  in the insoluble membrane increased earlier compared with that of PKC- $\alpha$  and was maximized at 2 minutes after the drug exposure. The partition of PKC- $\alpha$  in the soluble membrane fraction was prominent after 30-minute exposure to chryso splenol-C with a gradual decrease in the level of cytosolic PKC- $\alpha$  (Fig. 8A), although the signal changes were not statistically significant (Fig. 8C). The level of PKC- $\delta$  in the soluble membrane was also increased from 2 minutes to 5 minutes after exposure to the drug (Fig. 8, A and C). The expression of PKC- $\epsilon$  was detected in the soluble and membrane fractions, but the partition of PKC- $\epsilon$  in the membrane fractions was not observed after the treatment of chryso splenol-C (Fig. 8, B and C). This result suggests that the activity of PKC- $\alpha$  and PKC- $\delta$  increases after chryso splenol-C treatment. The time course of redistribution of each isoform

suggests that PKC- $\delta$  may play a role in chryso splenol-C-induced positive inotropy.


## Discussion

Our data provide a cellular mechanism for the previously reported positive inotropy by chryso splenol-C in rat ventricular myocytes. We found that chryso splenol-C increases  $\text{Ca}^{2+}$  transient magnitude (Fig. 1) with no change in  $\text{Ca}^{2+}$  influx through the L-type  $\text{Ca}^{2+}$  channel (Fig. 2) and that this chemical enhances resting  $\text{Ca}^{2+}$  spark occurrence and spark amplitude and duration with a significant increase in the SR  $\text{Ca}^{2+}$  content (Figs. 3–5). The stimulatory effects of chryso splenol-C on  $\text{Ca}^{2+}$  transients and  $\text{Ca}^{2+}$  sparks were eliminated by PKC inhibition and partially suppressed by CaMKII suppression (Fig. 6). Consistently, the blockade of PKC abolished chryso splenol-C-induced positive inotropic effects (Fig. 7). Pharmacological data suggested that PLC may not play a role in the positive inotropic effect exerted by chryso splenol-C (Fig. 7). Immunoblotting in the subcellular fractions of rat ventricular myocytes demonstrated significant translocation/activation of PKC- $\delta$  by chryso splenol-C with a time course similar to that of the positive inotropic effect (time-to-peak effect = 2 minutes; Son et al., 2011) and a delayed PKC- $\alpha$  translocation to the membrane fractions (Fig. 8). Our data suggest that chryso splenol-C may elicit a positive inotropic effect by enhancing SR  $\text{Ca}^{2+}$  releases on depolarizations with increasing SR  $\text{Ca}^{2+}$  loading via the activation of specific PKC isoform independently of PLC.

It is well known that enhancements of resting  $\text{Ca}^{2+}$  spark occurrence depend on SR  $\text{Ca}^{2+}$  loading (Satoh et al., 1997). Larger SR  $\text{Ca}^{2+}$  loading in chryso splenol-C-treated myocytes compared with cells under control conditions (Fig. 5) may explain the increased  $\text{Ca}^{2+}$  spark frequency as well as the larger sparks observed in the presence of chryso splenol-C (Fig. 4). Due to larger  $\text{Ca}^{2+}$  loading in the SR in the presence of chryso splenol-C,  $\text{Ca}^{2+}$  release on depolarizations may also increase (Cheng et al., 1993; Bassani et al., 1995; Satoh et al., 1997). Previous reports support the notion that SR  $\text{Ca}^{2+}$  load correlates with  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release gain function in cardiac myocytes (Cheng et al., 1993; Bassani et al., 1995). Although both depolarization-induced  $\text{Ca}^{2+}$  release and SR  $\text{Ca}^{2+}$  content were increased by chryso splenol-C, the fractional  $\text{Ca}^{2+}$  release was not altered (Fig. 5B). This result also supports the notion that the stimulatory effects of chryso splenol-C on  $\text{Ca}^{2+}$  releases involve a mode of action distinct from  $\beta_1$ -adrenergic signaling (Son et al., 2011) because  $\beta_1$ -adrenergic stimulation augments the fractional release as well as SR  $\text{Ca}^{2+}$  content (Ginsburg and Bers, 2004). Consistently, neither the decay of  $\text{Ca}^{2+}$  transient nor  $I_{\text{Ca}}$  was significantly altered in the presence of chryso splenol-C (Figs. 1 and 2).

It should be noted that fractional  $\text{Ca}^{2+}$  release measured as intracellular total  $\text{Ca}^{2+}$  concentrations would be slightly different from that assessed as free  $\text{Ca}^{2+}$  concentrations (Bassani et al., 1995). We estimated fractional release by the measurements of cytosolic free  $\text{Ca}^{2+}$  using a nonlinear  $\text{Ca}^{2+}$  indicator (fluo-4) as previously reported (Kim et al., 2015). It is suggested that  $\text{Ca}^{2+}$  buffering power decreases with increasing intracellular  $\text{Ca}^{2+}$  level (Eisner et al., 2000). Therefore, given that the caffeine-evoked rise is larger than that produced by electrical stimulation, a bigger effect of intracellular  $\text{Ca}^{2+}$  buffering at lower  $\text{Ca}^{2+}$  concentrations would have caused underestimation of the fractional release under drug stimulation. Nevertheless, using the current






**Fig. 7.** Role of PKC in chrysosplenol-C-induced positive inotropy in rat ventricular myocytes. Chrysosplenol-C (CP-C) (50  $\mu\text{M}$ )-induced positive inotropic effects under control conditions (A) and their modulations in cells pretreated with KN-93 (1  $\mu\text{M}$ , 6 minutes), the CaMKII inhibitor (B), KN-92 (1  $\mu\text{M}$ , 6 minutes), the inactive analog of KN-93 (C), chelerythrine (D; 2  $\mu\text{M}$ , Chel), PLC inhibitor (E; 5  $\mu\text{M}$  U73122), or U73343, the inactive analog of U73122 (F; 5  $\mu\text{M}$  U73343). *i* shows representative cell shortening traces recorded before (Con) and after each drug effect became stable. *ii* represents mean magnitude of cell shortening measured before and after the applications of CP-C in the absence (A;  $n = 8$ ) and presence of KN-93 (B;  $n = 5$ ), KN-92 (C;  $n = 5$ ), chelerythrine (D;  $n = 5$ ), U73122 (E;  $n = 6$ ), or U73343 (F;  $n = 5$ ), (B), KN-92 (C), U73122 (E), or U73343 (F) (paired *t* test). (G) Comparison of mean % increases in cell shortening by 50  $\mu\text{M}$  CP-C between control conditions and interventions indicated. GF109203X: 5  $\mu\text{M}$  ( $n = 6$ ). Considering the new *p* value after the Bonferroni correction ( $0.05/6 = 0.0083$ ), % change in contraction by CP-C was significantly different in the presence of chelerythrine and GF109203X compared with that under control condition (\*).

method, we have successfully observed the enhancement of fractional release by another positive inotropic agent murrayafoline-A that distinctly increases  $I_{\text{Ca}}$ , SR  $\text{Ca}^{2+}$  content, and the sensitivity of  $\text{Ca}^{2+}$  release sites (Kim et al., 2015).

Since SR  $\text{Ca}^{2+}$  loading was increased with no significant acceleration of  $\text{Ca}^{2+}$  transient decay in the presence of chrysosplenol-C, larger SR  $\text{Ca}^{2+}$  loading may be due to increased  $\text{Ca}^{2+}$  entry and reduced  $\text{Ca}^{2+}$  removal across the cell membrane. The decay rate of caffeine-induced  $\text{Ca}^{2+}$  transient that reflects the activity of forward mode  $\text{Na}^{+}$ - $\text{Ca}^{2+}$  exchanger was not altered by chrysosplenol-C (Fig. 5B). In this regard, we have observed that the removal of external  $\text{Na}^{+}$  and  $\text{Ca}^{2+}$  suppressed the chrysosplenol-C-mediated increase in the resting spark frequency (Q.A.Le and S.H.Woo, unpublished

observations). This suggests that the  $\text{Na}^{+}$  and  $\text{Ca}^{2+}$  transporter/channels in the cell membrane may be modulated by chrysosplenol-C-PKC signaling.

In fact, large SR  $\text{Ca}^{2+}$  leaks can result in a decrease in SR  $\text{Ca}^{2+}$  content. However, the maximal effect of chrysosplenol-C on the  $\text{Ca}^{2+}$  spark frequency was  $\sim 1.7$ – $1.8$  fold only, and chrysosplenol-C did not dramatically increase diastolic  $\text{Ca}^{2+}$  levels (Fig. 1). This mild increase in the resting  $\text{Ca}^{2+}$  spark occurrence is in contrast with the effect of isoproterenol, which increases spark frequency in ventricular myocytes by  $\sim 5$ -fold (Potenza et al., 2019). It should be noted that because chrysosplenol-C increases cytosolic  $\text{Ca}^{2+}$  and SR  $\text{Ca}^{2+}$  loading, it could generate adverse effects, such as arrhythmias, due to intracellular  $\text{Ca}^{2+}$  overload, as other  $\text{Ca}^{2+}$  mobilizing positive inotropic



**Fig. 8.** Redistributions of PKC subtypes by the treatment of chrysosplenol-C in rat ventricular myocytes. (A and B) Western blots with antibodies specific for PKC- $\alpha$  (A), PKC- $\delta$  (A), and PKC- $\epsilon$  (B) using cytosolic and particulate fractions [soluble membrane (mem, M) and insoluble membrane] of rat ventricular myocytes treated with control solutions and with 80  $\mu$ M chrysosplenol-C for 2, 5, and 30 minutes. As a standard for cytosolic protein and membrane protein, GAPDH and Na<sup>+</sup>-K<sup>+</sup> pump (NKP) were used, respectively. (C) Relative quantification of each PKC subtype Western bands (A and B) compared with standard protein expression (GAPDH or NKP). Four rats (four Western blotting) were used. Two-way ANOVA-related measures with post hoc testing to control for multiple comparisons were used to test statistical significance (\*). Namely, each ANOVA was run at a new *p* value of 0.05/3 (0.017).

agents (e.g., isoproterenol and digitalis) showed (Vassalle and Lin, 2004; Steinberg, 2018). In this regard, CaMKII that appears to be secondarily activated by PKC activity (Figs. 6 and 7) may induce SR Ca<sup>2+</sup> leak (Tani, 1990), thereby contributing to arrhythmogenesis.

Our findings indicate that PKC may play a key role in the chrysosplenol-C-induced positive inotropic effect by enhancing SR Ca<sup>2+</sup> releases in ventricular myocytes. In fact, there are significant differences in the inotropic responses under various first messengers (agonist/hormones) to activate PKC or PKC activators in terms of time course and effects. Involvement of PKC in the biphasic inotropic responses of cardiac myocytes has been previously reported under the stimulations of the endothelin receptor and  $\alpha_1$ -adrenergic receptor, although there have been controversial findings depending on species and experimental conditions (Capogrossi et al., 1991; Woo and Lee, 1999a,b; O-Uchi et al., 2008; Smyrniak et al., 2018). Direct activation of PKC using phorbol 12-myristate 13-acetate has also induced positive inotropic effects and enhanced Ca<sup>2+</sup> transients and I<sub>Ca</sub> in cardiac myocytes, but there were other inconsistent observations (decrease or no change) depending on the species and experimental conditions (Lacerda et al., 1988; Walsh and Kass, 1988; Capogrossi et al., 1990; MacLeod and Harding, 1991; Tseng and Boyden, 1991; Woo and Lee, 1999b). The positive inotropic effects of these stimulations appeared to be related to mild enhancement of Ca<sup>2+</sup> release and Ca<sup>2+</sup> current. In case of the effect of chrysosplenol-C, PKC does not seem to alter I<sub>Ca</sub>. Diverse effects of PKC under different agonists in cardiac myocytes may be caused by differences in the PKC isoforms involved and compartmentalization of relevant signaling molecules within the cells. According to our results, translocation of PKC- $\delta$  to the membrane fractions

was the most prominent after 2 minutes of the chrysosplenol-C exposure (Fig. 8) when we normally observed maximal stimulatory effects on contraction (Son et al., 2011), Ca<sup>2+</sup> transients, and sparks. Therefore, it is plausible to think that this isoform mediates the positive inotropic effect of chrysosplenol-C. There was a difference in the redistribution patterns between the soluble and insoluble membrane fractions of PKC- $\alpha$  and PKC- $\delta$ . PKC- $\alpha$  increased in the insoluble membrane after 5 minutes with a gradual decrease in its level in the cytosolic fraction, whereas PKC- $\delta$  was increased in both soluble and insoluble membrane fractions for 5 minutes with no significant reduction in its level in the cytosolic fractions (Fig. 8). It is known that various signaling molecules, including mitogen-activated protein kinases, extracellular signal-regulated kinases, and scaffolding proteins in the t-tubules, caveolae, and cytoskeleton are known to be contained in the Triton-insoluble membrane fraction in cardiac myocytes (Yan et al., 1996; Anderson, 1998; Ballard-Croft et al., 2008). Partitions of PKC- $\alpha$  and - $\delta$  into the insoluble membrane fractions suggest that interaction between the PKC isoforms and signaling molecules in those subcellular compartments.

Previous reports of the heart from PKC- $\alpha$  knockout mice, PKC- $\alpha$ -overexpressing mice, and cardiac cells with adenoviral PKC- $\alpha$  knockdown suggest that PKC- $\alpha$  reduces cardiac contraction via lowering Ca<sup>2+</sup> transient and Ca<sup>2+</sup> loading in the SR (Braz et al., 2004). Regarding the role of PKC- $\delta$  in cardiac contraction regulation, it has been demonstrated in myocytes expressing PKC- $\delta$ -GFP that phorbol dibutyrate (PDBu), the PKC activator, caused a transient negative inotropic response followed by a robust and sustained positive inotropic response that paralleled PKC- $\delta$  accumulation in the Golgi and other subcellular domain (Kang and Walker, 2005). The

cardiac cells expressing PKC- $\epsilon$ -GFP also showed similar stimulatory effects by PDBu on contraction (Kang and Walker, 2005). The PKC- $\epsilon$ , however, has been shown to mediate endothelin-1-induced negative inotropy in rat ventricular myocytes (Smyrniats et al., 2018). In addition, the PDBu-induced negative inotropic response has been suggested to be mediated by PKC- $\alpha$  (Kang and Walker, 2005). This previous evidence appears to be consistent with our hypothesis that PKC- $\delta$  that was early partitioned by chryso splenol-C to the soluble and insoluble membrane may be responsible for the positive inotropy. Such hypothesis warrants further investigations.

#### Authorship Contributions

Participated in research design: Woo.

Conducted experiments: Wang, Trinh, Vu, Ohk, Zhang.

Contributed new reagents or analytic tools: Hoang, Nguyen.

Performed data analysis: Wang, Trinh, Kim, Woo.

Wrote or contributed to the writing of the manuscript: Wang, Trinh, Vu, Kim, Woo.


#### References

- Anderson RGW (1998) The caveolae membrane system. *Annu Rev Biochem* **67**:199–225.
- Asai T, Shuba LM, Pelzer DJ, and McDonald TF (1996) PKC-independent inhibition of cardiac L-type  $Ca^{2+}$  channel current by phorbol esters. *Am J Physiol* **270**:H620–H627.
- Ballard-Croft C, Locklar AC, Keith BJ, Mentzer Jr RM, and Lasley RD (2008) Oxidative stress and adenosine  $A_1$  receptor activation differentially modulate subcellular cardiomyocyte MAPKs. *Am J Physiol Heart Circ Physiol* **294**:H263–H271.
- Bassani JWM, Bassani RA, and Bers DM (1994) Relaxation in rabbit and rat cardiac cells: species-dependent differences in cellular mechanisms. *J Physiol* **476**:279–293.
- Bassani JWM, Yuan W, and Bers DM (1995) Fractional SR  $Ca$  release is regulated by trigger  $Ca$  and SR  $Ca$  content in cardiac myocytes. *Am J Physiol* **268**:C1313–C1319.
- Beuckelmann DJ and Wier WG (1988) Mechanism of release of calcium from sarcoplasmic reticulum of guinea-pig cardiac cells. *J Physiol* **405**:233–255.
- Bovo E, Huke S, Blatter LA, and Zima AV (2017) The effect of PKA-mediated phosphorylation of ryanodine receptor on SR  $Ca^{2+}$  leak in ventricular myocytes. *J Mol Cell Cardiol* **104**:9–16.
- Braz JC, Gregory K, Pathak A, Zhao W, Sahin B, Kleivitsky R, Kimball TF, Lorenz JN, Nairn AC, Liggett SB, et al. (2004) PKC- $\alpha$  regulates cardiac contractility and propensity toward heart failure. *Nat Med* **10**:248–254.
- Cannell MB, Cheng H, and Lederer WJ (1994) Spatial non-uniformities in  $[Ca^{2+}]_i$  during excitation-contraction coupling in cardiac myocytes. *Biophys J* **67**:1942–1956.
- Capogrossi MC, Kaku T, Filburn CR, Peltó DJ, Hansford RG, Spurgeon HA, and Lakatta EG (1990) Phorbol ester and dioctanoylglycerol stimulate membrane association of protein kinase C and have a negative inotropic effect mediated by changes in cytosolic  $Ca^{2+}$  in adult rat cardiac myocytes. *Circ Res* **66**:1143–1155.
- Capogrossi MC, Kachadorian WA, Gambassi G, Spurgeon HA, and Lakatta EG (1991)  $Ca^{2+}$  dependence of  $\alpha$ -adrenergic effects on the contractile properties and  $Ca^{2+}$  homeostasis of cardiac myocytes. *Circ Res* **69**:540–550.
- Cheng H, Lederer WJ, and Cannell MB (1993) Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle. *Science* **262**:740–744.
- Eisner DA, Choi HS, Diaz ME, O'Neill SC, and Trafford AW (2000) Integrative analysis of calcium cycling in cardiac muscle. *Circ Res* **87**:1087–1094.
- Ginsburg KS and Bers DM (2004) Modulation of excitation-contraction coupling by isoproterenol in cardiomyocytes with controlled SR  $Ca^{2+}$  load and  $Ca^{2+}$  current trigger. *J Physiol* **556**:463–480.
- Goldberg M, Zhang HL, and Steinberg SF (1997) Hypoxia alters the subcellular distribution of protein kinase C isoforms in neonatal rat ventricular myocytes. *J Clin Invest* **99**:55–61.
- Guo T, Zhang T, Mestril R, and Bers DM (2006)  $Ca^{2+}$ /calmodulin-dependent protein kinase II phosphorylation of ryanodine receptor does affect calcium sparks in mouse ventricular myocytes. *Circ Res* **99**:398–406.
- Györke I and Györke S (1998) Regulation of the cardiac ryanodine receptor channel by luminal  $Ca^{2+}$  involves luminal  $Ca^{2+}$  sensing sites. *Biophys J* **75**:2801–2810.
- Hamill OP, Marty A, Neher E, Sakmann B, and Sigworth FJ (1981) Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch* **391**:85–100.
- Han S, Schiefer A, and Isenberg G (1994)  $Ca^{2+}$  load of guinea-pig ventricular myocytes determines efficacy of brief  $Ca^{2+}$  currents as trigger for  $Ca^{2+}$  release. *J Physiol* **480**:411–421.
- Huang DT, Kamperdick C, and Sung TV (2004) Homogenitac acid derivatives from *Milusa balansae*. *J Nat Prod* **67**:445–447.
- Janczewski AM, Spurgeon HA, Stern MD, and Lakatta EG (1995) Effects of sarcoplasmic reticulum  $Ca^{2+}$  load on the gain function of  $Ca^{2+}$  release by  $Ca^{2+}$  current in cardiac cells. *Am J Physiol* **268**:H916–H920.
- Kang M and Walker JW (2005) Protein kinase C delta and epsilon mediate positive inotropy in adult ventricular myocytes. *J Mol Cell Cardiol* **38**:753–764.
- Kim JC, Wang J, Son MJ, Cuong NM, and Woo SH (2015) Sensitization of cardiac  $Ca^{2+}$  release sites by protein kinase C signaling: evidence from action of murrayafoline A. *Pflügers Arch* **467**:1607–1621.
- Lacerda AE, Rampe D, and Brown AM (1988) Effects of protein kinase C activators on cardiac  $Ca^{2+}$  channels. *Nature* **335**:249–251.
- Lu S, Liao Z, Lu X, Katschinski DM, Mercola M, Chen J, Heller Brown J, Molkentin JD, Bossuyt J, and Bers DM (2020) Hyperglycemia acutely increases cytosolic reactive oxygen species via O-linked GlcNAcylation and CaMKII activation in mouse ventricular myocytes. *Circ Res* **126**:e80–e96.
- MacLeod KT and Harding SE (1991) Effects of phorbol ester on contraction, intracellular pH and intracellular  $Ca^{2+}$  in isolated mammalian ventricular myocytes. *J Physiol* **444**:481–498.
- Maier LS and Bers DM (2007) Role of  $Ca^{2+}$ /calmodulin-dependent protein kinase (CaMK) in excitation-contraction coupling in the heart. *Cardiovasc Res* **73**:631–640.
- Näbauer M, Callewaert G, Cleemann L, and Morad M (1989) Regulation of calcium release is gated by calcium current, not gating charge, in cardiac myocytes. *Science* **244**:800–803.
- Negretti N, O'Neill SC, and Eisner DA (1993) The relative contributions of different intracellular and sarcolemmal systems to relaxation in rat ventricular myocytes. *Cardiovasc Res* **27**:1826–1830.
- Niggli E and Lederer WJ (1990) Voltage-independent calcium release in heart muscle. *Science* **250**:565–568.
- Ogrodnik J and Niggli E (2010) Increased  $Ca^{2+}$  leak and spatiotemporal coherence of  $Ca^{2+}$  release in cardiomyocytes during  $\beta$ -adrenergic stimulation. *J Physiol* **588**:225–242.
- O-Uchi J, Sasaki H, Morimoto S, Kusakari Y, Shinji H, Obata T, Hongo K, Komukai K, and Kurihara S (2008) Interaction of alpha1-adrenoceptor subtypes with different G proteins induces opposite effects on cardiac L-type  $Ca^{2+}$  channel. *Circ Res* **102**:1378–1388.
- Pereira L, Métrich M, Fernández-Velasco M, Lucas A, Leroy J, Perrier R, Morel E, Fischmeister R, Richard S, Bénitah JP, et al. (2007) The cAMP binding protein Epac modulates  $Ca^{2+}$  sparks by a  $Ca^{2+}$ /calmodulin kinase signalling pathway in rat cardiac myocytes. *J Physiol* **583**:685–694.
- Potenza DM, Janicek R, Fernandez-Tenorio M, Camors E, Ramos-Mondragón R, Valdivia HH, and Niggli E (2019) Phosphorylation of the ryanodine receptor 2 at serine 2030 is required for a complete  $\beta$ -adrenergic response. *J Gen Physiol* **151**:131–145.
- Puglisi JL, Yuan W, Timofeyev V, Myers RE, Chiamvimonvat N, Samarel AM, and Bers DM (2011) Phorbol ester and endothelin-1 alter functional expression of  $Na^{+}/Ca^{2+}$  exchange,  $K^{+}$ , and  $Ca^{2+}$  currents in cultured neonatal rat myocytes. *Am J Physiol Heart Circ Physiol* **300**:H617–H626.
- Satoh H, Blatter LA, and Bers DM (1997) Effects of  $[Ca^{2+}]_i$ , SR  $Ca^{2+}$  load, and rest on  $Ca^{2+}$  spark frequency in ventricular myocytes. *Am J Physiol* **272**:H657–H668.
- Semple SJ, Nobbs SF, Pyke SM, Reynolds GD, and Flower RL (1999) Antiviral flavonoid from Pterocaulon sphaacelatum, an Australian Aboriginal medicine. *J Ethnopharmacol* **68**:283–288.
- Shacklock PS, Wier WG, and Balke CW (1995) Local  $Ca^{2+}$  transients ( $Ca^{2+}$  sparks) originate at transverse tubules in rat heart cells. *J Physiol* **487**:601–608.
- Simonis G, Honold J, Schwarz K, Braun MU, and Strasser RH (2002) Regulation of the isozymes of protein kinase C in the surviving rat myocardium after myocardial infarction: distinct modulation for PKC- $\alpha$  and for PKC- $\delta$ . *Basic Res Cardiol* **97**:223–231.
- Smyrniats I, Goodwin N, Wachten D, Skogestad J, Aronsen JM, Robinson EL, Demydenko K, Segonds-Pichon A, Oxley D, Sadayappan S, et al. (2018) Contractile responses to endothelin-1 are regulated by PKC phosphorylation of cardiac myosin binding protein-C in rat ventricular myocytes. *J Mol Cell Cardiol* **117**:1–18.
- Son MJ, Kim HK, Huong TT, Kim YH, Van Sung T, Cuong NM, and Woo SH (2011) Chryso splenol C increases contraction in rat ventricular myocytes. *J Cardiovasc Pharmacol* **57**:259–262.
- Steinberg SF (2018)  $\beta$ 1-adrenergic receptor regulation revisited. *Circ Res* **123**:1199–1201.
- Tani M (1990) Mechanisms of  $Ca^{2+}$  overload in reperfused ischemic myocardium. *Annu Rev Physiol* **52**:543–559.
- Tseng GN and Boyden PA (1991) Different effects of intracellular  $Ca$  and protein kinase C on cardiac T and L  $Ca$  currents. *Am J Physiol* **261**:H364–H379.
- Vassalle M and Lin CI (2004) Calcium overload and cardiac function. *J Biomed Sci* **11**:542–565.
- Walsh KB and Kass RS (1988) Regulation of a heart potassium channel by protein kinase A and C. *Science* **242**:67–69.
- Waxham MN and Aronowski J (1993)  $Ca^{2+}$ /calmodulin-dependent protein kinase II is phosphorylated by protein kinase C in vitro. *Biochemistry* **32**:2923–2930.
- Woo SH and Lee CO (1999a) Effects of endothelin-1 on  $Ca^{2+}$  signaling in guinea-pig ventricular myocytes: role of protein kinase C. *J Mol Cell Cardiol* **31**:631–643.
- Woo SH and Lee CO (1999b) Role of PKC in the effects of  $\alpha$ 1-adrenergic stimulation on  $Ca^{2+}$  transients, contraction and  $Ca^{2+}$  current in guinea-pig ventricular myocytes. *Pflügers Arch* **437**:335–344.
- Woo SH, Cleemann L, and Morad M (2003) Spatiotemporal characteristics of junctional and nonjunctional focal  $Ca^{2+}$  release in rat atrial myocytes. *Circ Res* **92**:e1–e11.
- Yan SR, Fumagalli L, and Berton G (1996) Activation of SRC family kinases in human neutrophils. Evidence that p58C-FGR and p53/56LYN redistributed to a Triton X-100-insoluble cytoskeletal fraction, also enriched in the caveolar protein caveolin, display an enhanced kinase activity. *FEBS Lett* **380**:198–203.
- Zhou P, Zhao YT, Guo YB, Xu SM, Bai SH, Lakatta EG, Cheng H, Hao XM, and Wang SQ (2009)  $\beta$ -adrenergic signaling accelerates and synchronizes cardiac ryanodine receptor response to a single L-type  $Ca^{2+}$  channel. *Proc Natl Acad Sci USA* **106**:18028–18033.

Address correspondence to: Sun-Hee Woo, Chungnam National University, 99 Daehak-ro, Daejeon, South Korea. E-mail: shwoo@cnu.ac.kr

“Chrysosplenol-C increases contraction by augmentation of sarcoplasmic reticulum  $\text{Ca}^{2+}$  loading and release via protein kinase C in rat ventricular myocytes” J. Wang, T.N. Trinh, A. T. V. Vu, J.C. Kim, A.T.N. Hoang, C.J. Ohk, Y.H. Zhang, C.M. Nguyen, S.H. Woo, Molecular Pharmacology, MOLPHARM-AR-2021-000365R1

### Supplemental material



**Fig. S1.** Effects of KN-93 and chelerythrine on SR  $\text{Ca}^{2+}$  loading and fractional release in rat ventricular myocytes. (A and C)  $\text{Ca}^{2+}$  signal traces measured on field stimulation (arrowheads) followed by 10 mM-caffeine ("caff") exposure in representative ventricle cells under control conditions and after application of 1  $\mu\text{M}$  KN-93 for 5 min (A) or 2  $\mu\text{M}$  chelerythrine for 10 min (C). (B and D) Summary of the effects of KN-93 (B;  $n = 8$ ) and chelerythrine ("Chel"; D;  $d = 8$ ) on the magnitude of caffeine-induced  $\text{Ca}^{2+}$  release (SR  $\text{Ca}^{2+}$  content) and fractional release, showing a significant decrease in SR  $\text{Ca}^{2+}$  content by KN-93 only. Paired Student's  $t$  tests followed by Bonferroni correction.  $P < (0.05/2)$  was considered to be significant.