

Nucleosome Positioning and Gene Regulation of the *SGLT2* Gene in the Renal Proximal Tubular Epithelial Cells [Ⓢ]

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, and Ichiro Ieiri

Department of Clinical Pharmacokinetics, Graduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka, Japan

Received January 16, 2018; accepted June 27, 2018

ABSTRACT

Filtered glucose is mostly reabsorbed by sodium-glucose cotransporter 2 (SGLT2) in the proximal tubules. SGLT2 is predominantly expressed in the human kidney. However, the regulatory mechanisms for *SGLT2* gene expression in the human kidney remain unclear. We in this work elucidated the transcriptional regulatory mechanisms for the *SGLT2* gene by nucleosome occupancy in the *SGLT2* promoter region. Expressions of *SGLT2* mRNA and protein were markedly weaker in human kidney-derived HK-2 cells than the human kidney. The nucleosome occupancy level in the *SGLT2* promoter region was low in the kidney, but high in HK-2 cells. A treatment with a histone deacetylase inhibitor trichostatin A (TSA) decreased nucleosome occupancy in the promoter region and increased *SGLT2*

expression levels in HK-2 cells. The upregulation of *SGLT2* expression by histone acetylation was accompanied by a higher binding frequency of hepatocyte nuclear factor (HNF) 1 α , a transcriptional modulator of *SGLT2* in the human kidney, to the promoter region. The transfection of a HNF1 α expression plasmid into HK-2 cells resulted in the upregulation of *SGLT2* mRNA expression in the presence of TSA, but not in the treatment of dimethylsulfoxide as a control. Nucleosome occupancy in the promoter region was markedly higher in the liver and small intestine than the kidney. Our results indicate that tissue-specific nucleosome occupancy plays an important role in the regulation of *SGLT2* gene expression via HNF1 α binding at the *SGLT2* promoter region.

Introduction

The kidney plays an important role in the reabsorption of glucose from the glomerular filtrate. Approximately 180 g glucose is typically filtered by the kidneys each day and reabsorbed in proximal tubule epithelial cells (PTECs) by sodium-glucose cotransporters (SGLTs) (Wright, 2001; Chen et al., 2010; Gerich, 2010; Pérez López et al., 2010). In the S1 and S2 segments of the proximal tubule, approximately 90% of filtered glucose is reabsorbed through SGLT2 (Gerich, 2010). Previous studies demonstrated that a deficiency in SGLT2 expression decreased blood glucose levels. The knockout of *Sgt2* in diabetic mice reduced blood glucose levels and prevented glomerular hyperfiltration (Vallon et al., 2013). Therefore, the inhibition of SGLT2 activity in the kidney is regarded as an effective strategy for glycemic control in diabetic patients. Selective SGLT2 inhibitors have been developed as a new class of oral medication for the treatment of type 2 diabetes mellitus. These drugs reduce glucose reabsorption by SGLT2 in the proximal tubules, thereby increasing urinary

glucose excretion and lowering plasma glucose levels (Jabbour and Goldstein, 2008; Boldys and Okopień, 2009; Misra, 2013).

Previous studies focused on the regulation of *SGLT2* expression by transcription factors. Hepatocyte nuclear factor (HNF) 1 α , which has been characterized as a modulator of *SGLT2* expression, was shown to directly bind to the *SGLT2* 5'-flanking region (5'-FR) and control *SGLT2* gene expression (Pontoglio et al., 2000; Freitas et al., 2008; Zhao et al., 2016). Additionally, HNF4 α and SP1 have been reported to play an important role in *SGLT2* expression (Kothinti et al., 2010; Bonner et al., 2015). Although these transcription factors are expressed in various tissues such as the liver and intestines, *SGLT2* is not. Therefore, another mechanism for the regulation of *SGLT2* expression appears to contribute to tissue-specific gene expression. The nucleosome is a basic unit of chromatin and consists of approximately 147 bp of DNA wrapped around histones. Nucleosome occupancy plays an important role in the epigenetic regulation of gene expression by inhibiting the initiation of transcription at gene promoters (Lorch et al., 1987; Workman, 2006). Previous studies demonstrated that nucleosomes control tissue-specific transcription factor binding to gene promoters, and, thus, contribute to tissue-specific gene expression (Hoffman et al., 2010; Hu et al., 2011; Tsui et al., 2011; Yang et al., 2013; Zhang et al., 2015). However, the relationship between nucleosome occupancy and tissue-specific *SGLT2* expression remains unknown. Therefore, a

This work was supported by Japan Society for the Promotion of Science KAKENHI [Grant JP17K08954].

<https://doi.org/10.1124/mol.118.111807>.

[Ⓢ] This article has supplemental material available at molpharm.aspetjournals.org.

ABBREVIATIONS: 5'-FR, 5'-flanking region; ChIP, chromatin immunoprecipitation; DMSO, dimethylsulfoxide; HNF, hepatocyte nuclear factor; NOMe-Seq, nucleosome occupancy and methylome sequencing; NuSA, nucleosome-scanning assay; PCR, polymerase chain reaction; PTEC, proximal tubule epithelial cell; SGLT2, sodium-glucose cotransporter 2; siHNF1 α , HNF1 α siRNA; siRNA, small interfering RNA; TSA, trichostatin A; TSS, transcription start site.




Fig. 1. Analysis of transcriptional activity of *SGLT2* 5'-FR and *SGLT2* expression levels in HK-2 cells. (A) Luciferase activity of a series of reporter constructs containing *SGLT2* 5'-FR in HK-2 cells. Results are expressed as fold increases in pGL4.10. (B) *SGLT2* mRNA levels in HK-2 cells and the human kidney were measured by quantitative PCR and normalized to RPL13 mRNA levels. Results are expressed relative to *SGLT2* mRNA levels in the human kidney. (C) Western blot analyses of *SGLT2* and β -actin protein expression in HK-2 cells and the human kidney. *SGLT2* protein levels represent the density ratio for the bands of *SGLT2* and β -actin. Results are expressed relative to *SGLT2* protein levels in the human kidney. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$.

clearer understanding of nucleosome occupancy at the *SGLT2* promoter may provide significant insights into transcriptional regulatory mechanisms for the *SGLT2* gene in the human kidney.

In the present study, we focused on the transcriptional regulation of the *SGLT2* gene by nucleosome occupancy in *SGLT2* 5'-FR in the human kidney. We demonstrated that decreasing nucleosome occupancy in the *SGLT2* promoter plays a critical role in the upregulation of *SGLT2* expression. We also revealed that HNF1 α binds to the *SGLT2* promoter in the human kidney, and the activation of *SGLT2* transcription by HNF1 α requires low nucleosome occupancy through histone acetylation in the *SGLT2* promoter.

Materials and Methods

Human Tissue. A human kidney cortex and small intestine obtained from surgical specimens were purchased from Integrated Laboratory Services-Biotech (Chestertown, MD). A human liver sample was obtained from a Caucasian donor at the National Disease Research Interchange (Philadelphia, PA). All assays using human tissue samples were performed by three independent experiments ($n = 3$). The use of human tissues was approved by the Kyushu University Institutional Review Board for Human Genome/Genome Research (Fukuoka, Japan).

Cell Culture. HK-2 cells were obtained from the American Type Culture Collection (Manassas, VA). Cells were cultured in DMEM/F12 medium (Thermo Fisher Scientific, Waltham, MA) supplemented with 10% fetal bovine serum at 37°C in an incubator with 5% CO₂.

Construction of *SGLT2* Reporter Plasmids. The DNA fragment of *SGLT2* 5'-FR (-3185/+18) was amplified by a polymerase chain reaction (PCR) using primers (described as -3185 and +18 in Supplemental Table 1) and human genomic DNA as a template. The pGL4.10 vector (Promega, Madison, WI) was digested by KpnI and HindIII restriction enzymes and then ligated with the PCR product using the In-Fusion HD Cloning kit (TaKaRa, Shiga, Japan). Deletion constructs [-2320/+18, -1587/+18, -485/+18, -154/+18, -44/+18, and del(-51/-37)] were generated by In-Fusion cloning or the QuikChange II site-directed mutagenesis method (Supplemental Table 1) using the -3185/+18 construct as a template. The sequences of these plasmids were confirmed by direct sequencing using primers (Supplemental Table 2) described previously (Nishimura et al., 2014).

Construction of a HNF1 α Expression Plasmid. The coding region of HNF1 α was amplified by PCR using cDNA synthesized from the human kidney and primers (Supplemental Table 3). The pcDNA3.1(+) vector (Invitrogen, Carlsbad, CA) was digested by EcoRI

and EcoRV restriction enzymes and then ligated with the PCR product using the In-Fusion HD Cloning kit. The sequence of the plasmid was confirmed by directed sequencing using primers (Supplemental Table 3).

Luciferase Assay. HK-2 cells were seeded on 24-well culture plates at 0.2×10^5 cells/well and incubated for 24 hours. A series of reporter constructs [equimolar amounts of the empty pGL4.10 plasmid (250 ng), 500 ng HNF1 α expression plasmid or the empty pcDNA3.1(+) plasmid, and 25 ng internal standard pGL4.70 (Promega)] were transfected into HK-2 cells using FuGENE HD Transfection Reagent (Promega), according to the manufacturer's instructions. After 48 hours, luciferase activities were measured using the Dual-Luciferase Reporter Assay System (Promega) and a TD-20/20 Luminometer (Promega). Transcriptional activities were expressed as the ratio of *Firefly* luciferase to *Renilla* luciferase activity.

RNA Interference. The transfection mix was prepared in Opti-MEM (Invitrogen) with HNF1 α small interfering RNA (siRNA) (siHNF1 α ; Sigma-Aldrich, St. Louis, MO) or control siRNA (MISSION siRNA Universal Negative Control; SIC-001; Sigma-Aldrich) and Lipofectamine RNAiMAX (Invitrogen), according to the manufacturer's directions. The final concentrations of siRNA and lipofectamine added to the cells were 20 nM and 2 μ l/ml, respectively. HK-2 cells were cultured in the presence of the transfection mixture for 24 hours, and cells were treated with dimethylsulfoxide (DMSO) or 1.0 μ M trichostatin A (TSA; Wako, Osaka, Japan) for 24 hours. siHNF1 α sequences were as follows: 5'-CAGUGAGACUGCAGAA-GUAtt-3' (sense sequence) and 5'-TACTTCTGCAGTCTCACTGtt-3' (antisense sequence).

Total RNA Isolation and Capture of Nascent RNAs. HK-2 cells were seeded on a six-well plate at 1.5×10^5 cells/well and incubated for 24 hours, and cells were then transfected with 500 ng HNF1 α expression plasmid or pcDNA3.1(+) plasmid using the FuGENE HD Transfection Reagent (Promega). After 48 hours, cells were treated with DMSO or 1.0 μ M TSA, together with 0.2 mM 5-ethynyl uridine for 24 hours. Total RNA was isolated from the human kidney and HK-2 cells using ISOSPIN Cell & Tissue RNA (Nippon Gene, Tokyo, Japan). Ethynyl uridine-labeled nascent RNA was biotinylated and captured using a Click-iT Nascent RNA Capture Kit (Invitrogen), according to manufacturer's instructions.

cDNA Synthesis and Quantitative PCR. Total RNA (500 ng) or nascent RNA (200 ng) was transcribed using the Verso cDNA Synthesis Kit (Thermo Fisher Scientific). Quantitative PCR was performed on a StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA) with SYBR Premix Ex *Taq* (TaKaRa) under the following conditions: 95°C for 2 minutes, followed by 40 cycles at 95°C for 3 seconds and 60°C for 30 seconds. *SGLT2* and HNF1 α mRNA levels were normalized to 60S ribosomal protein L13 (RPL13) mRNA levels. The primers used in this study are listed in Supplemental Table 4.




Fig. 2. Analysis of epigenetic regulation in *SGLT2* 5'-FR in HK-2 cells and the human kidney. (A and B) NOME-Seq data of *SGLT2* 5'-FR in HK-2 cells and the human kidney. The arrow indicates the *SGLT2* TSS. (A) Vertical lines indicate CpG sites. White circles represent unmethylated CpG sites, and black circles represent methylated CpG sites. (B) Vertical lines indicate GpC sites. White circles represent unmethylated GpC sites, and black circles represent methylated GpC sites. Deep gray bars represent nucleosome occupancy, which is the region of consecutive unmethylated GpC sites over 147 bp. (C) NuSA data of 5'-FR in HK-2 cells (black circles) and the human kidney (white circles). Relative nucleosome occupancy is expressed relative to the level of nucleosome occupancy in each sample without the micrococcal nuclease treatment and indicated by the midpoints of each amplicon. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$ for HK-2 cells versus the human kidney.

Nucleosome Occupancy and Methylome Sequencing. Nucleosome occupancy and methylome sequencing (NOME-Seq) were performed according to the manufacturer's instructions (Active Motif, Carlsbad, CA), with slight modifications. Approximately 50 mg frozen human tissues and 7.5×10^5 of HK-2 cells were cross-linked by 1% formaldehyde at room temperature for 15 and 5 minutes, respectively, and cross-linking was quenched by adding 1.25 M glycine. Cells were homogenized in lysis buffer with 40 strokes of a Potter-Elvehjem homogenizer (Wheaton Industries, Millville, NJ). Homogenized kidney tissues were filtered through an 80- μ m Nylon Woven Net Filter (Merck Millipore, Darmstadt, Germany) to remove tissue debris. Chromatin was sheared to fragments larger than 1 kb. Sheared

chromatin was treated with 0 or 20 U GpC methyltransferase at 37°C for 4 hours. Reactions were stopped by the addition of stop solution and incubated at 95°C for 15 minutes. Bisulfite conversion was then performed on DNA. Bisulfite-converted DNA was used for PCR amplification with primers for *SGLT2* 5'-FR (Supplemental Table 5). PCR products were cloned into the pGEM-T easy vector (Promega). Twenty colonies in GpC methyltransferase-treated DNA and 10 colonies in nontreated DNA were subjected to colony PCR, followed by sequencing with primers (Supplemental Table 5) for the assessment of nucleosome occupancy and CpG methylation, respectively. GpCpG sites were removed from the analysis of nucleosome occupancy as they are both a GpC and CpG context.




Fig. 3. Influence of the TSA treatment on *SGLT2* expression and histone acetylation in *SGLT2* 5'-FR in HK-2 cells. HK-2 cells were treated with DMSO or TSA for 24 hours. (A) *SGLT2* mRNA levels were measured by quantitative PCR and normalized to RPL13 mRNA levels. Results are expressed relative to *SGLT2* mRNA levels in DMSO-treated cells (control). (B) Scheme of the ChIP assay in 5'-FR. The upper arrow indicates the *SGLT2* TSS. Lower arrows indicate two sets of PCR primers targeting 5'-FR, designated as distal and proximal. (C) ChIP analyses targeting 5'-FR in DMSO- or TSA-treated cells using antibodies against normal IgG (white bars) and acetylated histone H3 (black bars). (D) Semiquantitative analyses of acetylated histone H3 enrichment in two regions in ChIP analyses in (C). Results are expressed as the percentage of the immunoprecipitate over total input DNA. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$.

Nucleosome Scanning Assay. A nucleosome-scanning assay (NuSA) was performed using the EpiScope Nucleosome Preparation Kit (TaKaRa), according to the manufacturer's protocol, with slight modifications. Approximately 20 mg frozen human kidney or 2.0×10^6 of HK-2 cells was washed with ice-cold phosphate saline buffer and incubated in lysis buffer for 10 minutes. Tissue samples were homogenized and filtered, as described above. The nuclei released were treated with 0 or 4 U micrococcal nuclease for 30 minutes at 37°C and purified using NucleoSpin Gel and PCR Clean-up (TaKaRa). Nucleosomal DNA enrichment was evaluated by quantitative PCR with overlapping primer pairs (Supplemental Table 6), producing a high-resolution, quantitative map of the nucleosome location and occupancy. Relative nucleosome occupancy was presented by the ratio to the nucleosomal DNA enrichment of LINE-1 as a reference gene using the Pfaffl method (Pfaffl, 2001).

Chromatin Immunoprecipitation Assay. Chromatin immunoprecipitation (ChIP) assays were performed using the EpiQuik ChIP or Tissue ChIP kit (Epigentek, Farmingdale, NY), according to manufacturer's instructions. Briefly, approximately 50 mg human kidney or 1.0×10^6 of HK-2 cells were cross-linked, homogenized, and filtered, as described above. Chromatin was sheared to fragments ranging from 300 to 600 bp. The sonicated lysate ($5 \mu\text{l}$) was used to quantify the total amount of DNA (input). Immunoprecipitation was performed for 90 minutes with $2 \mu\text{g}$ antibodies: anti-acetylated histone H3 (Merck), anti-HNF1 α (sc-393925; Santa Cruz Biotechnology, Santa Cruz, CA), or normal mouse IgG as the negative control. Immunoprecipitated DNA and input DNA were analyzed by semi-quantitative PCR or quantitative PCR using primers for *SGLT2* 5'-FR (Supplemental Table 7). Semiquantitative PCR was performed using Gene RED PCR Mix Plus (Nippon Gene). PCR products were resolved by electrophoresis on a 2% agarose gel and stained with ethidium bromide. The intensity of bands was quantified with the Lumino Image Analyzer LAS-3000 and Multi Gauge (Fujifilm, Tokyo, Japan).

Western Blot Analysis. The human kidney was pulverized and dissolved in radioimmunoprecipitation assay buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% sodium deoxycholate, 1% Nonidet P-40, and 0.1% SDS). HK-2 cells were harvested and homogenized with radioimmunoprecipitation assay buffer 24 hours after the TSA treatment. Deglycosylation was performed by incubation with peptide-N-glycosidase F (New England Biolabs, Ipswich, MA). Lysate samples were separated on 9% SDS-polyacrylamide gels and transferred to polyvinylidene difluoride membranes by Trans-Blot SD Semi-Dry Electrophoretic Transfer Cell (Bio-Rad, Richmond, CA). Membranes were blocked in phosphate-buffered saline with 0.1% Tween 20 and 5% nonfat dry milk at room temperature for 20 minutes and incubated with the primary anti- β -actin (ab6276, diluted at 1:5000; Abcam, Cambridge, UK), anti-SGLT2 (sc-393350, diluted at 1:200; Santa Cruz Biotechnology), and anti-HNF1 α (sc-6547, diluted at 1:1000; Santa Cruz Biotechnology) antibodies at 4°C overnight. After washing, the membrane was incubated at room temperature for 1 hour with the ECL Mouse IgG, horseradish peroxidase-linked whole Ab (NA931, diluted at 1:10,000; GE Healthcare, Little Chalfont, UK), or donkey anti-goat IgG-horseradish peroxidase (sc-2020, diluted at 1:10,000; Santa Cruz Biotechnology). After washing the membrane, immune complexes were visualized using a chemiluminescence reagent (Amersham ECL Select; GE Healthcare) and captured with the Lumino Image Analyzer LAS-3000.

Statistical Analysis. Results from three independent experiments are expressed as the mean \pm S.D. Statistical analyses were performed with the statistical software program R. The means of two groups were compared with an unpaired Student's *t* test (two-tailed). Comparisons of the means of multiple groups were performed with the Tukey-Kramer test. Two-way analysis of variance was used to examine the effects of two factors on an experimental response. All tests were considered significant at $P < 0.05$.

Results

Analysis of *SGLT2* mRNA Levels and Nucleosome Occupancy in HK-2 Cells and Human Kidney Tissue. *SGLT2* is primarily expressed in the cortex of the kidney and located in the S1 and S2 segments of the proximal tubule. The HK-2 cell line, an immortalized PTEC line from a normal adult human kidney, is available as a human PTEC model (Ryan et al., 1994; Wilmer et al., 2010). However, a luciferase assay with HK-2 cells showed no significant changes in the luciferase activity of *SGLT2* 5'-FR reporter constructs (Fig. 1A). Quantitative PCR with cDNA synthesized from total RNA and Western blot analyses demonstrated that *SGLT2* mRNA and protein levels were markedly lower in HK-2 cells than in the human kidney (Fig. 1, B and C). These results suggest different regulatory mechanisms for *SGLT2* gene




Fig. 4. Influence of the TSA treatment on nucleosome occupancy in *SGLT2* 5'-FR in HK-2 cells. (A) NOME-Seq data of *SGLT2* 5'-FR in DMSO- or TSA-treated cells. Vertical lines indicate GpC sites. White circles represent unmethylated GpC sites, and black circles represent methylated GpC sites. Deep gray bars represent nucleosome occupancy, which is the region of consecutive unmethylated GpC sites over 147 bp. (B) NuSA data of 5'-FR in DMSO (black circles)- or TSA (white circles)-treated cells. Relative nucleosome occupancy is expressed relative to the level of nucleosome occupancy in each sample without the micrococcal nuclease treatment and indicated by the midpoints of each amplicon. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$ for DMSO group versus TSA group.

expression between HK-2 cells and the human kidney. Therefore, we performed NOME-Seq to identify endogenous DNA methylation and the distribution of nucleosome occupancy in 5'-FR in HK-2 cells and the human kidney. NOME-Seq results of CpG methylation profiles showed that 5'-FR was highly methylated in HK-2 and the human kidney (Fig. 2A). In NOME-Seq, GpC methyltransferase may artificially methylate GpC sites that are not protected by nucleosomes or DNA-binding proteins. GpC methylation patterns represent the sites of a genome occupied by nucleosomes. To confirm whether GpC sites in GpC methyltransferase-treated samples were successfully methylated, we assessed GpC sites in the *GAPDH* promoter. These GpC sites were highly methylated in extracted DNA samples (Supplemental Fig. 1). The 5'-FR was occupied by more nucleosomes in HK-2 cells than in the human kidney (Fig. 2B; Supplemental Fig. 2A). We then performed NuSA to confirm nucleosome occupancy in *SGLT2* 5'-FR. Relative nucleosome occupancy was markedly higher in HK-2 cells than in the human kidney (Fig. 2C). These results indicated that *SGLT2* 5'-FR in HK-2 cells is highly occupied by nucleosomes.

Influence of the TSA Treatment on *SGLT2* Expression and Nucleosome Occupancy in *SGLT2* 5'-FR in HK-2 Cells. Histone acetylation is a well-known mechanism

for epigenetic regulation that upregulates gene expression. Histone deacetylase inhibitors stimulate gene expression by enhancing histone acetylation. In the present study, HK-2 cells were treated with TSA, the most common histone deacetylase inhibitor, to evaluate the effects of histone acetylation on *SGLT2* expression and nucleosome occupancy in *SGLT2* 5'-FR. Quantitative PCR with cDNA synthesized from total RNAs showed that the TSA treatment significantly upregulated *SGLT2* mRNA expression (Fig. 3A). The ChIP assay showed that the TSA treatment induced the acetylation of histone H3 in the region between -145 and $+135$ bp upstream of the *SGLT2* transcription start site (TSS), whereas histone H3 acetylation between -547 and -212 bp upstream of the TSS was not significantly induced (Fig. 3, B–D). We then performed NOME-Seq and NuSA to assess changes in nucleosome occupancy in 5'-FR. The nucleosome occupancy level within approximately 300 bp upstream of the TSS was decreased by the TSA treatment, whereas nucleosome occupancy was not changed in the region between -500 and -300 bp upstream of the TSS (Fig. 4; Supplemental Fig. 2B). NOME-Seq results also showed that the TSA treatment had no effect on endogenous DNA methylation in 5'-FR (Supplemental Fig. 3). These results indicate that




Fig. 5. Analysis of HNF1 α expression and HNF1 α -binding frequency after the TSA treatment in HK-2 cells. (A) ChIP analyses with quantitative PCR targeting the HNF1 α binding site in *SGLT2* 5'-FR in DMSO- or TSA-treated cells and the human kidney using antibodies against normal IgG (white bars) and HNF1 α (black bars). The relative binding frequency of HNF1 α was measured by quantitative PCR and normalized to that of the input. (B) HK-2 cells were reverse transfected with negative control or siHNF1 α for 24 hours and then treated with DMSO or TSA for 24 hours. Western blot analyses representing HNF1 α and β -actin protein expression in HK-2 cells. (C) Protein levels represent the density of the ratio for the bands of HNF1 α and β -actin in (B). Results were normalized to HNF1 α protein levels in DMSO-treated cells (control). (D) *SGLT2* mRNA levels were measured by quantitative PCR and normalized to RPL13 mRNA levels. Results are expressed relative to *SGLT2* mRNA levels in DMSO-treated cells. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$.

histone acetylation by the TSA treatment upregulates *SGLT2* expression and decreases nucleosome occupancy levels within approximately 300 bp upstream of the TSS.

Analysis of HNF1 α Expression and HNF1 α -Binding Frequency After the TSA Treatment in HK-2 Cells.

Transcription factors capable of binding to *SGLT2* 5'-FR were screened by the Database Center for Life Science Galaxy (Institute in Research Organization of Information and Systems, Chiba, Japan). A binding site of HNF1 α , which is a transcriptional modulator of *SGLT2* in the human kidney, was found between -51 and -37 bp ($-51/-37$) upstream of the *SGLT2* TSS. Chromatin immunoprecipitated (ChIP) assays with quantitative PCR demonstrated that HNF1 α binding was detected in the region between -144 and $+26$ bp upstream, including the predicted HNF1 α binding site ($-51/-37$), in the human kidney (Fig. 5A). TSA-treated HK-2 cells showed a significantly higher binding frequency than DMSO-treated cells (Fig. 5A). We next performed RNA interference with siHNF1 α to evaluate a role for HNF1 α in regulating *SGLT2* gene expression in HK-2 cells. Quantitative PCR and Western blot analyses showed that the TSA treatment caused the upregulated HNF1 α mRNA (Supplemental Fig. 4) and protein expression (Fig. 5, B and C). The transfection of siHNF1 α in DMSO- or TSA-treated HK-2 cells significantly decreased HNF1 α mRNA (Supplemental Fig. 4) and protein expression (Fig. 5, B and C). The knockdown of HNF1 α decreased *SGLT2* mRNA expression in TSA-treated HK-2 cells, whereas no significant changes were observed in DMSO-treated HK-2 cells (Fig. 5D). These results suggest that the TSA treatment upregulates *SGLT2* gene expression by increased HNF1 α binding in *SGLT2* 5'-FR in HK-2 cells.

Deletion Analysis of *SGLT2* 5'-FR in HNF1 α -Expressing HK-2 Cells. We performed luciferase assays using HNF1 α -expressing HK-2 cells to clarify whether the predicted HNF1 α binding site contributes to the transcription of the *SGLT2* gene (Fig. 6). The transfection of the HNF1 α -negative plasmid (pcDNA3.1) showed that luciferase activity levels were not significantly different in the four reporter constructs containing *SGLT2* 5'-FR from the control reporter plasmid (pGL4.10). The transfection of the HNF1 α expression plasmid significantly enhanced the luciferase activities of the $-154/+18$ construct and $-485/+18$ construct, whereas no significant differences were observed in luciferase activities between $-154/+18$ and $-485/+18$. These constructs contained the predicted HNF1 α binding site ($-51/-37$). We then measured the luciferase activity of the HNF1 α binding site-deleted construct [del($-51/-37$)]. Cells transfected with del($-51/-37$) demonstrated that the deletion of the binding site resulted in loss of HNF1 α -mediated luciferase activity. These results suggest that the predicted HNF1 α binding site ($-51/-37$) is essential for HNF1 α -mediated transcription of the *SGLT2* gene.

Quantitative Analysis of Effects of the TSA Treatment on *SGLT2* mRNA Levels in HNF1 α -Expressing HK-2 Cells. Quantitative PCR with cDNA synthesized from total RNA and nascent RNA was performed to evaluate the effects of the TSA treatment on *SGLT2* mRNA expression in HNF1 α -expressing HK-2 cells (Fig. 7). In the present study, the nascent RNA was defined as RNA transcribed after the DMSO or TSA treatment. Total and nascent *SGLT2* mRNA expression levels were significantly higher in HNF1 α -expressing cells treated with TSA than in pcDNA3.1-transfected cells. Nascent




Fig. 6. Deletion analysis of *SGLT2* 5'-FR in HNF1 α -expressing HK-2 cells. HK-2 cells were transfected with a series of reporter constructs or a control reporter plasmid (pGL4.10), together with the HNF1 α -negative plasmid (pcDNA3.1, white bars) or HNF1 α expression plasmid (black bars). The position of the deleted region is indicated with Del(-51/-37). Relative luciferase activity is expressed as a fold increase in pGL4.10 and represents the mean \pm S.D. of three independent experiments. * $P < 0.05$.

SGLT2 mRNA levels were 1.8-fold higher than total *SGLT2* mRNA in HNF1 α -expressing cells treated with TSA. In contrast, no significant changes in *SGLT2* mRNA expression levels were observed in HNF1 α -expressing cells treated with DMSO. These results indicate that histone acetylation induced by the TSA treatment plays an important role in the HNF1 α -mediated upregulation of the *SGLT2* gene.

Analysis of Nucleosome Occupancy in Human Kidney, Liver, and Small Intestine Tissues. HNF1 α is expressed in various tissues, such as the liver and intestines, whereas *SGLT2* is not. We attempted to identify nucleosome occupancy in 5'-FR containing the HNF1 α binding site in the human liver and small intestine. The region within approximately 300 bp upstream of the *SGLT2* TSS showed markedly higher nucleosome occupancy in the liver and small intestine than in the kidney (Fig. 8). These results suggest that tissue-specific nucleosome occupancy in 5'-FR plays an important role in the regulation of *SGLT2* expression via HNF1 α binding.

Discussion

In the present study, we focused on the transcriptional regulatory mechanisms of the *SGLT2* gene in the human




Fig. 7. Quantitative analysis of effects of the TSA treatment on *SGLT2* mRNA levels in HNF1 α -expressing HK-2 cells. HK-2 cells were transfected with the HNF1 α -negative plasmid (pcDNA3.1, white circles) or HNF1 α expression plasmid (black circles) and treated with DMSO or TSA for 24 hours. Total and nascent *SGLT2* mRNA levels were measured by quantitative PCR and normalized to RPL13 mRNA levels. Results are expressed relative to *SGLT2* mRNA levels in empty pcDNA3.1-transfected cells treated with DMSO. Results represent the mean \pm S.D. of three independent experiments. * $P < 0.05$ for HNF1 α group versus pcDNA3.1 group.

kidney. *SGLT2* is known to be expressed in PTECs of the human kidney. Some PTEC lines, such as LLC-PK1, KPT2, and MDCK cells, have been used to analyze expression and regulation of the *SGLT2* gene in the kidney (Da Costa-Pessoa et al., 2014; Jiang et al., 2014; Zapata-Morales et al., 2014). The human kidney-derived HK-2 cell line, which is available as a human PTEC model (Ryan et al., 1994; Wilmer et al., 2010), has been used to evaluate the regulatory mechanism for the *SGLT2* gene in PTECs of the human kidney (Panchapakesan et al., 2013). The principal portion of the kidney was divided into two distinct regions: renal cortex and renal medulla. The expression level was extremely higher in renal cortex than the renal medulla (You et al., 1995; Chen et al., 2010). The proximal tubule, which contains PTECs, is a portion of the renal cortex. Therefore, we performed in vitro analyses using the HK-2 cells derived from PTECs and the renal cortex to reveal the transcriptional profiles of the *SGLT2* gene in the human kidney.

The regulatory mechanisms for *SGLT2* gene expression in the human kidney remain unclear. Our results showed that HK-2 cells exhibited weak transcriptional activity for the *SGLT2* gene (Fig. 1A). We demonstrated that the levels of *SGLT2* mRNA and protein expression were extremely lower in HK-2 cells than the human kidney (Fig. 1, B and C). These results suggest that *SGLT2* expression is not activated in HK-2 cells, which is derived from the PTEC of the human kidney. DNA methylation, which is a well-known epigenetic mechanism for transcriptional silencing (Jones and Baylin, 2007), plays an important role in the regulation of tissue-specific gene expression by inhibiting the binding of transcription factors to DNA (Song et al., 2005; Moore et al., 2013; Medvedeva et al., 2014). Previous studies have reported that kidney-specific expression of various genes is associated with DNA methylation of the promoter region around the TSS (Aoki et al., 2008; Kikuchi et al., 2010; Azuma et al., 2012). Our results of the assessment of CpG methylation using NOME-Seq demonstrated that all CpG sites in *SGLT2* 5'-FR were highly methylated in HK-2 cells and the human kidney (Fig. 2A), suggesting that endogenous DNA methylation is not involved in the regulation of *SGLT2* expression. In contrast, we found that 5'-FR in HK-2 cells was occupied by more nucleosomes than in the human kidney (Fig. 2B). The highly occupied 5'-FR in HK-2 cells was confirmed by NuSA (Fig. 2C). Nucleosome occupancy may regulate transcription by inhibiting the access of transcription factors to promoters (Lorch et al., 1987; Kornberg and Lorch, 1999). These results suggest that the weak *SGLT2* transcriptional activity of HK-2 cells is due to high nucleosome occupancy in *SGLT2* 5'-FR.




Fig. 8. Analysis of nucleosome occupancy in human kidney, liver, and small intestine tissues. NOME-Seq data of *SGLT2* 5'-FR in the human kidney (white squares), liver (white circles), and small intestine (black circles). Graph represents the proportion of nucleosome-occupied GpC sites in 5'-FR in the three tissues. The arrow indicates the *SGLT2* TSS. Vertical lines indicate GpC sites.

Our results showed that the TSA treatment upregulated *SGLT2* mRNA expression, induced the acetylation of histone H3, and decreased nucleosome occupancy in 5'-FR within approximately 300 bp upstream of the *SGLT2* TSS (Figs. 3 and 4). TSA strongly increases the acetylation of the N-terminal tails of histone H3, which is a well-established acetylation target in nucleosome dynamics and gene activation (Choi and Howe, 2009; Hansen et al., 2010). Previous studies reported higher levels of histone H3 acetylation and lower nucleosome occupancy at active genes in promoter regions (Strenkert et al., 2011). Histone acetylation alters the nucleosomal conformation, which may increase the accessibility of transcriptional factors (Norton et al., 1989; Lee et al., 1993). Most transcribed genes have low nucleosome occupancy at promoters (Bell et al., 2011). These findings suggest that decreased nucleosome occupancy in *SGLT2* 5'-FR altered by the TSA-induced acetylation of histone H3 is involved in the upregulation of *SGLT2* expression via the binding of transcription factors to 5'-FR.

Previous studies demonstrated that the TSA-mediated upregulation of gene expression is associated with histone H3 acetylation and the recruitment of HNF1 α to the promoter (Yang et al., 2010). Our results using TSA-treated HK-2 cells showed an enhanced HNF1 α -binding frequency in *SGLT2* 5'-FR (Fig. 5A). The knockdown of HNF1 α decreased HNF1 α protein levels in DMSO- or TSA-treated HK-2 cells (Fig. 5, B and C), but *SGLT2* mRNA expression level was significantly decreased in TSA-treated cells, not in DMSO-treated cells (Fig. 5D). Furthermore, the HNF1 α overexpression, which was confirmed by Western blot analysis (Supplemental Fig. 5), significantly enhanced the transcriptional activity of 5'-FR (Fig. 6). These results suggest that the upregulation of *SGLT2* mRNA expression after the TSA treatment is associated with the enhanced HNF1 α -binding frequency in 5'-FR. In contrast, the loss of HNF1 α -mediated transcriptional activity was caused by the deletion of the predicted HNF1 α binding site (Fig. 6). Previous studies reported that a promoter region of the rodent *Sglt2* gene is located in 5'-FR around the HNF1 α binding site (Pontoglio et al., 2000; Freitas et al., 2008). In addition, reduction in the HNF1 α -binding frequency at the binding site suppresses *SGLT2* mRNA expression (Zhao et al., 2016). Collectively, our results suggest that 5'-FR around the HNF1 α binding site is a promoter region of the *SGLT2* gene.

The upregulation of *SGLT2* mRNA expression by HNF1 α was observed in TSA-treated HK-2 cells (Fig. 7). The TSA treatment also caused a decrease in nucleosome occupancy in 5'-FR containing the HNF1 α binding site (Fig. 4). Histone acetylation decreases nucleosome occupancy and enhances the accessibility of transcription factors to promoter regions

(Li et al., 2005; Zampetaki et al., 2007; Haberland et al., 2009). These results indicate that nucleosome occupancy in 5'-FR plays a critical role in interfering with the binding of HNF1 α , thereby regulating *SGLT2* expression.

Previous studies reported that *SGLT2* is predominantly expressed in the human kidney (Chen et al., 2010; Wright et al., 2011), whereas HNF1 α is strongly expressed in the human kidney, liver, and small intestine (Cereghini, 1996; Harries et al., 2009; D'Angelo et al., 2010). Our results revealed markedly higher nucleosome occupancy in the human liver and small intestine than in the kidney (Fig. 8). In contrast, the results of the assessment of CpG methylation demonstrated that all CpG sites in *SGLT2* 5'-FR were highly methylated in the human kidney, liver, and small intestine (Fig. 2A; Supplemental Fig. 3B). These results suggest that the different nucleosome occupancy levels in these tissues may affect tissue-specific HNF1 α -binding frequency and, thus, lead to tissue-specific *SGLT2* expression. This hypothesis is supported by previous findings demonstrating that tissue-specific gene expression is associated with tissue-specific nucleosome occupancy in the upstream region of the TSS (Kikuchi et al., 2010; Chen et al., 2017).

The inhibition of *SGLT2* has been reported as a new therapeutic strategy for diabetes (Chao and Henry, 2010; Terami et al., 2014). Previous studies demonstrated that *SGLT2* expression levels are increased in patients with type 2 diabetes (Rahmoune et al., 2005; Wang et al., 2017). In contrast, the recent study reported that *SGLT2* expression was significantly higher in healthy controls than participants with type 2 diabetes (Solini et al., 2017). Decreased *SGLT2* expression levels were observed in patients with maturity onset diabetes of the young type 3 (Pontoglio et al., 2000), caused by mutations in the *HNF1 α* gene (Menzel et al., 1998). However, the mechanisms responsible for the regulation of *SGLT2* expression in these diabetic patients remain unclear. Our results demonstrated that *SGLT2* expression levels are associated with the HNF1 α -binding frequency in 5'-FR. An increased HNF1 α -binding frequency in the *Sglt2* promoter region is involved in the diabetes-induced overexpression of *Sglt2* in the rat kidney (Freitas et al., 2008). In addition, a high sodium intake suppresses *SGLT2* expression by reducing the HNF1 α -binding frequency in 5'-FR (Zhao et al., 2016). Therefore, a clearer understanding of the HNF1 α -binding frequency in 5'-FR in the human kidney will provide new insights into how *SGLT2* expression is regulated in diabetic patients.

In conclusion, our results clearly demonstrate that the transcriptional regulatory mechanisms for the *SGLT2* gene are controlled by nucleosome occupancy and HNF1 α binding

in the *SGLT2* promoter. This is the first study on epigenetic mechanisms to regulate *SGLT2* expression in human tissues. Further studies are required to investigate in more detail the histone modifications causing alterations in nucleosome occupancy in the *SGLT2* gene related to the regulation of *SGLT2* expression.

Acknowledgments

We thank the Research Support Center, Graduate School of Medical Sciences, Kyushu University, for technical support.

Authorship Contributions

Participated in research design: Takesue, Hirota, Ieiri.

Conducted experiments: Takesue, Hirota, Tachimura, Tokashiki.

Performed data analysis: Takesue, Hirota.

Wrote or contributed to the writing of the manuscript: Takesue, Hirota, Ieiri.

References

- Aoki M, Terada T, Kajiwara M, Ogasawara K, Ikai I, Ogawa O, Katsura T, and Inui K (2008) Kidney-specific expression of human organic cation transporter 2 (OCT2/SLC22A2) is regulated by DNA methylation. *Am J Physiol Renal Physiol* **295**:F165–F170.
- Azuma M, Koyama D, Kikuchi J, Yoshizawa H, Thasinas D, Shiizaki K, Kuro-o M, Furukawa Y, and Kusano E (2012) Promoter methylation confers kidney-specific expression of the *Klotho* gene. *FASEB J* **26**:4264–4274.
- Bell O, Tiwari VK, Thomä NH, and Schübeler D (2011) Determinants and dynamics of genome accessibility. *Nat Rev Genet* **12**:554–564.
- Boldys A and Okopień B (2009) Inhibitors of type 2 sodium glucose co-transporters—a new strategy for diabetes treatment. *Pharmacol Rep* **61**:778–784.
- Bonner C, Kerr-Conte J, Gmyr V, Queniat G, Moerman E, Thévenet J, Beaucamps C, Delalleau N, Popescu I, Malaisse WJ, et al. (2015) Inhibition of the glucose transporter SGLT2 with dapagliflozin in pancreatic alpha cells triggers glucagon secretion. *Nat Med* **21**:512–517.
- Cereghini S (1996) Liver-enriched transcription factors and hepatocyte differentiation. *FASEB J* **10**:267–282.
- Chao EC and Henry RR (2010) SGLT2 inhibition—a novel strategy for diabetes treatment. *Nat Rev Drug Discov* **9**:551–559.
- Chen J, Li E, Zhang X, Dong X, Lei L, Song W, Zhao H, and Lai J (2017) Genome-wide nucleosome occupancy and organization modulates the plasticity of gene transcriptional status in maize. *Mol Plant* **10**:962–974.
- Chen J, Williams S, Ho S, Loraine H, Hagan D, Whaley JM, and Feder JN (2010) Quantitative PCR tissue expression profiling of the human SGLT2 gene and related family members. *Diabetes Ther* **1**:57–92.
- Choi JK and Howe LJ (2009) Histone acetylation: truth of consequences? *Biochem Cell Biol* **87**:139–150.
- da Costa-Pessoa JM, Damasceno RS, Machado UF, Beloto-Silva O, and Oliveira-Souza M (2014) High glucose concentration stimulates NHE-1 activity in distal nephron cells: the role of the Mek/Erk1/2/p90RSK and p38MAPK signaling pathways. *Cell Physiol Biochem* **33**:333–343.
- D'Angelo A, Bluteau O, Garcia-Gonzalez MA, Gresh L, Doyen A, Garbay S, Robine S, and Pontoglio M (2010) Hepatocyte nuclear factor 1alpha and beta control terminal differentiation and cell fate commitment in the gut epithelium. *Development* **137**:1573–1582.
- Freitas HS, Anhé GF, Melo KFS, Okamoto MM, Oliveira-Souza M, Bordin S, and Machado UF (2008) Na⁽⁺⁾-glucose transporter-2 messenger ribonucleic acid expression in kidney of diabetic rats correlates with glycemic levels: involvement of hepatocyte nuclear factor-1α expression and activity. *Endocrinology* **149**:717–724.
- Gerich JE (2010) Role of the kidney in normal glucose homeostasis and in the hyperglycaemia of diabetes mellitus: therapeutic implications. *Diabet Med* **27**:136–142.
- Haberland M, Montgomery RL, and Olson EN (2009) The many roles of histone deacetylases in development and physiology: implications for disease and therapy. *Nat Rev Genet* **10**:32–42.
- Hansen JC, Nyborg JK, Luger K, and Stargell LA (2010) Histone chaperones, histone acetylation, and the fluidity of the chromogenome. *J Cell Physiol* **224**:289–299.
- Harries LW, Brown JE, and Gloy AL (2009) Species-specific differences in the expression of the HNF1A, HNF1B and HNF4A genes. *PLoS One* **4**:e7855.
- Hoffman BG, Robertson G, Zavaglia B, Beach M, Cullum R, Lee S, Soukhatcheva G, Li L, Wederell ED, Thiessen N, et al. (2010) Locus co-occupancy, nucleosome positioning, and H3K4me1 regulate the functionality of FOXA2-, HNF4A-, and PDX1-bound loci in islets and liver. *Genome Res* **20**:1037–1051.
- Hu G, Schones DE, Cui K, Ybarra R, Northrup D, Tang Q, Gattinoni L, Restifo NP, Huang S, and Zhao K (2011) Regulation of nucleosome landscape and transcription factor targeting at tissue-specific enhancers by BRG1. *Genome Res* **21**:1650–1658.
- Jabbour SA and Goldstein BJ (2008) Sodium glucose co-transporter 2 inhibitors: blocking renal tubular reabsorption of glucose to improve glycaemic control in patients with diabetes. *Int J Clin Pract* **62**:1279–1284.
- Jiang M, Wang Q, Karasawa T, Koo JW, Li H, and Steyger PS (2014) Sodium-glucose transporter-2 (SGLT2; SLC5A2) enhances cellular uptake of aminoglycosides. *PLoS One* **9**:e108941.
- Jones PA and Baylín SB (2007) The epigenomics of cancer. *Cell* **128**:683–692.
- Kikuchi R, Yagi S, Kusuhara H, Imai S, Sugiyama Y, and Shiota K (2010) Genome-wide analysis of epigenetic signatures for kidney-specific transporters. *Kidney Int* **78**:569–577.
- Kornberg RD and Lorch Y (1999) Twenty-five years of the nucleosome, fundamental particle of the eukaryote chromosome. *Cell* **98**:285–294.
- Kothinti RK, Blodgett AB, Petering DH, and Tabatabai NM (2010) Cadmium down-regulation of kidney Sp1 binding to mouse SGLT1 and SGLT2 gene promoters: possible reaction of cadmium with the zinc finger domain of Sp1. *Toxicol Appl Pharmacol* **244**:254–262.
- Lee DY, Hayes JJ, Pruss D, and Wolffe AP (1993) A positive role for histone acetylation in transcription factor access to nucleosomal DNA. *Cell* **72**:73–84.
- Li B, Pattenden SG, Lee D, Gutiérrez J, Chen J, Seidel C, Gerton J, and Workman JL (2005) Preferential occupancy of histone variant H2AZ at inactive promoters influences local histone modifications and chromatin remodeling. *Proc Natl Acad Sci USA* **102**:18385–18390.
- Lorch Y, LaPointe JW, and Kornberg RD (1987) Nucleosomes inhibit the initiation of transcription but allow chain elongation with the displacement of histones. *Cell* **49**:203–210.
- Medvedeva YA, Khamis AM, Kulakovskiy IV, Ba-Alawi W, Bhuyan MSI, Kawaji H, Lassmann T, Harbers M, Forrest ARR, and Bajic VB; FANTOM Consortium (2014) Effects of cytosine methylation on transcription factor binding sites. *BMC Genomics* **15**:119.
- Menzel R, Kaisaki PJ, Rjasanowski I, Heinke P, Kerner W, and Menzel S (1998) A low renal threshold for glucose in diabetic patients with a mutation in the hepatocyte nuclear factor-1α (HNF-1α) gene. *Diabet Med* **15**:816–820.
- Misra M (2013) SGLT2 inhibitors: a promising new therapeutic option for treatment of type 2 diabetes mellitus. *J Pharm Pharmacol* **65**:317–327.
- Moore LD, Le T, and Fan G (2013) DNA methylation and its basic function. *Neuro-psychopharmacology* **38**:23–38.
- Nishimura K, Ide R, Hirota T, Kawazu K, Kodama S, Takesue H, and Ieiri I (2014) Identification and functional characterization of novel nonsynonymous variants in the human multidrug and toxin extrusion 2-K. *Drug Metab Dispos* **42**:1432–1437.
- Norton VG, Imai BS, Yau P, and Bradbury EM (1989) Histone acetylation reduces nucleosome core particle linking number change. *Cell* **57**:449–457.
- Panchapakesan U, Pegg K, Gross S, Komala MG, Mudaliar H, Forbes J, Pollock C, and Mather A (2013) Effects of SGLT2 inhibition in human kidney proximal tubular cells—renoprotection in diabetic nephropathy? *PLoS One* **8**:e54442.
- Pérez López G, González Albarrán O, and Cano Megías M (2010) Sodium-glucose cotransporter type 2 inhibitors (SGLT2): from familial renal glycosuria to the treatment of type 2 diabetes mellitus. *Nefrologia* **30**:618–625.
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* **29**:e45.
- Pontoglio M, Prié D, Cheret C, Doyen A, Leroy C, Froguel P, Velho G, Yaniv M, and Friedlander G (2000) HNF1alpha controls renal glucose reabsorption in mouse and man. *EMBO Rep* **1**:359–365.
- Rahmoune H, Thompson PW, Ward JM, Smith CD, Hong G, and Brown J (2005) Glucose transporters in human renal proximal tubular cells isolated from the urine of patients with non-insulin-dependent diabetes. *Diabetes* **54**:3427–3434.
- Ryan MJ, Johnson G, Kirk J, Fuerstenberg SM, Zager RA, and Torok-Storb B (1994) HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney. *Kidney Int* **45**:48–57.
- Solini A, Rossi C, Mazzanti CM, Proietti A, Koepsell H, and Ferrannini E (2017) Sodium-glucose co-transporter (SGLT2) and SGLT1 renal expression in patients with type 2 diabetes. *Diabetes Obes Metab* **19**:1289–1294.
- Song F, Smith JF, Kimura MT, Morrow AD, Matsuyama T, Nagase H, and Held WA (2005) Association of tissue-specific differentially methylated regions (TDMs) with differential gene expression. *Proc Natl Acad Sci USA* **102**:3336–3341.
- Strenkert D, Schmollinger S, Sommer F, Schulz-Raffelt M, and Schroda M (2011) Transcription factor-dependent chromatin remodeling at heat shock and copper-responsive promoters in *Chlamydomonas reinhardtii*. *Plant Cell* **23**:2285–2301.
- Terami N, Ogawa D, Tachibana H, Hatanaka T, Wada J, Nakatsuka A, Eguchi J, Horiguchi CS, Nishii N, Yamada H, et al. (2014) Long-term treatment with the sodium glucose cotransporter 2 inhibitor, dapagliflozin, ameliorates glucose homeostasis and diabetic nephropathy in db/db mice. *PLoS One* **9**:e100777.
- Tsui K, Dubuis S, Gebbia M, Morse RH, Barkai N, Tirosh I, and Nislow C (2011) Evolution of nucleosome occupancy: conservation of global properties and divergence of gene-specific patterns. *Mol Cell Biol* **31**:4348–4355.
- Vallon V, Rose M, Gerasimova M, Satriano J, Platt KA, Koepsell H, Cunard R, Sharma K, Thomson SC, and Rieg T (2013) Knockout of Na-glucose transporter SGLT2 attenuates hyperglycemia and glomerular hyperfiltration but not kidney growth or injury in diabetes mellitus. *Am J Physiol Renal Physiol* **304**:F156–F167.
- Wang XX, Levi J, Luo Y, Myakala K, Herman-Edelstein M, Qiu L, Wang D, Peng Y, Grenz A, Lucia S, et al. (2017) SGLT2 protein expression is increased in human diabetic nephropathy: SGLT2 protein inhibition decreases renal lipid accumulation, inflammation, and the development of nephropathy in diabetic mice. *J Biol Chem* **292**:5335–5348.
- Wilmer MJ, Saleem MA, Masereeuw R, Ni L, van der Velden TJ, Russel FG, Mathieson PW, Mommens LA, van den Heuvel LP, and Levchenko EN (2010) Novel conditionally immortalized human proximal tubule cell line expressing functional influx and efflux transporters. *Cell Tissue Res* **339**:449–457.
- Workman JL (2006) Nucleosome displacement in transcription. *Genes Dev* **20**:2009–2017.
- Wright EM (2001) Renal Na⁽⁺⁾-glucose cotransporters. *Am J Physiol Renal Physiol* **280**:F10–F18.
- Wright EM, Luo DDF, and Hirayama BA (2011) Biology of human sodium glucose transporters. *Physiol Rev* **91**:733–794.
- Yang H, Nie Y, Li Y, and Wan Y-JY (2010) Histone modification-mediated CYP2E1 gene expression and apoptosis in HepG2 cells. *Exp Biol Med (Maywood)* **235**:32–39.

- Yang Z, Yoshioka H, and McCarrey JR (2013) Sequence-specific promoter elements regulate temporal-specific changes in chromatin required for testis-specific activation of the *Pgk2* gene. *Reproduction* **146**:501–516.
- You G, Lee WS, Barros EJG, Kanai Y, Huo TL, Khawaja S, Wells RG, Nigam SK, and Hediger MA (1995) Molecular characteristics of Na(+)-coupled glucose transporters in adult and embryonic rat kidney. *J Biol Chem* **270**:29365–29371.
- Zampetaki A, Zeng L, Xiao Q, Margariti A, Hu Y, and Xu Q (2007) Lacking cytokine production in ES cells and ES-cell-derived vascular cells stimulated by TNF-alpha is rescued by HDAC inhibitor trichostatin A. *Am J Physiol Cell Physiol* **293**:C1226–C1238.
- Zapata-Morales JR, Galicia-Cruz OG, Franco M, and Martinez Y Morales F (2014) Hypoxia-inducible factor-1 α (HIF-1 α) protein diminishes sodium glucose transport 1 (SGLT1) and SGLT2 protein expression in renal epithelial tubular cells (LLC-PK1) under hypoxia. *J Biol Chem* **289**:346–357.
- Zhang T, Zhang W, and Jiang J (2015) Genome-wide nucleosome occupancy and positioning and their impact on gene expression and evolution in plants. *Plant Physiol* **168**:1406–1416.
- Zhao Y, Gao P, Sun F, Li Q, Chen J, Yu H, Li L, Wei X, He H, Lu Z, et al. (2016) Sodium intake regulates glucose homeostasis through the PPAR δ /adiponectin-mediated SGLT2 pathway. *Cell Metab* **23**:699–711.

Address correspondence to: Dr. Ichiro Ieiri, Department of Clinical Pharmacokinetics, Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka, 812-8582, Japan. E-mail: ieiri-ttr@umin.ac.jp

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri

Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology

Supplemental Table 1

Primer sequences for the cloning of *SGLT2* reporter plasmids

Primer		Sequence (5' to 3') ^a	Annealing (°C)	Restriction enzymes
Cloning of the <i>SGLT2</i> 5'-flanking region				
-3185	Forward	TGGCCTAACTGGCCGGTACCTTCCCGACCGCCT	60.0	-
+18	Reverse	AGTACCGGATTGCCACTCCCCAGGATCTGCCCC		
In-Fusion cloning and site-directed mutagenesis				
Del-(-51/-37)	Forward	GGCTCAGTGCCCCTGCTTCCCCTGGGGGAATCC	60.0	-
	Reverse	CAGGGGCACTGAGCCGACAAGTCCCCCAGGTCT		
-2320	Forward	GTTTGTAAATGAAGGAAGGT <u>ACCAGGAAGGAAGAAAGA</u>	60.0	Kpn□
	Reverse	TCTTTCCTTCCTTCCTGGT <u>TACCTTCCTTCATTAACAAAC</u>		
-1587	Forward	CCAAGTCTCTTTGTGGT <u>TACCTGACAAATGACACAC</u>	60.0	Kpn□
	Reverse	GTGTGTCATTTGTCAGGGT <u>ACCACAAAGAGCAGTTGG</u>		
-485	Forward	CAAAAATCTGGGCTGGT <u>TACCTTAAAGGAGTGGGAAAGGA</u>	60.0	Kpn□
	Reverse	TCCTTTCCTTCCTTTAAGGT <u>ACCAGCCCAGATTTTGT</u>		
-154	Forward	TGGAAGGGCCCAGGT <u>ACCCAAGACCAGCC</u>	60.0	Kpn□
	Reverse	GGCTGGTCTTGGT <u>TACCTGGGCCCTTCCA</u>		
-44	Forward	GGCTCAGTGCCCCTGAGGT <u>ACCCATTAATCCTTC</u>	60.0	Kpn□
	Reverse	GAAGGATTAATGGT <u>TACCTCAGGGGCACTGAGCC</u>		

^aThe restriction site is underlined, and nucleotide changes are marked in bold letters.

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri

Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology

Supplemental Table 2

Direct sequence oligonucleotides for the cloning of *SGLT2* reporter plasmids

Location		Sequence (5' to 3')
SGLT2	Antisense	TGAGAGAAATCCAGTGCCAAGT
	Antisense	CCTGAGATGAGAATTTGTGTGC
	Sense	GCTTTGTTGGTTTTTCTCCTTGT
	Sense	CCACACCCAGCCAGTCCTAC
	Sense	GGAAGGATGAGCGGGAATTG
pGL4.10	Sense	GCAGGTGCCAGAACATTTCT

Supplemental Table 3

Primer sequences and oligonucleotides for the cloning of the HNF1 α expression plasmid

Primer		Sequence (5' to 3')
Cloning of the HNF1 α coding region		
	Forward	CAGTGTGGTGAATTATGGTTTCTAAACTGAGCCA
	Reverse	GCCACTGTGCTGGATTTACTGGGAGGAAGAGGC
Direct sequencing		
HNF1 α	Sense	AGCAGTTCACCCATGCAGG
T7	Sense	TTGTAATACGACTCACTATAG
BGH	Antisense	TAGAAGGCACAGTCGAGG

Supplemental Table 4

Primer sequences for quantitative PCR

Gene		Sequence (5' to 3')	Position
SGLT2	Forward	TTCAGTCTCCGGCATAGCAA	1700 to 1719
	Reverse	CATCTCCATGGCACTCTCTGG	1807 to 1787
HNF1 α	Forward	CCCTGGGTCCTACGTTCA	1463 to 1480
	Reverse	GGGTCACATGGCTCTGCA	1657 to 1640
RPL13	Forward	GAGACAGTTCTGCTGAAGAACTGAA	486 to 510
	Reverse	TCCGGACGGGCATGAC	551 to 536

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri

Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology

Supplemental Table 5

Primer sequences for NOME-Seq

Primer		Sequence (5' to 3')	Position
Bisulfite PCR			
SGLT2	Forward	TGGGAAAGGATTTTTGATTTTTTTT	-477 to -454
	Reverse	CCCCTAAATCCCCCAAAAA	-14 to -33
GAPDH	Forward	GGGTTTTTGTTTTTGATTTTTTAGTGTTT	-220 to -192
	Reverse	CAATCCCAGCCCAAATCTTAAA	+25 to +3
Colony PCR			
T7	Forward	TTCAGTCTCCGGCATAGCAA	
SP6	Reverse	CATCTCCATGGCACTCTCTGG	

Supplemental Table 6

Primer sequences for NuSA

Position		Sequence (5' to 3')	Midpoint
-547 to -522	Forward	TTTGGTGGGGATAAAATATCTGGTCA	-462
-377 to -400	Reverse	TCTTCAGCCTGATTTCCAATCCTG	
-508 to -487	Forward	GCAAAAATCTGGGCTGGGTAGG	-445
-383 to -405	Reverse	GCCTGATTTCCAATCCTGGTCAT	
-454 to -430	Forward	CTAGATTTGGTTTGGAGAAGCAGGG	-381
-309 to -334	Reverse	TTTCAAATCCAAGTCTGACAGGGTC	
-419 to -402	Forward	GCGGGAATTGGGGCATGA	-353
-287 to -318	Reverse	TTTAACTAATCCAGAGGAATCATTTTCAAATC	
-365 to -342	Forward	GAGCTATGGAGGGTTCCTGAGGAG	-289
-214 to -237	Reverse	TGCTCCAGGCTCAAATCACTCTT	
-316 to -287	Forward	TTTGAAAATGATTCCTCTGGATTAGTTAAA	-258
-200 to -217	Reverse	CGCCCTCTCCCCTGTGCT	
-316 to -287	Forward	TTTGAAAATGATTCCTCTGGATTAGTTAAA	-236
-157 to -179	Reverse	GCCCTTCCAAGTTCAAGAGCACT	
-237 to -214	Forward	AAGAGTGATTTTGAGCCTGGAGCA	-170
-104 to -131	Reverse	TGTTTAGCTGAATCAGGTCATATCAAGG	
-181 to -158	Forward	AGAGTGCTCTTGAACCTGGAAGGG	-123
-65 to -84	Reverse	CCGACAAGTCCCCCAGGTCT	
-144 to -120	Forward	GACCAGCCTTCAGCCTTGATATGAC	-69
+6 to -13	Reverse	CCCCATCCAGGAACCAGCC	
-91 to -71	Forward	GGGAATGAGACCTGGGGGACT	-25
+41 to +20	Reverse	CTGCCTCTGTGTGCTCCTCCAT	
-17 to +1	Forward	GGGGCTGGTTCCTGGATG	+51
+119 to +95	Reverse	GGAAATATGCAGCAATGACTAGGAT	
+3 to +23	Forward	GGCAGATCCTGGGGAGAATGG	+74
+144 to +124	Reverse	CACAAGCCAACGCCAATGACC	

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri

Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology

Supplemental Table 7


Primer sequences for the ChIP assay

Name		Sequence (5' to 3')	Position	Annealing (°C)
Semi-quantitative PCR				
Distal	Forward	TTTGGTGGGGATAAAATATCTGGTCAA	-547 to -521	60.0
	Reverse	TGTGCTCCAGGCTCAAATCACTC	-212 to -235	
Proximal	Forward	AGACCAGCCTTCAGCCTTGATATGA	-145 to -121	60.0
	Reverse	ACGCCAATGACCAGCAGGAAATA	+135 to +113	
Quantitative PCR				
	Forward	GACCAGCCTTCAGCCTTGATATGACC	-144 to -119	
	Reverse	CCTCCATTCTCCCCAGGATCTGC	+26 to +4	

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri


Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology




Supplemental Figure 1 NOME-Seq analysis of nucleosome occupancy in *GAPDH* 5'-FR in HK-2 cells and human tissues.

The arrow indicates the *GAPDH* TSS. Vertical lines indicate GpC sites. White circles represent unmethylated GpC sites and black circles represent methylated GpC sites. Gray bars represent nucleosome occupancy, which is the region of consecutive unmethylated GpC sites over 147 bp.




Supplemental Figure 2 Analysis of nucleosome occupancy in HK-2 cells and the human kidney

NOME-Seq data of *SGLT2* 5'-FR in (A) HK-2 cells and the human kidney, and (B) DMSO- and TSA-treated HK-2 cells. Graph represents the proportion of nucleosome-occupied GpC sites in 5'-FR in the three tissues. The arrow indicates the *SGLT2* TSS. Vertical lines indicate GpC sites.



Supplemental Figure 3 Analysis of endogenous DNA methylation in HK-2 cells and human tissues.

NOMe-Seq data of *SGLT2* 5'-FR in (A) DMSO- or TSA-treated HK-2 cells and (B) human liver and small intestine. The arrow indicates the *SGLT2* TSS. Vertical lines indicate CpG sites. White circles represent unmethylated CpG sites and black circles represent methylated CpG sites.




Supplemental Figure 4 Analysis of HNF1α mRNA expression in siHNF1α-transfected HK-2 cells

HK-2 cells were reverse-transfected with negative control (white bars) or siHNF1α (black bars) for 24 hours, and then treated with DMSO or TSA for 24 hours. HNF1α mRNA levels were measured by quantitative PCR and normalized to RPL13 mRNA levels. Results are expressed relative to HNF1α mRNA levels in the control group. Results represent the mean ± SD of three independent experiments. *P < 0.05.

Hiroaki Takesue, Takeshi Hirota, Mami Tachimura, Ayane Tokashiki, Ichiro Ieiri

Nucleosome positioning and gene regulation of the *SGLT2* gene in the human kidney

Molecular Pharmacology



Supplemental Figure 5 Analysis of HNF1 α expression in HNF1 α -expressing HK-2 cells

Western blot analyses representing HNF1 α and β -actin protein expression in HNF1 α -expressing HK-2 cells with DMSO or TSA treatment.