

**MOLPHARM 2004 0497**

## **Mini-review**

### **G protein-coupled receptor dimerization: function and ligand pharmacology**

Graeme Milligan

Molecular Pharmacology Group

Division of Biochemistry and Molecular Biology

Institute of Biomedical and Life Sciences

University of Glasgow

Glasgow G12 8QQ

Scotland, U.K.

MOLPHARM 2004 0497

Running title: GPCR dimerization

### Manuscript Information

Text pages 32

Tables 0

Figures 0

References 85

Total word count 4,548

Abstract 87

Introduction: Not relevant

Discussion Not relevant

Address all correspondence to:

Graeme Milligan, Molecular Pharmacology Group, Davidson Building, University of  
Glasgow, Glasgow G12 8QQ Scotland, U.K.

Tel: 44 (0) 141 330 5557

FAX: 44 (0) 141 330 4620

e-mail: [g.milligan@bio.gla.ac.uk](mailto:g.milligan@bio.gla.ac.uk)

<sup>1</sup> Abbreviations: FRET; fluorescence resonance energy transfer, BRET;  
bioluminescence resonance energy transfer, GPCR; G protein–coupled receptor.

MOLPHARM 2004 0497

## **Abstract**

It is now generally accepted that G protein-coupled receptors (GPCRs) can exist as dimers or as part of larger oligomeric complexes. Increasing evidence suggests that a dimer is the minimal functional structure but considerable variation exists between reports of the effects of agonist ligands on quaternary structure. Many studies have intimated the existence of hetero-dimeric GPCR pairings. Key questions that remain to be addressed effectively include the prevalence and relevance of these in native tissues and the implications of hetero-dimerization for pharmacology and, potentially, for drug design.

## **Introduction**

Although the generation of functional complexes via dimerization and oligomerization is a common theme for polypeptides including membrane proteins (Woolf and Linderman, 2003), for many years G protein-coupled receptors (GPCRs<sup>1</sup>) were assumed to exist and function as monomeric species. In many regards this was understandable. Both the expression of single DNA species encoding GPCRs and functional reconstitution studies employing artificial lipid vesicles and purified proteins indicated that the product of a single gene and single polypeptides chains were able to generate the anticipated pharmacology and function. Of course such studies did not address, nor indeed seek to address, whether the GPCR functioned as a monomer or if function required quaternary structure. In the recent past the concept that GPCRs can exist as dimers or higher-order oligomers has advanced rapidly (see Angers et al., 2002, Lee et al., 2003, Milligan et al., 2003, for reviews). There is also growing appreciation that GPCR dimerization may be a requisite for function. A

MOLPHARM 2004 0497

wide range of approaches has contributed to the view that individual GPCRs can exist as homo-dimers. These include the analysis of ligand binding studies (Armstrong and Strange, 2001), the tendency of GPCRs to migrate, even in SDS-polyacrylamide gels, as complexes with the molecular mass anticipated for dimers and/or higher-order structures (Nimchinsky et al., 1997), the ability to co-immunoprecipitate co-expressed but differentially epitope-tagged forms of a GPCR (Hebert et al., 1996, Cjevic and Devi, 1997) and the capacity to reconstitute function following the co-expression of distinct pairs of mutants that are both individually non-functional (Lee et al., 2002, Carrillo et al., 2003). As well as reductionist techniques, a range of studies have employed methods such as fluorescence resonance energy transfer (FRET) (Overton and Blumer 2000, Rocheville et al., 2000a, McVey et al., 2001, Dinger et al., 2003, Stanasila et al., 2003), bioluminescence resonance energy transfer (BRET) (Angers et al., 2000, McVey et al., 2001, Cheng and Miller, 2001, Kroeger et al., 2001, Ayoub et al., 2002, Issafras et al., 2002, Babcock et al., 2003) and imaging of functional reconstitution (Carrillo et al., 2003) to explore such interactions in living cells, either at the population level or in single cells. Essentially in parallel with these efforts, studies using similar techniques have noted the capacity of many pairs of GPCRs to form hetero-dimers or -oligomers (Marshall, 2001, George et al., 2002). This is clearly not a random process (Milligan et al., 2004) and in certain cases, for example the GABA<sub>B</sub> receptor (Marshall et al., 1999, Mohler and Fritschy, 1999) and taste receptors (Zhao et al., 2003, Matsunami and Amrein, 2004), selective dimerization defines ligand pharmacology and response. However, although both direct experiments (Ramsay et al., 2002) and computational strategies (Filizola et al., 2002, Soyer et al., 2003) have been employed to analyse such interactions, understanding of

MOLPHARM 2004 0497

the molecular basis of hetero-dimerization and of the relative propensity of GPCRs to hetero-dimerize is currently in its infancy.

One key question that remains contentious is how, or indeed if, receptor ligands alter the dimerization or organizational structure of GPCRs. The growing literature on GPCR hetero-dimerization has also resulted in attempts to address if ligand pharmacology or function at hetero-dimer pairs is distinct from that at the corresponding homo-dimers. It is possible this might provide molecular explanations for tissue pharmacology that is not easily explained by heterologous expression of individual GPCRs. Neither of these questions can be addressed fully at this stage. However, in this commentary I review data pertinent to these points and try to encapsulate key issues that require further study. As many of the approaches used cannot effectively discriminate between dimers and higher-order oligomers, I have used the term 'dimer' except in situations in which higher-order organization appears likely.

### **Are all GPCRs dimers?**

The most convincing demonstration that members of the rhodopsin-related, family A GPCRs exist as dimers *in situ*, and may indeed be further arranged into oligomeric arrays, derives from the application of atomic force microscopy to the organization of rhodopsin in rod outer segment membranes (Liang et al., 2003). Although some reservations have been expressed about the preparation procedures employed and the implications for molecular organization of biophysical experiments conducted a number of years ago (Chabre et al., 2003) these studies demonstrated

MOLPHARM 2004 0497

rhodopsin to be arranged in para-crystalline arrays. Models based on these pictures were inferred to indicate that intra-dimer links might be provided by contacts between transmembrane helices IV and V with further organizational structure derived from interactions involving transmembrane helices I and II. Only occasional rhodopsin monomers were observed. It could be argued that the high density of rhodopsin in the rod outer segment might impose regularity and organization above and beyond that required for GPCRs expressed at much lower density. However, virtually all of the basic functions and mechanisms of rhodopsin have subsequently been found to have parallels in other family A GPCRs and as such, it has provided an excellent template to explore and understand other members of this family.

It is difficult to ascertain the fraction of any family A GPCR that exists as dimers. Although GPCR aggregates with molecular mass corresponding to dimers often survive resolution by SDS-PAGE, the relative fraction of apparent monomers to dimers observed in such studies may indicate more about the relative interaction affinities of GPCR pairs than the fraction present as dimers in intact cells. Attempts at quantitation derived from FRET-based studies on neuropeptide Y receptor subtypes suggested that some 26% of NPY Y2 receptor and 44% of NPY Y1 receptor constructs were able to generate effective FRET signals. These numbers are likely to represent minimum estimates of dimerization as they were calculated relative to an NPY Y2 receptor construct with a FRET positive control attached to its sequence (Dinger et al., 2003). Studies employing saturation BRET, in which various ratios of *Renilla* luciferase and GFP-tagged forms of the  $\beta_2$ -adrenoceptor were co-expressed

MOLPHARM 2004 0497

(Mercier et al., 2002), estimated that more than 80% of this GPCR was present in dimers. Because GPCR dimerization is difficult to analyse in native cells and tissues it has also been suggested that some examples of GPCR dimerization may, at least in part, be artefacts derived from the high levels of expression that can often be achieved in heterologous cell systems. Although Mass-Action implies this issue must be taken seriously, Mercier et al., (2002) estimated that the extent of dimerization of the  $\beta_2$ -adrenoceptor was unchanged over a 20 fold range of GPCR expression levels.

Equally, although it is often convenient to monitor signals in resonance energy transfer experiments with higher levels of expression, constitutive dimerization of the CCR5 chemokine/HIV co-entry receptor was observed at levels comparable to those expressed endogenously (Issafras et al., 2002).

Although the preceding discussion has concentrated on the family A GPCRs because they are by far the largest grouping, key information on constitutive dimerization and the inherent requirement of this for membrane delivery has been gleaned from study of the metabotropic glutamate-like family C GPCRs that all possess long N-terminal sequences. Although these share no homology with the family A GPCRs, both the recognition that the GABA<sub>B</sub>R1 GPCR is not trafficked effectively to the cell surface in the absence of expression of the GABA<sub>B</sub>R2 (Marshall et al., 1999) and that the extracellular domain of the metabotropic glutamate receptor 1a forms a constitutive dimer (Kunishima et al., 2000) were integral to more widespread appreciation of the general nature of GPCR dimerization. The intrinsic requirement of dimerization of the family C GPCRs for function is so firmly established that this is no longer a point of contention.

MOLPHARM 2004 0497

### **Why should GPCRs be dimers?**

There is growing evidence that GPCR-GPCR interactions occur initially during biosynthesis. Maturing forms of most family A GPCRs become terminally N-glycosylated and this is likely a key ‘quality control’ step prior to export from the Golgi apparatus. At least for the family C GABA<sub>B</sub> receptor (White et al, 1998) this seems to require protein-protein interactions provided by dimerization. In studies of oxytocin and vasopressin receptors, Terrillon et al., (2003) demonstrated immature forms of these GPCRs were present as dimers whilst still present in the endoplasmic reticulum. This again suggests that dimerization may be an integral aspect of GPCR maturation and that dimerization is an early commitment step in the production of functional GPCRs. It is also noteworthy that both artificially produced and naturally occurring GPCR truncation and splice variants can act to limit cell surface delivery of GPCRs (Coge et al., 1999, Lee et al., 2000, Karpa et al., 2000). This is likely to reflect interactions that do not result in final maturation of the dimer.

Models generated from the structural organization of rhodopsin and the actual molecular size of both this GPCR and G proteins indicate that a dimer would be required to provide an appropriate ‘footprint’ for the binding of a single heterotrimeric G protein consisting of transducin  $\alpha$  and the  $\beta_1 + \gamma_1$  complex (Liang et al., 2003). It is clear that GPCRs are able to bind G protein  $\alpha$  and  $\beta/\gamma$  subunits independently and that  $\beta/\gamma$  interactions contribute substantially to the ability of agonists to activate G protein  $\alpha$  subunits (Liu et al., 2002). Biophysical studies following expression in *E. Coli*, denaturation and refolding of the leukotriene B<sub>4</sub> BLT1 receptor followed by reconstitution of the GPCR with a heterotrimeric G protein consisting of  $G\alpha_{i2}$ ,  $\beta_1$  and



MOLPHARM 2004 0497

$\gamma_2$  also provided compelling evidence for the overall assembly of a pentameric (BLT1)<sub>2</sub> G $\alpha_{i2}$ ,  $\beta_1$  and  $\gamma_2$  complex (Baneres and Parello, 2003). Interestingly, the structure of arrestins (Han et al., 2001) is also consistent with the binding of a single arrestin across the cytoplasmic face of a GPCR dimer. Thus, binding of a single arrestin might be expected to desensitize a GPCR dimer.

One early attempt to disrupt GPCR dimerization involved the use of a peptide corresponding to transmembrane helix VI of the  $\beta_2$ -adrenoceptor. As monitored by the relative proportions of immunodetected GPCR migrating through SDS-PAGE in positions consistent with monomer and dimer, the proportion of dimer was decreased by this peptide in a concentration-dependent manner (Hebert et al., 1996). This correlated with a reduction in agonist (but not forskolin or NaF)-mediated adenylyl cyclase activity (Hebert et al., 1996). Such observations would certainly be consistent with the idea that high affinity binding of G protein, and therefore effective G protein-mediated signal transduction, would require the GPCR dimer. A key dimer interface in the *Saccharomyces cerevisiae*  $\alpha$ -factor GPCR involves the glycoporphin A-like GXXXG dimerization sequence that is present in transmembrane helix I (Overton et al., 2003). Mutation of this sequence inhibited protein-protein interactions as monitored in FRET studies and also resulted in a marked reduction in ligand-mediated signal generation. This was argued not simply to reflect poor membrane delivery of the mutated receptor and shown not to reflect poor agonist affinity. Early studies on dimerization of the dopamine D2 receptor indicated that as well as a peptide corresponding to transmembrane region VI, an equivalent peptide corresponding to transmembrane region VII was also able to reduce the proportion of dimer (Ng et al., 1996). Few subsequent studies or models of GPCR dimers have suggested a likely

MOLPHARM 2004 0497

role for transmembrane region VII as a dimer interface. However, a recent study directly introduced peptides corresponding to this region from a variety of GPCRs into rats and recorded a remarkable range of loss of physiological functions that would be anticipated to reflect activation of the corresponding receptors (George et al., 2003). Both these authors (Lee et al., 2003) and others (Guo et al., 2003) have also indicated a key role for transmembrane domain IV in dimerization of the dopamine D2 receptor. Thus, although it is difficult to rationalize the molecular basis of these results they would be consistent with reduction or loss of GPCR function resulting from disruption of specific GPCR homo-dimers. Such studies would certainly benefit from independent replication.

### **Are GPCRs permanent dimers?**

As discussed above, emerging evidence indicates dimerization to occur early in the biogenesis of at least certain GPCRs. This supports a model in which GPCRs are delivered to the plasma membrane as dimers. Reports indicating that ligands can regulate GPCR dimerization do not seem consistent with this scenario and will be discussed later. The genetic tractability of *Saccharomyces cerevisiae* resulted in early evidence that the  $\alpha$ -factor GPCR is internalized as a dimeric complex (Yesilaltay and Jenness, 2000). The intracellular complement of GPCRs, such as the  $\alpha_{1a}$ -adrenoceptor, that constitutively internalize at a rapid rate in the absence of ligands, also appears to be composed of dimers (Ramsay et al., 2004). There is therefore growing evidence that the entire life-cycle of some GPCRs takes place within a dimeric complex.

MOLPHARM 2004 0497

The concept that GPCRs internalize as dimers has been utilized to attempt to explore the selectivity of GPCR hetero-dimerization. For example, following co-expression of the  $\alpha_{1a}$ - and  $\alpha_{1b}$ -adrenoceptors, the  $\alpha_{1a}$ -selective agonist oxymetazoline was able to induce co-internalization of the two GPCRs. However, this agonist did not produce co-internalization of either the NK1 tachykinin receptor or the CCR5 chemokine receptor when they were co-expressed with the  $\alpha_{1a}$ -adrenoceptor (Stanasila et al., 2003). This suggests that the  $\alpha_{1a}$ - and  $\alpha_{1b}$ -adrenoceptors form a high affinity hetero-dimer but that the other pairs do not. In a similar vein, co-expression of the NK1 receptor with the MOP opioid receptor resulted in co-internalization of these two GPCRs in response to addition of selective ligands for either receptor (Pfeiffer et al., 2003). It has also been reported that when the  $\beta_2$ -adrenoceptor is co-expressed with the DOP opioid receptor, internalization of the opioid receptor is seen in response to isoproterenol (Jordan et al., 2001). However, when the  $\beta_2$ -adrenoceptor was co-expressed with the KOP opioid receptor, isoproterenol treatment failed to induce  $\beta_2$ -adrenoceptor internalization (Jordan et al., 2001). These observations are of considerable interest but also pose problems of interpretation. The  $\beta_2$ -adrenoceptor-DOP opioid receptor data indicate a capacity of these two GPCRs to interact and that agonist-occupation of one element of the hetero-dimer is sufficient to induce internalization. The KOP opioid receptor is much more resistant to homologous ligand-induced internalization than the DOP opioid receptor. A reasonable interpretation of these results is that the production of a KOP opioid receptor- $\beta_2$ -adrenoceptor hetero-dimer restrains or blocks ligand-induced internalization. Interestingly, however, direct measurements indicate the propensity of the  $\beta_2$ -adrenoceptor and KOP opioid receptors to form a hetero-dimer is much less than for

MOLPHARM 2004 0497

the KOP opioid receptor to form a homo-dimer or a hetero-dimer with the DOP opioid receptor (Ramsay et al., 2002). KOP opioid receptor- $\beta_2$ -adrenoceptor interactions can be measured, but only at high expression levels (Ramsay et al., 2002). This raises questions about the physiological relevance of such a dimer. If two co-expressed GPCRs, A and B, have equivalent propensity to form homo and hetero-dimers, then 50% of the population should be represented by A-A and B-B. Therefore, if the KOP opioid receptor and the  $\beta_2$ -adrenoceptor are co-expressed but have only relatively low mutual affinity then it must be anticipated that the majority of dimers formed would be  $\beta_2$ -adrenoceptor- $\beta_2$ -adrenoceptor and KOP opioid receptor-KOP opioid receptor homo-dimers. There is no reason to suppose that a  $\beta_2$ -adrenoceptor- $\beta_2$ -adrenoceptor homo-dimer would not internalize in response to isoproterenol. Although at the current time this is pure speculation, the data of Jordan et al. (2001) may therefore indicate the production of a larger oligomeric complex containing copies of both GPCRs. When the  $\beta_2$ -adrenoceptor and the DOP opioid receptor were stably co-expressed at relatively modest levels in HEK293 cells, agonists at both GPCRs were required to produce internalization of the two polypeptides (Tsao and von Zastrow, 2000). Interestingly, the trafficking patterns of the two GPCRs could be resolved (Tsao and von Zastrow, 2000). These observations suggest that if the GPCRs are permanent dimers, then the homo-dimers predominate or if there is a significant hetero-dimer population these are separated into monomers during intracellular trafficking. Other data, discussed later, would allow the second hypothesis. Interactions with  $\beta$ -arrestins to allow clathrin-mediated internalization are consistent with a trimeric (GPCR) $_2$ - $\beta$ -arrestin complex (and therefore the GPCR dimer) persisting at least until removal of  $\beta$ -arrestin within acidified endosomal

MOLPHARM 2004 0497

vesicles. This concept also raises intriguing questions about the fate of GPCR hetero-dimers that incorporate so called type A and type B GPCRs. These are separated according to their affinity of interactions with  $\beta$ -arrestins. The type A GPCRs have relatively poor  $\beta$ -arrestin affinity and are rapidly separated from it following internalization. The type B GPCRs have substantially higher affinity for, and generally co-internalise with,  $\beta$ -arrestins into endocytic vesicles (Shenoy and Lefkowitz, 2003). Co-expression of pairs of related GPCRs such the thyrotropin releasing hormone receptors 1 and 2 resulted in altered patterns of  $\beta$ -arrestin subtype trafficking than seen for either GPCR alone (Hanyaloglu et al., 2002). Terrillon et al., (2004) have recently reported similar data following co-expression of the type B vasopressin V2 receptor and the type A vasopressin V1a receptor. This clearly is an area requiring further detailed analysis. However, the capacity of  $\beta$ -arrestin-2 to act as a scaffolding protein for some pathways of GPCR-mediated ERK MAP kinase activation (Shenoy and Lefkowitz, 2003) may underlie observations of altered regulation of this pathway following co-expression of pairs of GPCRs. This has been a popular endpoint measure used to support potential distinct functions of GPCR hetero-dimers.

### **What is the propensity of GPCRs to form hetero-dimers?**

Before we can attempt to rationalize potential variations in ligand function and pharmacology with GPCR hetero-dimerization we must understand the expression pattern of the GPCR family. The best estimates of the number of functional GPCR genes in man are for some 339 encoding olfactory receptors (Malnic et al., 2004) and

MOLPHARM 2004 0497

in the region of 400 others (Fredriksson et al., 2003, Vassilatis et al., 2003).

Assuming that olfactory GPCRs do indeed form dimers (and difficulties in expressing these in heterologous systems have restricted analysis of this issue) then at least in the olfactory system they are restricted to homo-dimers because each olfactory neurone expresses only one member of the family. Although not yet fully confirmed, microarray and related studies have, however, indicated a surprisingly widespread distribution of a number of olfactory GPCRs (Hakak et al., 2003). Microarray-based information has confirmed general suspicions that individual organs and small regions of tissue may express up to 100 distinct GPCRs (Hakak et al., 2003). However, at least in the public domain, such information is lacking for individual cells. Despite this, it is clear from the sensitivity of widely used cell culture lines to chemicals used in high throughput screens that a substantial number of GPCRs may be co-expressed in detectable amounts. Although transport of mRNAs to different cellular locations prior to translation in terminally differentiated and polarized cells may prevent co-temporal and spatial synthesis of some potential GPCR dimer pairings, the above information does suggest the possibility of a potentially bewildering set of combinations.

One enduring and frustrating aspect of studies of GPCR dimerization selectivity is the difficulty in producing clear and unambiguous negative controls. For example, in co-immunoprecipitation studies Salim et al., (2002) reported the capacity of all GPCRs tested to be co-immunoprecipitated with the serotonin 5-HT<sub>1A</sub> receptor. In studies of a number of family A GPCRs negative controls have utilised family C GPCRs, such as the GABA<sub>B</sub>R1, that have no inherent homology. However, even this may not be

MOLPHARM 2004 0497

absolute. Interactions have been reported between the family A adenosine A<sub>2A</sub> receptor and the family C metabotropic Glu<sub>5</sub> receptor (Ferre et al., 2002), although the molecular basis for this interaction remains unknown. The chemokine CCR5 receptor has also been a popular negative control for non-chemokine GPCRs. Relatively few studies have tried to assess relative interaction affinities directly but saturation BRET techniques (Mercier et al., 2002) offer one approach. The most obvious hypothesis is that more homologous GPCRs would show higher interaction affinities. This is clearly too simplistic. Saturation BRET studies indicate the  $\beta_1$ -adrenoceptor and  $\beta_2$ -adrenoceptor can form hetero-dimers with similar affinity as the corresponding homo-dimers (Mercier et al., 2002). Equally, KOP opioid receptor-DOP opioid hetero-dimers form at least as effectively as KOP opioid receptor homo-dimers (Ramsay et al., 2002). However, the inability of the  $\alpha_{1a}$ -adrenoceptor and  $\alpha_{1d}$ -adrenoceptors to form hetero-dimers (Uberty et al., 2003), the reported lack of interactions between KOP and MOP opioid receptors (Jordan and Devi, 1999) and the ability of only some somatostatin receptor subtypes to generate hetero-dimers (Rocheville et al., 2000b) clearly points to greater complexity. Informatic studies have suggested a basis for the opioid receptor results (Filizola et al., 2002) but this remains to be validated and expanded to a general case model. In the medium term it is likely that relative GPCR interaction affinities will have be measured directly on a case by case basis.

MOLPHARM 2004 0497

### **Do ligands alter GPCR dimerization?**

Although this might seem a relatively easy question to answer, the range of observations on this topic is wide and confusing. Early studies on the  $\beta_2$ -adrenoceptor indicated a substantial level of constitutive dimerization and small but significant increases in this with addition of isoproterenol (Hebert et al., 1996, Angers et al., 2000). In contrast, initial studies on the DOP opioid receptor suggested that ligands capable of inducing receptor internalization also caused monomerization of the receptor (Cvejic and Devi, 1997). This conclusion was in marked contrast to the case of the *Saccharomyces cerevisiae*  $\alpha$ -factor receptor that showed a lack of effect of ligand (Overton and Blumer, 2000) and, where as noted above, the dimer is internalized. A substantial number of RET-based experiments have shown small effects of agonist ligands above substantial signals corresponding to constitutive dimers/oligomers (see Milligan, 2004 for review). Despite this, an emerging consensus favours family A GPCRs with small endogenous ligands that bind within the seven transmembrane domains predominantly as constitutive dimers. Observed effects of ligands are then most likely due to conformational alterations within the complex resulting in small changes in the orientation or distance between the RET reporters (Ayoub et al., 2002).

This generalization cannot currently be extended to family A GPCR groupings in which larger ligands bind, at least partially, to elements of the N-terminal domain and the extracellular loops. One of the most interesting groupings is the chemokine receptors. Early studies indicated the chemokine SDF-1 $\alpha$  to produce dimerization of the CXCR4 receptor that was almost undetectable in the absence of the ligand (Vila-



MOLPHARM 2004 0497

Coro et al., 1999). In complete contrast, combinations of BRET and sedimentation studies indicated the CXCR4 receptor to be a constitutive dimer that was completely unaffected by the presence of SDF-1 $\alpha$  (Babcock et al., 2003). Ligand-independent dimerization of both the CXCR2 receptor (Trettel et al., 2003) and the CCR5 receptor (Issafras et al., 2002) has also been reported recently. Ligand-induced dimerization was, however, also recorded for the CCR2 receptor on addition of monocyte chemoattractant protein-1 (Rodriguez-Frade et al., 1999). As well as the effect of agonist, function was induced by addition of a bivalent anti-chemokine receptor monoclonal antibody but not by Fab fragments derived from the antibody. These authors drew direct parallels between the ability of chemokine receptor agonists to induce GPCR dimerization and the effects of cytokines on their, non-GPCR, receptor targets. Fascinatingly, this group have also reported that the capacity of HIV-1 to utilise the CCR5 receptor as a 'co-receptor' for cell entry was blocked by dimerization of the receptor by an antibody or by chemokine agonists at the receptor (Vila-Coro et al., 2000). Although not directly related, in the same studies that reported the CXCR4 receptor as a constitutive dimer, the CCR5 receptor was indicated to be a monomer (Babcock et al., 2003). Furthermore, in the studies that showed ligand-independent CXCR2 dimerization, the closely related CXCR1 receptor, which also uses interleukin 8 as a ligand, was reported not to dimerize (Trettel et al., 2003). Hetero-dimerization has also been reported between some (Mellado et al., 2001, Rodriguez-Frade et al., 2004) but not other (Babcock et al., 2003, Trettel et al., 2003) chemokine receptor pairs and at this time a clear pattern is difficult to discern. One issue may be the comparison of the effects of chemokine ligands and monoclonal antibodies. It is possible that the reported effects of

MOLPHARM 2004 0497

antibodies relate to clustering and capping of receptors rather than alterations in dimerization *per se* (Issafras et al., 2002).

It is not only the chemokine receptors at which substantial and divergent effects of ligands have been reported. Each of neuropeptide Y Y4 receptor (Berglund et al., 2003), TSH receptor (Latif et al., 2001) and type A cholecystokinin receptor (Cheng and Miller, 2001) dimers have been reported to be constitutively formed but dissociated by agonists. In light of the discussion of the capacity of arrestins to bridge GPCRs dimers it might be expected that ligand-induced dissociation of a GPCR dimer would limit or prevent interactions with arrestins and therefore both desensitization and arrestin-dependent internalization. Direct analysis of this issue would be of interest. By contrast, the GnRH (Cornea et al., 2001), lutropin (Tao et al., 2004) and a number of other receptors, have been reported to increase in aggregation state in response to agonists whereas the C5a complement receptor is a constitutive dimer that is unaffected by agonist (Floyd et al., 2003).

### **Does GPCR dimerization alter ligand pharmacology or function?**

With the caveats noted above that suggest some reported GPCR hetero-dimer pairs may have limited physiological significance, perhaps the most important questions in terms of relevance of GPCR hetero-dimers relate to whether they generate novel receptors in terms of either ligand pharmacology or function. This is a highly complex area, but at least for certain hetero-dimer pairs, this does appear likely. This issue has been addressed most directly for opioid receptor pairings. This reflects a combination of the large number of subtype-selective ligands available and the long appreciated view that opioid tissue pharmacology cannot be explained only by the

MOLPHARM 2004 0497

molecularly defined and characterized receptor subtypes. Many of the key experiments have been reviewed in detail (Jordan et al., 2000, Devi, 2001) and more recent studies have expanded this (Pan et al., 2002). Equally, the idea of opioid receptor interactions has influenced thinking on the molecular basis of opioid tolerance (He et al., 2002). Despite clear variations in opioid pharmacology following co-expression of opioid receptor pairs, it has been difficult to fully replicate opioid tissue pharmacology using this approach. It remains unclear if this is simply because receptor expression ratios need to be more fully titrated and controlled because, as discussed earlier co-expression of, for example, the MOP and DOP opioid receptors will not result in production of a pure population of MOP-DOP hetero-dimers. Ligands able to selectively target hetero-dimer pairs need to be identified and then their function analysed in physiological settings. The greatest progress in identifying such ligands undoubtedly stems from work on GPCRs that have historically attracted great efforts in medicinal chemistry. For example, a number of anti-parkinsonian agents may have significantly higher affinity at dopamine D3/D2 hetero-dimers than at the corresponding homo-dimers (Maggio et al., 2003). A substantial number of other studies have reported unique pharmacology associated with the co-expression of pairs of GPCRs (e.g. Yoshioka et al., 2001). Equally, the ability of  $\beta$ -blockers to interfere with angiotensin AT1-mediated signalling and of the AT1 receptor blocker valsartan to reduce catecholamine-induced elevation in heart rate (Barki-Harrington et al., 2003) may indicate functional angiotensin AT1- $\beta$ -adrenoceptor interactions *in vivo*. Such studies although fascinating are indirect and may reflect cross-talk between the signalling pathways at a level downstream of receptor activation. Such

MOLPHARM 2004 0497

interactions between  $G_s$  and  $G_q$ -mediated signals are well established to occur (Cordeaux and Hill, 2002). Strategies that allow the detection of ligand binding or function only by a hetero-dimer pair in the presence of the corresponding homo-dimers are required to allow rapid and effective screening and detection of ligands with these characteristics. Only with such ligands to hand will it be possible to tease out the physiological relevance of GPCR hetero-dimerization.

### References

- Angers S, Salahpour A and Bouvier M (2002) Dimerization: an emerging concept for G protein-coupled receptor ontogeny and function. *Annu Rev Pharmacol Toxicol* **42**:409-435.
- Angers S, Salahpour A, Joly E, Hilairat S, Chelsky D, Dennis M and Bouvier M (2000) Detection of beta 2-adrenergic receptor dimerization in living cells using bioluminescence resonance energy transfer (BRET). *Proc Natl Acad Sci U S A* **97**:3684-3689.
- Armstrong D and Strange PG (2001) Dopamine D2 receptor dimer formation: evidence from ligand binding. *J Biol Chem* **276**:22621-22629.
- Ayoub MA, Couturier C, Lucas-Meunier E, Angers S, Fossier P, Bouvier M and Jockers R (2002) Monitoring of ligand-independent dimerization and ligand-induced conformational changes of melatonin receptors in living cells by bioluminescence resonance energy transfer. *J Biol Chem* **277**: 21522-21528.
- Babcock GJ, Farzan M and Sodroski J (2003) Ligand-independent dimerization of CXCR4, a principal HIV-1 coreceptor. *J Biol Chem* **278**:3378-3385.

MOLPHARM 2004 0497

Barki-Harrington L, Luttrell LM and Rockman HA (2003) Dual inhibition of beta-adrenergic and angiotensin II receptors by a single antagonist: a functional role for receptor-receptor interaction in vivo. *Circulation* **108**:1611-1618.

Baneres JL and Parello J (2003) Structure-based analysis of GPCR function: evidence for a novel pentameric assembly between the dimeric leukotriene B4 receptor BLT1 and the G-protein. *J Mol Biol* **329**:815-829.

Berglund MM, Schober DA, Esterman MA and Gehlert DR (2003) Neuropeptide Y Y4 receptor homodimers dissociate upon agonist stimulation. *J Pharmacol Exp Ther* **307**:1120-1126.

Carrillo JJ, Pediani J and Milligan G (2003) Dimers of class A G protein-coupled receptors function via agonist-mediated trans-activation of associated G proteins. *J Biol Chem* **278**:42578-42587.

Chabre M, Cone R and Saibil H (2003) Biophysics: is rhodopsin dimeric in native retinal rods? *Nature* **426**:30-31.

Cheng ZJ and Miller LJ (2001) Agonist-dependent dissociation of oligomeric complexes of G protein-coupled cholecystinin receptors demonstrated in living cells using bioluminescence resonance energy transfer. *J Biol Chem* **276**:48040-48047.

Coge F, Guenin SP, Renouard-Try A, Rique H, Ouvry C, Fabry N, Beauverger P, Nicolas JP, Galizzi JP, Boutin JA and Canet E (1999) Truncated isoforms inhibit [<sup>3</sup>H]prazosin binding and cellular trafficking of native human alpha1A-adrenoceptors. *Biochem J* **343**:231-239.

MOLPHARM 2004 0497

Cordeaux Y and Hill SJ (2002) Mechanisms of cross-talk between G-protein-coupled receptors. *Neuro-Signals* **11**:45-57.

Cornea A, Janovick JA, Maya-Nunez G and Conn PM (2001) Gonadotropin-releasing hormone receptor microaggregation. Rate monitored by fluorescence resonance energy transfer. *J Biol Chem* **276**:2153-2158.

Cvejjic S and Devi LA (1997) Dimerization of the delta opioid receptor: implication for a role in receptor internalisation. *J Biol Chem* **272**:26959-26964.

Devi LA (2001) Heterodimerization of G-protein-coupled receptors: pharmacology, signaling and trafficking. *Trends Pharmacol Sci* **22**:532-537.

Dinger MC, Bader JE, Kobor AD, Kretzschmar AK and Beck-Sickingler AG (2003) Homodimerization of neuropeptide y receptors investigated by fluorescence resonance energy transfer in living cells. *J Biol Chem* **278**:10562-10571.

Ferre S, Karcz-Kubicha M, Hope BT, Popoli P, Burgueno J, Gutierrez MA, Casado V, Fuxe K, Goldberg SR, Lluís C, Franco R and Ciruela F (2002) Synergistic interaction between adenosine A2A and glutamate mGlu5 receptors: implications for striatal neuronal function. *Proc Natl Acad Sci U S A* **99**:11940-11945.

Filizola M, Olmea O and Weinstein H (2002) Prediction of heterodimerization interfaces of G-protein coupled receptors with a new subtractive correlated mutation method. *Prot Engineering* **15**:881-885.

Floyd DH, Geva A, Bruinsma SP, Overton MC, Blumer KJ and Baranski TJ (2003) C5a receptor oligomerization. II. Fluorescence resonance energy transfer studies of a human G protein-coupled receptor expressed in yeast. *J Biol Chem* **278**:35354-35361.

MOLPHARM 2004 0497

Fredriksson R, Lagerstrom MC, Lundin LG and Schioth HB (2003) The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol* **63**:1256-1272.

George SR, Ng GY, Lee SP, Fan T, Varghese G, Wang C, Deber CM, Seeman P and O'Dowd BF (2003) Blockade of G protein-coupled receptors and the dopamine transporter by a transmembrane domain peptide: novel strategy for functional inhibition of membrane proteins in vivo. *J Pharmacol Exp Ther* **307**:481-489.

George SR, O'Dowd BF and Lee SP (2002) G-protein-coupled receptor oligomerization and its potential for drug discovery. *Nature Reviews Drug Discovery* **1**:808-820.

Guo W, Shi L and Javitch JA (2003) The fourth transmembrane segment forms the interface of the dopamine D2 receptor homodimer. *J Biol Chem* **278**:4385-4388.

Hakak Y, Shrestha D, Goegel MC, Behan DP and Chalmers DT (2003) Global analysis of G-protein-coupled receptor signaling in human tissues. *FEBS Lett* **550**:11-17.

Han M, Gurevich VV, Vishnivetskiy SA, Sigler PB and Schubert C (2001) Crystal structure of beta-arrestin at 1.9 Å: possible mechanism of receptor binding and membrane translocation. *Structure* **9**:869-880.

Hanyaloglu AC, Seeber RM, Kohout TA, Lefkowitz RJ and Eidne KA (2002) Homo- and hetero-oligomerization of thyrotropin-releasing hormone (TRH) receptor subtypes. Differential regulation of beta-arrestins 1 and 2. *J Biol Chem* **277**:50422-50430.

MOLPHARM 2004 0497

He L, Fong J, von Zastrow M and Whistler JL (2002) Regulation of opioid receptor trafficking and morphine tolerance by receptor oligomerization. *Cell* **108**:271-282.

Hebert TE, Moffett S, Morello JP, Loisel TP, Bichet DG, Barret C and Bouvier M (1996) A peptide derived from a beta2-adrenergic receptor transmembrane domain inhibits both receptor dimerization and activation. *J Biol Chem* **271**:16384-16392.

Issafras H, Angers S, Bulenger S, Blanpain C, Parmentier M, Labbe-Jullie C, Bouvier M and Marullo S (2002) Constitutive agonist-independent CCR5 oligomerization and antibody-mediated clustering occurring at physiological levels of receptors. *J Biol Chem* **277**:34666-24673.

Jordan BA and Devi LA (1999) G-protein-coupled receptor heterodimerization modulates receptor function. *Nature* **399**:697-700.

Jordan BA, Cvejic S and Devi LA (2000) Opioids and their complicated receptor complexes. *Neuropsychopharmacology* **23**:S5-S18.

Jordan BA, Trapaidze N, Gomes I, Nivarthi R and Devi LA (2001) Oligomerization of opioid receptors with beta 2-adrenergic receptors: a role in trafficking and mitogen-activated protein kinase activation. *Proc Natl Acad Sci U S A* **98**:343-348.

Karpa KD, Lin R, Kabbani N and Levenson R (2000) The dopamine D3 receptor interacts with itself and the truncated D3 splice variant d3nf: D3-D3nf interaction causes mislocalization of D3 receptors. *Mol Pharmacol* **58**:677-683.

Kroeger KM, Hanyaloglu AC, Seeber RM, Miles LE, and Eidne KA (2001) Constitutive and agonist-dependent homo-oligomerization of the thyrotropin-releasing hormone receptor. Detection in living cells using bioluminescence resonance energy transfer. *J Biol Chem* **276**:12736-12743.



MOLPHARM 2004 0497

Kunishima N, Shimada Y, Tsuji Y, Sato T, Yamamoto M, Kumasaka T, Nakanishi S, Jingami H and Morikawa K (2000) Structural basis of glutamate recognition by a dimeric metabotropic glutamate receptor. *Nature* **407**:971-977.

Latif R, Graves P and Davies TF (2001) Ligand-dependent inhibition of oligomerization at the human thyrotropin receptor. *J Biol Chem* **277**:45059-45067.

Lee C, Ji I, Ryu K, Song Y, Conn PM and Ji TH (2002) Two defective heterozygous luteinizing hormone receptors can rescue hormone action. *J Biol Chem* **277**:15795-15800.

Lee SP, O'Dowd BF and George SR (2003) Homo- and hetero-oligomerization of G protein-coupled receptors. *Life Sci* **74**:173-180.

Lee SP, O'Dowd BF, Ng GY, Varghese G, Akil H, Mansour A, Nguyen T and George SR (2000) Inhibition of cell surface expression by mutant receptors demonstrates that D2 dopamine receptors exist as oligomers in the cell. *Mol Pharmacol* **58**:120-128.

Lee SP, O'Dowd BF, Rajaram RD, Nguyen T and George SR (2003) D2 dopamine receptor homodimerization is mediated by multiple sites of interaction, including an intermolecular interaction involving transmembrane domain 4. *Biochemistry* **42**:11023-11031.

Liang Y, Fotiadis D, Filipek S, Saperstein DA, Palczewski K and Engel A (2003) Organization of the G protein-coupled receptors rhodopsin and opsin in native membranes. *J Biol Chem* **278**:21655-21662.

Liu S, Carrillo JJ, Pediani J and Milligan G (2002) Effective information transfer from the  $\alpha_{1b}$ -adrenoceptor to G<sub>11</sub>  $\alpha$  requires both  $\beta/\gamma$  interactions and an aromatic group 4 amino acid from the C-terminus of the G protein. *J Biol Chem* **277**:25707-25714.

MOLPHARM 2004 0497

McVey M, Ramsay D, Kellett E, Rees S, Wilson S, Pope AJ, and Milligan G (2001)

Monitoring receptor oligomerization using time-resolved fluorescence resonance energy transfer and bioluminescence resonance energy transfer: The human  $\delta$  opioid receptor displays constitutive oligomerization at the cell surface which is not regulated by receptor occupancy. *J Biol Chem* **276**: 14092-14099.

Maggio R, Scarselli M, Novi F, Millan MJ and Corsini GU(2003) Potent activation of dopamine D3/D2 heterodimers by the antiparkinsonian agents, S32504, pramipexole and ropinirole. *J Neurochem* **87**:631-641.

Malnic B, Godfrey PA and Buck LB (2004) The human olfactory receptor gene family. *Proc Natl Acad Sci U S A* **101**:2584-2589.

Marshall FH (2001) Heterodimerization of G-protein-coupled receptors in the CNS. *Curr Opin Pharmacol* **1**:40-44.

Marshall FH, Jones KA, Kaupmann K and Bettler B (1999) GABAB receptors - the first 7TM heterodimers. *Trends Pharmacol Sci* **20**:396-399.

Matsunami H and Amrein H (2004) Taste perception: how to make a gourmet mouse. *Curr Biol* **14**:118-120.

Mellado M, Rodriguez-Frade JM, Vila-Coro AJ, Fernandez S, Martin de Ana A,

Jones DR, Toran JL and Martinez-A C (2001) Chemokine receptor homo- or heterodimerization activates distinct signaling pathways. *EMBO J* **20**:2497-2507.

Mercier JF, Salahpour A, Angers S, Breit A and Bouvier M (2002) Quantitative assessment of beta 1- and beta 2-adrenergic receptor homo- and heterodimerization by bioluminescence resonance energy transfer. *J Biol Chem* **277**:44925-44931.

MOLPHARM 2004 0497

Milligan G (2004) Applications of bioluminescence- and fluorescence resonance energy transfer to drug discovery at G protein-coupled receptors. *Eur J Pharm Sci.* **21**:397-405.

Milligan G, Lopez-Gimenez JF, Wilson S and Carrillo J (2004) Selectivity in the oligomerisation of G protein-coupled receptors. *Sem Cell Develop. Biol. (in press)*.

Milligan G, Ramsay D, Pascal G, and Carrillo JJ (2003) GPCR dimerisation. *Life Sci* **74**:181-188.

Mohler H and Fritschy JM (1999) GABAB receptors make it to the top--as dimers. *Trends Pharmacol Sci* **20**:87-89.

Ng GY, O'Dowd BF, Lee SP, Chung HT, Brann MR, Seeman P and George SR (1996) Dopamine D2 receptor dimers and receptor-blocking peptides. *Biochem Biophys Res Commun* **227**:200-204.

Nimchinsky EA, Hof PR, Janssen WG, Morrison JH and Schmauss C (1997) Expression of dopamine D3 receptor dimers and tetramers in brain and in transfected cells. *J Biol Chem* **272**: 29229-29237.

Overton MC and Blumer KJ (2000) G-protein-coupled receptors function as oligomers *in vivo*. *Curr Biol* **10**:341-344.

Overton MC, Chinault SL and Blumer KJ (2003) Oligomerization, biogenesis, and signaling is promoted by a glycoporphin A-like dimerization motif in transmembrane domain 1 of a yeast G protein-coupled receptor. *J Biol Chem* **278**:49369-49377.

Pan YX, Bolan E and Pastmak GW (2002) Dimerization of morphine and orphanin/nociceptin receptors: generation of a novel opioid receptor subtype. *Biochem Biophys Res Comm* **297**:659-663.

MOLPHARM 2004 0497

Pfeiffer M, Kirscht S, Stumm R, Koch T, Wu D, Laugsch M, Schroder H, Holtt V and Schulz S (2003) Heterodimerization of substance P and mu-opioid receptors regulates receptor trafficking and resensitization. *J Biol Chem* **278**:51630-51637.

Ramsay D, Carr IC, Padiani J, Lopez-Gimenez JF, Thurlow R, Fidock M and Milligan G (2004) High affinity interactions between human  $\alpha_{1A}$ -adrenoceptor C-terminal splice variants produce homo and hetero-dimers but do not generate the  $\alpha_{1L}$ -adrenoceptor. *Mol. Pharmacol.* (submitted for publication). Manuscript 2003/3152

Ramsay D, Kellett E, McVey M, Rees S and Milligan G (2002) Homo- and hetero-oligomeric interactions between G-protein-coupled receptors in living cells monitored by two variants of bioluminescence resonance energy transfer (BRET): hetero-oligomers between receptor subtypes form more efficiently than between less closely related sequences. *Biochem J* **365**:429-440.

Rocheville M, Lange DC, Kumar U, Patel SC, Patel RC and Patel YC (2000a) Receptors for dopamine and somatostatin: formation of hetero-oligomers with enhanced functional activity. *Science* **288**:154-157.

Rocheville M, Lange DC, Kumar U, Sasi R, Patel RC and Patel YC (2000b) Subtypes of the somatostatin receptor assemble as functional homo- and heterodimers. *J Biol Chem* **275**:7862-7869.

Rodriguez-Frade JM, Del Real G, Serrano A, Hernanz-Falcon P, Soriano SF, Vila-Coro AJ, De Ana AM, Lucas P, Prieto I, Martinez-A C and Mellado M (2004) Blocking HIV-1 infection via CCR5 and CXCR4 receptors by acting in trans on the CCR2 chemokine receptor. *EMBO J* **23**:66-76.

MOLPHARM 2004 0497

Rodriguez-Frade JM, Vila-Coro AJ, de Ana AM, Albar JP, Martinez-A C and Mellado M (1999) The chemokine monocyte chemoattractant protein-1 induces functional responses through dimerization of its receptor CCR2. *Proc Natl Acad Sci U S A* **96**:3628-3633.

Salim K, Fenton T, Bacha J, Urien-Rodriguez H, Bonnert T, Skynner HA, Watts E, Kerby J, Heald A, Beer M, McAllister G and Guest PC (2002) Oligomerization of G-protein-coupled receptors shown by selective co-immunoprecipitation. *J Biol Chem* **277**:15482-15485.

Shenoy SK and Lefkowitz RJ (2003) Multifaceted roles of beta-arrestins in the regulation of seven-membrane-spanning receptor trafficking and signalling. *Biochem J* **375**:503-515.

Soyer OS, Dimmic MW, Neubig RR and Goldstein RA (2003) Dimerization in aminergic G-protein-coupled receptors: application of a hidden-site class model of evolution. *Biochemistry* **42**:14522-14531.

Stanasila L, Perez J-B, Vogel H and Cotecchia S (2003) Oligomerization of the  $\alpha_{1a}$ - and  $\alpha_{1b}$ -adrenergic receptor subtypes. *J. Biol. Chem.* **278**: 40239-40251.

Tao YX, Johnson NB and Segaloff DL (2004) Constitutive and agonist-dependent self-association of the cell surface human lutropin receptor. *J Biol Chem* **279**:5904-5914.

Terrillon S, Barberis C and Bouvier M (2004) Heterodimerization of V1a and V2 vasopressin receptors determines the interaction with beta-arrestin and their trafficking patterns. *Proc Natl Acad Sci U S A* **101**:1548-1553.

Terrillon S, Durroux T, Mouillac B, Breit A, Ayoub MA, Taulan M, Jockers R, Barberis C and Bouvier M (2003) Oxytocin and vasopressin V1a and V2 receptors

MOLPHARM 2004 0497

form constitutive homo- and heterodimers during biosynthesis. *Mol Endocrinol* **17**:677-691.

Trettel F, Di Bartolomeo S, Lauro C, Catalano M, Ciotti MT, Limatola C (2003) Ligand-independent CXCR2 dimerization. *J Biol Chem* **278**: 40980-40988.

Tsao PI and von Zastrow M (2000) Type-specific sorting of G protein-coupled receptors after endocytosis. *J Biol Chem* **275**:11130-11140.

Uberti MA, Hall RA and Minneman KP (2003) Subtype-specific dimerization of alpha 1-adrenoceptors: effects on receptor expression and pharmacological properties. *Mol Pharmacol* **64**:1379-1390.

Vassilatis DK, Hohmann JG, Zeng H, Li F, Ranchalis JE, Mortrud MT, Brown A, Rodriguez SS, Weller JR, Wright AC, Bergmann JE and Gaitanaris GA (2003) The G protein-coupled receptor repertoires of human and mouse. *Proc Natl Acad Sci U S A* **100**:4903-4908.

Vila-Coro AJ, Mellado M, Martin de Ana A, Lucas P, del Real G, Martinez-A C and Rodriguez-Frade JM (2000) HIV-1 infection through the CCR5 receptor is blocked by receptor dimerization. *Proc Natl Acad Sci U S A* **97**:3388-3393.

Vila-Coro AJ, Rodriguez-Frade JM, Martin De Ana A, Moreno-Ortiz MC, Martinez-A C and Mellado M (1999) The chemokine SDF-1alpha triggers CXCR4 receptor dimerization and activates the JAK/STAT pathway. *FASEB J* **13**:1699-1710.

White JH, Wise A, Main MJ, Green A, Fraser NJ, Disney GH, Barnes AA, Emson P, Foord SM and Marshall FH (1998) Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature* **396**:679-682.

Woolf PJ and Linderman JJ (2003) Self organization of membrane proteins via dimerization. *Biophys Chem* **104**:217-227.

MOLPHARM 2004 0497

Yesilaltay A and Jenness DD (2000) Homo-oligomeric complexes of the yeast alpha-factor pheromone receptor are functional units of endocytosis. *Mol Biol Cell* **11**:2873-2884.

Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ and Zuker CS (2003) The receptors for mammalian sweet and umami taste. *Cell* **115**:255-266.

Yoshioka K, Saitoh O and Nakata H (2001) Heteromeric association creates a P2Y-like adenosine receptor. *Proc Natl Acad Sci U S A* **98**:7617-7622.

MOLPHARM 2004 0497

### **Footnotes**

Work on GPCR dimerization within my group is supported by the Biotechnology and Biosciences Research Council, The Medical Research Council and the Wellcome Trust.

Address all correspondence to:

Graeme Milligan

Molecular Pharmacology Group

Davidson Building

University of Glasgow

Glasgow G12 8QQ

Scotland, U.K.

Tel: 44 (0) 141 330 5557

FAX: 44 (0) 141 330 4620

e-mail: [g.milligan@bio.gla.ac.uk](mailto:g.milligan@bio.gla.ac.uk)