

MOL #50534

## Title Page

# A DOPAMINE D<sub>2</sub> RECEPTOR MUTANT CAPABLE OF G PROTEIN-MEDIATED SIGNALING BUT DEFICIENT IN ARRESTIN BINDING

**Hongxiang Lan, Yong Liu, Michal I. Bell, Vsevolod V. Gurevich, and Kim A. Neve**

*Departments of Behavioral Neuroscience (Y.L., M.I.B., K.A.N.) and Physiology & Pharmacology (H.L.), Oregon Health & Science University, and Veterans Affairs Medical Center (K.A.N.), Portland, Oregon 97239; Department of Pharmacology (V.V.G.), Vanderbilt University School of Medicine, Nashville, Tennessee 37232*

MOL #50534

## Running Title Page

**Running Title:** Arrestin Binding to the D<sub>2</sub> Receptor Third Cytoplasmic Loop

**Address correspondence to:** Dr. Kim A. Neve, Portland VA Medical Center (R&D-30),  
3710 SW US Veterans Hospital Road, Portland, OR 97239-2999, Telephone: 503-721-7911, Fax:  
503-721-7839, E-mail: [nevek@ohsu.edu](mailto:nevek@ohsu.edu)

The number of text pages: 38

The number of figures: 9

The number of references: 45

The number of words in the Abstract: 252

The number of words in the Introduction: 490

The number of words in the Discussion: 1587

**ABBREVIATIONS:** CMF-PBS, calcium-, magnesium-free phosphate-buffered saline; GPCR, G protein-coupled receptor; GRK, GPCR kinase; GST, glutathione-S-transferase; ERK, extracellular signal-regulated kinase; D<sub>2</sub>-A4, IYIV212-215A4 mutant of the rat dopamine D<sub>2L</sub> receptor; IC3, third intracellular loop of a GPCR; D<sub>2</sub>- or D<sub>3</sub>-IC3, third intracellular loop of the rat dopamine D<sub>2L</sub> or D<sub>3</sub> receptor; PTX, pertussis toxin; TBS, Tris-buffered saline

MOL #50534

## ABSTRACT

Arrestins mediate G protein-coupled receptor desensitization, internalization, and signaling. Dopamine D<sub>2</sub> and D<sub>3</sub> receptors have similar structures, but distinct characteristics of interaction with arrestins. The goals of this study were to compare arrestin-binding determinants in D<sub>2</sub> and D<sub>3</sub> receptors other than phosphorylation sites and to create a D<sub>2</sub> receptor that is deficient in arrestin binding. We first assessed the ability of purified arrestins to bind to glutathione S-transferase (GST) fusion proteins containing the receptor third intracellular loops (IC3). Arrestin3 bound to IC3 of both D<sub>2</sub> and D<sub>3</sub> receptors, with the affinity and localization of the binding site indistinguishable between the receptor subtypes. Mutagenesis of the GST-IC3 fusion proteins identified an important determinant of the binding of arrestin3 in the N-terminal region of IC3. Alanine mutations of this determinant (IYIV212-215) in the full-length D<sub>2</sub> receptor generated a signaling-biased receptor with intact ligand binding and G-protein coupling and activation, but deficient in receptor-mediated arrestin3 translocation to the membrane, agonist-induced receptor internalization, and agonist-induced desensitization in human embryonic kidney 293 cells. This mutation also decreased arrestin-dependent activation of extracellular signal-regulated kinases. The finding that non-phosphorylated D<sub>2</sub>-IC3 and D<sub>3</sub>-IC3 have similar affinity for arrestin is consistent with previous suggestions that the differential effects of D<sub>2</sub> and D<sub>3</sub> receptor activation on membrane translocation of arrestin and receptor internalization are at least in part due to differential phosphorylation of the receptors. In addition, these results imply that the sequence IYIV212-215 at the N-terminus of IC3 of the D<sub>2</sub> receptor is a key element of the arrestin binding site.

MOL #50534

## Introduction

The non-visual arrestins arrestin2 and 3 (also termed  $\beta$ -arrestin1 and 2) are cytosolic proteins involved in homologous desensitization and resensitization of G protein-coupled receptors (GPCRs), and serve as adaptors to link GPCRs to the endocytotic machinery. Arrestins also redirect GPCRs to alternative, G protein-independent signaling pathways (Pierce and Lefkowitz, 2001). Although most or all GPCRs bind arrestins, and phosphorylated serine and threonine residues often comprise part of the arrestin binding site, little information is available concerning other structural features of receptors that cause them to be recognized by arrestins.

Three D2-like receptors, D<sub>2</sub>, D<sub>3</sub>, and D<sub>4</sub>, have been identified. D2-like receptors are the major targets of antipsychotic drugs. When activated, they couple to the G<sub>i/o</sub> family of G proteins and regulate adenylate cyclases, potassium channels, calcium channels, and other effectors (Neve *et al.*, 2004). Despite close sequence homology between D<sub>2</sub> and D<sub>3</sub> receptors, agonist-induced receptor phosphorylation, arrestin translocation to the plasma membrane, and receptor internalization for the two subtypes differ dramatically. These differences can be attributed primarily to IC2 and IC3 of the two receptors (Kim *et al.*, 2001).

Numerous studies have examined the role of IC2, IC3, and the carboxyl-terminus of GPCRs in receptor internalization. In the classic model of G protein-coupled receptor kinase- (GRK-) and arrestin-mediated intracellular trafficking of GPCRs, GRK phosphorylates residues in the intracellular segments of receptors, recruiting arrestin to bind to the phosphorylated residues and to other residues whose accessibility is regulated by receptor phosphorylation and by the activation state of the receptor (Pierce and Lefkowitz, 2001; Lee *et al.*, 2000; Kim *et al.*, 2004; Gurevich and Gurevich, 2006). One way to quantify the contribution of non-phosphorylated residues to the binding of arrestin is to use peptides representing intracellular receptor domains,

MOL #50534

either measuring their ability to inhibit arrestin binding to receptors or measuring direct binding of arrestin to the receptor fragments. Such studies have demonstrated that arrestin binds to multiple unphosphorylated intracellular domains of GPCRs (Wu *et al.*, 1997; Gelber *et al.*, 1999; Cen *et al.*, 2001; Han *et al.*, 2001; DeGraff *et al.*, 2002; Macey *et al.*, 2004; Macey *et al.*, 2005).

To identify non-phosphorylated arrestin-binding sites of D<sub>2</sub> and D<sub>3</sub> receptors, we generated glutathione S-transferase (GST) fusion proteins of intracellular loops of the receptors and used them in direct binding assays with purified arrestins. We now report that arrestin3 bound avidly to IC3 of both receptors. Binding of arrestin3 to D<sub>2</sub>-IC3 required 4-5 residues at the N-terminus of the loop. Simultaneous substitution of alanine for four of these residues in the full-length D<sub>2</sub> receptor abolished receptor-mediated recruitment of arrestin3 to the membrane, agonist-induced receptor internalization, and agonist-induced desensitization of cyclic AMP accumulation in HEK 293 cells and decreased arrestin-dependent activation of extracellular signal-regulated kinases (ERKs), without altering high- or low-affinity agonist binding to the receptor and the potency of agonist for inhibition of cyclic AMP accumulation. Our data suggest that the D<sub>2</sub>-A4 mutant differentiates between G protein- and arrestin-dependent responses to receptor activation.

MOL #50534

## Materials and Methods

**Materials.** [<sup>3</sup>H]spiperone (83 Ci/mmol) was purchased from GE Healthcare (Little Chalfont, Buckinghamshire, UK), and [<sup>3</sup>H]sulpiride (77.7 Ci/mmol) from PerkinElmer Life and Analytical Sciences (Boston, MA). Serum was purchased from Hyclone Laboratories (Logan, UT).

Dopamine, (+)-butaclamol, haloperidol, and most reagents, including culture medium, were purchased from Sigma-Aldrich (St. Louis, MO). Antibodies used include: mouse anti-rat arrestin2 (A47520 from BD Transduction Laboratories, Lexington, KY), mouse anti-human arrestin3 (sc-13140 from Santa Cruz Biotechnology, Santa Cruz, CA), rabbit anti-dually phosphorylated (i.e., activated) ERKs (44-680G from Biosource International, Camarillo, CA), p44/42 MAP kinase antibody (i.e., total ERKs; #9102 from Cell Signaling Technology, Danvers, MA), and secondary antibodies horseradish peroxidase conjugated goat anti-mouse IgG (31430, from Pierce Biotechnology, Rockford, IL) and horseradish peroxidase conjugated goat anti-rabbit IgG (#1858415 from Pierce Biotechnology). The cyclic AMP EIA kit was from Cayman Chemical (Ann Arbor, MI). Arrestin-3-pCMV5 was a generous gift from Dr. Marc Caron.

**Generation of GST Fusion Proteins and D<sub>2</sub> Receptor Constructs.** For construction of the GST fusion proteins the IC3 of the rat dopamine D<sub>2L</sub> receptor (D<sub>2</sub>-IC3), amino acids 211-371, and IC3 of the rat dopamine D<sub>3</sub> receptor (D<sub>3</sub>-IC3), amino acids 210-372, were amplified by polymerase chain reaction, subcloned into SpeI-XhoI sites in pET-41a(+) (Novagen, Madison, Wisconsin), transformed into NovaBlue competent cells, sequenced, and subsequently transformed into Rosetta 2(DE3) competent cells (Novagen, Madison, Wisconsin). For purification of GST fusion proteins, Rosetta 2(DE3) cells were grown in 2X YTK medium containing kanamycin (50 µg/ml) at 37 °C to A<sub>600</sub> = 0.8 and induced with 1 mM isopropyl-β-D-thiogalactopyranoside for 2 h at 32 °C. Cells were pelleted, resuspended in lysis buffer (50 mM

MOL #50534

Tris, 1 mM EDTA, 0.5 mg/ml lysozyme, pH 8.0) containing Complete protease inhibitor tablet (Roche Diagnostics, Mannheim, Germany), and incubated for 20 min with gentle rotation at room temperature. The homogenates were clarified by centrifugation, and supernatants were applied to microcentrifuge tubes containing Glutathione Sepharose 4B beads (GE Healthcare), and purified as described by the manufacturer. To quantify the amounts of fusion proteins, SDS sample loading buffer was applied to beads bound with purified proteins, samples were separated by SDS-PAGE and the gel was stained with Gel Code Blue (Pierce, Rockford, IL). BSA was used as a standard.

The D<sub>2</sub>-IC3 substitution mutants, the A4 mutant (myc-D<sub>2</sub>-IYIV212-215A4), and the IC3ΔMID mutant (myc-D<sub>2</sub>-IC3ΔMID) were constructed using the QuikChange mutagenesis kit (Stratagene, La Jolla, CA) through one or more mutagenesis steps, with GST-D<sub>2</sub>-IC3 or the myc-D<sub>2</sub> receptor as a template. The IC3 truncation mutants were generated using a QuikChange method modified to introduce large truncations (Makarova *et al.*, 2000) with GST-D<sub>2</sub>-IC3 or GST-D<sub>3</sub>-IC3 plasmid as template. The myc-D<sub>2</sub> receptor (referred to herein as wild type D<sub>2</sub> to differentiate it from the D<sub>2</sub>-A4 mutant) was constructed by cloning a rat D<sub>2L</sub> receptor cDNA into SfiI-XhoI sites in the pCMV-Myc vector (BD Biosciences Clontech, Mountain View, CA).

**Purified Arrestin Binding to GST Fusion Proteins.** To obtain purified arrestins, plasmids were expressed in BL21 cells and arrestins purified using heparin-Sepharose chromatography, followed by Q-Sepharose chromatography (Han *et al.*, 2001). GST fusion proteins bound to glutathione Sepharose 4B beads were incubated with purified bovine arrestin2 or arrestin3 in arrestin binding buffer (25 mM Tris-HCl, 150 mM NaCl, pH 7.2, Complete protease inhibitor tablet, 0.1% Triton X-100) for 30 min at room temperature. Incubation mixtures were washed four times in wash buffer (arrestin binding buffer without protease inhibitors), and the proteins

MOL #50534

were released with SDS sample loading buffer. Proteins were separated by SDS-PAGE, transferred to PVDF membranes that were then blocked with 5% nonfat-dry milk in Tris-buffered saline (TBS), and detected by immunoblotting using anti-arrestin2 (1:300 dilution in TBS) or anti-arrestin3 (1:400 dilution in TBS) antibody, with horseradish peroxidase-conjugated goat anti-mouse IgG (1:10,000 dilution in TBS) as secondary antibody. Visualization of the secondary antibody was performed using the SuperSignal West Pico Chemiluminescence kit (Pierce, Rockford, IL) and quantified by IP Lab (Scanalytics, Fairfax, VA). The amount of bound arrestin2 or arrestin3 was calculated from linear regression of a standard curve generated using background optical density (i.e., no arrestin) and 3-6 concentrations of arrestin2 or arrestin3 varying between 0.25 and 20 ng. Saturation analysis of the binding of arrestin was carried out by incubating various concentrations of arrestin2 or -3 with a fixed concentration of GST alone (150 or 175 ng) or GST-IC3 (300 ng) for 30 min at room temperature. The resulting concentration-response curves were analyzed by nonlinear regression using Prism 3.0 (Graphpad Software) and statistical comparisons of the curves were made using two-way ANOVA followed by Bonferroni post test analysis. In experiments where only one concentration of arrestin was used, statistical significance was evaluated using a paired t test.

**Internalization Assay.** Internalization was measured using the intact cell [<sup>3</sup>H]sulpiride binding assay described by Itokawa et al. (Itokawa *et al.*, 1996). HEK 293 cells grown to 80% confluency were co-transfected with 30 ng D<sub>2</sub> wild type, 10 μg D<sub>2</sub>-A4 mutant receptor DNA (the A4 mutant was expressed on the membrane at a lower density so a greater amount of DNA was used to achieve similar expression levels), or 100 ng D<sub>2</sub>-IC3ΔMID mutant receptor DNA and 3 μg arrestin3-pCMV5 using Lipofectamine2000 (Invitrogen, Carlsbad, CA). Cells were split into 2 plates after 12 h, and 2 days later rinsed once with pre-warmed, calcium- and magnesium-free



MOL #50534

phosphate-buffered saline (CMF-PBS; 138 mM NaCl, 4.1 mM KCl, 5.1 mM sodium phosphate, 5 mM potassium phosphate, and 0.2% glucose, pH 7.4), and pre-incubated for 15 min with pre-warmed, CO<sub>2</sub>-saturated serum-free Dulbecco's modified Eagle's medium containing 20 mM HEPES, pH 7.4, at 37 °C. Cells were stimulated with 10 μM dopamine in the same HEPES-buffered medium at 37 °C for 20 min. Stimulation was terminated by quickly cooling the plates on ice and washing the cells three times with ice-cold calcium-, magnesium-free phosphate-buffered saline (CMF-PBS), after which cells were removed from the plate in 2 ml of ice-cold CMF-PBS assay buffer (CMF-PBS containing 2 mM EDTA and 0.001% bovine serum albumin). Cells were gently mixed, added to assay tubes in a final volume of 250 μl with [<sup>3</sup>H]sulpiride (5 nM final concentration), and incubated at 4 °C for 150 min in the absence and presence of unlabeled haloperidol (10 μM final concentration). The assay was terminated by filtration through Whatman GF/C filters presoaked with 0.05% polyethylenimine using a 96-well Tomtec cell harvester (Orange, CT) and ice-cold wash buffer (10 mM Tris-HCl, pH 7.4, 0.9% NaCl). Filters were allowed to dry, and BetaPlate scintillation fluid (50 μl) was added to each sample. Radioactivity on the filters was determined using a Wallac 1205 BetaPlate scintillation counter (Gaithersburg, MD). To confirm the expression of arrestin3, the remaining cells were pelleted and resuspended with ice-cold lysis buffer (20 mM HEPES, 20 mM NaCl, 5 mM EDTA, 0.5% 3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate (CHAPS) and Complete protease inhibitor tablet). The suspensions were gently rocked on an orbital shaker at 4 °C for 60 min and then were centrifuged at 100,000 x g for 30 min at 4°C. The supernatant was saved and immunoblotting of overexpressed arrestin3 was performed as described below.

**Arrestin3 Translocation.** Transient expression of wild type D<sub>2</sub> or D<sub>2</sub>-A4 mutant receptor with arrestin3 and dopamine stimulation of cells were performed as described above, except that each

MOL #50534

plate was split into 3 plates after transfection (one plate for the intact cell [<sup>3</sup>H]sulpiride binding to detect the levels of receptor expression, performed as described above but without dopamine treatment). Stimulation was terminated by quickly cooling the plates on ice and washing the cells once with ice-cold CMF-PBS. Cells were lysed with 1 ml ice-cold lysis buffer (20 mM HEPES, 20 mM NaCl, 5 mM EDTA, and Complete protease inhibitor tablet), scraped, collected, homogenized with a glass-Teflon homogenizer, and sonicated for 8-10 s. Samples were centrifuged at 1,000 x g for 10 min at 4 °C. Supernatants were transferred to new centrifuge tubes and centrifuged at 100,000 x g for 30 min at 4 °C. Supernatants were collected; pellets were rinsed carefully with ice-cold CMF-PBS and then resuspended with 100 µl CMF-PBS. The abundance of arrestin3 in both pellet and supernatant fractions was quantified by immunoblotting as described above, except with a 1:100 dilution of the anti-arrestin3 antibody and the addition of 0.1% Tween 20 and 5% dry milk to the incubations with primary and secondary antibodies.

**Radioligand Binding Assays.** Cells expressing wild type D<sub>2</sub> receptor or the D<sub>2</sub>-A4 mutant were lysed in ice-cold hypotonic buffer (1 mM Na<sup>+</sup>HEPES, pH 7.4, 2 mM EDTA) for 15 min, scraped from the plate, and centrifuged at 17,000 x g for 20 min. The resulting crude membrane fraction was resuspended with a Brinkmann Polytron homogenizer (Brinkmann Instruments, Westbury, NY) at setting 6 for 8 to 10 s in TBS for saturation assays of the binding of [<sup>3</sup>H]spiperone, or resuspended in preincubation buffer (50 mM Tris-HCl, pH 7.4, 0.9% NaCl, 5 mM MgCl<sub>2</sub>, 1 mM dithiothreitol), preincubated for 30 min at 37 °C, centrifuged at 17,000 x g for 10 min, and resuspended again in Tris assay buffer (50 mM Tris-HCl, pH 7.4, 6 mM MgCl<sub>2</sub>, 1 mM EDTA, 1 mM dithiothreitol, 0.001% BSA, 0.002% ascorbic acid) for competition binding studies in which dopamine displacement of the binding of [<sup>3</sup>H]spiperone was assessed. Membranes (40-100 µg protein) were incubated in duplicate in a total reaction volume of 1 ml with [<sup>3</sup>H]spiperone at

MOL #50534

concentrations ranging from 0.01-0.6 nM for saturation binding or ~0.1 nM with the appropriate concentration of the competing drug dopamine for competition binding. (+)-Butaclamol (2  $\mu$ M final) was used to define nonspecific binding. Reactions were incubated at 37 °C for 45 min and terminated by filtration as described above. Data for saturation and competition binding were analyzed by nonlinear regression using the computer program Prism 3.0 (GraphPad, San Diego, CA) to determine  $K_d$  and  $IC_{50}$  values. Apparent affinity ( $K_i$ ) values were calculated from the  $IC_{50}$  values by the method of Cheng and Prusoff (Cheng and Prusoff, 1973). In all assays, the free concentration of radioligand was calculated as the concentration added minus the concentration specifically bound.

**Cyclic AMP Accumulation Assay.** Wild-type or mutant  $D_2$  receptors in pCMV-Myc were cotransfected with pcDNA3.1 (for G418 resistance) into HEK 293 cells using Lipofectamine2000, and clonal cell line stably expressing the receptors were isolated after selection with G418 (800  $\mu$ g/ml). Cell lines were maintained in Dulbecco's modified Eagle's medium supplemented with 5% iron-supplemented calf bovine serum, 5% fetal bovine serum and 600  $\mu$ g/ml G418 at 37 °C and 10%  $CO_2$ . For production of cell lines overexpressing arrestin3, stable  $D_2$  cell lines were transiently transfected with 3  $\mu$ g arrestin3-pCMV5. The cyclic AMP accumulation assay was performed as described by Liu et al. (Liu *et al.*, 2007), except that the agonist dopamine was used and the assay was performed two days after transient transfection. The expression of cell surface receptor was determined by intact cell [ $^3H$ ]sulpiride binding. The overexpression of arrestin3 was confirmed by immunoblotting as described above. In some experiments, cells were pre-treated with 10  $\mu$ M dopamine for 20 min at 37°C, followed by 3 rinses with ice-cold Earle's balanced salt solution, prior to measuring dopamine inhibition of forskolin-stimulated cyclic AMP accumulation for 10 min at 37°C.

MOL #50534

**Immunodetection of ERKs.** Stable expression of D<sub>2</sub> receptors and transient expression of arrestin3 were performed as described above. For detection of phosphospecific ERKs, PVDF membranes were probed with rabbit anti-dually phosphorylated ERKs (1/100 dilution in TBST (TBS + 0.1% Tween 20) with 5% dry milk), followed by horseradish peroxidase conjugated goat anti-rabbit IgG (1/200 dilution in TBST with 1% dry milk). Phospho-ERKs were visualized and quantified as described above for arrestins. Multiple dilutions of sample WT+arr-DA were used to verify that the concentration of phospho-ERKs varied linearly with optical density. For detection of total ERKs, PVDF membranes were blocked with 5% dry milk in TBST, and detected by immunoblotting using p44/42 MAP kinase antibody (1/1000 dilution in TBST), with horseradish peroxidase conjugated goat anti-rabbit IgG (1/1000 dilution in TBST) as secondary antibody.

MOL #50534

## Results

**Robust Binding of Arrestin3 to IC3.** GST-D<sub>2</sub>-IC3 and GST-D<sub>3</sub>-IC3 were constructed and the binding of arrestin determined using an *in vitro* GST pull-down assay. To identify conditions for equilibrium binding, the rate of association of arrestin3 with GST-D<sub>2</sub>-IC3 was determined. The half-time for binding was approximately 2 min and the binding approached equilibrium within 15 min (data not shown). Therefore, GST binding assays were carried out for 30 min. Arrestin3 bound avidly to both GST-D<sub>2</sub>-IC3 and GST-D<sub>3</sub>-IC3, showing no apparent difference between the two IC3 fusion proteins (Fig. 1, Table 1). Arrestin2 bound weakly to both fusion proteins (Fig. 1).

### **Identification of Arrestin3 Binding Sites within IC3 of the D<sub>2</sub> and D<sub>3</sub> Receptors.**

Intracellular residues in proximity to the transmembrane domains are critical for GPCR interactions with many cytosolic proteins, so we made truncation/deletion mutants to identify regions of arrestin3 binding to IC3, focusing on residues close to those transmembrane domains (Fig. 2A). GST pull-down assays revealed that deletion of the first 20 residues of D<sub>2</sub>-IC3 (D<sub>2</sub>-IC3ΔNT20) dramatically decreased its arrestin-binding capability; when the last 20 amino acids were deleted (D<sub>2</sub>-IC3ΔCT20), the binding capability also decreased, though to a lesser extent; when both the first and last 20 residues were deleted (D<sub>2</sub>-IC3Δ(NT20+CT20)), little binding was observed (Fig. 2B). In contrast, deletion of ~120 residues comprising all of IC3 except the 20 residues at each terminus (D<sub>2</sub>-IC3ΔMID) had little effect on the binding of arrestin. The arrestin binding patterns of the truncation/deletion mutants of the D<sub>3</sub> receptor were the same as for the D<sub>2</sub> receptor (Fig. 2C).

Having localized arrestin3 binding to the first and last 20 residues of IC3, with the N-terminal segment playing a more important role, we constructed additional N-terminal truncation

MOL #50534

mutants of D<sub>2</sub>-IC3 in which the first 5 or 10 or the second 10 amino acids were deleted (Fig. 3A). In the GST pull-down assay, deletion of either the first 5 (D<sub>2</sub>-IC3ΔNT5) or 10 (D<sub>2</sub>-IC3ΔNT10) residues decreased arrestin binding by about 60%, whereas deletion of the second 10 residues (D<sub>2</sub>-IC3ΔNT10-2) had little effect (Fig. 3B). We also made smaller C-terminal truncation/deletion mutants, deleting the last 5 or 10 or the penultimate 10 residues, but none of these deletions caused a significant reduction in the binding of arrestin (data not shown). This is consistent with the lesser role of the C-terminus suggested by the 20-residue truncation, and suggests that any contiguous 10 residues in the C-terminus are sufficient for the smaller contribution of the C-terminus to the binding of arrestin.

Because most of the effect of deleting the N-terminal 20 residues of D<sub>2</sub>-IC3 could be attributed to the first 5 residues (D<sub>2</sub>-IC3ΔNT5), we made additional mutations within this segment (Fig. 4A). Deletion of the first 3 residues (D<sub>2</sub>-IC3ΔNT3) had no effect on the binding of arrestin3 (Fig. 4B). On the other hand, deletion of D<sub>2</sub>-IC3 residues 2-5 (IYIV212-215 in the D<sub>2</sub> receptor, D<sub>2</sub>-IC3ΔIYIV) was as deleterious to the binding of arrestin as deletion of the first 5 residues (D<sub>2</sub>-IC3ΔNT5) (Fig. 4B).

Within the cytosolic domains of many integral membrane proteins, the motif YXXϕ (where Y is tyrosine, X is any amino acid, and ϕ is an amino acid with a bulky hydrophobic group) mediates endocytosis and other intracellular trafficking events (Collins *et al.*, 2002; Vogt *et al.*, 2005; Boll *et al.*, 1996). A copy of this motif, YIVL213-216, is at positions 3-6 of D<sub>2</sub>-IC3, and a deletion mutant of residues 2-6 that removes the entire motif (D<sub>2</sub>-IC3ΔIYIVL) caused a 90% loss of arrestin binding that was slightly greater than the loss of binding to D<sub>2</sub>-IC3ΔIYIV (Fig. 4A and B) and similar to the 83% loss of binding caused by the N-terminal 20-residue truncation (D<sub>2</sub>-IC3ΔNT20; Fig. 3). Finally, we mutated IYIV212-215 or IYIVL212-216 to alanines,

MOL #50534

creating the substitution mutants D<sub>2</sub>-IC3IYIV212-215A4 and D<sub>2</sub>-IC3IYIVL212-216A5 that correspond to the deletion mutants D<sub>2</sub>-IC3ΔIYIV and D<sub>2</sub>-IC3ΔIYIVL (Fig. 4A). Alanine substitutions were at least as effective as corresponding deletion mutations at decreasing the binding of arrestin (Fig. 4B).

**The A4 Mutation had no Effect on D<sub>2</sub> Receptor Affinity for Ligands.** The effect of the A4 (IYIV212-215A4) mutation in the context of the full-length D<sub>2</sub> receptor was evaluated by analysis of radioligand binding to membranes prepared from HEK 293 cells transiently expressing wild type D<sub>2</sub> and D<sub>2</sub>-A4 mutant receptors. Saturation analysis of the binding of the D<sub>2</sub>-selective antagonist radioligand, [<sup>3</sup>H]spiperone, yielded *K<sub>d</sub>* values of 83 ± 13 pM and 86 ± 4 pM for D<sub>2</sub> and D<sub>2</sub>-A4 receptors, respectively (Table II; Fig. 5A). Competition binding analysis of the ability of dopamine to decrease the binding of [<sup>3</sup>H]spiperone indicated that high- and low-affinity binding of dopamine to the A4 mutant was indistinguishable from binding to the wild type D<sub>2</sub> receptor (Table II; Fig. 5B), suggesting that the A4 mutation did not alter D<sub>2</sub> receptor affinity for ligands or coupling to G proteins.

**Inhibition of Cyclic AMP by D<sub>2</sub>-A4.** The ability of the A4 mutant to inhibit adenylate cyclase was also characterized. The potency of the agonist was unchanged by the mutation, either in the absence or presence of overexpressed arrestin3 (Fig. 6A and B). Similarly, maximal inhibition of cyclic AMP accumulation did not differ significantly between wild type D<sub>2</sub> and D<sub>2</sub>-A4 in the absence of added arrestin (Fig. 6C), indicating that the A4 mutation did not affect receptor activation of the G proteins mediating inhibition of adenylate cyclase. Overexpression of arrestin3 significantly decreased D<sub>2</sub> receptor-mediated maximum inhibition of cyclic AMP accumulation (57 ± 1%), whereas the maximum inhibition mediated by the A4 mutant was not

MOL #50534

altered ( $69 \pm 4\%$ ), perhaps reflecting the consequences of differential arrestin translocation and receptor desensitization and internalization (but see below).

**The A4 Mutation Abolished Arrestin3 Translocation and Receptor Internalization.** The functional interaction of D<sub>2</sub> and D<sub>2</sub>-A4 receptors with arrestin was evaluated by quantifying agonist-induced translocation of arrestin to the cell membrane and agonist-induced receptor internalization in HEK 293 cells transiently co-expressing wild type or mutant D<sub>2</sub> receptor and arrestin3. The amount of arrestin3 in the membrane fraction of cells transfected with the wild type D<sub>2</sub> receptor was doubled from  $1.0 \pm 0.2$  ng of arrestin3 in untreated cells to  $2.1 \pm 0.3$  ng after treatment with 10  $\mu$ M dopamine for 20 min (Fig. 7A). In contrast, in cells coexpressing D<sub>2</sub>-A4 and arrestin3, the levels of arrestin3 in the membrane preparations were equal in vehicle- and dopamine-treated cells ( $1.2 \pm 0.1$  vs.  $1.1 \pm 0.1$ , respectively), indicating that the A4 mutation prevented dopamine-induced arrestin3 translocation to the plasma membrane (Fig. 7A). Neither D<sub>2</sub> nor D<sub>2</sub>-A4 mediated detectable translocation of arrestin2 in HEK 293 cells co-transfected with arrestin2 (data not shown).

D<sub>2</sub> receptor internalization was assessed by quantifying the agonist-induced loss of cell-surface binding of the hydrophilic ligand [<sup>3</sup>H]sulpiride in an intact cell binding assay. In cells transiently expressing the wild type D<sub>2</sub> receptor and arrestin3, treatment with 10  $\mu$ M dopamine for 20 min decreased binding of [<sup>3</sup>H]sulpiride by  $32 \pm 2\%$  (Fig. 7B). In contrast, co-transfection with arrestin2 supported little agonist-induced internalization (data not shown). Consistent with the lack of arrestin3 translocation mediated by the A4 mutant and the role of arrestin binding in D<sub>2</sub> receptor internalization (Macey *et al.*, 2004), the A4 mutation abolished dopamine-induced receptor internalization. In contrast, the D<sub>2</sub>-IC3 $\Delta$ MID mutation (deletion of 121 residues comprising all of IC3 except the 20 residues at each terminus; see Fig. 2A) had little effect on



MOL #50534

receptor internalization (Fig. 7B). Together, these data suggest that the motif IYIV212-215 at the N-terminus of the D<sub>2</sub> receptor IC3 is required for a functional interaction between the receptor and arrestin3.

**The A4 Mutation Abolished Desensitization of Cyclic AMP Accumulation.** Because of the role of arrestin in desensitization of GPCRs, we evaluated the effect of the A4 mutation on dopamine-induced desensitization of D<sub>2</sub> receptor-mediated inhibition of cyclic AMP accumulation. HEK 293 cells stably expressing wild type or mutant D<sub>2</sub> receptors and transiently overexpressing arrestin3 were treated with 10 μM dopamine for 20 min prior to measuring acute inhibition of forskolin-stimulated activity (Fig. 8). Dopamine pretreatment caused a decrease in the potency of dopamine at the wild type D<sub>2</sub> receptor from 3.7 nM (pEC<sub>50</sub> = 8.43 ± 0.05) to 16 nM (pEC<sub>50</sub> = 7.78 ± 0.09; *p* < 0.01), whereas the potency at the D<sub>2</sub>-A4 mutant receptor was unaffected (pEC<sub>50</sub> = 7.97 ± 0.03 for vehicle-treated cells vs. 8.12 ± 0.33 for dopamine-treated cells). Maximal inhibition of cyclic AMP accumulation was not altered by dopamine treatment, but in this set of experiments the maximal inhibition by D<sub>2</sub>-A4 was slightly but not significantly enhanced compared to wild type D<sub>2</sub> (Fig. 8C).

**The A4 Mutation Decreased Arrestin-dependent Activation of ERKs.** Because the D<sub>2</sub>-A4 mutation abolished receptor internalization/desensitization and arrestin3 translocation, one intriguing issue is whether this mutation inhibits ERK activation. Receptor-mediated activation of ERKs (ERK1, 44 kDa and ERK2, 42 kDa) was measured by quantifying the abundance of dually phosphorylated ERKs. Treatment with dopamine induced rapid and robust activation of ERKs in HEK 293 cells expressing either the wild type or the mutant receptor (Fig. 9). The levels of dopamine-stimulated phospho-ERKs for D<sub>2</sub> wild type receptor and D<sub>2</sub>-A4 receptor in the absence of overexpressed arrestin were not significantly different (Fig. 9C), which suggested

MOL #50534

that D<sub>2</sub> receptor stimulation of ERKs in HEK 293 cells was independent of arrestin binding to the receptor in the absence of arrestin overexpression. For wild type D<sub>2</sub> receptor, overexpression of arrestin3 dramatically increased basal (12 fold) and dopamine-induced (2.9 fold) ERK phosphorylation, whereas for the A4 mutant the corresponding increments were smaller (3.2 and 1.5 fold, respectively; Fig. 9A and C). Thus, the A4 mutation decreased arrestin-dependent activation of ERKs. Similar results were observed with the D<sub>2</sub>-selective agonist quinpirole (data not shown).

To assess the G protein-dependence of D<sub>2</sub> receptor-mediated activation of ERKs, some cells were pretreated with pertussis toxin (PTX; 50 ng/ml overnight) prior to agonist treatment. This treatment regimen abolishes D<sub>2</sub> receptor inhibition of adenylate cyclase activity and coupling to G proteins in a variety of cell lines, including HEK 293 cells (Neve *et al.*, 1989; Watts and Neve, 1996; Watts and Neve, 1997; Watts *et al.*, 1998). PTX treatment greatly decreased activation of ERKs by D<sub>2</sub> and D<sub>2</sub>-A4 receptors (Fig. 9), indicating that a major component of ERK activation by the D<sub>2</sub> receptor in HEK 293 cells requires PTX-sensitive G proteins G $\alpha_{i/o}$ . PTX treatment also decreased the activation of ERKs in cells overexpressing arrestin3, but a substantial PTX-insensitive component may be mediated by arrestin rather than G $\alpha_{i/o}$ . This PTX-insensitive and arrestin-dependent component is smaller but still present in cells expressing D<sub>2</sub>-A4, suggesting that the mutant receptor retains residual ability to bind arrestins.

MOL #50534

## Discussion

Studies of receptor-mediated translocation of GFP-tagged arrestin to the cell membrane have identified IC2 and IC3 as being particularly important for the interaction between D2-like receptors and arrestin (Kim *et al.*, 2001). In the present study, we constructed GST fusion proteins as “bait” to identify arrestin-binding subdomains. We determined that IC3 from both D<sub>2</sub> and D<sub>3</sub> receptors bound arrestin3 more avidly than arrestin2. Our data also indicated that D<sub>2</sub>-IC3 and D<sub>3</sub>-IC3 bound arrestin with similar affinities; this is in contrast to the lower and different affinities of D<sub>2</sub>-IC2 and D<sub>3</sub>-IC2 for arrestin (Lan *et al.*, 2008). Four residues at the N-terminus of D<sub>2</sub>-IC3 were critical for arrestin binding. Alanine mutations of those four residues abolished D<sub>2</sub> receptor-mediated translocation of arrestin and receptor internalization and decreased arrestin-dependent activation of ERKs, without altering receptor coupling to G proteins and the potency of agonist for inhibition of cyclic AMP accumulation, suggesting that the D<sub>2</sub>-A4 mutant is a receptor that is biased towards G protein-mediated signaling pathways and away from arrestin-dependent pathways.

Accumulating evidence suggests that multiple mechanisms contribute to GPCR-arrestin interactions. Phosphorylation of receptor intracellular domains by GRKs may promote arrestin binding due to electrostatic interactions between negatively charged phosphates on the receptor and positively charged arrestin residues that serve as a phosphorylation sensor (Gurevich and Gurevich, 2006). These electrostatic interactions may also induce a conformational change in arrestin that enhances its binding to the receptor (Gurevich and Gurevich, 2004). Phosphorylation of the receptor may also contribute indirectly to arrestin binding by initiating conformational changes of the intracellular domains that expose binding sites for arrestin (Kim *et al.*, 2004). Receptor activation is also accompanied by conformational changes, exposing

MOL #50534

receptor sites that interact with a proposed activation sensor in arrestin (Gurevich and Gurevich, 2006). Because the GST fusion proteins are not phosphorylated, this method does not identify receptor phosphorylation sites that bind arrestin, but identifies only determinants of binding that are revealed by activation- or phosphorylation-dependent conformational changes of the receptor intracellular domains; the determinants are presumably occluded in the inactive and/or unphosphorylated full-length receptor but exposed when the receptor fragments are expressed as GST fusion proteins, free from constraints imposed by other domains of the intact receptor. Thus, this *in vitro* binding assay may identify both phosphorylation-independent and some phosphorylation-dependent determinants of the interaction between receptor and arrestin.

Arrestin-binding domains/residues vary among GPCRs, based on results from binding studies with purified arrestins along with recombinant or synthetic peptides representing receptor intracellular fragments. Both arrestin2 and arrestin3 bind to IC3 of the 5-HT<sub>2A</sub> receptor and the  $\delta$ -opioid receptor, and to the C-terminal domain of  $\delta$ - and  $\kappa$ -opioid receptors, with certain serine/threonine residues in these receptor regions being important for binding (Gelber *et al.*, 1999; Cen *et al.*, 2001). Binding of arrestin2 to the M<sub>3</sub>-muscarinic receptor requires both N- and C-terminal regions of the IC3 of the M<sub>3</sub>-muscarinic receptor subdomains, whereas for arrestin3 the C-terminal region of the IC3 is sufficient for binding (Wu *et al.*, 1997). Attempts to define more precisely non-phosphorylated residues that are sites of arrestin binding have identified the highly conserved DRY sequence at the N-terminus of IC2 (Hüttenrauch *et al.*, 2002), an aspartate residue in IC3 of the luteinizing hormone/choriogonadotropin receptor that is thought to mimic a phosphorylated residue (Mukherjee *et al.*, 2002), and BXXBB (B = basic residue, X = any) motifs present in the N- and C-terminal portions of IC3 of the  $\alpha_2$ -adrenoceptors (DeGraff

MOL #50534

*et al.*, 2002). Our studies on basic residues in similar locations in the D<sub>2</sub> receptor suggest that they are not involved in the binding of arrestin3 to this receptor (unpublished observations).

Our GST pull-down studies demonstrated the ability of D<sub>2</sub>-like receptor IC3 to bind arrestin3 with high affinity ( $K_d = 70\text{-}80$  nM). These values are lower (*i.e.*, higher affinity) than the micromolar affinities derived from surface plasmon resonance studies (Cen *et al.*, 2001; Liu *et al.*, 2004), closer to the sub-nanomolar affinity of arrestin for the intact phosphorylated active receptors (Gurevich *et al.*, 1995), and consistent with estimates of arrestin concentrations in neurons (30-200 nM) (Gurevich *et al.*, 2004), suggesting that arrestin affinities for the receptor loops are biologically relevant. Consistent with the view that stronger agonist-induced phosphorylation of the D<sub>2</sub> receptor than of the D<sub>3</sub> receptor contributes to differential patterns of translocation of arrestin to cell membrane (Kim *et al.*, 2001), we observed that arrestin bound with equal affinity to non-phosphorylated IC3 from both receptors.

Studies with full-length receptors have used direct binding of arrestin, arrestin translocation, and receptor internalization as assays to explore receptor determinants of arrestin binding (Gurevich and Gurevich, 2006), although with these assays it is sometimes difficult to distinguish between phosphorylated or non-phosphorylated determinants of binding. These approaches revealed that receptor elements involved in arrestin binding can be localized almost anywhere on the intracellular surface of the receptor, including in the C-terminal tail (Qian *et al.*, 2001), IC3 (Kim *et al.*, 2001; Lee *et al.*, 2000; DeGraff *et al.*, 2002; Namkung and Sibley, 2004), IC2 (Raman *et al.*, 1999; Kim *et al.*, 2001), and IC1 (Raman *et al.*, 1999; Kishi *et al.*, 2002). The present work describes a significant non-phosphorylated determinant of arrestin binding at the N terminus of D<sub>2</sub>-IC3.

MOL #50534

The predominant pathway for GPCR endocytosis/internalization involves arrestin-dependent recruitment into clathrin-coated vesicles (Ferguson, 2001). A number of motifs have been proposed to be required for receptor endocytosis, including NP(X<sub>2,3</sub>)Y, DRYXXV/IXXPL, BXXBB, and a dileucine motif, all located within or close to the transmembrane proximal domains of IC2, IC3, and the carboxyl-terminus of GPCRs (DeGraff *et al.*, 2002; Barak *et al.*, 1994; Moro *et al.*, 1994; Arora *et al.*, 1995; Gabilondo *et al.*, 1997). Interestingly, a membrane-proximal YXX $\phi$  motif, where  $\phi$  is any bulky hydrophobic residue, was reported to interact with  $\mu$ 2 subunit of the AP2 complex and to be important for clathrin-mediated endocytosis of many integral membrane proteins (Ohno *et al.*, 1995; Boll *et al.*, 1996; Royle *et al.*, 2005; Owen and Evans, 1998; Collins *et al.*, 2002; Vogt *et al.*, 2005). In our study, via GST pull-down assays, we identified the 5-residue sequence IYIVL at the N-terminus of D<sub>2</sub>-IC3 that incorporates the YXX $\phi$  motif and that is required for high-affinity binding of arrestin. Mutation of four of these residues to alanine in the full-length receptor (D<sub>2</sub>-A4) prevented D<sub>2</sub> receptor-mediated translocation of arrestin to the membrane, agonist-induced receptor internalization, and desensitization of cyclic AMP accumulation associated with overexpression of arrestin, indicating that this sequence is required for a functional interaction between the D<sub>2</sub> receptor and arrestin. Importantly, the A4 mutation did not alter either high-affinity binding of dopamine to the D<sub>2</sub> receptor or the potency of dopamine for D<sub>2</sub>-mediated inhibition of cyclic AMP accumulation, suggesting that coupling of the mutant receptor to G proteins and the activation of G proteins were not affected by the mutation; thus, distinct structural determinants of the D<sub>2</sub> receptor mediate its interactions with arrestins and G proteins.

The lack of evidence for binding of arrestin2 to the D<sub>2</sub> receptor is surprising in light of an earlier publication from this laboratory reporting that the D<sub>2</sub> receptor preferentially interacts with

MOL #50534

arrestin2 over arrestin3 in neostriatal neurons (Macey *et al.*, 2004). Much of the discrepancy could potentially be explained by the type of cell in which the receptor is expressed. In the earlier publication we reported that the preferential interaction with arrestin2 was observed in neostriatal neurons, but not in NS20Y neuroblastoma cells, and we concluded that the preferential interaction with arrestin2 in neurons was related to compartmentalization of the receptor, arrestins, or other interacting proteins rather than to an inherent preference for arrestin3. The preferential interaction with arrestin3 in HEK293 cells may be due to the influence of similar cell type-dependent factors. In addition, the preference for arrestin3 described here is consistent with work indicating that arrestin3, but not arrestin2, mediates D<sub>2</sub> receptor inhibition of the protein kinase Akt and methamphetamine-induced locomotor activity (Beaulieu *et al.*, 2005).

As a signaling-biased GPCR, capable of activating G protein-regulated signaling pathways but with reduced ability to bind arrestin, the D<sub>2</sub>-A4 mutant receptor is a useful experimental tool. Stimulation of some GPCRs induces the assembly of a protein complex in which arrestin acts as a receptor-regulated scaffold, recruiting mitogen-activated protein kinase cascades to the agonist-occupied GPCR and facilitating GPCR-dependent kinase activation (Luttrell *et al.*, 1999; McDonald *et al.*, 2000). That the D<sub>2</sub> receptor-mediated activation of ERKs was not affected by the A4 mutation but almost entirely prevented by PTX in HEK 293 cells indicates that when the abundance of arrestins is low, D<sub>2</sub> receptor activation of ERKs is independent of arrestin binding to the receptor and mediated predominantly by PTX-sensitive G proteins. In contrast, overexpression of arrestin3 greatly increased the D<sub>2</sub> receptor-mediated activation of ERKs, in particular enhancing a PTX-insensitive component of ERK activation, but had a much smaller effect in cells expressing the arrestin-insensitive mutant D<sub>2</sub>-A4. Similarly, parathyroid hormone

MOL #50534

receptor-mediated activation of ERK in HEK 293 cells is comprised of distinct arrestin- and G protein-dependent pathways (Gesty-Palmer *et al.*, 2006).

Arrestin3 is also a scaffold for Akt and protein phosphatase 2A, and disruption of signaling through this protein complex reduces dopamine-dependent behaviors in mice (Beaulieu *et al.*, 2005). If arrestin3 binding to the D<sub>2</sub> receptor is required for activation of this signaling pathway, we predict that the D<sub>2</sub>-A4 mutant would also be deficient in signaling via Akt. The use of arrestin3 null mutant mice is one way in which investigators have differentiated between G protein- and arrestin-dependent responses to receptor activation (Beaulieu *et al.*, 2005; Raehal *et al.*, 2005); the use of arrestin-insensitive mutant receptors represents an alternative approach.



MOL #50534

## References

- Arora KK, Sakai A, and Catt KJ (1995) Effects of second intracellular loop mutations on signal transduction and internalization of the gonadotropin-releasing hormone receptor. *J Biol Chem* **270**:22820-22826.
- Barak LS, Tiberi M, Freedman NJ, Kwatra MM, Lefkowitz RJ, and Caron MG (1994) A highly conserved tyrosine residue in G protein-coupled receptors is required for agonist-mediated  $\beta_2$ -adrenergic receptor sequestration. *J Biol Chem* **269**:2790-2795.
- Beaulieu JM, Sotnikova TD, Marion S, Lefkowitz RJ, Gainetdinov RR, and Caron MG (2005) An Akt/ $\beta$ -arrestin 2/PP2A signaling complex mediates dopaminergic neurotransmission and behavior. *Cell* **122**:261-273.
- Boll W, Ohno H, Songyang Z, Rapoport I, Cantley LC, Bonifacino JS, and Kirchhausen T (1996) Sequence requirements for the recognition of tyrosine-based endocytic signals by clathrin AP-2 complexes. *EMBO J* **15**:5789-5795.
- Cen B, Xiong Y, Ma L, and Pei G (2001) Direct and differential interaction of  $\beta$ -arrestins with the intracellular domains of different opioid receptors. *Mol Pharmacol* **59**:758-764.
- Cheng Y-C and Prusoff WH (1973) Relationship between the inhibition constant ( $K_I$ ) and the concentration of inhibitor which causes 50 per cent inhibition ( $I_{50}$ ) of an enzymatic reaction. *Biochem Pharmacol* **22**:3099-3108.
- Collins BM, McCoy AJ, Kent HM, Evans PR, and Owen DJ (2002) Molecular architecture and functional model of the endocytic AP2 complex. *Cell* **109**:523-535.
- DeGraff JL, Gurevich VV, and Benovic JL (2002) The third intracellular loop of  $\alpha_2$ -adrenergic receptors determines subtype specificity of arrestin interaction. *J Biol Chem* **277**:43247-43252.

MOL #50534

Ferguson SSG (2001) Evolving concepts in G protein-coupled receptor endocytosis: The role in receptor desensitization and signaling. *Pharmacol Rev* **53**:1-24.

Gabilondo AM, Hegler J, Krasel C, Boivin-Jahns V, Hein L, and Lohse MJ (1997) A dileucine motif in the C terminus of the  $\beta_2$ -adrenergic receptor is involved in receptor internalization. *Proc Natl Acad Sci USA* **94**:12285-12290.

Gelber EI, Kroeze WK, Willins DL, Gray JA, Sinar CA, Hyde EG, Gurevich V, Benovic J, and Roth BL (1999) Structure and function of the third intracellular loop of the 5-hydroxytryptamine<sub>2A</sub> receptor: The third intracellular loop is  $\alpha$ -helical and binds purified arrestins. *J Neurochem* **72**:2206-2214.

Gesty-Palmer D, Chen M, Reiter E, Ahn S, Nelson CD, Wang S, Eckhardt AE, Cowan CL, Spurney RF, Luttrell LM, and Lefkowitz RJ (2006) Distinct  $\beta$ -arrestin- and G protein-dependent pathways for parathyroid hormone receptor-stimulated ERK1/2 activation. *J Biol Chem* **281**:10856-10864.

Gurevich EV, Benovic JL, and Gurevich VV (2004) Arrestin2 expression selectively increases during neural differentiation. *J Neurochem* **91**:1404-1416.

Gurevich VV, Dion SB, Onorato JJ, Ptasienski J, Kim CM, Sterne-Marr R, Hosey MM, and Benovic JL (1995) Arrestin interactions with G protein-coupled receptors. Direct binding studies of wild type and mutant arrestins with rhodopsin,  $\beta_2$ -adrenergic, and m2 muscarinic cholinergic receptors. *J Biol Chem* **270**:720-731.

Gurevich VV and Gurevich EV (2004) The molecular acrobatics of arrestin activation. *Trends in Pharmacological Sciences* **25**:105-111.

Gurevich VV and Gurevich EV (2006) The structural basis of arrestin-mediated regulation of G-protein-coupled receptors. *Pharmacol Ther* **110**:465-502.

MOL #50534

Han M, Gurevich VV, Vishnivetskiy SA, Sigler PB, and Schubert C (2001) Crystal structure of  $\beta$ -arrestin at 1.9 Å: Possible mechanism of receptor binding and membrane translocation.

*Struct Fold Des* **9**:869-880.

Hüttenrauch F, Nitzki A, Lin FT, Höning S, and Oppermann M (2002)  $\beta$ -arrestin binding to CC chemokine receptor 5 requires multiple C-terminal receptor phosphorylation sites and involves a conserved Asp-Arg-Tyr sequence motif. *J Biol Chem* **277**:30769-30777.

Itokawa M, Toru M, Ito K, Tsuga H, Kameyama K, Haga T, Arinami T, and Hamaguchi H (1996) Sequestration of the short and long isoforms of dopamine D<sub>2</sub> receptors expressed in Chinese hamster ovary cells. *Mol Pharmacol* **49**:560-566.

Kim KM, Valenzano KJ, Robinson SR, Yao WD, Barak LS, and Caron MG (2001) Differential regulation of the dopamine D<sub>2</sub> and D<sub>3</sub> receptors by G protein-coupled receptor kinases and  $\beta$ -arrestins. *J Biol Chem* **276**:37409-37414.

Kim OJ, Gardner BR, Williams DB, Marinec PS, Cabrera DM, Peters JD, Mak CC, Kim KM, and Sibley DR (2004) The role of phosphorylation in D<sub>1</sub> dopamine receptor desensitization: evidence for a novel mechanism of arrestin association. *J Biol Chem* **279**:7999-8010.

Kishi H, Krishnamurthy H, Galet C, Bhaskaran RS, and Ascoli M (2002) Identification of a short linear sequence present in the C-terminal tail of the rat follitropin receptor that modulates arrestin-3 binding in a phosphorylation-independent fashion. *J Biol Chem* **277**:21939-21946.

Lan H, Gurevich VV, and Neve KA (2008) An intracellular loop 2 amino acid residue that contributes to the binding of arrestin to the dopamine D<sub>2</sub> receptor. *Mol Pharmacol*

Submitted for publication.

MOL #50534

- Lee KB, Ptasienski JA, Pals-Rylaarsdam R, Gurevich VV, and Hosey MM (2000) Arrestin binding to the M<sub>2</sub> muscarinic acetylcholine receptor is precluded by an inhibitory element in the third intracellular loop of the receptor. *J Biol Chem* **275**:9284-9289.
- Liu P, Roush ED, Bruno J, Osawa S, and Weiss ER (2004) Direct binding of visual arrestin to a rhodopsin carboxyl terminal synthetic phosphopeptide. *Mol Vis* **10**:712-719.
- Liu Y, Buck DC, Macey TA, Lan H, and Neve KA (2007) Evidence that calmodulin binding to the dopamine D<sub>2</sub> receptor enhances receptor signaling. *J Recept Signal Transduct Res* **27**:47-65.
- Luttrell LM, Ferguson SSG, Daaka Y, Miller WE, Maudsley S, Della Rocca GJ, Lin FT, Kawakatsu H, Owada K, Luttrell DK, Caron MG, and Lefkowitz RJ (1999)  $\beta$ -arrestin-dependent formation of  $\beta_2$  adrenergic receptor Src protein kinase complexes. *Science* **283**:655-661.
- Macey TA, Gurevich VV, and Neve KA (2004) Preferential interaction between the dopamine D<sub>2</sub> receptor and arrestin2 in neostriatal neurons. *Mol Pharmacol* **66**:1635-1642.
- Macey TA, Liu Y, Gurevich VV, and Neve KA (2005) Dopamine D<sub>1</sub> receptor interaction with arrestin3 in neostriatal neurons. *J Neurochem* **93**:128-134.
- Makarova O, Kamberov E, and Margolis B (2000) Generation of deletion and point mutations with one primer in a single cloning step. *Biotechniques* **29**:970-972.
- McDonald PH, Chow CW, Miller WE, Laporte SA, Field ME, Lin FT, Davis RJ, and Lefkowitz RJ (2000)  $\beta$ -Arrestin 2: A receptor-regulated MAPK scaffold for the activation of JNK3. *Science* **290**:1574-1577.

MOL #50534

- Moro O, Shockley MS, Lameh J, and Sadee W (1994) Overlapping multi-site domains of the muscarinic cholinergic Hm1 receptor involved in signal transduction and sequestration. *J Biol Chem* **269**:6651-6655.
- Mukherjee S, Gurevich VV, Preninger A, Hamm HE, Bader MF, Fazleabas AT, Birnbaumer L, and Hunzicker-Dunn M (2002) Aspartic acid 564 in the third cytoplasmic loop of the luteinizing hormone/choriogonadotropin receptor is crucial for phosphorylation-independent interaction with arrestin2. *J Biol Chem* **277**:17916-17927.
- Namkung Y and Sibley DR (2004) Protein kinase C mediates phosphorylation, desensitization, and trafficking of the D<sub>2</sub> dopamine receptor. *J Biol Chem* **279**:49533-49541.
- Neve KA, Henningsen RA, Bunzow JR, and Civelli O (1989) Functional characterization of a rat dopamine D-2 receptor cDNA expressed in a mammalian cell line. *Mol Pharmacol* **36**:446-451.
- Neve KA, Seamans JK, and Trantham-Davidson H (2004) Dopamine receptor signaling. *J Recept Signal Transduct Res* **24**:165-205.
- Ohno H, Stewart J, Fournier MC, Bosshart H, Rhee I, Miyatake S, Saito T, Gallusser A, Kirchhausen T, and Bonifacino JS (1995) Interaction of tyrosine-based sorting signals with clathrin-associated proteins. *Science* **269**:1872-1875.
- Owen DJ and Evans PR (1998) A structural explanation for the recognition of tyrosine-based endocytotic signals. *Science* **282**:1327-1332.
- Pierce KL and Lefkowitz RJ (2001) Classical and new roles of  $\beta$ -arrestins in the regulation of G-protein-coupled receptors. *Nat Rev Neurosci* **2**:727-733.

MOL #50534

Qian H, Pipolo L, and Thomas WG (2001) Association of  $\beta$ -arrestin 1 with the type 1A

angiotensin II receptor involves phosphorylation of the receptor carboxyl terminus and correlates with receptor internalization. *Mol Endocrinol* **15**:1706-1719.

Raehal KM, Walker JK, and Bohn LM (2005) Morphine side effects in  $\beta$ -arrestin 2 knockout mice. *J Pharmacol Exp Ther* **314**:1195-1201.

Raman D, Osawa S, and Weiss ER (1999) Binding of arrestin to cytoplasmic loop mutants of bovine rhodopsin. *Biochemistry* **38**:5117-5123.

Royle SJ, Qureshi OS, Bobanovic LK, Evans PR, Owen DJ, and Murrell-Lagnado RD (2005) Non-canonical YXXG $\phi$  endocytic motifs: recognition by AP2 and preferential utilization in P2X4 receptors. *J Cell Sci* **118**:3073-3080.

Vogt C, Eickmann M, Diederich S, Moll M, and Maisner A (2005) Endocytosis of the Nipah virus glycoproteins. *J Virol* **79**:3865-3872.

Watts VJ and Neve KA (1996) Sensitization of endogenous and recombinant adenylylase by activation of D<sub>2</sub> dopamine receptors. *Mol Pharmacol* **50**:966-976.

Watts VJ and Neve KA (1997) Activation of type II adenylylase by D<sub>2</sub> and D<sub>4</sub> but not D<sub>3</sub> dopamine receptors. *Mol Pharmacol* **52**:181-186.

Watts VJ, Wiens BL, Cumbay MG, Vu MN, Neve RL, and Neve KA (1998) Selective activation of G $\alpha_o$  by D<sub>2L</sub> dopamine receptors in NS20Y neuroblastoma cells. *J Neurosci* **18**:8692-8699.

Wu GY, Krupnick JG, Benovic JL, and Lanier SM (1997) Interaction of arrestins with intracellular domains of muscarinic and  $\alpha_2$ -adrenergic receptors. *J Biol Chem* **272**:17836-17842.

MOL #50534

### **Footnotes**

This work was supported by United States Public Health Service grants MH045372 (K.A.N.), GM63097 (V.V.G.), and EY11500 (V.V.G.), and the VA Merit Review and Career Scientist programs (K.A.N.).

MOL #50534

## Figure Legends

**Fig. 1.** Binding of arrestins to GST-D<sub>2</sub>-IC3 and GST-D<sub>3</sub>-IC3 fusion proteins. GST alone (GST, 150 ng) or receptor third intracellular loop GST fusion proteins (GST-D<sub>2</sub>-IC3 and GST-D<sub>3</sub>-IC3, 300 ng) were incubated with the indicated amount of arrestin2 or arrestin3. The amount of arrestin that co-eluted with GST or the GST fusion proteins was determined by immunoblotting with anti-arrestin antibodies. Results were quantified using standard curves constructed with known amounts of arrestin2 and arrestin3. **A**, Immunoblots are shown from an experiment representative of 4 independent experiments. Arrestin standards are shown on the right of each blot. **B**, The protein-stained gel demonstrates that equal amounts of GST-D<sub>2</sub>-IC3 and GST-D<sub>3</sub>-IC3 were included in the reactions. **C**, Results shown are the mean  $\pm$  S.E.M. for the binding of arrestins to IC3 fusion proteins. The amount of arrestin bound is plotted against the amount of arrestin included in the pull-down assay.

**Fig. 2.** Binding of arrestin3 to D<sub>2</sub> and D<sub>3</sub> receptor IC3 truncation mutants. **A**, Alignment of IC3 from the rat D<sub>2</sub> and D<sub>3</sub> dopamine receptors. Each IC3 is divided into NT (20 residues), MID (121 residues not shown for D<sub>2</sub> and 123 residues for D<sub>3</sub>) and CT (20 residues). **B** and **C**, Purified GST or GST fusion proteins (ranging from 500 to 800 ng to keep the molar concentration constant) were incubated with 100 ng of purified arrestin3, and the amount of arrestin bound was determined by quantitative immunoblotting. **Upper panels**, representative arrestin immunoblots are shown. The amount of bound arrestin was quantified using a standard curve constructed with known amounts of arrestin as shown on the right of each blot. **Middle panels**, protein-stained gels used to verify the size of the truncation mutants and to quantify the amount of each fusion protein are shown. **Lower panels**, the results shown are the mean  $\pm$  S.E.M. from 4 independent



MOL #50534

experiments (\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  versus the respective wild type fusion protein by paired t test).

**Fig. 3.** Binding of arrestin3 to truncation mutants of the amino-terminal region of the D<sub>2</sub> receptor IC3. **A**, Sequence of the 20 amino acids in the amino-terminal region of D<sub>2</sub>-IC3. The first 5 or 10 or the second 10 amino acids were deleted to form the indicated mutants. **B**, Purified GST or GST fusion proteins (400 ng) were incubated with 100 ng of purified arrestin3, and the amount of arrestin bound was determined by quantitative immunoblotting. The results shown are the mean  $\pm$  S.E.M. from 4 independent experiments (\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  versus D<sub>2</sub>-IC3 by paired t test).

**Fig. 4.** Binding of arrestin3 to 3-, 4-, and 5-residue deletion mutants and 4- and 5-residue substitution mutants of the amino-terminal region of D<sub>2</sub>-IC3. **A**, Sequence of amino acids in the amino-terminal region of D<sub>2</sub>-IC3, along with the name of each construct where  $\Delta$  denotes the residues that were deleted. **B**, Purified GST or GST fusion proteins (600 ng) were incubated with 100 ng of purified arrestin3, and the amount of arrestin bound was determined by quantitative immunoblotting. Results shown are the mean  $\pm$  S.E.M. from 4-5 independent experiments. (\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  versus D<sub>2</sub>-IC3 by paired t test).

**Fig. 5.** Ligand binding properties of the D<sub>2</sub>-A4 mutant receptor. **A**, Saturation analysis of the binding of the D<sub>2</sub> receptor antagonist [<sup>3</sup>H]spiperone to membrane preparations from HEK 293 cells expressing wild type D<sub>2</sub> or D<sub>2</sub>-A4 receptors. Data are plotted as specific binding (fmol/mg protein) versus the free ligand concentration of radioligand. **B**, Inhibition of the binding of [<sup>3</sup>H]spiperone by dopamine. Data are plotted as a percentage of specific binding in the absence

MOL #50534

of dopamine versus the logarithm of the concentration of dopamine. In both A and B, the experiment shown is representative of three or more independent experiments.

**Fig. 6.** Dopamine inhibition of forskolin-stimulated cyclic AMP accumulation. HEK293 cells overexpressing wild type D<sub>2</sub> (WT) or D<sub>2</sub>-A4 (A4) receptors in the absence or presence of exogenous arrestin3 (arr) were incubated with 30 μM forskolin and increasing concentrations of dopamine, and cyclic AMP accumulation was determined. **A**, data are plotted as a percentage of forskolin-stimulated cyclic AMP accumulation in the absence of dopamine. **B** and **C**, the mean ± S.E.M. values for EC<sub>50</sub> and maximum inhibition from four experiments are shown. (\*\*\*)  $p < 0.001$  versus wild type by paired t test). The expression levels of cell surface D<sub>2</sub> receptors were similar in all cell lines (data not shown).

**Fig 7.** Agonist-induced translocation of arrestin3 and receptor internalization in HEK 293 cells coexpressing arrestin3 and wild type or mutant D<sub>2</sub> receptors. **A**, Cells were treated with 10 μM dopamine for 20 min, membranes were prepared, and levels of arrestin3 were determined by quantitative immunoblotting. Results were quantified using standard curves constructed with known amounts of purified arrestin3. The expression of cell surface receptor for the A4 mutant is similar to or higher than that for D<sub>2</sub> wild type, as determined by intact cell [<sup>3</sup>H]sulpiride binding (data not shown). **Upper panel**, an immunoblot is shown from an experiment representative of 4 independent experiments. Supernatants were loaded to show similar expression levels of arrestin3. **Lower panel**, Results from all four experiments are shown as the mean ± S.E.M. **B**, Cells were treated with 10 μM dopamine for 20 min, and were subjected to the intact cell [<sup>3</sup>H]sulpiride binding assay. Results are expressed as the percentage by which the binding of [<sup>3</sup>H]sulpiride to cell surface receptors decreased after agonist stimulation, and are shown as the

MOL #50534

mean  $\pm$  S.E.M. from 4 independent experiments (\*  $p < 0.05$ , \*\*  $p < 0.01$  versus wild type by paired t test). The expression levels of arrestin3 were similar in cells transfected with wild type or mutant receptors (data not shown).

**Fig. 8.** Desensitization of dopamine inhibition of forskolin-stimulated cyclic AMP accumulation. HEK293 cells stably overexpressing wild type D<sub>2</sub> (WT) or D<sub>2</sub>-A4 (A4) receptors and transiently overexpressing arrestin3 were treated with 10  $\mu$ M dopamine (DA) or vehicle (Con) for 20 min. After washing, cells were incubated with 30  $\mu$ M forskolin and increasing concentrations of dopamine for 10 min, and cyclic AMP accumulation was determined. **A**, results from a representative experiment are plotted as a percentage of forskolin-stimulated cyclic AMP accumulation in the absence of dopamine. **B** and **C**, the mean  $\pm$  S.E.M. values for EC<sub>50</sub> and maximum inhibition from four independent experiments are shown. (\*\*  $p < 0.01$  versus vehicle-treated cells by paired t test).

**Fig. 9.** Interaction of arrestin3 with the dopamine D<sub>2</sub> receptor modulates receptor activation of ERKs. HEK293 cells overexpressing wild type D<sub>2</sub> (WT) or D<sub>2</sub>-A4 (A4) receptors in the absence or presence of heterologously expressed arrestin3 (arr) were treated with 10  $\mu$ M dopamine for 5 min prior to quantification of activated ERKs in cell lysates. Some cells were pretreated with pertussis toxin (PTX) overnight. **A, Upper panel**, a representative immunoblot is shown for phospho-ERKs in lysates of cells without (CTL) or with (DA) dopamine treatment. **Middle panel**, an immunoblot shows similar expression levels of total ERKs. **Lower panel**, an immunoblot shows that the band optical density varies linearly with the concentration of phospho-ERKs from sample WT+arr-DA. **B**, the expression levels of arrestin3 are shown. **C**, results from

MOL #50534

four independent experiments are shown as the mean  $\pm$  S.E.M. (\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , versus the same drug/receptor combination without overexpressed arrestin, paired t test).

MOL #50534

## Tables

TABLE 1

Saturation analysis of binding of arrestin3 to receptor fragments

$K_d$  values were calculated based on the molecular weight of arrestin3 (~54 kDa).  $B_{max}$  values represent the amount of arrestin3 bound to 300 ng of GST-IC3 (~44 kDa). Each value represents the mean  $\pm$  S.E. of 3 or more independent experiments.

Fusion Protein	Arrestin3	
	$K_d$ (nM)	$B_{max}$ (ng)
GST-D <sub>2</sub> -IC3	72 $\pm$ 14	10.1 $\pm$ 0.6
GST-D <sub>3</sub> -IC3	79 $\pm$ 14	10.2 $\pm$ 0.6

MOL #50534


TABLE 2

Binding characteristics of the wild type and A4 mutant dopamine D<sub>2</sub> receptors

Affinity values for [<sup>3</sup>H]spiperone ( $K_d$ ) were determined by saturation analyses. The apparent affinity values for dopamine were determined by inhibition of the binding of [<sup>3</sup>H]spiperone, followed by nonlinear regression analysis of the competition curves. Two-site fits generated values for binding sites with high affinity ( $K_{i\text{high}}$ ) and low affinity ( $K_{i\text{low}}$ ) for dopamine. Each value represents the mean  $\pm$  S.E. of 3 or more independent experiments.

Receptor	$K_d$ [ <sup>3</sup> H]Spiperone ( <i>pM</i> )	$K_{i\text{high}}$ for dopamine ( <i>nM</i> )	$K_{i\text{low}}$ for dopamine ( $\mu\text{M}$ )	Receptor in high affinity state (%)
D <sub>2</sub> wild type	83 $\pm$ 13	9.0 $\pm$ 2.9	4.3 $\pm$ 0.5	22.1 $\pm$ 3.5
D <sub>2</sub> -A4	86 $\pm$ 4	11.2 $\pm$ 3.2	3.6 $\pm$ 0.7	22.7 $\pm$ 3.4

**Figure 1**





**Figure 2**




Figure 3

**A**



**B**



**Figure 4**

**A**

D2-IC3:  $^{211}$ KIYIVLRKRRKRVNTKRSSR $^{230}$

D2-IC3 $\Delta$ NT5 =  $\Delta$ KIYIV    D2-IC3 $\Delta$ NT3 =  $\Delta$ KIY

D2-IC3 $\Delta$ IYIV212-215    D2-IC3 $\Delta$ IYIVL212-216

D2-IC3IYIV212-215A4    D2-IC3IYIVL212-216A5

**B**




Figure 5








Figure 6






**Figure 7**

Figure 8






Figure 9

