

MOL #55897

Diphosphoinositol Polyphosphates: Metabolic Messengers?

Stephen B. Shears

Inositide Signaling Group, National Institute of Environmental Health Sciences, NIH,
DHHS, Research Triangle Park, PO Box 12233, NC 27709, USA

MOL #55897

Running title: Diphosphoinositol Polyphosphates; Metabolic Messengers?

Author for correspondence: Stephen Shears, Inositide Signaling Group, National Institute of Environmental Health Sciences, NIH, DHHS, Research Triangle Park, PO Box 12233, NC 27709, USA E-MAIL shears@niehs.nih.gov

Abbreviations: Inositol phosphates are designated according to IUPAC nomenclature, i.e., InsP₅; inositol pentakisphosphate, InsP₆; inositol hexakisphosphate, PP-InsP₄, diphosphoinositol tetrakisphosphate; PP-InsP₅, diphosphoinositol pentakisphosphate ("InsP₇"); (PP)₂-InsP₄, bis-diphosphoinositol tetrakisphosphate ("InsP₈"); AICAR, 5-aminoimidazole-4-carboxamide ribonucleoside; EPAC, exchange protein directly activated by cyclic AMP; PLC, phosphoinositide-specific phospholipase C; PPIP5K, PP-InsP₅ kinase; TNF, tumor necrosis factor; TNP, N²-(*m*-trifluoromethyl)benzyl), N⁶-(*p*-nitrobenzyl)purine; ZMP, 5-amino-4-imidazolecarboxamide riboside monophosphate.

MOL #55897

Abstract

The diphosphoinositol polyphosphates ("inositol pyrophosphates") are a specialized subgroup of the inositol phosphate signaling family. This review proposes that many of the current data concerning the metabolic turnover and biological effects of the diphosphoinositol polyphosphates are linked by a common theme, namely, that these polyphosphates act as metabolic messengers. This review will also discuss the latest proposals concerning possible molecular mechanisms of action of this intriguing class of molecules.

MOL #55897

Introduction

It was the discovery of cyclic AMP (Rall and Sutherland, 1958; Sutherland and Rall, 1958) that introduced us to the concept of a “second messenger” (Robison et al., 1968): a diffusible molecule (or ion) that, in response to an extracellular stimulus, is rapidly generated at (or released from) a particular subcellular site and then regulates particular effector proteins within the cell so as to elicit a cellular response. Of course, evolution has a remarkable tendency to repeat a good idea, so many different second messengers are now known. The inositol phosphate family represents the convergence of several “good signaling ideas”, most notably in their use of a recurring theme in the cell-signaling genre, namely, phosphate groups. Phosphates have two especially prominent features that facilitate specificity of interactions between cell signaling entities. First, the bulky nature of the phosphate group imposes geometric constraints on ligand-protein and protein-protein interactions. Second, the phosphate’s negative charge at physiological pH bestows specificity on its interactions with target proteins through multiple ionic and hydrogen bonds. The negative charges on the phosphate group also make soluble, phosphorylated molecules lipid-impermeant, so that they can be retained inside cells.

Inositol offers several additional assets for a signaling molecule. It is chemically stable, it is small (hence it diffuses through cytosol quickly), and it is only a short synthetic offshoot from the glycolytic pathway (Sherman et al., 1977). There is also a functionally significant plane of symmetry across the 2/5-axis of the inositol ring (Fig. 1). That symmetry permits one inositol phosphate to imitate another’s three-dimensional phosphate recognition pattern, when the orientation of the inositol ring changes in relation to the protein’s ligand-binding site (Wilcox et al., 1994), although in addition the binding site itself has to be somewhat flexible (Ho et al., 2002). This phenomenon

MOL #55897

provides a molecular explanation for the metabolic promiscuity of certain inositol phosphate kinases (see (Shears, 2004) for a review) and it also facilitates functionally important metabolic interactions between different inositol phosphates (Chamberlain et al., 2007). With all of these advantageous features, it is not surprising that the inositol phosphates are a phylogenetically-widespread family of second messengers.

There are also a large number of different inositol phosphates, thereby creating opportunities for substantial structural and functional diversity within this class molecules. A mere handful of these molecules is shown in Fig. 2. In fact, the inositol ring -- “a six-bit signaling scaffold” (York, 2006) – can, theoretically, host 63 different combinations of one-to-six monoester phosphate groups; more than half of these possible molecules are now acknowledged to exist in Nature (Majerus et al., 2008; Irvine and Schell, 2001; Michell, 2008). Yet, even that understates their abundance: in the early 1990s it became clear that the presupposed upper limit of six phosphates per inositol ring was going to be exceeded, following reports that cells contain inositol-based molecules that are even more polar than InsP_6 (Oliver et al., 1992; Stephens et al., 1991; Wong et al., 1992; Europe-Finner et al., 1991). Two groups working independently (Stephens et al., 1993; Menniti et al., 1993) determined that these molecules are diphosphorylated derivatives of InsP_5 and InsP_6 (Fig. 3). More recently, evidence has arisen of diphosphorylated derivatives of InsP_3 and InsP_4 (Seeds et al., 2005). The threat that cells might also synthesize triphosphates of InsP_6 (Shears, 2004; Draskovic et al., 2008) foreshadows a further potential expansion of this family.

The initial discovery of diphosphorylated inositol phosphates immediately generated an anticipation that these molecules have some special functional significance (Stephens et al., 1993; Menniti et al., 1993). This argument goes well beyond the obvious “cells

MOL #55897

would not make them unless they are important” category. Instead, it was noted that these molecules undergo exceptionally high rates of metabolic turnover through coupled kinase/phosphatase substrate cycles (Menniti et al., 1993;Stephens et al., 1993). The investment of cellular energy in the maintenance of these cycles must be quite substantial and, presumably, biologically significant. Moreover, substrate cycles are themselves tell-tale signs of regulatory steps in a metabolic pathway (Hers and Hue, 1983), from which we can infer that the levels of the substrates and/or products must be worth regulating. The diphosphoinositol polyphosphates are also molecules with severe electrostatic and steric congestion. The relief of these constraints following hydrolysis of the diphosphate groups has long-been viewed as contributing to a “high energy” reaction that ought to have biological significance. Hence, for example, the origin of the idea that diphosphoinositol polyphosphates might transfer phosphate to proteins (Stephens et al., 1993;Voglmaier et al., 1996;Hand and Honek, 2007;Laussmann et al., 1996).

Nevertheless, despite their pedigree, the diphosphoinositol polyphosphates have been slow to achieve the status of being widely-accepted intracellular signals (Burton et al., 2009). One significant problem has been that their global intracellular concentrations do not change in response to the activation, by any naturally-occurring extracellular agent, of a defined signal transduction pathway. This factor has been a major impediment to our attempts to place in a cell-signaling context the diversity of proposed cellular activities that have been attributed to these compounds. A major goal of this review is to assimilate evidence that points to a resolution of this problem. The key is not to consider "second messengers" as just being mediators of extracellular stimuli, as originally envisaged (Robison et al., 1968). There is little evidence that diphosphoinositol polyphosphates fulfil this requirement. Instead, we need to look at other signaling systems that respond to biochemical information that is generated from

MOL #55897

within the cell, rather than from outside of it. An example from within this genre are the “metabolic messengers” (Rolland et al., 2001). These particular entities are bio-sensors that first detect variations in energy balance and subsequently communicate this information to other cellular networks, which then initiate adaptive responses. This ensures the maintenance of tight energy homeostasis, which is a fundamental necessity for cell survival (Hardie, 2004). Some recent developments ((Bennett et al., 2006; Lee et al., 2007; Choi et al., 2008), for example) have led to the suggestion that diphosphoinositol polyphosphates might serve this role of coupling signaling pathways to the energetic status of the cell. This evidence is discussed here. I will also expand this new concept, by arguing that it provides a common theme that rationalizes many of the apparently disparate functions of the diphosphoinositol polyphosphates. This review will also discuss the latest proposals concerning possible molecular mechanisms of action of this intriguing class of molecules.

Nomenclature

Some comments on nomenclature are warranted. This review takes an approach that might at first seem both unwieldy and unnecessarily pedantic, but it is hoped that the reader will ultimately appreciate that the field will benefit from the use of terminology that is unambiguous. Some of the current concerns with nomenclature originate from the initial discovery of these compounds (Stephens et al., 1993; Menniti et al., 1993). Stephens and colleagues (Stephens et al., 1993) described these molecules as “diphosphoinositol polyphosphates”, which is the IUPAC-preferred terminology (IUPAC-IUB Commission on Biochemical Nomenclature, 1977). We, however, classified these molecules as “inositol pyrophosphate polyphosphates” - *mea culpa* - and subsequently it has become common practice to abbreviate this too, simply, “inositol pyrophosphates” (e.g. (Saiardi et al., 2002)). The reason that IUPAC frowns on this vernacular is because

MOL #55897

“pyrophosphate” is a term that was originally introduced to define a diphosphate that was produced non-enzymatically, by heating (the prefix “pyro-“ is derived from Greek, meaning “fire” in this context). Another, perhaps less finicky argument against using “inositol pyrophosphate” as a designation is that it is inadequate for describing a molecule that contains *both* phosphates and diphosphates. Thus, in this review, “diphosphoinositol polyphosphates” is the preferred nomenclature. However, the thesis in this review that these molecules are metabolic messengers applies, for now, only to the canonical members of this family: PP-InsP₄, PP-InsP₅ and (PP)₂-InsP₄ (Fig. 3). Some caution is warranted as we await further evidence of the significance of (PP)₂-InsP₃ (Ingram et al., 2003; Caffrey et al., 2000; Draskovic et al., 2008), and the diphosphorylated derivatives of InsP₃ and InsP₄ (Seeds et al., 2005). IPMK (see Fig. 3) can also add a diphosphate to Ins(1,3,4,5,6)P₅; the position of the diphosphate is not known, and it remains unclear if this reaction has any biological importance (Zhang et al., 2001; Saiardi et al., 2001a).

There is also the issue of triphosphates. Some time ago, following prolonged incubation of InsP₆ with an InsP₆ kinase, we noticed that PP-InsP₅ was not the only product. In addition, small amounts (5-10%) of more polar material were observed (Saiardi et al., 2000). We (Saiardi et al., 2000) initially proposed that this novel molecule was an isomer of (PP)₂-InsP₄; only later (Shears, 2004) was the possibility raised that it might instead be a PPP-InsP₅, i.e. a triphosphate. This alternate option was shown to be correct, following the publication of an impressive analytical study (Draskovic et al., 2008). As if this subject were not complicated enough, PPP-InsP₅ can be even further phosphorylated, perhaps to PPPP-InsP₅ (Draskovic et al., 2008). However, these are all data acquired from experiments that were performed *in vitro*. So, it has to be asked,

MOL #55897

do these more esoteric tri- and tetra-phosphate derivatives occur *in vivo*, or are they merely by-products of the availability of excessive quantities of recombinant enzyme and over-indulgent incubation times *in vitro*? This question has not yet been answered satisfactorily. In part this may reflect technical issues; we have found that we cannot quantitatively recover PPP-InsP₅ from a strong anion exchange HPLC column using our standard elution conditions (Shears, unpublished data). It is possible, then, that this material is being lost from our HPLC assays of cell extracts. Nevertheless, some [³H]inositol-labeled molecules of approximately the appropriate polarity have been observed upon HPLC analysis of a strain of *S. cerevisiae* lacking its endogenous InsP₆ kinase but transformed with a human InsP₆ kinase gene (Draskovic et al., 2008). It still needs to be directly determined if these compounds are actually tri- and tetra-phosphates, as opposed to some of the multiply diphosphorylated derivatives that the InsP₆ kinase can also synthesize, from InsP₅ for example (Draskovic et al., 2008). More importantly, we must also be cautious in attributing biological relevance to an experiment involving supra-physiological, heterologous gene expression, and the deletion of an endogenous gene, InsP₆ kinase, that has many phenotypic consequences (see below).

To return to the subject of decrees from IUPAC, in this review I will agree with their contention that "Ins" is a more intuitive abbreviation for inositol than is "I" (IUPAC-IUB Commission on Biochemical Nomenclature, 1969), although both are widely deployed elsewhere (that is, inositol hexakisphosphate is written here as "InsP₆" rather than "IP₆"). Another point to emphasize is that the diphosphorylated derivatives of InsP₆, namely, PP-InsP₅ and (PP)₂-InsP₄, are frequently abbreviated in the literature to "IP₇" and "IP₈", respectively. This terminology is avoided here because it, too, can introduce

MOL #55897

confusion. For example, the aforementioned PPP-InsP₅ contains eight phosphates, but is clearly different from (PP)₂-InsP₄ (Fig. 3). Furthermore, one might argue that it is the inherent inadequacies of "IP₇" as a nomenclature that have led it occasionally - and incorrectly - to be defined as either a "heptaphosphate" (Mishra and Bhalla, 2002) (i.e. an entirely mythical beast endowed with a chain of seven phosphates) or a "heptakisphosphate" (Lee et al., 2007) (which would require seven phosphates to each be individually attached to a carbon - impossible of course because inositol has only six carbons). The usage of "IP₇/IP₈" should also be avoided because it can obscure the fact that InsP₆ is not the only precursor for this class of compounds; PP-InsP₄ is a member of this family, but it is synthesized from InsP₅ (Fig. 3).

Another issue to be cognizant of are the different names that have been used at various times to describe the enzymes that metabolize the diphosphoinositol polyphosphates (Table 1). For example, although the phosphatases that degrade these molecules are known in the field as DIPPs ("Diphosphoinositol Polyphosphate Phosphohydrolases" (Safrany et al., 1998) the Human Genome Organization Gene Nomenclature Committee (HGNC) has allocated them to the NUDT family (NUDT = "Nudix [Nucleoside Diphosphate attached moiety 'x']-Type motif), in honor of their eponymous catalytic site (G_x₅E_x₅[UA]_xR_E_x₂E_E_xGU, or similar, in which U is an aliphatic, hydrophobic residue (McLennan, 2006)). The yeast gene is named Ddp1.

The HGNC originally proposed that "*IHPK*" (for "inositol hexakisphosphate kinase") be used to designate the three members of the human gene family that phosphorylate InsP₆ and InsP₅ to 5-PP-InsP₅ and 5-PP-InsP₄ respectively (Fig. 3). However, in deference to more common usage, the HGNC has now adopted "*IP6K*" as the name for this gene family (Table 1). The orthologous yeast gene is *Kcs1* (Saiardi et al., 1999).

MOL #55897

The mammalian IP6Ks -- there are three isoforms -- have a molecular weight of 46 to 49 kDa (Saiardi et al., 2001b; Saiardi et al., 1999; Schell et al., 1999). These kinases possess a PxxxDxKxG catalytic domain, and a remote, catalytically-essential SSL (or similar) tetrapeptide. Some homologues of these kinases are rather larger. Kcs1 from *S. cerevisiae* is a 120 kDa protein (Saiardi et al., 1999). It contains leucine zippers near its C-terminus which do not contribute to catalytic activity but instead are important for cell wall integrity and vacuolar biogenesis (Dubois et al., 2002). Thus, Kcs1 has other functions, and so its deletion from cells can give misleading information concerning the roles of diphosphoinositol polyphosphates.

There are separate, unrelated kinases which further phosphorylate 5-PP-InsP₅ to (PP)₂-InsP₄ (Shears et al., 1995). These enzymes are around 120-160 kDa in size, and, unlike the IP6Ks, they utilize an ATP-grasp catalytic domain to support kinase activity. These enzymes can also phosphorylate InsP₆ (Mulugu et al., 2007; Choi et al., 2007; Fridy et al., 2007), but they are not typically called "InsP₆ kinases", which conveniently avoids confusion with the canonical IP6K family. A recent study unequivocally determined that these particular enzymes phosphorylate the 1/3-position of the inositol ring (Lin et al., 2009). The latter work corrected an earlier tentative proposal (Mulugu et al., 2007) that the enzyme is a 4/6-kinase. York's laboratory (Mulugu et al., 2007) was the first to describe the identify of this gene, in a study with *S. cerevisiae*, so this group retained the name (*Vip1*) that it had already been christened with, well before it was known to encode a kinase. While the latter study was in progress, our laboratory independently purified and sequenced an PP-InsP₅ kinase from rat brain, and we also identified the two human genes (Choi et al., 2007). York's group have also described the mammalian homologues (Fridy et al., 2007). These proteins were already listed in the genome database as HISPPD2A and HISPPD1

MOL #55897

because they contain an acid phosphatase-like domain. The HGNC will not accept “VIP” as an alternative name for these genes, because that has already been allocated to a different human gene. We have proposed that they be known as PPIP5K1 and PPIP5K2 (Choi et al., 2007). This decision reflects on our conclusion that the catalytic preference of this kinase *in vitro* - in mammals at least - is to phosphorylate 5-PP-InsP₅ rather than InsP₆ (Choi et al., 2007). A recent pharmacological study is also consistent with 5-PP-InsP₅ being the main substrate *in vivo* (again, in mammalian cells) (Padmanabhan et al., 2009). Additionally, when we over-expressed a kinase-dead mutant version of PPIP5K1 in HEK cells, we did not see any decrease in the cellular levels of any inositol phosphate (Choi et al., 2007), which argues against the phosphatase domain having significant hydrolytic activity against inositol phosphatases *in vivo*.

Despite the substrate preference of the human enzyme that is explicitly implied by the PPIP5K1/2 nomenclature, it should be noted that the phosphorylation of InsP₆ to 1/3-PP-InsP₅ is in itself functionally significant, at least in yeasts and plants. For example, in *S. cerevisiae* 1/3-PP-InsP₅ regulates cyclin kinase activity (Lee et al., 2008; Lee et al., 2007). *Arabidopsis* appears to encode homologues of Vip1/PPIP5K in its genome, but not homologues of IP6K. Thus, Vip/PPIP5K may be the *only* source of PP-InsP₅ in some organisms.

Incidentally, the uncertainty over whether the diphosphate in this isomer of PP-InsP₅ is positioned at either the 1 or 3 position (hence the “1/3” designation, see Fig. 3) reflects an analytical impediment created by the axis of symmetry in the inositol ring (see Fig. 1). Thus, 1-PP-InsP₅ and 3-PP-InsP₅ are an enantiomeric pair, so a stereoselective

MOL #55897

technique is required to distinguish between them.

Cellular Levels, Metabolic Turnover and the Issue of Compartmentalization.

If the diphosphoinositol polyphosphates are to act as cellular signals, then the levels of one or more of this family of molecules should respond to some defined intra- or extra-cellular stimulus. The slime-molds, and in particular *Dictyostelium* (Watts, 1984), provide a singularly dramatic example of such a phenomenon. The bacteria-munching, amoeboid form of *Dictyostelium* contains around 10 μM of both PP-InsP₅ and (PP)₂-InsP₄ (Laussmann et al., 2000). When the cells are starved of their bacteria food supply, the amoeboid cells aggregate, the first of a series of morphogenetic stages that culminate in the formation of a fruiting body (Watts, 1984). During the initial aggregation process, the levels of PP-InsP₅ and (PP)₂-InsP₄ each increase to 100 and 250 μM , respectively (Laussmann et al., 2000; Luo et al., 2003). Perhaps slime molds use these exceptionally high levels of diphosphoinositol polyphosphates as “high-energy” phosphate donors. However, these organisms are not a representative eukaryotic cell model for studying the turnover and biological functions of the diphosphoinositol polyphosphates. For a start, slime molds synthesize both 6-PP-InsP₅ and 5,6-(PP)₂-InsP₄ (Laussmann et al., 1997; Laussmann et al., 1996), which are not found in yeast and mammalian cells, which instead predominantly contain 1/3-PP-InsP₅, 5-PP-InsP₅, and 1/3,5-[PP]₂-InsP₄ (Lin et al., 2009; Albert et al., 1997). Additionally, there are much lower concentrations of diphosphoinositol polyphosphates in yeast and mammalian cells. Levels of PP-InsP₅ usually lie in the 1 to 5 μM range (Ingram et al., 2003; Fisher et al., 2002; Barker et al., 2004; Illies et al., 2007; Bennett et al., 2006). The levels of PP-InsP₄ and (PP)₂-InsP₄ are even lower in yeast and mammalian cells, each only about

MOL #55897

10-20% of those of PP-InsP₅ (Glennon and Shears, 1993;Choi et al., 2008;Choi et al., 2005).

The fact that most eukaryotic cells normally contain low levels of diphosphoinositol polyphosphates does not preclude these molecules from being cellular signals. Similar concentrations are found for other bio-active inositol phosphates, such as Ins(1,4,5)P₃ (Streb et al., 1983) and Ins(1,3,4,5)P₄ (Huang et al., 2007b). The issue is that such low levels add an additional layer of difficulty to the technical challenges involved in accurately measuring stimulus-dependent changes in the turnover of the diphosphoinositol polyphosphates. Currently, diphosphoinositol polyphosphates can only be quantified following their individual separation by HPLC analysis of cell extracts. There is a non-radioactive, automated, in-line HPLC assay for inositol phosphates that relies on a post-column formation of a metal-dye complex (Mayr, 1988); this technique has been used to assay PP-InsP₅ in mammalian cells (Albert et al., 1997), but it is not quite sensitive enough to record (PP)₂-InsP₄ turnover (Lin et al., 2009). Thus, most groups assay cellular turnover of the diphosphoinositol polyphosphates after they are first radiolabeled, by incubating cells with [³H]inositol until isotopic equilibrium is reached (Shears, 1997). Unfortunately, this procedure takes several days (Shears, 1997). Moreover, unless appropriate precautions are taken, the diphosphoinositol polyphosphates may not be quantitatively recovered from cell lysates, which obviously affects the data that are obtained. For example, we add 1 mM EDTA to the media used to quench and extract the cells, and also to the HPLC elution buffers, so as to chelate trace quantities of trivalent cations that otherwise can cause the diphosphoinositol polyphosphates to precipitate out of solution and be lost from the analysis. When counting the radioactivity in the HPLC eluate, some laboratories routinely use in-line scintillation counters (e.g., see (Fridy et al., 2007)), but our experience has been that

MOL #55897

the short count-times that are an inherent aspect of this equipment is a hindrance to the accurate assessment of the especially low cellular levels of $(PP)_2$ - $[^3H]$ InsP₄. We are also concerned that a homogeneous fluid phase may not be achieved immediately upon in-line mixing of the scintillation fluid with the high-salt HPLC buffers, which could decrease counting efficiency around the point in the gradient where $(PP)_2$ - $[^3H]$ InsP₄ is normally eluted. Thus, we (Safrany and Shears, 1998) and others (Azevedo and Saiardi, 2006), believe that it is more accurate to direct the HPLC eluate to a fraction collector, manually add scintillant to each fraction, mix vigorously, and then use a traditional counter. Anyone who doubts the value of our precautions should note how difficult it was to detect changes in PP-InsP₅ and $(PP)_2$ -InsP₄ levels when an in-line counter was used to assess the effects of overexpression of full-length human PP-InsP₅ kinase in HEK cells (Fridy et al., 2007). In contrast, in similar experiments with the same cell line, we (Choi et al., 2007) collected and analyzed individual fractions of HPLC eluate and we reported that $(PP)_2$ - $[^3H]$ InsP₄ levels increased 10-fold following overexpression of the PP-InsP₅ kinase. This point is emphasized in order to illustrate how varying interpretations of the biological function of an enzyme could result from practical differences in the analytical procedures that are used.

Another approach that can assist the analysis of turnover of diphosphoinositol polyphosphates in an intact mammalian model system is to use the DDT₁-MF₂ hamster vas deferens smooth muscle cell-line (Safrany and Shears, 1998) because these cells contain about tenfold higher levels of diphosphoinositol polyphosphates than are present in other mammalian cells. Others (Fridy et al., 2007; Otto et al., 2007) have successfully used genetic approaches to elevate the cellular levels of diphosphoinositol polyphosphates and thereby facilitate the analysis of their cellular turnover; this group achieved that goal by overexpression of both InsP₅ kinase and a G-protein (G α_q) that

MOL #55897

activates PLC.

As noted above, and in Fig. 3, two classes of enzymes (IP6K and VIP/PPIP5K) cooperate to provide two routes by which $(PP)_2\text{-InsP}_4$ can be synthesized from InsP_6 in yeast and mammalian cells. These two pathways use either 5-PP- InsP_5 or 1/3-PP- InsP_5 as an intermediate (Fig. 3). Which of these is quantitatively the most important route? This is not an easy question to answer with certainty. HPLC analyses (Albert et al., 1997) indicate that 5-PP- InsP_5 is the major PP- InsP_5 isomer to accumulate in mammalian cells. The situation is probably the same in yeast, since deletion of *Kcs1*, which synthesizes 5-PP- InsP_5 (Draskovic et al., 2008), is associated with >80% reduction in PP- InsP_5 levels (Saiardi et al., 2000). Therefore, even though most of the previously published assays of PP- InsP_5 in intact yeast and mammalian cells did not distinguish which isomer was being studied, we can safely assume it was predominantly 5-PP- InsP_5 . However, the steady-state levels of a metabolic intermediate are not necessarily reflective of its rate of metabolism. Additionally, the nature of the PP- InsP_5 isomers that accumulate in cells reflects upon both $(PP)_2\text{-InsP}_4$ synthesis *and* its metabolism. For example, when presented with 1/3,5- $(PP)_2\text{-InsP}_4$ as a substrate, DIPPs prefer to hydrolyze the diphosphate that is added by PPIP5K (Shears et al., 1995), which we now know to be the 1/3-diphosphate (Lin et al., 2009). A positional preference of DIPPs for one of the two diphosphate groups will clearly influence which PP- InsP_5 isomers are formed following $(PP)_2\text{-InsP}_4$ dephosphorylation (see Fig. 3). This phenomenon may help explain why the activities of the DIPPs can differentially “mask” the rates of flux through the two different PP- InsP_5 isomers during $(PP)_2\text{-InsP}_4$ synthesis (Padmanabhan et al., 2009). Nevertheless, there are two good reasons for

MOL #55897

proposing that the major pathway of $(PP)_2$ -InsP₄ synthesis goes through 5-PP-InsP₅ in mammalian cells. First, kinetic data inform us that, *in vitro*, the human VIP/PPIP5Ks prefer to phosphorylate 5-PP-InsP₅ over InsP₆ (Choi et al., 2007). Second, there are pharmacological data that support this proposal: TNP, or N²-(*m*-trifluoromethyl)benzyl), N⁶-(*p*-nitrobenzyl)purine, has been used as a cell-permeant and selective inhibitor of IP6Ks without affecting the VIP/PPIP5K enzymes (Padmanabhan et al., 2009). The addition of TNP to either HeLa cells or to *S. cerevisiae* lowered the cellular levels of both $(PP)_2$ -InsP₄ and PP-InsP₅ (Padmanabhan et al., 2009). These data are consistent with 5-PP-InsP₅ (synthesized by the TNP-sensitive IP6Ks) being the most important precursor for $(PP)_2$ -InsP₄ (see Fig. 3). At least in “resting” cells. The situation may be different after one or more of the enzymes is either down- or up-regulated, such as occurs after hyperosmotic stress (Pesesse et al., 2004), for example.

The DIPPs are very active enzymes: four mammalian genes have been cloned, all of which encode proteins of around 20 kDa in size: type 1 (Safrany et al., 1998; Chu et al., 2004), types 2 α /2 β (Caffrey et al., 2000; Hua et al., 2001) and types 3 α /3 β (Leslie et al., 2002; Hidaka et al., 2002; Hua et al., 2003). The specificity constants (k_{cat}/K_m) for these enzymes range in value from 2×10^5 to $5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$, the latter being close to the limit for diffusion-controlled encounter between enzyme and substrate (Fersht, 1985). With DIPP activities such as these, we should perhaps wonder why the cellular concentrations of diphosphoinositol polyphosphates are actually as *high* as they are! This high metabolic turnover of the diphosphoinositol polyphosphates becomes evident when cells are incubated with 1 mM fluoride, which inhibits DIPP activities (Menniti et al., 1993; Safrany and Shears, 1998). This pharmacological maneuver causes rapid, many-fold increases in the levels of diphosphoinositol polyphosphates (Menniti et al., 1993; Safrany and Shears, 1998). In the absence of fluoride, high ongoing metabolism

MOL #55897

may impede our ability to detect subtle changes in its rate, especially if metabolic regulation were to be compartmentalized to certain regions of the cell. However, fluoride is of limited practical use as a pharmacological tool, because it is so non-specific. It is a well-known protein phosphatase inhibitor and, by perturbing the activities of numerous enzymes regulated by phosphorylation/dephosphorylation cycles, fluoride affects many signaling processes.

Since the DIPPs are so active, their acute spatiotemporal regulation could greatly impact upon the turnover of the diphosphoinositol polyphosphates. However, there is as yet no evidence that this is exploited as a short-term signaling mechanism; no covalent modification of DIPPs has been found (Safrany et al., 1998). Furthermore, studies with GFP-DIPP constructs offer no evidence of their compartmentalization; the fusion proteins were uniformly distributed throughout the cell ((Leslie et al., 2002); Choi, K. and Shears, S.B., unpublished data). As for the intracellular distribution of *endogenous* DIPPs, we do not currently have antibodies with sufficient sensitivity for such studies.

This current lack of evidence that DIPPS might be compartmentalized impacts upon the idea, noted later in this review and elsewhere (Burton et al., 2009), that a particular action of a diphosphoinositol polyphosphate could be facilitated if there were to be a localized elevation in its concentration. So as to explain the latter point, we can turn to the example shown by another intracellular signal, cAMP. There seems little doubt that the non-uniform distribution of cAMP through the cell (i.e. its compartmentalization) is strictly dependent upon the physical separation of the “on-switch” (adenylyl cyclase) from the “off-switch” (cAMP phosphodiesterase) (Chen et al., 2008). By analogy, it is arguable that there is little possibility of there being locally elevated concentrations of diphosphoinositol polyphosphates, if the highly-active DIPPs are uniformly distributed

MOL #55897

throughout the cell. This last observation also makes it unclear whether a significant localized accumulation of any diphosphoinositol polyphosphates can arise from IP6K2 being compartmentalized in the nucleus (as indicated upon over-expression of a GFP-IP6K2 fusion construct; (Saiardi et al., 2001b)). On the other hand, the VIP/PP1P5Ks are excluded from the nucleus (Choi et al., 2007; Fridy et al., 2007) indicating that the products of these particular kinases are likely present in lower amounts in the nucleus compared to the cytoplasm.

One possibility that has yet to be addressed is whether there might be longer-term regulation of the levels of diphosphoinositol polyphosphates through regulated changes in the degree of *DIPP* expression. For example, there are several DIPP2 mRNA transcripts, which utilize multiple and non-canonical polyadenylation signals (Caffrey et al., 2000). This situation is often an indicator of cells utilizing translational control mechanisms in order to regulate gene expression (Edwards-Gilbert et al., 1997). The exceptionally high G/C content of the 5'-UTRs of the DIPP1 and DIPP2 mRNAs (Caffrey et al., 2000; Safrany et al., 1998) may also influence mRNA stability (Kozak, 1996) and hence gene expression. Incidentally, the expression of DIPP2 α and DIPP2 β from a single gene relies upon an unusual mode of alternate splicing that we christened "intron boundary skidding" (Caffrey and Shears, 2001). Finally, the expression of two *DIPP3* genes (Hua et al., 2003) is of some interest because it appears to represent rare evidence in support of the concept (Force et al., 1999) of subfunctionalization following gene duplication.

Another factor that might contribute to the cellular turnover of the diphosphoinositol polyphosphates is their ability to directly phosphorylate proteins (Bhandari et al., 2007; Saiardi et al., 2004). However, current evidence indicates that there is a relatively slow-acting "off-switch" for this process, namely, phosphatase-directed cleavage of the

MOL #55897

phosphorylated protein (Bhandari et al., 2008; Burton et al., 2009). It has been suggested that this is a manifestation of a long-lived signaling process (Burton et al., 2009). If this is the case, it would greatly limit the availability of protein substrate for phosphorylation, so that the rate of consumption of diphosphoinositol polyphosphates for this process might not be significant in relation to their ongoing high rate of metabolism by the DIPPs.

The Relationship Between Cellular Bioenergetic Status and Cellular Levels of Diphosphoinositol Polyphosphates.

As mentioned above, some cellular signals do not respond exclusively to extrinsic agents. Instead, there are signaling activities that originate from within the cell. For example, cells utilize bio-sensors to detect variations in energy balance and subsequently communicate this information to other cellular networks which then initiate adaptive responses. One of the most sensitive indicators of bioenergetic balance is the AMP-activated protein kinase (AMPK), which is activated by an elevated [AMP]/[ATP] ratio (Hardie, 2004). Increased [AMP] renders AMPK susceptible to activation by phosphorylation, whereupon AMPK quickly inhibits ATP-consuming anabolic processes (protein synthesis, gluconeogenesis, fatty acid synthesis) and simultaneously activates ATP-generating, catabolic pathways (glycolysis, fatty acid oxidation) (Hardie, 2004). In this way, AMPK acts to restore an appropriate cellular "energy balance".

Phosphorylation and activation of AMPK are largely driven by the kinases's ultrasensitivity to just small increases in [AMP] (Hardie, 2004), which in turn are considered to represent a state of "mild energy depletion" (Inoki et al., 2003). This terminology usefully distinguishes this situation from the more severe energy stress that arises following a substantial decrease in cellular ATP levels (Inoki et al., 2003; Browne and Proud, 2002). The latter would typically only be expected to occur under rather extreme conditions such as during muscle fatigue, for example, or during an acute

MOL #55897

stress such as ischaemia (Hochachka, 1999). These more significant decreases in [ATP] also prompt additional stress sentinels to become activated (Zhuo et al., 2005).

There is a pharmacological tool for studying the actions of cellular sentinels that react to increases in [AMP]: 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR). Upon its uptake into cells, AICAR is phosphorylated to 5-amino-4-imidazolecarboxamide riboside monophosphate (ZMP), an AMP-mimetic which, for example, activates AMPK (Merrill et al., 1997; Sabina et al., 1985). It should be emphasized that this experimental approach is designed to invoke relatively mild energy stress, that is, an increase in [AMP], without there being a significant effect upon [ATP]. We (Choi et al., 2008) have recently reported that the treatment of cells with AICAR causes a reduction in the cellular levels of $(PP)_2$ -InsP₄. This effect was most easily detected in experiments with the DDT₁-MF₂ smooth muscle cell line (Choi et al., 2008), in which “resting” levels of $(PP)_2$ -InsP₄ are unusually high (Safrany and Shears, 1998). In other cell types the levels of $(PP)_2$ -InsP₄ are often too low to accurately assess inhibition of its synthesis. In these other cell types, this inhibitory effect of AICAR was most readily observed when the drug was used to antagonize the several-fold elevation in $(PP)_2$ -InsP₄ levels that normally accompanies hyperosmotic stress (Pesesse et al., 2004; Choi et al., 2008). Incidentally, this osmotic response is an area of study that could benefit from further research, as we do not yet understand the mechanisms that are involved, nor the biological significance. We have previously argued that the degree of osmotic stress that activates $(PP)_2$ -InsP₄ synthesis lies within the biologically relevant range of conditions to which cells are normally exposed (Yang et al., 2008).

Unexpectedly, we have found that the inhibition of $(PP)_2$ -InsP₄ synthesis following the incubation of cells with AICAR is independent of its canonical target, AMPK (Choi et al.,

MOL #55897

2008). So, clearly, there must be other proteins that can respond to the changes in [AMP] that AICAR - through its conversion to ZMP - is considered to mimic. If we are to look for candidates, we should consider that the AMP-binding cystathionine- β -synthase (CBS) module that is present in AMPK also occurs in a large range of diverse proteins, including ATP-binding cassette (ABC) transporters, voltage-gated chloride channels and transporters, a variety of other transporter families, and a number of enzymes (Biemans-Oldehinkel et al., 2006). We (Choi et al., 2008) have also perturbed [AMP] by incubating cells with the mitochondrial poison oligomycin, but in the presence of glucose to sustain glycolytic ATP synthesis. Again, the relatively mild energy-stress that ensued was associated with a decline in $(PP)_2$ -InsP₄ synthesis (Choi et al., 2008). These observations led us (Choi et al., 2008) to propose that the levels and hence the signaling strength of $(PP)_2$ -InsP₄ might provide a novel link between signaling and bioenergetic networks. In other words, $(PP)_2$ -InsP₄ was hypothesized to be a metabolic messenger.

We have not yet been able to determine the mechanism by which increases in [AMP] inhibits $(PP)_2$ -InsP₄ synthesis. It seems likely that the activities of the VIP/PPIP5K proteins may be regulated, and now that these enzymes have been cloned protein (Mulugu et al., 2007; Fridy et al., 2007; Choi et al., 2007), it is hoped that mechanistic information will shortly be forthcoming. The regulation of VIP/PPIP5K activity may not only involve the kinase domain by itself, which only accounts for the N-terminal one-third of the entire protein (Mulugu et al., 2007; Fridy et al., 2007; Choi et al., 2007). In addition, the VIP/PPIP5Ks also possess an apparently catalytically-inactive phosphatase domain followed by long C-termini (Mulugu et al., 2007; Fridy et al., 2007; Choi et al., 2007). Perhaps these are regulatory domains? Since regulation of protein function by phosphorylation is such a widespread regulatory process, recent

MOL #55897

advances in phosphoproteomic methodology (Blackburn and Goshe, 2009) may help us gain insight into the regulation of VIP/PPIP5K activity. For example, such techniques have already demonstrated that EGF treatment of HeLa cells increases the degree of phosphorylation of Ser475 in the phosphatase domain of PPIP5K1 (Olsen et al., 2006).

The cellular levels of $(PP)_2\text{-InsP}_4$ are reduced following cell treatment with either of three MEK inhibitors: U0126, PD98059 or PD184352 (Choi et al., 2008;Choi et al., 2005;Pesesse et al., 2004). It is important to note that this inhibition of $(PP)_2\text{-InsP}_4$ synthesis is not a result of the MEK/ERK cascade being blocked. We can make this assertion because the efficient “knock-down” of either MEK or ERK using RNA interference had no impact upon $(PP)_2\text{-InsP}_4$ levels (Choi et al., 2008). Instead, it has turned out that all three of the MEK inhibitors have a puzzling “off-target” effect of somehow impairing cellular energy homeostasis (Dokladda et al., 2005;Choi et al., 2008). Again, these effects upon energetic status are relatively mild, with the MEK inhibitors primarily elevating [AMP] and not altering [ATP]. While we cannot yet explain this phenomenon, it does provide additional pharmacological evidence of the close association between $(PP)_2\text{-InsP}_4$ levels and cellular energy status (Choi et al., 2008). This is also a remarkable turn of events for three such structurally dissimilar drugs that, until recently, were considered to be quite specific MEK inhibitors (Bain et al., 2007;Dokladda et al., 2005).

This idea that the cellular levels of the diphosphoinositol polyphosphates might be dependent upon the metabolic health of the cell can help us view in a new light an otherwise puzzling observation: the inhibition of cellular synthesis of $(PP)_2\text{-InsP}_4$ following elevations in intracellular [cAMP] (Safrany and Shears, 1998). By using several commercially available cAMP analogues as pharmacological tools, we have

MOL #55897

excluded both PKA (Safrany and Shears, 1998) and EPAC (Hidaka, K., Choi, K. and Shears, S.B., unpublished data) from mediating the inhibition of $(PP)_2$ -InsP₄ synthesis by cAMP. We are still searching for another mechanistic explanation. Perhaps $(PP)_2$ -InsP₄ metabolism is controlled by a metabolic sensor that poorly discriminates between cAMP and AMP? This sensor, if it were to exist, could not be AMPK, because as noted above, this particular kinase does not affect $(PP)_2$ -InsP₄ synthesis. In any case, it is known that cAMP cannot activate AMPK directly (Henin et al., 1996; Carling et al., 1989). Despite that, increases in cellular cAMP do lead to an activation of AMPK in adipocytes (Daval et al., 2005; Yin et al., 2003). We have found a similar effect to occur in DDT₁-MF₂ cells (Choi, K. And Shears, S.B., unpublished data). This puzzling but reproducible phenomenon has apparently received little subsequent attention. Is it possible that a sustained elevation in cAMP might somehow stress the energy status of the cell? For example, others (Epperson et al., 2005) have hypothesized that, in those cells that have high levels of cAMP generation and metabolism, this might generate sufficient AMP to activate a sensor of energy-stress; Epperson *et al* (2005) of course thought that this sensor would be AMPK, but we could speculate further that maybe $(PP)_2$ -InsP₄ synthesis might also be regulated by some other entity that reacts to the AMP that is inevitably produced from cAMP. In any case, these are reasons to suggest that the response of $(PP)_2$ -InsP₄ levels to cAMP reflects the roles of diphosphoinositol polyphosphates as metabolic signals.

While an increased [AMP] may be considered symptomatic of a relatively mild bioenergetic challenge, a decline in [ATP] places a considerably greater metabolic stress upon cells. A number of energy-homeostasis processes respond to changes in cellular ATP levels (for example, see (Zhuo et al., 2005) and (Ruiz-Stewart et al., 2004)). Recent work in our laboratory suggests that this list should now include

MOL #55897

diphosphoinositol polyphosphates: we (Choi, K. and Shears, S.B., unpublished data) have found that an oligomycin-induced, 15% decrease in cellular [ATP] elicits a 50% drop in (PP)₂-InsP₄ levels in the DDT₁ MF-2 smooth muscle cell-line (we do not yet understand the mechanism underlying this effect). Additionally, in the same experiments, there was also a 15% decrease in PP-InsP₅ levels (Choi, K. and Shears, S.B., unpublished data; as discussed above, almost all of the PP-InsP₅ that accumulates in mammalian cells is the 5-PP-InsP₅ isomer). In earlier experiments in which [ATP] levels were perturbed by cell treatment with antimycin, the concentration of PP-InsP₅ decreased much faster ($t_{1/2} = 5$ min) than did the levels of all of the other inositol phosphates (Oliver et al., 1992). The explanation for these rapid responses of PP-InsP₅ to a drop in cellular [ATP] may lie in the IP6Ks exhibiting a relatively high K_m value of approximately 1 mM for ATP (Voglmaier et al., 1996; Saiardi et al., 1999), which lies within the range of values (1-5 mM) usually accredited to the cytosolic [ATP] in mammalian cells (Soboll et al., 1978). Just to emphasize how unusual is this situation, it should be noted that the InsP₄ and InsP₅ kinases display a K_m for ATP which is less than 100 μ M (Tan et al., 1997; Verbsky et al., 2002). In other words, IP6Ks have an atypically low affinity for ATP that could make 5-PP-InsP₅ synthesis especially sensitive to fluctuations in cytosolic [ATP].

The high K_m of the IP6Ks for ATP might also explain an otherwise puzzling observation that the synthesis of both PP-InsP₄ and PP-InsP₅ synthesis is inhibited 20-50% following short-term (<2 hr) treatment of liver cells with thapsigargin (Glennon and Shears, 1993; Padmanabhan et al., 2009). Perhaps IP6K activity decreases in direct response to the decline in [ATP] levels that others have shown can accompany cell-treatment with thapsigargin (Waring and Beaver, 1996). Furthermore, others (Tamas et

MOL #55897

al., 2006) have shown thapsigargin to activate the AMPK cascade. This phenomenon was suggested to represent a homeostatic adjustment to an increased energy-demand following activation of Ca^{2+} -dependent signaling pathways (Tamas et al., 2006). In any case, the latter study provides a context for understanding why the levels of a metabolic messenger might respond to thapsigargin. Yet again, a previously inexplicable behavior of a diphosphoinositol polyphosphate can be rationalized by considering the molecule to be a metabolic messenger.

What might be the biological consequences of this decrease in the levels of 5-PP-InsP₅ and (PP)₂-InsP₄ that seems to accompany a decline in the bioenergetic health of the cell? It should first be noted that a reduction in the rate of synthesis of diphosphoinositol polyphosphates will, just by itself, reduce cellular ATP consumption. This in itself might help defend cellular energy balance. We also know from genetic work with yeasts that when the synthesis of the diphosphoinositol polyphosphates is blocked, the cells grow more slowly (Saiardi et al., 2000), and there are defects in endocytic trafficking (Saiardi et al., 2002) and DNA repair (Luo et al., 2002). These are biological processes that normally consume significant quantities of cellular energy. If these cellular activities are constrained in response to reduced levels of diphosphoinositol polyphosphates, that might provide a mechanism for energy conservation when the cell is bioenergetically challenged. Clearly, these are speculative ideas at present. We need to obtain more precise information on the biological actions of diphosphoinositol polyphosphates, particularly in mammalian cells. This is key to future progress in this field of research.

Since the levels of diphosphoinositol polyphosphates seem to be so closely tied to the bioenergetic status of the cell (see above), we should also consider what might be the biological impact if the levels of the polyphosphates were to be elevated following an

MOL #55897

increase in the cellular [ATP]. For one possible explanation, we next turn our attention to pancreatic β -cells. These cells are metabolically specialized so that cellular [ATP]/[ADP] ratios increase dramatically in response to elevations in serum glucose levels (Detimary et al., 1998). The unusual sensitivity of IP6Ks to fluctuations in cellular [ATP] would appear to be an appropriate mechanism by which cellular 5-PP-InsP₅ concentration in β -cells could be tied to serum glucose levels. This could be an important topic to pursue because 5-PP-InsP₅ has been shown to stimulate insulin release, in an impressive series of electrophysiological experiments (Illies et al., 2007). More recently, mice in which the *IP6K1* was deleted were found to have reduced levels of circulating insulin (Bhandari et al., 2008) and TNP-mediated inhibition of IP6K was shown to reduce insulin secretion from the Min6 pancreatoma cell line (Padmanabhan et al., 2009). The 5-PP-InsP₅ enhances secretion of the hormone by increasing the size of the rapidly-releasable pool of insulin vesicles (Illies et al., 2007). This subset of granules is in close proximity with the plasma membrane, where they are poised to undergo exocytosis as soon as the cell receives the appropriate stimulus (e.g. elevated glucose). This process, which drives the initial phase of insulin secretion, was enhanced either by over-expression of IP6K, or by the direct application of PP-InsP₅, in a dose-dependent manner over the physiologically-relevant concentration range of 1 - 10 μ M (Illies et al., 2007). Any isomer of PP-InsP₅ was able to elicit an increase in insulin secretion (Illies et al., 2007) which has led to speculation that PP-InsP₅ acts by phosphorylating some (as yet undefined) target protein, since, as discussed below, such a molecular action also promises not to be restricted to any particular isomer of PP-InsP₅. The lack of any specificity for PP-InsP₅ suggests that (PP)₂-InsP₄ might also stimulate insulin secretion, but this has yet to be tested. In any case, we can conclude that organismal energy metabolism is regulated by the diphosphoinositol polyphosphates.

MOL #55897

The idea that cellular levels of the diphosphoinositol polyphosphates might reflect cellular bioenergetic status adds a new context to recent observations that PP-InsP₄ levels influence telomere length, in yeast at least (Saiardi et al., 2005; York et al., 2005). The shortening of telomeres is a topic of interest because it accompanies the inevitable process of cellular senescence in somatic cells (Feldser and Greider, 2007; Herbig et al., 2006). This quite normal biological process is beneficial in an animal's early life because it has tumour-suppressive properties (Sedivy, 2007). However, cellular senescence is detrimental in older animals because it contributes to the functional decline of various organ systems (i.e. the ageing process) by inexorably depleting tissues of the functional cells that are required to maintain organ homeostasis. The timing of this telomere-dependent phenomenon of "good things going bad" (Campisi and d'Adda di Fagagna, 2007) or, more technically, "antagonistic pleiotropy" (Shayman and Kirkwood, 1987), is increasingly recognized to be influenced by environmental stress (Rando, 2006; Sedivy, 2007) and, notably, cellular energy status (Narala et al., 2008); cellular senescence is associated with a decline in bioenergetic health (Wang et al., 2003; Zwerschke et al., 2003). The observation that telomere length in yeast is also correlated with the synthesis of very long-chain fatty acids (Ponnusamy et al., 2008) makes this subject even more intriguing because, in mammals, systemic energy balance is influenced by circulating long-chain fatty acids (Lam et al., 2005). It could be productive to study if there is a connection between PP-InsP₄ levels and telomere length in animal cells which reflects an interplay between cellular bioenergetic homeostasis and the diphosphoinositol polyphosphates.

It is also worth noting that some of the proteins that participate in telomere homeostasis, for example the highly-conserved REM (MRE11/RAD50/NBN) complex, also function to regulate DNA repair (Czornak et al., 2008). This observation suggests that it is not

MOL #55897

coincidental that both telomere maintenance (Saiardi et al., 2005; York et al., 2005) and DNA repair (Luo et al., 2002) each require input from the diphosphoinositol polyphosphates; perhaps the polyphosphates act upon both processes through a common mechanism. This could be an important topic, especially as DNA repair is an energy-intensive process that has to be co-ordinated with the mechanisms that maintain cellular energy homeostasis (Olovnikov et al., 2008).

So far, we have discussed circumstances in which the levels of diphosphoinositol polyphosphates positively correlate with the metabolic health of the cell, as described by the levels of ATP and AMP. Interestingly, apoptosis is an important circumstance in which the relationship between adenine nucleotides and diphosphoinositol polyphosphates can sometimes be reversed. This observation comes in part from experiments in which an ovarian carcinoma cell line was treated with cisplatin (Nagata et al., 2005), a platinum-based chemotherapeutic agent that not only cross-links DNA, but also impairs cellular bioenergetic health (Rodriguez-Enriquez et al., 2009). Yet, following cisplatin treatment, the rate of synthesis of 5-PP-InsP₅ initially increased, apparently due to persistent activation of IP6K2 (Nagata et al., 2005). If an increased synthesis of PP-InsP₅ is normally a signal of bioenergetic health (see above), is it possible that the cisplatin-mediated activation of IP6K2 “misleads” the cell, preventing it from countering the bioenergetic stress that cisplatin has induced? In such an event, it can be appreciated that the metabolic crisis could be exacerbated; a sort-of “double-whammy” for cellular energy-balance. Apoptosis is frequently the ultimate outcome of a sustained failure to adapt to metabolic stress (Jin et al., 2007). Is this why cells in which IP6K is over-expressed then succumb to apoptosis? A recent study (Morrison et al., 2009) indicated that apoptosis could even be induced by microinjection into a cell of 25 μM 5-PP-InsP₅ (the product of IP6K activity; Fig. 3).

MOL #55897

Since apoptosis is recognized to be a means by which emerging cancer cells can be purged (Jin et al., 2007), perhaps IP6K activity can be exploited as an anti-cancer defense mechanism. Indeed, the over-expression of IP6K augments the pro-apoptotic actions of not only cisplatin (Nagata et al., 2005), but also other cellular stressors -- etoposide, hydrogen peroxide, hypoxia (Nagata et al., 2005) -- which, again, compromise bioenergetic homeostasis (Rodriguez-Enriquez et al., 2009; Lambotte, 1977; Hara and Abiko, 1995). IP6K2 also enhances the pro-apoptotic actions of interferon- β (Morrison et al., 2002; Morrison et al., 2001). It might be worth studying if the latter effect is related to the decrease in cellular [ATP] reportedly induced by interferon- β (Lewis et al., 1996).

The reason why the type 2 IP6K is selected for this pro-apoptotic process might be associated with its unique, stress-dependent relocation from the nucleus to Bax-positive (i.e., damaged) mitochondria; the type 1 and 3 IP6Ks do not show this effect (Nagata et al., 2005). However, a subsequent study failed to detect increased mitochondrial IP6K2 during apoptosis (Morrison et al., 2005). Perhaps instead the activity of the type 2 enzyme is unique among the IP6K family in being activated by certain pro-apoptotic stimuli, through covalent modification for example (Nagata et al., 2005), or by impeding the association of IP6K2 with HSP90, which normally inhibits the kinase activity (Chakraborty et al., 2008). IP6K2 also has the ability to bind to TNF receptor-associated factor-2, thereby attenuating the latter's influence over the anti-apoptotic NF- κ B signaling pathway (Morrison et al., 2007).

In the discussion of PP-InsP₅ turnover in the preceding paragraphs, we have largely discussed the relationship that the 5-PP-InsP₅ isomer has with cellular energy balance. In yeasts at least, the quantitatively minor 1/3-PP-InsP₅ isomer also has a biologically important role that is directly relevant to metabolic well-being; this PP-InsP₅ participates

MOL #55897

in phosphate homeostasis (Lee et al., 2007). Inorganic phosphate is, of course, an essential nutrient that is required in large amounts for nucleic acid and phospholipid biosynthesis, but it is also essential for cellular energy metabolism. O'Shea and colleagues (Lee et al., 2007) have shown that the synthesis of total PP-InsP₅ levels are up-regulated in yeast cells grown for 1 to 2 hr in low-phosphate media. Genetic evidence indicates that it is the 1/3-isomer of PP-InsP₅ that must have been elevated (Lee et al., 2007), although this was not directly confirmed. In any case (see above), the levels of the other PP-InsP₅ isomer, 5-PP-InsP₅, would not be expected to rise when cells are bioenergetically-stressed, which is the expected outcome of phosphate starvation (Martinez et al., 1998). As discussed in more detail below, elevated 1/3-PP-InsP₅ levels inhibit a cyclin kinase activity that regulates the expression of phosphate responsive (*Pho*) gene products that are important for extracellular phosphate generation and assimilation, such as a phosphate transporter and a secreted acid phosphatase (Springer et al., 2003). Thus, the data published by O'Shea and colleagues (Lee et al., 2007) can be considered to have provided us with a classical demonstration of the actions of a metabolic messenger: a bio-sensor (1/3-PP-InsP₅) that helps the cell avoid perturbations in energy balance when extracellular inorganic phosphate is limiting. As the turnover of diphosphoinositol polyphosphates is quite high, the yeast cells must invest a significant amount of energy to sustain the 10-fold increase in the levels of 1/3-PP-InsP₅ observed during phosphate deprivation (Lee et al., 2007). However, yeasts have significant energy reserves in the form of inorganic polyphosphate, which is mobilized during the initial stages of phosphate starvation (Martinez et al., 1998). Thus, a short-term investment of cellular energy in the increased synthesis of 1/3-PP-InsP₅ might be justified when, hopefully, it can subsequently be repaid by improved scavenging of extracellular organic phosphates. Nevertheless, the window of opportunity for this adaptation will likely be open only for a

MOL #55897

limited time; both ATP and inorganic polyphosphate levels decline substantially after a few hours phosphate starvation (Martinez et al., 1998), which would be expected (see above) to put severe strain upon the ability of yeasts to sustain elevated levels of diphosphoinositol polyphosphates.

As noted by others (Bennett et al., 2006), there are earlier, albeit less-characterized observations, that also speak to there being lines of communication between cellular phosphate status and diphosphoinositol polyphosphates. For example, one group uncovered the molecular identity of IP6K1 while pursuing an explanation for its ability to stimulate cellular uptake of inorganic phosphate (Schell et al., 1999). Consistent with this observation, the *kcs1Δ* strain of *S. cerevisiae* was found to exhibit reduced inorganic phosphate uptake (Saiardi et al., 2004). Deletion of *Kcs1* has also been shown to activate the PHO system (Auesukaree et al., 2005), though in retrospect one might now suggest that this reflects compensatory up-regulation of *Vip1* expression (for example, see (York et al., 2005)), and elevation of 1/3-PP-InsP₅ levels.

Protein Phosphorylation by Diphosphoinositol Polyphosphates?

It is only relatively recently that solid information has emerged that offers possible mechanisms of action of the diphosphoinositol polyphosphates. One proposal has drawn from the a long-standing recognition (Stephens et al., 1993; Hand and Honek, 2007; Laussmann et al., 1996) that the hydrolysis of one of their phosphate groups must be associated with a significant free-energy change. It was consideration of this point that led Stephens and colleagues (Stephens et al., 1993) to hypothesize that the diphosphoinositol polyphosphates might act as phosphate donors in phosphotransferase reactions. Snyder and colleagues have actively pursued this idea. This group (Saiardi et al., 2004; Bhandari et al., 2007) has shown, at least *in vitro*, that

MOL #55897

all of the diphosphoinositol polyphosphates can phosphorylate certain proteins. The consensus phosphorylation site is a serine that is surrounded by acidic residues (Saiardi et al., 2004). The appropriate target sequence is especially well-represented in proteins such as Nsr1 (yeast nucleolin), NOPP140 and TCOF1, which are all nucleolar residents (Saiardi et al., 2004). The primary function of the nucleolus is to synthesize ribosomes (Boisvert et al., 2007). It is therefore of interest that there is evidence of genetic interactions between ribosomal biogenesis and the synthesis of diphosphoinositol polyphosphates (Horigome et al., 2009). Moreover, ribosomal biogenesis consumes 80% of the energy expenditure of a proliferating cell (Thomas, 2000). Thus, any regulation of the rate of ribosome synthesis that might occur following changes in the phosphorylation status of nucleolar proteins could have a significant impact upon cellular bioenergetic homeostasis. This may be one means by which the actions of diphosphoinositol polyphosphates can be linked to cellular energy conservation, which, the reader may recall, I have speculated to be the *raison d'être* for their roles as metabolic messengers (see above).

The transfer of the phosphate group from the diphosphoinositol polyphosphate to a protein substrate occurs independently of protein kinase activity (Saiardi et al., 2004). This phenomenon is particularly remarkable because in the absence of enzymatic assistance, phosphoric anhydrides are considered to be chemically stable, protected by their negative charges from rapid attack by water and other nucleophiles (Westheimer, 1987). It seems that it is the co-ordination of the negative charge by Mg^{2+} that enables the phosphate transfer to occur (Bhandari et al., 2007; Saiardi et al., 2004). There is also a requirement that the target proteins must first be "primed" by an initial casein kinase 2 (CK2)-dependent phosphorylation event (Bennett et al., 2006; Bhandari et al., 2007). Furthermore, the evidence now points to the diphosphoinositol polyphosphates actually further phosphorylating the serine that is initially phosphorylated by CK2

MOL #55897

(Bhandari et al., 2007). These data provide a provocative and novel idea concerning the molecular action of diphosphoinositol polyphosphates while also unveiling evidence of a novel mechanism of covalent modification: nonenzymic diphosphorylation of serine. But does this occur *in vivo*?

The last question is difficult to answer directly, in no small part because of considerable technical challenges. Despite ongoing improvements in the sensitivity of mass spectrometry, it remains a daunting task to identify changes in phosphorylation status of a particular protein in cell lysates (Blackburn and Goshe, 2009). Let alone when the goal is to distinguish between serine diphosphates and serine monophosphates in a peptide fragment. Perhaps in the future it might be possible to develop antibodies against diphospho-serine that can achieve the same goal. Until such a time, only indirect approaches have been possible. For example, Snyder and colleagues used yeast cells in which the InsP₆ kinase (Kcs1) that makes 5-PP-InsP₅ was genetically eliminated (Saiardi et al., 2004). They found that the degree of phosphorylation of endogenous Nsr1 was substantially reduced in these cells. This is a promising observation, but it should be noted that the deletion of Kcs1 impairs cell-wall integrity (Dubois et al., 2002) and compromises a number of yeast's normal biological processes: vacuolar biogenesis (Saiardi et al., 2002; Saiardi et al., 2000), endocytosis (Saiardi et al., 2002), stress responses (Dubois et al., 2002), and DNA recombination (Luo et al., 2002). Moreover, the expression of at least 20 yeast genes changes in response to the elimination of Kcs1 (El Alami et al., 2003). In such circumstances, a change in the phosphorylation status of Nsr1 could be argued to arise independently of PP-InsP₅ synthesis *per se*, and might instead reflect the cell's adjustments to the many molecular responses to the *kcs1Δ* genotype.

Another point to discuss is that the deletion of the *Nsr1* gene in *S. cerevisiae* has been

MOL #55897

observed to be associated with a doubling of intracellular levels of PP-InsP₅ and (PP)₂-InsP₄ (Saiardi et al., 2004). These increases were proposed to reflect a reduced demand for diphosphoinositol polyphosphate turnover, once this proposed target of phosphorylation was eliminated (Saiardi et al., 2004). However, the validity of this proposal might now be questioned by the expansion of the number of proteins now put forward as substrates for protein phosphorylation by the diphosphoinositol polyphosphates (Bhandari et al., 2007). If there really are such a large number of protein substrates, removing just one of them would not be expected to significantly impact the cellular levels of diphosphoinositol polyphosphates. Especially if the putative serine-diphosphate is long-lived (Burton et al., 2009), since this also limits the impact of the phosphorylation process upon the turnover of the phosphate donors. Once again, we should perhaps consider that the changes in levels of PP-InsP₅ and (PP)₂-InsP₄ in the *nsr1Δ* strain might be an indirect effect, perhaps a consequence of the accompanying growth-impaired phenotype.

Another issue to consider is that InsP₆ is quite an effective inhibitor of protein phosphorylation by diphosphoinositol polyphosphates (Saiardi et al., 2004). This might not always be an obstacle in the slime-molds, particularly the aggregated form of *Dictyostelium*, in which levels of (PP)₂-InsP₄ match those of InsP₆ (Laussmann et al., 2000). Moreover, *Dictyostelium* are apparently unique in their synthesis of the 5,6-diphosphate isomer of (PP)₂-InsP₄ (Laussmann et al., 2000; Lin et al., 2009); the hydrolysis of one of these vicinal diphosphates is an especially “high-energy” reaction that could facilitate phosphotransfer to proteins (Hand and Honek, 2007). Thus, slime molds might be a singularly appropriate model in which to study the biological relevance of protein phosphorylation by diphosphoinositol polyphosphates. In contrast, since in all other eukaryotic cells the cellular levels of InsP₆ are at least 25-fold higher than the

MOL #55897

diphosphoinositol polyphosphates, the latter will likely only be capable of phosphorylating proteins in an a microenvironment from which InsP₆ is relatively excluded. This scenario is plausible. There is certainly evidence that some InsP₆ is divided into metabolically-separated “pools” (Otto et al., 2007). Other data showing a punctate distribution of the InsP₅ 2-kinase within certain cellular structures such as nucleoli and stress-granules also indicates that intracellular InsP₆ synthesis is compartmentalized (Brehm et al., 2007). On the other hand, the apparently tight metabolic equilibrium between [³H]-inositol labeled pools of InsP₆ and PP-InsP₅ (Menniti et al., 1993) argues strongly that InsP₆ is not actually physically separated from the diphosphoinositol polyphosphates *in vivo*.

Using the human homologue of Nsr1 - nucleolin - as a model, we (Yang et al., 2008) have searched for evidence that its phosphorylation by diphosphoinositol polyphosphates might be physiologically relevant. We made the assumption that, if Snyder and colleagues (Bhandari et al., 2007; Saiardi et al., 2004) are correct, the degree of nucleolin phosphorylation should increase as the cellular levels of (PP)₂-InsP₄ and/or PP-InsP₅ are elevated. We also noted previous experiments demonstrating that the phosphorylation of nucleolin is associated with its transfer from the nucleolus into the nucleoplasm (Kim et al., 2005). Thus, the extent to which nucleolin accumulates in the nucleoplasm can be anticipated to provide a readout of its degree of phosphorylation. We therefore manipulated cellular levels of diphosphoinositol polyphosphates in an osteosarcoma cell line using a combination of hyperosmotic stress, and some pharmacological tricks (Yang et al., 2008). We found that a hyperosmotic challenge did indeed cause nucleolin to accumulate in the nucleoplasm -- suggesting its degree of phosphorylation was increased -- but this response occurred independently of changes in levels of diphosphoinositol polyphosphates (Yang et al.,

MOL #55897

2008). Nevertheless, our experiments with nucleolin can only be considered an indirect test of the hypothesis put forward by Snyder and colleagues. Further progress in this area requires new methodologies to be developed that can directly detect protein diphosphorylation by diphosphoinositol polyphosphates *in vivo*.

One might anticipate that if diphosphoinositol polyphosphates were indeed to phosphorylate proteins *in vivo*, then the reverse reaction - dephosphorylation of the protein - might also be a regulated event. Yet, so far, no such phosphatase activity has been observed, and in fact, the diphosphorylated proteins are notably resistant to dephosphorylation when added to cell lysates (Bhandari et al., 2007). This metabolic stability has been argued to be biologically significant by ensuring that signaling through this process is long-lived (Burton et al., 2009). Nevertheless, the identification of the requisite phosphatase, even if it is not very active, is key to bolstering the credentials of this hypothesis.

Finally, it is also a little puzzling from a signaling perspective that each of the individual diphosphoinositol polyphosphates have similar abilities to phosphorylate proteins *in vitro* (Bhandari et al., 2007). Why should the cell invest resources in synthesizing several highly-phosphorylated molecules that all have an identical mechanism of action? And, if all of the different diphosphoinositol polyphosphates act in the same way, why does the cell sometimes independently regulate their turnover (see, for example: (Lee et al., 2007;Choi et al., 2007;Choi et al., 2008;Choi et al., 2005;Pesesse et al., 2004))? These are questions which suggest that there must be other mechanisms of action of the diphosphoinositol polyphosphates. In fact, in the case of regulation by 1/3-PP-InsP₅ of yeast cyclin kinase activity (see below), it is almost certain that 1/3-PP-InsP₅ does not act by directly phosphorylating a protein. We can make that statement with some confidence because any of the diphosphoinositol polyphosphates can donate a

MOL #55897

phosphate to a protein, at least *in vitro* (Bhandari et al., 2007). In contrast, O'Shea and colleagues (Lee et al., 2007) described how the kinase inhibition was relatively specific for the 1/3-isomer of PP-InsP₅, while 5-PP-InsP₅ was considerably less effective. Such specificity suggests the participation of a more classical receptor-based signaling mechanism, as discussed below.

Do Cells contain “Receptors” for Diphosphoinositol Polyphosphates?

A traditional role for an intracellular signal is for it to selectively bind to an intracellular “receptor” and thereby alter some inherent property of the protein. In a group of mid-1990 publications, we identified several PP-InsP₅-binding proteins, and at that time it seemed of potential interest that they all have in common a role in regulating vesicular traffic: Coatamer, AP2, and AP180 (previously sometimes called “AP3”) (Ye et al., 1995; Shears et al., 1995; Fleischer et al., 1994). The affinity of AP180 for PP-InsP₅ was determined to be 3 to 5-fold greater than it is for InsP₆ (Ye et al., 1995; Saiardi et al., 2002) and these observations were considered to support the idea that PP-InsP₅ plays a role in endocytosis (Saiardi et al., 2002; Ye et al., 1995). However, these ligand-binding assays were performed before we fully appreciated the biological importance of Mg²⁺ coordinating the negative charge of these highly electronegative polyphosphates (Torres et al., 2005). Since our earlier binding assays did not include divalent cations, the values of the binding affinities that we obtained must now be interpreted cautiously, as previously discussed (Shears, 2001; Bennett et al., 2006). Moreover, we subsequently demonstrated that the domain in AP180 that binds PP-InsP₅ also binds inositol lipids (Ye et al., 1995). The current consensus in the field is that the lipids are the physiologically-relevant ligands, and most notably for PtdIns(4,5)P₂, they are viewed as *promoting* endocytosis (Legendre-Guillemain et al., 2004). This has rather muddled

MOL #55897

the waters with regards to understanding the significance of our *in vitro* clathrin assembly assays, which indicated that endocytosis should be *inhibited* when AP180 binds either the lipids (Hao et al., 1997) or the diphosphoinositol polyphosphates (Ye et al., 1995).

In later ligand-binding experiments that also did not contain divalent cations, Snyder and colleagues (Luo et al., 2003) reported that 5-PP-InsP₅ bound to the PH domain of *Dictyostelium* CRAC (cytosolic regulator of adenylyl cyclase) (Luo et al., 2003). The affinity of PP-InsP₅ was reportedly similar to that of Ins(1,3,4,5)P₄, which is the "headgroup" of PtdIns(3,4,5)P₃ (Luo et al., 2003). This and other evidence suggested that competition between 5-PP-InsP₅ and PtdIns(3,4,5)P₃ for binding to this PH domain regulated the intracellular distribution of the CRAC, which is a key factor for determining the directionality of chemotaxis of *Dictyostelium* (Luo et al., 2003). The fact that *Dictyostelium* contains such exceptionally high levels of diphosphoinositol polyphosphates (see above) makes this hypothesis of special interest. However, it is harder to justify the biological relevance of the apparent binding of PP-InsP₅ to PH domains in several mammalian proteins: PKB, PIKE and TIAM (Luo et al., 2003). The latter argument is made because, in mammalian cells, the levels of PP-InsP₅ are 100-fold lower than they are in *Dictyostelium*. In any case, others (Downes et al., 2005) have been unable to reproduce the observation that PP-InsP₅ binds to PKB, at least. A separate, detailed study of the PH domain in PDK1 could not detect any binding of PP-InsP₅ (Komander et al., 2004).

Arguably the most promising - and selective - intracellular "receptor" for a diphosphoinositol polyphosphate was recently identified by O'Shea and colleagues (Lee et al., 2008; Lee et al., 2007) in experiments with *S. cerevisiae*. We should note at the

MOL #55897

outset that these binding assays were not only performed in the presence of Mg^{2+} , thereby making the data more reliable (see above), but in addition the ligand binding was even found to depend upon divalent cations being present (Lee et al., 2007). In this work, 1/3-PP-InsP₅ was shown to bind to the Pho80/Pho85/Pho81 cyclin-dependent kinase/cyclin kinase inhibitor complex. (In an effort to spare the reader some confusion, we should note that in the O'Shea studies (Lee et al., 2008; Lee et al., 2007) the isomer in question is said to be 4/6-PP-InsP₅. Only subsequently (Lin et al., 2009) was it determined to actually be the 1/3-isomer; also see above for an explanation).

The biological significance of 1/3-PP-InsP₅ binding to the Pho80/Pho85/Pho81 complex is as follows: When *S. cerevisiae* has limited access to inorganic phosphate, Pho81 inhibits cyclin kinase activity, so that it no longer hyperphosphorylates the transcription factor Pho4 (Kaffman et al., 1994), which then enters the nucleus to drive the transcription of genes important for phosphate generation and assimilation, such as a phosphate transporter and a secreted acid phosphatase (Springer et al., 2003). O'Shea and colleagues have demonstrated that 1/3-PP-InsP₅ augments the inhibitory activity of Pho81 (Lee et al., 2007; Lee et al., 2008). This is a relatively specific effect, as 5-PP-InsP₅ is much less effective (Lee et al., 2007). Additionally, from the key perspective of physiological relevance, the inhibitory effect of 1/3-PP-InsP₅ was unaffected by a 50-fold excess of InsP₆ (Lee et al., 2008). Thus, InsP₆ is unlikely to interfere with this action of 1/3-PP-InsP₅ *in vivo*, even though cellular InsP₆ levels are 25-fold greater than those of PP-InsP₅ (Lee et al., 2007). It appears that 1/3-PP-InsP₅ either interacts with both Pho81 and Pho80/Pho85, or it stabilizes the inhibitory association of Pho81 with Pho80/Pho85 (Lee et al., 2008). There are also strong indications of a more complex relationship between Vip1 and the Pho80/Pho85/Pho81 system that deserves further

MOL #55897

attention. For example, the expression of *Vip1* is regulated by Pho4 (Maerkl and Quake, 2007). Of further interest is the observation that the Vip1 protein is phosphorylated by Pho80-Pho85, at least *in vitro* (Dephoure et al., 2005). Pho85 – and its human homologue, Cdk5 – are also important for cell cycle progression, polarized cell growth and cytoskeletal dynamics (Huang et al., 1999;Huang et al., 2007a), thus raising the possibility that such activities might also be regulated by the products of Vip/PPIP5K activity.

The 1/3-PP-InsP₅ inhibits cyclin kinase activity *in vitro* with an IC₅₀ value of 55 μM (Lee et al., 2007). O'Shea and colleagues (Lee et al., 2007) conclude that this is physiologically-relevant in phosphate-restricted yeast, in which they estimated that total PP-InsP₅ levels can rise to 10-30 μM (a 10-fold increase over basal levels). It is useful to look closely at how this estimate was obtained. To arrive at this value, total intracellular PP-InsP₅ concentration in [³H]inositol-labeled cells was estimated from the ratio, PP-[³H]InsP₅ / [³H]InsP₆, and the value of the denominator was *assumed* to be 100 μM (Lee et al., 2007). Is this a realistic assumption? It does have some precedents in eukaryotic cells: *Dictyostelium* synthesizes 300 μM InsP₆ (Laussmann et al., 2000), but this organism appears uniquely adapted in this respect. Some transformed haemopoietic cell lines have been reported to contain 90-100 μM InsP₆ (Bunce et al., 1993). However, another species of yeast, *Schizosaccharomyces pombe* contains only 36 μM InsP₆ (Ingram et al., 2003). All of the other published estimates of cellular InsP₆ levels – including some direct mass assays – are in the 15 - 60 μM range (Barker et al., 2004;Szwergold et al., 1987;Irvine and Schell, 2001;Pittet et al., 1989;Letcher et al., 2008). So 100 μM InsP₆, while not an implausible concentration, would certainly be an unusually high value. Which means that the levels of PP-InsP₅ in

MOL #55897

S. cerevisiae may also have been overestimated.

Additionally, we should note that the O'Shea group calculated the concentration of 1/3-PP-InsP₅ from two more-or-less equally sized [³H]inositol-labeled peaks that eluted after InsP₆ during HPLC analysis (Lee et al., 2007). If these two peaks are actually two different isomers of PP-InsP₅ (Fig. 3), only one-half of this "total PP-[³H]InsP₅" can represent the one specific isomer, 1/3-PP-InsP₅, that inhibits cyclin kinase activity. In this scenario, cellular levels of 1/3-PP-InsP₅ are over-estimated by a factor of two. If instead one of these two "post-InsP₆" peaks were to comprise (PP)₂-InsP₄, then "total PP-[³H]InsP₅" would actually include (PP)₂-InsP₄ and 5-PP-InsP₅, as well as genuine 1/3-PP-InsP₅, leading to an even larger over-estimation of the actual 1/3-PP-InsP₅ levels. At the very least, therefore, we should reduce by half the concentration of 1/3-PP-InsP₅ that should be calculated from the two "post-InsP₆" peaks. Once this correction is made, and after also taking a more conservative value for the intracellular levels of InsP₆ (15-60 μM instead of 100μM), the actual levels of 1/3-PP-InsP₅ after phosphate starvation could lie in the 3-9 μM range, well below the 55 μM required to half-maximally inhibit cyclin kinase activity *in vitro*.

On the other hand, we should not forget that very persuasive genetic data indicate that the cyclin kinase activity of Pho80/Pho85 is inhibited by the catalytic activity of Vip1, which synthesizes 1/3-PP-InsP₅ (Lee et al., 2007; Mulugu et al., 2007). One of the key experiments involved a *vip1Δ* strain of *S. cerevisiae*. In these cells, Pho4 was constitutively cytoplasmic and, therefore, incompetent to activate transcription, even in phosphate-starved yeast. Remember that Pho4 is inactive when it is hyperphosphorylated by the Pho80-Pho85 cyclin/cyclin-dependent kinase (Kaffman et

MOL #55897

al., 1994). Thus, in the *vip1Δ* strain, and, therefore, in the absence of 1/3-PP-InsP₅, the cyclin kinase could not be inactivated by low phosphate conditions (Lee et al., 2007). Data as telling as these indicate that we should offer suggestions as to how the genetic results might be reconciled with the biochemical observations. For example, it may be productive to determine the effects of 1/3,5-(PP)₂-InsP₄ upon cyclin kinase activity. Alternately, perhaps other regulatory factors remain to be discovered that sensitize the cyclin kinase so that 1/3-PP-InsP₅ acts with greater potency *in vivo*. These and other possible explanations are especially worth pursuing, because the isomer-specific association of 1/3-PP-InsP₅ with Pho81 and/or Pho80/Pho85 *in vitro* is, to date, the best characterized example of a specific "receptor" for a diphosphoinositol polyphosphate. In addition to further confirming that this interaction is biologically significant, it will also be important to explicitly define the amino-acid residues that mediate these interactions, and to search for similar motifs in other proteins, such as Cdk5, the mammalian homologue of Pho85 (Huang et al., 1999). In this way, we might uncover additional actions of 1/3-PP-InsP₅.

Having said all that, a couple of additional comments that relate to the aforementioned "post-InsP₆" peaks are warranted. Recall (see above) that the O'Shea data (Lee et al., 2007) can be interpreted to indicate that, in addition to 1/3-PP-InsP₅, the cellular levels of 5-PP-InsP₅ and/or 1/3,5-[PP]₂-InsP₄ may also be elevated in *S. cerevisiae* during phosphate starvation. If this interpretation is correct, then clearly the initial response of the yeast cells to impending bioenergetic stress is different from that which would be expected in mammalian cells. As discussed above, in mammalian cells the levels of 5-PP-InsP₅ and/or 1/3,5-[PP]₂-InsP₄ often appear to *decrease* when bioenergetic health is compromised. One group (Burton et al., 2009) has even indicated that they have

MOL #55897

unpublished data showing that phosphate-starved *S. cerevisiae* do indeed exhibit a decrease in PP-InsP₅ levels. Presumably it is the 5-PP-InsP₅ isomer that these authors are referring to, as this is the predominant isomer in *S. cerevisiae* (see above). Thus, it would be useful to clarify what are the responses of each of the individual diphosphoinositol polyphosphates to phosphate starvation in yeast. It seems that much more information is required before we can fully appreciate the molecular actions of diphosphoinositol polyphosphates *in vivo*.

Concluding Statement

The idea that diphosphoinositol polyphosphates are second messengers - as envisaged in its original definition (Robison et al., 1968), is hard to justify when their cellular levels do not respond to extracellular agents. This situation has prompted the proposal in this review that these molecules respond to an intrinsic change in cellular circumstances: the cellular bioenergetic status. This hypothesis does have the advantage of providing a context for the conditions in which changes in the levels of these molecules has actually been observed. However, even if the idea that diphosphoinositol polyphosphates are metabolic messengers makes a convincing treatise, this review has shown that many questions remain unanswered. Finding the answers is a task of considerable importance. Adaptation to energy-stress is key to the very survival of the cell. A fuller understanding of cellular energy-sensing machinery is potentially also of use as an exploitable target for cancer therapy (Swinnen et al., 2005; Sofer et al., 2005). Finally, our data raise the possibility of a new phenomenon associated with aging: attenuation of inositol pyrophosphate signaling, because of its hypersensitivity to the slight, but progressive decline in cellular energy homeostasis that others have noted during the approach to cellular senescence (Miyoshi et al., 2006; Wang et al., 2003). Pharmacological or genetic intervention in the pathways of inositol pyrophosphate

MOL #55897

signaling may therefore ultimately prove to be of benefit to human health.

Acknowledgements

I would like to thank Drs Erin O'Shea, John York, Adolfo Saiardi and Robin Irvine for their insightful comments. I also thank present and past members of the Inositide Signaling Group for their dedication and contributions to this field of research.

MOL #55897

References

Agranoff BW (1978) Textbook Errors: Cyclitol Confusion. *Trends Biochem Sci* **3**:N283-N285.

Albert C, Safrany S T, Bembenek M E, Reddy K M, Reddy K K, Falck J R, Bröker M, Shears S B and Mayr G W (1997) Biological Variability in the Structures of Diphosphoinositol Polyphosphates in *Dictyostelium Discoideum* and Mammalian Cells. *Biochem J* **327**:553-560.

Auesukaree C, Tochio H, Shirakawa M, Kaneko Y and Harashima S (2005) Plc1p, Arg82p and Kcs1p, Enzymes Involved in Inositol Pyrophosphate Synthesis, Are Essential for Phosphate Regulation and Polyphosphate Accumulation in *Saccharomyces Cerevisiae*. *J Biol Chem* **280**:25127-25133.

Azevedo C and Saiardi A (2006) Extraction and Analysis of Soluble Inositol Polyphosphates From Yeast. *Nat Protoc* **1**:2416-2422.

Bain J, Plater L, Elliott M, Shpiro N, Hastie J, McLauchlan H, Klevernic I, Arthur S, Alessi D and Cohen P (2007) The Selectivity of Protein Kinase Inhibitors; a Further Update. *Biochem J*.

Barker CJ, Wright J, Hughes P J, Kirk C J and Michell R H (2004) Complex Changes in Cellular Inositol Phosphate Complement Accompany Transit Through the Cell Cycle. *Biochem J* **380** :465-473.

Bennett M, Onnebo S M, Azevedo C and Saiardi A (2006) Inositol Pyrophosphates: Metabolism and Signaling. *Cell Mol Life Sci* **63**:552-564.

Bhandari R, Juluri K R, Resnick A C and Snyder S H (2008) Gene Deletion of Inositol

MOL #55897

Hexakisphosphate Kinase 1 Reveals Inositol Pyrophosphate Regulation of Insulin Secretion, Growth, and Spermiogenesis. *Proc Natl Acad Sci U S A*.

Bhandari R, Saiardi A, Ahmadibeni Y, Snowman A M, Resnick A C, Kristiansen T Z, Molina H, Pandey A, Werner J K, Jr., Juluri K R, Xu Y, Prestwich G D, Parang K and Snyder S H (2007) Protein Pyrophosphorylation by Inositol Pyrophosphates Is a Posttranslational Event. *Proc Natl Acad Sci U S A* **104**:15305-15310.

Biemans-Oldehinkel E, Mahmood N A and Poolman B (2006) A Sensor for Intracellular Ionic Strength. *Proc Natl Acad Sci U S A* **103**:10624-10629.

Blackburn K and Goshe M B (2009) Challenges and Strategies for Targeted Phosphorylation Site Identification and Quantification Using Mass Spectrometry Analysis. *Brief Funct Genomic Proteomic* **8**:90-103.

Boisvert FM, van Koningsbruggen S, Navascues J and Lamond A I (2007) The Multifunctional Nucleolus. *Nat Rev Mol Cell Biol* **8**:574-585.

Brearley CA and Hanke D E (1996) Metabolic Evidence for the Order of Addition of Individual Phosphate Esters to the Myo-Inositol Moiety of Inositol Hexakisphosphate in the Duckweed *Spirodela Polrhiza* L. *Biochem J* **314**:227-233.

Brehm MA, Schenk T M, Zhou X, Fanick W, Lin H, Windhorst S, Nalaskowski M M, Kobras M, Shears S B and Mayr G W (2007) Intracellular Localization of Human Inositol 1,3,4,5,6-Pentakisphosphate 2-Kinase. *Biochem J* **408**:335-345.

Browne GJ and Proud C G (2002) Regulation of Peptide-Chain Elongation in Mammalian Cells. *Eur J Biochem* **269**:5360-5368.

Bunce CM, French P J, Allen P, Mountford J C, Moor B, Greaves M F, Michell R H and Brown G (1993) Comparison of the Levels of Inositol Metabolites in Transformed

MOL #55897

Haemopoietic Cells and Their Normal Counterparts. *Biochem J* **289**:667-673.

Burton A, Hu X and Saiardi A (2009) Are Inositol Pyrophosphates Signalling Molecules? *J Cell Physiol* **in press**.

Caffrey JJ, Safrany S T, Yang X and Shears S B (2000) Discovery of Molecular and Catalytic Diversity Among Human Diphosphoinositol Polyphosphate Phosphohydrolases: An Expanding NUDT Family. *J Biol Chem* **275**:12730-12736.

Caffrey JJ and Shears S B (2001) Genetic Rationale for Microheterogeneity of Human Diphosphoinositol Polyphosphate Phosphohydrolase Type 2. *Gene* **269**:53-60.

Campisi J and d'Adda di Fagagna F (2007) Cellular Senescence: When Bad Things Happen to Good Cells. *Nat Rev Mol Cell Biol* **8**:729-740.

Carling D, Clarke P R, Zammit V a and Hardie D G (1989) Purification and Characterization of the AMP-Activated Protein Kinase. Copurification of Acetyl-CoA Carboxylase Kinase and 3-Hydroxy-3-Methylglutaryl-CoA Reductase Kinase Activities. *Eur J Biochem* **186**:129-136.

Chakraborty A, Koldobskiy M A, Sixt K M, Juluri K R, Mustafa A K, Snowman A M, van Rossum D B, Patterson R L and Snyder S H (2008) HSP90 Regulates Cell Survival Via Inositol Hexakisphosphate Kinase-2. *Proc Natl Acad Sci U S A* **105**:1134-1139.

Chamberlain PP, Qian X, Stiles A R, Cho J, Jones D H, Lesley S A, Grabau E A, Shears S B and Spraggon G (2007) Integration of Inositol Phosphate Signaling Pathways Via Human ITPK1. *J Biol Chem* **282**:28117-28125.

Chen W, LeVine H and Rappel W J (2008) A Mathematical Analysis of Second

MOL #55897

Messenger Compartmentalization. *Phys Biol* **5**:46006.

Choi JH, Williams J, Cho J, Falck J R and Shears S B (2007) Purification, Sequencing, and Molecular Identification of a Mammalian PP-InsP5 Kinase That Is Activated When Cells Are Exposed to Hyperosmotic Stress. *J Biol Chem* **282**:30763-30775.

Choi, K., Mollapour, E., Choi, J. H., Falck, J. R., and Shears, S. B. Signaling by inositol pyrophosphates is supervised by "high-energy" phosphate balance, independently of AMP-activated protein kinase. submitted . 2008.

Ref Type: Journal (Full)

Choi K, Mollapour E and Shears S B (2005) Signal Transduction During Environmental Stress: InsP₈ Operates Within Highly Restricted Contexts. *Cell Signal* **17**:1533-1541.

Chu C, Alapat D, Wen X, Timo K, Burstein D, Lisanti M, Shears S and Kohtz D S (2004) Ectopic Expression of Murine Diphosphoinositol Polyphosphate Phosphohydrolase 1 Attenuates Signaling Through the ERK1/2 Pathway. *Cell Signal* **16**:1045-1059.

Czornak K, Chughtai S and Chrzanowska K H (2008) Mystery of DNA Repair: the Role of the MRN Complex and ATM Kinase in DNA Damage Repair. *J Appl Genet* **49**:383-396.

Daval M, Diot-Dupuy F, Bazin R, Hainault I, Viollet B, Vaulont S, Hajduch E, Ferre P and Foufelle F (2005) Anti-Lipolytic Action of AMP-Activated Protein

MOL #55897

Kinase in Rodent Adipocytes. *J Biol Chem* **280**:25250-25257.

Dephoure N, Howson R W, Blethrow J D, Shokat K M and O'Shea E K (2005) Combining Chemical Genetics and Proteomics to Identify Protein Kinase Substrates. *Proc Natl Acad Sci U S A* **102**:17940-17945.

Detimary P, Dejonghe S, Ling Z, Pipeleers D, Schuit F and Henquin J C (1998) The Changes in Adenine Nucleotides Measured in Glucose-Stimulated Rodent Islets Occur in Beta Cells but Not in Alpha Cells and Are Also Observed in Human Islets. *J Biol Chem* **273**:33905-33908.

Dokladda K, Green K A, Pan D A and Hardie D G (2005) PD98059 and U0126 Activate AMP-Activated Protein Kinase by Increasing the Cellular AMP:ATP Ratio and Not Via Inhibition of the MAP Kinase Pathway. *FEBS Lett* **579**:236-240.

Downes CP, Gray A, Fairservice A, Safrany S T, Batty I H and Fleming I (2005) The Regulation of Membrane to Cytosol Partitioning of Signalling Proteins by Phosphoinositides and Their Soluble Headgroups. *Biochem Soc Trans* **33**:1303-1307.

Draskovic P, Saiardi A, Bhandari R, Burton A, Ilc G, Kovacevic M, Snyder S H and Podobnik M (2008) Inositol Hexakisphosphate Kinase Products Contain Diphosphate and Triphosphate Groups. *Chem Biol* **15**:274-286.

Dubois E, Scherens B, Vierendeels F, Ho M W Y, Messenguy F and Shears S B (2002) In *Saccharomyces Cerevisiae*, the Inositol Polyphosphate Kinase Activity of Kcs1p Is Required for Resistance to Salt Stress, Cell Wall Integrity and

MOL #55897

Vacuolar Morphogenesis. *J Biol Chem* **277**:23755-23763.

Edwards-Gilbert G, Veraldi K L and Milcarek C (1997) Alternative Poly(A) Site Selection in Complex Transcription Units: Means to an End? *Nucleic Acids Research* **25**:2547-2561.

El Alami M, Messenguy F, Scherens B and Dubois E (2003) Arg82p Is a Bifunctional Protein Whose Inositol Polyphosphate Kinase Activity Is Essential for Nitrogen and PHO Gene Expression but Not for Mcm1p Chaperoning in Yeast. *Mol Microbiol* **49**:457-468.

Epperson S, Gustafsson A, Gonzalez A, Villegas S, Meszaros J and Brunton L (2005) Pharmacology of G-Protein-linked Signaling in Cardiac Fibroblasts, in *Interstitial Fibrosis in Heart Failure* pp 83-97, New York, Springer.

Europe-Finner GN, Gammon B and Newell P C (1991) Accumulation of [3H]-Inositol into Inositol Polyphosphates During Development of Dictyostelium. *Biochem Biophys Res Commun* **181**:191-196.

Feldser DM and Greider C W (2007) Short Telomeres Limit Tumor Progression in Vivo by Inducing Senescence. *Cancer Cell* **11**:461-469.

Fersht A (1985) *Enzyme Structure and Mechanism*. W.H.Freeman Co., San Francisco.

Fisher DI, Safrany S T, McLennan A G and Cartwright J L (2002) Nudix Hydrolases That Degrade Dinucleoside and Diphosphoinositol Polyphosphates Also Have 5-Phosphoribosyl 1-Pyrophosphate (PRPP) Pyrophosphatase Activity

MOL #55897

That Generates the Glycolytic Activator Ribose 1,5-Bisphosphate. *J Biol Chem* **277**:47313-47317.

Fleischer B, Xie J, Mayrleitner M, Shears S B, Palmer D J and Fleischer S (1994) Golgi Coatomer Binds, and Forms K⁺-Selective Channels Gated by, Inositol Polyphosphates. *J Biol Chem* **269**:17826-17832.

Force A, Lynch M, Pickett F B, Amores A, Yan Y and Postlethwait J (1999) Preservation of Duplicate Genes by Complementary, Degenerative Mutations. *Genetics* **151**:1531-1545.

Fridy PC, Otto J C, Dollins D E and York J D (2007) Cloning and Characterization of Two Human VIP1-Like Inositol Hexakisphosphate and Diphosphoinositol Pentakisphosphate Kinases. *J Biol Chem* **282**:30754-30762.

Glennon MC and Shears S B (1993) Turnover of Inositol Pentakisphosphates, Inositol Hexakisphosphate and Diphosphoinositol Polyphosphates in Primary Cultured Hepatocytes. *Biochem J* **293**:583-590.

Hand CE and Honek J F (2007) Phosphate Transfer From Inositol Pyrophosphates InsP₅PP and InsP₄(PP)₂: a Semi-Empirical Investigation. *Bioorg Med Chem Lett* **17**:183-188.

Hao W, Tan Z, Prasad K, Reddy K K, Chen J, Prestwich G D, Falck J R, Shears S B and Lafer E M (1997) Regulation of AP-3 Function by Inositides. Identification of Phosphatidylinositol 3,4,5-Trisphosphate As a Potent Ligand. *J Biol Chem* **272**:6393-6398.

MOL #55897

Hara A and Abiko Y (1995) Protective Effect of Hypoxia on Mechanical and Metabolic Changes Induced by Hydrogen Peroxide in Rat Hearts. *Am J Physiol* **268**:H614-H620.

Hardie DG (2004) The AMP-Activated Protein Kinase Pathway--New Players Upstream and Downstream. *J Cell Sci* **117**:5479-5487.

Henin N, Vincent M F and Van den B G (1996) Stimulation of Rat Liver AMP-Activated Protein Kinase by AMP Analogues. *Biochim Biophys Acta* **1290**:197-203.

Herbig U, Ferreira M, Condel L, Carey D and Sedivy J M (2006) Cellular Senescence in Aging Primates. *Science* **311**:1257.

Hers HG and Hue L (1983) Gluconeogenesis and Related Aspects of Glycolysis. *Ann Rev Biochem* **52**:617-653.

Hidaka K, Caffrey J J, Hua L, Zhang T, Falck J R, Nickel G C, Carrel L, Barnes L D and Shears S B (2002) An Adjacent Pair of Human NUDT Genes on Chromosome X Are Preferentially Expressed in Testis and Encode Two New Isoforms of Diphosphoinositol Polyphosphate Phosphohydrolase. *J Biol Chem* **277**:32730-32738.

Ho MWY, Yang X, Carew M A, Zhang T, Hua L, Kwon Y-U, Chung S-K, Adelt S, Vogel G, Riley A M, Potter B V L and Shears S B (2002) Regulation of Ins(3456)P4 Signaling by a Reversible Kinase/Phosphatase. *Current Biology* **12**:477-482.

MOL #55897

Hochachka PW (1999) The Metabolic Implications of Intracellular Circulation. *Proc Natl Acad Sci U S A* **96**:12233-12239.

Horigome C, Ikeda R, Okada T, Takenami K and Mizuta K (2009) Genetic Interaction Between Ribosome Biogenesis and Inositol Polyphosphate Metabolism in *Saccharomyces Cerevisiae*. *Biosci Biotechnol Biochem* **73**:443-446.

Hua LV, Green M, Warsh J J and Li P P (2001) Molecular Cloning of a Novel Isoform of Diphosphoinositol Polyphosphate Phosphohydrolase: A Potential Target for Lithium Therapy. *Neuropsychopharmacology* **24**:640-651.

Hua LV, Hidaka K, Pesesse X, Barnes L D and Shears S B (2003) Paralogous Murine Nudt10 and Nudt11 Genes Have Differential Expression Patterns but Encode Identical Proteins That Are Physiologically Competent Diphosphoinositol Polyphosphate Phosphohydrolases. *Biochem J* **373**:81-89.

Huang D, Friesen H and Andrews B (2007a) Pho85, a Multifunctional Cyclin-Dependent Protein Kinase in Budding Yeast. *Mol Microbiol* **66**:303-314.

Huang D, Patrick G, Moffat J, Tsai L H and Andrews B (1999) Mammalian Cdk5 Is a Functional Homologue of the Budding Yeast Pho85 Cyclin-Dependent Protein Kinase. *Proc Natl Acad Sci U S A* **96**:14445-14450.

Huang YH, Grasis J A, Miller A T, Xu R, Soonthornvacharin S, Andreotti A H, Tsoukas C D, Cooke M P and Sauer K (2007b) Positive Regulation of Itk PH Domain Function by Soluble IP4. *Science* **316**:886-889.

MOL #55897

Illies C, Gromada J, Fiume R, Leibiger B, Yu J, Juhl K, Yang S-N, Barma D K, Falck J R, Saiardi A, Barker C J and Berggren P-O (2007) Inositol Pyrophosphates Determine Exocytic Capacity. *Science* **318**:1299-1302.

Ingram SW, Safrany S T and Barnes L D (2003) Disruption and Overexpression of the *Schizosaccharomyces Pombe Aps1* Gene and the Effects on Growth Rate, Morphology, and Intracellular Diadenosine 5', 5'''-P¹, P⁵-Pentaphosphate and Diphosphoinositol Polyphosphate Concentrations. *Biochem J* **369**:519-528.

Inoki K, Zhu T and Guan K L (2003) TSC2 Mediates Cellular Energy Response to Control Cell Growth and Survival. *Cell* **115**:577-590.

Irvine RF and Schell M (2001) Back in the Water: the Return of the Inositol Phosphates. *Nature Reviews Molecular Cell Biology* **2**:327-338.

IUPAC-IUB Commission on Biochemical Nomenclature (1969) [IUPAC Commission on the Nomenclature of Organic Chemistry (CON) and IUPAC-IUB Commission on Biochemical Nomenclature (CBN). Nomenclature of Cyclitol. Tentative Rules]. *Hoppe Seylers Z Physiol Chem* **350**:523-535.

IUPAC-IUB Commission on Biochemical Nomenclature (1977) Nomenclature of Phosphorus-Containing Compounds of Biochemical Importance (Recommendations 1976). *Proc Nat Acad Sci USA* **74** :2222-2230.

Jin S, DiPaola R S, Mathew R and White E (2007) Metabolic Catastrophe As a Means to Cancer Cell Death. *J Cell Sci* **120**:379-383.

Kaffman A, Herskowitz I, Tjian R and O'Shea E K (1994) Phosphorylation of the

MOL #55897

Transcription Factor PHO4 by a Cyclin-CDK Complex, PHO80-PHO85. *Science* **263**:1153-1156.

Kim K, Dimitrova D D, Carta K M, Saxena A, Daras M and Borowiec J A (2005) Novel Checkpoint Response to Genotoxic Stress Mediated by Nucleolin-Replication Protein a Complex Formation. *Mol Cell Biol* **25**:2463-2474.

Komander D, Fairservice A, Deak M, Kular G S, Prescott A R, Peter D C, Safrany S T, Alessi D R and van Aalten D M (2004) Structural Insights into the Regulation of PDK1 by Phosphoinositides and Inositol Phosphates. *EMBO J* **23**:3918-3928.

Kozak M (1996) Interpreting CDNA Sequences: Some Insights From Studies on Translation. *Mammalian Genome* **7**:563-574.

Lam TK, Pocai A, Gutierrez-Juarez R, Obici S, Bryan J, guilar-Bryan L, Schwartz G J and Rossetti L (2005) Hypothalamic Sensing of Circulating Fatty Acids Is Required for Glucose Homeostasis. *Nat Med* **11**:320-327.

Lambotte L (1977) Effect of Anoxia and ATP Depletion on the Membrane Potential and Permeability of Dog Liver. *J Physiol* **269**:53-76.

Laussmann T, Eujen R, Weisshuhn C M, Thiel U, Falck J R and Vogel G (1996) Structures of Diphospho-Myo-Inositol Pentakisphosphate and Bisdiphospho-Myo-Inositol Tetrakisphosphate From *Dictyostelium* by NMR Analysis. *Biochem J* **315**:715-725.

Laussmann T, Pikzack C, Thiel U, Mayr G W and Vogel G (2000) Diphospho-

MOL #55897

Myo-Inositol Phosphates During the Life Cycle of *Dictyostelium* and *Polysphondylium*. *Eur J Biochem* **267**:2447-2451.

Laussmann T, Reddy K M, Reddy K K, Falck J R and Vogel G (1997) Diphospho-*Myo*-Inositol Phosphates From *Dictyostelium* Identified As D-6-Diphospho-*Myo*-Inositol Pentakisphosphate and D-5,6-Bisdiphospho-*Myo*-Inositol Tetrakisphosphate. *Biochem J* **322**:31-33.

Lee YS, Huang K, Quioco F A and O'Shea E K (2008) Molecular Basis of Cyclin-CDK-CKI Regulation by Reversible Binding of an Inositol Pyrophosphate. *Nat Chem Biol* **4**:25-32.

Lee YS, Mulugu S, York J D and O'Shea E K (2007) Regulation of a Cyclin-CDK-CDK Inhibitor Complex by Inositol Pyrophosphates. *Science* **316**:109-112.

Legendre-Guillemain V, Wasiak S, Hussain N K, Angers A and McPherson P S (2004) ENTH/ANTH Proteins and Clathrin-Mediated Membrane Budding. *J Cell Sci* **117**:9-18.

Leslie NR, McLennan A G and Safrany S T (2002) Cloning and Characterization of HAp1 and HAp2, Human Diadenosine Polyphosphate-Metabolizing Nudix Hydrolases. *BMC Biochemistry* **3**:20.

Letcher AJ, Schell M J and Irvine R F (2008) Do Mammals Make All Their Own Inositol Hexakisphosphate? *Biochem J* **416**:263-270.

Lewis JA, Huq A and Najarro P (1996) Inhibition of Mitochondrial Function by Interferon. *J Biol Chem* **271**:13184-13190.

MOL #55897

Lin H, Fridy P C, Ribeiro A A, Choi J H, Barma D K, Vogel G, Falck J R, Shears S B, York J D and Mayr G W (2009) Structural Analysis and Detection of Biological Inositol Pyrophosphates Reveals That the VIP/PPIP5K Family Are 1/3-Kinases. *J Biol Chem* **284**:1863-1872.

Luo HR, Huang Y E, Chen J C, Saiardi A, Iijima M, Ye K, Huang Y, Nagata E, Devreotes P and Snyder S H (2003) Inositol Pyrophosphates Mediate Chemotaxis in Dictyostelium Via Pleckstrin Homology Domain-PtdIns(3,4,5)P₃ Interactions. *Cell* **114**:559-572.

Luo HR, Saiardi A, Yu H, Nagata E, Ye K and Snyder S H (2002) Inositol Pyrophosphates Are Required for DNA Hyperrecombination in Protein Kinase C1 Mutant Yeast. *Biochemistry* **41**:2509-2515.

Maerkl SJ and Quake S R (2007) A Systems Approach to Measuring the Binding Energy Landscapes of Transcription Factors. *Science* **315**:233-237.

Majerus PW, Zou J, Marjanovic J, Kisseleva M V and Wilson M P (2008) The Role of Inositol Signaling in the Control of Apoptosis. *Adv Enzyme Regul* **48**:10-17.

Martinez P, Zvyagilskaya R, Allard P and Persson B L (1998) Physiological Regulation of the Derepressible Phosphate Transporter in *Saccharomyces Cerevisiae*. *J Bacteriol* **180**:2253-2256.

Mayr GW (1988) A Novel Metal-Dye Detection System Permits Picomolar-Range H.p.l.c. Analysis of Inositol Polyphosphates From Non-Radioactively Labelled

MOL #55897

Cell or Tissue Specimens. *Biochem J* **254**:585-591.

McLennan AG (2006) The Nudix Hydrolase Superfamily. *Cell Mol Life Sci* **63**:123-143.

Menniti FS, Miller R N, Putney J W, Jr. and Shears S B (1993) Turnover of Inositol Polyphosphate Pyrophosphates in Pancreatoma Cells. *J Biol Chem* **268**:3850-3856.

Merrill GF, Kurth E J, Hardie D G and Winder W W (1997) AICA Riboside Increases AMP-Activated Protein Kinase, Fatty Acid Oxidation, and Glucose Uptake in Rat Muscle. *Am J Physiol* **273**:E1107-E1112.

Michell RH (2008) Inositol Derivatives: Evolution and Functions. *Nat Rev Mol Cell Biol* **9**:151-161.

Mishra J and Bhalla U S (2002) Simulations of Inositol Phosphate Metabolism and Its Interaction With InsP(3)-Mediated Calcium Release. *Biophysical Journal* **83**:1298-1316.

Miyoshi N, Oubrahim H, Chock P B and Stadtman E R (2006) Age-Dependent Cell Death and the Role of ATP in Hydrogen Peroxide-Induced Apoptosis and Necrosis. *Proc Natl Acad Sci U S A* **103**:1727-1731.

Morrison BH, Bauer J A, Hu J, Grane R W, Ozdemir A M, Chawla-Sarkar M, Gong B, Almasan A, Kalvakolanu D V and Lindner D J (2002) Inositol Hexakisphosphate Kinase 2 Sensitizes Ovarian Carcinoma Cells to Multiple Cancer Therapeutics. *Oncogene* **21**:1882-1889.

MOL #55897

Morrison BH, Bauer J A, Kalvakolanu D V and Lindner D J (2001) Inositol Hexakisphosphate Kinase 2 Mediates Growth Suppressive and Apoptotic Effects of Interferon- β in Ovarian Carcinoma Cells. *J Biol Chem* **276**:24965-24970.

Morrison BH, Bauer J A, Lupica J A, Tang Z, Schmidt H, DiDonato J A and Lindner D J (2007) Effect of Inositol Hexakisphosphate Kinase 2 on Transforming Growth Factor Beta-Activated Kinase 1 and NF-KappaB Activation. *J Biol Chem* **282**:15349-15356.

Morrison BH, Tang Z, Jacobs B S, Bauer J A and Lindner D J (2005) Apo2L/TRAIL Induction and Nuclear Translocation of Inositol Hexakisphosphate Kinase 2 During IFN-Beta-Induced Apoptosis in Ovarian Carcinoma. *Biochem J* **385**:595-603.

Morrison BH, Haney R, Lamarre E, Drazba J, Prestwich GD and Lindner, DJ (2009) *Oncogene* doi:10.1038/onc.2009.113

Mulugu S, Bai W, Fridy P C, Bastidas R J, Otto J C, Dollins D E, Haystead T A, Ribeiro A A and York J D (2007) A Conserved Family of Enzymes That Phosphorylate Inositol Hexakisphosphate. *Science* **316**:106-109.

Nagata E, Luo H R, Saiardi A, Bae B I, Suzuki N and Snyder S H (2005) Inositol Hexakisphosphate Kinase-2, a Physiologic Mediator of Cell Death. *J Biol Chem* **280**:1634-1640.

Narala SR, Allsopp R C, Wells T B, Zhang G, Prasad P, Coussens M J, Rossi D J, Weissman I L and Vaziri H (2008) SIRT1 Acts As a Nutrient-Sensitive Growth Suppressor and Its Loss Is Associated With Increased AMPK and Telomerase

MOL #55897

Activity. *Mol Biol Cell* **19**:1210-1219.

Oliver KG, Putney J W, Jr., Obie J F and Shears S B (1992) The Interconversion of Inositol 1,3,4,5,6-Pentakisphosphate and Inositol Tetrakisphosphates in AR4-2J Cells. *J Biol Chem* **267**:21528-21534.

Olovnikov IA, Kravchenko J E and Chumakov P M (2008) Homeostatic Functions of the P53 Tumor Suppressor: Regulation of Energy Metabolism and Antioxidant Defense. *Semin Cancer Biol.*

Olsen JV, Blagoev B, Gnäd F, Macek B, Kumar C, Mortensen P and Mann M (2006) Global, in Vivo, and Site-Specific Phosphorylation Dynamics in Signaling Networks. *Cell* **127**:635-648.

Otto JC, Kelly P, Chiou S T and York J D (2007) Alterations in an Inositol Phosphate Code Through Synergistic Activation of a G Protein and Inositol Phosphate Kinases. *Proc Natl Acad Sci U S A* **104**:15653-15658.

Padmanabhan U, Dollins D E, Fridy P C, York J D and Downes C P (2009) Characterization of a Selective Inhibitor of Inositol Hexakisphosphate Kinases: Use in Defining Biological Roles and Metabolic Relationships of Inositol Pyrophosphates. *J Biol Chem.*

Pesesse X, Choi K, Zhang T and Shears S B (2004) Signalling by Higher Inositolpolyphosphates: Synthesis of Bis-Diphosphoinositol Tetrakisphosphate ("InsP8") Is Selectively Activated by Hyperosmotic Stress. *J Biol Chem* **279**:43378-43381.

MOL #55897

Pittet D, Schlegel W, Lew D P, Monod A and Mayr G W (1989) Mass Changes in Inositol Tetrakis- and Pentakisphosphate Isomers Induced by Chemotactic Peptide Stimulation in HL-60 Cells. *J Biol Chem* **264**:18489-18493.

Ponnusamy S, Alderson N L, Hama H, Bielawski J, Jiang J C, Bhandari R, Snyder S H, Jazwinski S M and Ogretmen B (2008) Regulation of Telomere Length by Fatty Acid Elongase 3 in Yeast. Involvement of Inositol Phosphate Metabolism and Ku70/80 Function. *J Biol Chem* **283**:27514-27524.

Rall TW and Sutherland E W (1958) Formation of a Cyclic Adenine Ribonucleotide by Tissue Particles. *J Biol Chem* **232**:1065-1076.

Rando TA (2006) Prognostic Value of Telomere Length: the Long and Short of It. *Ann Neurol* **60**:155-157.

Robison GA, Butcher R W and Sutherland E W (1968) Cyclic AMP. *Annu Rev Biochem* **37**:149-174.

Rodriguez-Enriquez S, Marin-Hernandez A, Gallardo-Perez J C, Carreno-Fuentes L and Moreno-Sanchez R (2009) Targeting of Cancer Energy Metabolism. *Mol Nutr Food Res* **53**:29-48.

Rolland F, Winderickx J and Thevelein J M (2001) Glucose-Sensing Mechanisms in Eukaryotic Cells. *Trends Biochem Sci* **26**:310-317.

Ruiz-Stewart I, Tiyyagura S R, Lin J E, Kazerounian S, Pitari G M, Schulz S, Martin E, Murad F and Waldman S A (2004) Guanylyl Cyclase Is an ATP Sensor Coupling Nitric Oxide Signaling to Cell Metabolism. *Proc Natl Acad Sci U S A*

MOL #55897

101:37-42.

Sabina RL, Patterson D and Holmes E W (1985) 5-Amino-4-Imidazolecarboxamide Riboside (Z-Riboside) Metabolism in Eukaryotic Cells. *J Biol Chem* **260**:6107-6114.

Safrany ST, Caffrey J J, Yang X, Bembenek M E, Moyer M B, Burkhart W A and Shears S B (1998) A Novel Context for the "MutT" Module, a Guardian of Cell Integrity, in a Diphosphoinositol Polyphosphate Phosphohydrolase. *EMBO J* **17**:6599-6607.

Safrany ST and Shears S B (1998) Turnover of Bis-Diphosphoinositol Tetrakisphosphate in a Smooth Muscle Cell Line Is Regulated by B₂- Adrenergic Receptors Through a CAMP-Mediated, A-Kinase-Independent Mechanism. *EMBO J* **17**:1710-1716.

Saiardi A, Bhandari A, Resnick R, Cain A, Snowman A M and Snyder S H (2004) Inositol Pyrophosphate: Physiologic Phosphorylation of Proteins. *Science* **306**:2101-2105.

Saiardi A, Caffrey J J, Snyder S H and Shears S B (2000) The Inositol Hexakisphosphate Kinase Family: Catalytic Flexibility, and Function in Yeast Vacuole Biogenesis. *J Biol Chem* **275**:24686-24692.

Saiardi A, Erdjument-Bromage H, Snowman A, Tempst P and Snyder S H (1999) Synthesis of Diphosphoinositol Pentakisphosphate by a Newly Identified Family of Higher Inositol Polyphosphate Kinases. *Current Biology* **9**:1323-1326.

MOL #55897

Saiardi A, Nagata E, Luo H R, Sawa A, Luo X, Snowman A M and Snyder S H (2001a) Mammalian Inositol Polyphosphate Multikinase Synthesizes Inositol 1,4,5-Trisphosphate and an Inositol Pyrophosphate. *Proc Natl Acad Sci U S A* **98**:2306-2311.

Saiardi A, Nagata E, Luo H R, Snowman A M and Snyder S H (2001b) Identification and Characterization of a Novel Inositol Hexakisphosphate Kinase. *J Biol Chem* **276**:39179-39185.

Saiardi A, Resnick A C, Snowman A M, Wendland B and Snyder S H (2005) Inositol Pyrophosphates Regulate Cell Death and Telomere Length Via PI3K-Related Protein Kinases. *Proc Nat Acad Sci USA* **102**:1911-1914.

Saiardi A, Sciambi C, McCaffery J M, Wendland B and Snyder S H (2002) Inositol Pyrophosphates Regulate Endocytic Trafficking. *Proc Nat Acad Sci USA* **99**:14206-14211.

Schell MJ, Letcher A J, Brearley C A, Biber J, Murer H and Irvine R F (1999) PiUS (Pi Uptake Stimulator) Is an Inositol Hexakisphosphate Kinase. *FEBS Lett* **461**:169-172.

Sedivy JM (2007) Telomeres Limit Cancer Growth by Inducing Senescence: Long-Sought in Vivo Evidence Obtained. *Cancer Cell* **11**:389-391.

Seeds AM, Bastidas R J and York J D (2005) Molecular Definition of a Novel Inositol Polyphosphate Metabolic Pathway Initiated by Inositol 1,4,5-Trisphosphate 3-Kinase Activity in *Saccharomyces Cerevisiae*. *J Biol Chem*.

MOL #55897

Seeds AM, Sandquist J C, Spana E P and York J D (2004) A Molecular Basis for Inositol Polyphosphate Synthesis in *Drosophila Melanogaster*. *J Biol Chem*.

Shayman JA and Kirkwood M T (1987) Bradykinin-Stimulated Changes in Inositol Phosphate in Renal Papillary Collecting Tubule Cells. *Biochem Biophys Res Commun* **145**:1119-1125.

Shears SB (1997) Measurement of inositol phosphate turnover in intact cells and cell-free systems, in *Signalling by Inositides: A Practical Approach*. (Shears SB ed) pp 33-52, Oxford University Press, Oxford, U.K.

Shears SB (2001) Assessing the Omnipotence of Inositol Hexakisphosphate. *Cell Signal* **13**:151-158.

Shears SB (2004) How Versatile Are Inositol Phosphate Kinases? *Biochem J* **377**:265-280.

Shears SB, Ali N, Craxton A and Bembenek M E (1995) Synthesis and Metabolism of *Bis*-Diphosphoinositol Tetrakisphosphate *in Vitro* and *in Vivo*. *J Biol Chem* **270**:10489-10497.

Sherman WR, Rasheed A, Mauck L A and Wiecko J (1977) Incubations of Testis Myo-Inositol-1-Phosphate Synthase With D-(5-18O)Glucose 6-Phosphate and With H₂¹⁸O Show No Evidence of Schiff Base Formation. *J Biol Chem* **252**:5672-5676.

Soboll S, Scholz R and Heldt H W (1978) Subcellular Metabolite Concentrations. Dependence of Mitochondrial and Cytosolic ATP Systems on the Metabolic State

MOL #55897

of Perfused Rat Liver. *Eur J Biochem* **87**:377-390.

Sofer A, Lei K, Johannessen C M and Ellisen L W (2005) Regulation of MTOR and Cell Growth in Response to Energy Stress by REDD1. *Mol Cell Biol* **25**:5834-5845.

Springer M, Wykoff D D, Miller N and O'Shea E K (2003) Partially Phosphorylated Pho4 Activates Transcription of a Subset of Phosphate-Responsive Genes. *PLoS Biol* **1**:E28.

Stephens LR, Hawkins P T, Stanley A F, Moore T, Poyner D R, Morris P J, Hanley M R, Kay R R and Irvine R F (1991) Myo-Inositol Pentakisphosphates. Structure, Biological Occurrence and Phosphorylation to Myo-Inositol Hexakisphosphates. *Biochem J* **275**:485-499.

Stephens LR and Irvine R F (1990) Stepwise Phosphorylation of Myo-Inositol Leading to Myo-Inositol Hexakisphosphate in Dictyostelium. *Nature* **346**:580-583.

Stephens LR, Radenberg T, Thiel U, Vogel G, Khoo K-H, Dell A, Jackson T R, Hawkins P T and Mayr G W (1993) The Detection, Purification, Structural Characterization and Metabolism of Diphosphoinositol Pentakisphosphate(s) and Bisdiphosphoinositol Tetrakisphosphate(s). *J Biol Chem* **268**:4009-4015.

Streb H, Irvine R F, Berridge M J and Schulz I (1983) Release of Ca²⁺ From a Nonmitochondrial Store in Pancreatic Cells by Inositol-1,4,5-Trisphosphate. *Nature* **306**:67-68.

Sutherland EW and Rall T W (1958) Fractionation and Characterization of a

MOL #55897

Cyclic Adenine Ribonucleotide Formed by Tissue Particles. *J Biol Chem* **232**:1077-1091.

Swinnen JV, Beckers A, Brusselmans K, Organe S, Segers J, Timmermans L, Vanderhoydonc F, Deboel L, Derua R, Waelkens E, De Schrijver E, Van de S T, Noel A, Fougelle F and Verhoeven G (2005) Mimicry of a Cellular Low Energy Status Blocks Tumor Cell Anabolism and Suppresses the Malignant Phenotype. *Cancer Res* **65**:2441-2448.

Szwegold BS, Graham R A and Brown T R (1987) Observation of Inositol Pentakis- and Hexakis-Phosphates in Mammalian Tissues by ^{31}P NMR. *Biochem Biophys Res Commun* **149**:874-881.

Tamas P, Hawley S A, Clarke R G, Mustard K J, Green K, Hardie D G and Cantrell D A (2006) Regulation of the Energy Sensor AMP-Activated Protein Kinase by Antigen Receptor and Ca^{2+} in T Lymphocytes. *J Exp Med* **203**:1665-1670.

Tan Z, Bruzik K S and Shears S B (1997) Properties of the Inositol 3,4,5,6-Tetrakisphosphate 1-Kinase Purified From Rat Liver. Regulation of Enzyme Activity by Inositol 1,3,4-Trisphosphate. *J Biol Chem* **272**:2285-2290.

Thomas G (2000) An Encore for Ribosome Biogenesis in the Control of Cell Proliferation. *Nat Cell Biol* **2**:E71-E72.

Torres J, Domínguez S, Cerdá F M, Obal G, Mederos A, Irvine R F, Díaz A and Kremer C (2005) Solution Behaviour of Myo -Inositol Hexakisphosphate in the Presence of Multivalent Cations. Prediction of a Neutral Pentamagnesium

MOL #55897

Species Under Cytosolic/Nuclear Conditions. *Journal of Inorganic Biochemistry* **99**:828-840.

Verbsky JW, Wilson M P, Kisseleva M V, Majerus P W and Wentz S R (2002) The Synthesis of Inositol Hexakisphosphate: Characterization of Human Inositol 1,3,4,5,6-Pentakisphosphate 2-Kinase. *J Biol Chem* **277**:31857-31862.

Voglmaier SM, Bembenek M E, Kaplin A I, Dormán G, Olszewski J D, Prestwich G D and Snyder S H (1996) Purified Inositol Hexakisphosphate Kinase Is an ATP Synthase: Diphosphoinositol Pentakisphosphate As a High-Energy Phosphate Donor. *Proc Nat Acad Sci USA* **93**:4305-4310.

Wang W, Yang X, Lopez d S, I, Carling D and Gorospe M (2003) Increased AMP:ATP Ratio and AMP-Activated Protein Kinase Activity During Cellular Senescence Linked to Reduced HuR Function. *J Biol Chem* **278**:27016-27023.

Waring P and Beaver J (1996) Cyclosporin A Rescues Thymocytes From Apoptosis Induced by Very Low Concentrations of Thapsigargin: Effects on Mitochondrial Function. *Exp Cell Res* **227**:264-276.

Watts DJ (1984) Protein Synthesis During Development and Differentiation in the Cellular Slime Mould Dictyostelium Discoideum. *Biochem J* **220**:1-14.

Westheimer FH (1987) Why Nature Chose Phosphates. *Science* **235**:1173-1178.

Wilcox RA, Safrany S T, Lampe D, Mills S J, Nahorski S R and Potter B V L (1994) Modification at C2 of Myo-Inositol 1,4,5-Trisphosphate Produces Inositol Trisphosphates and Tetrakisphosphates With Potent Biological Activities. *Eur J*

MOL #55897

Biochem **223**:115-124.

Wong NS, Barker C J, Morris A J, Craxton A, Kirk C J and Michell R H (1992) The Inositol Phosphates of WRK1 Rat Mammary Tumour Cells. *Biochem J* **286**:459-468.

Yang L, Reece J M, Cho J, Bortner C D and Shears S B (2008) The Nucleolus Exhibits an Osmotically Regulated Gatekeeping Activity That Controls the Spatial Dynamics and Functions of Nucleolin. *J Biol Chem* **283**:11823-11831.

Ye W, Ali N, Bembenek M E, Shears S B and Lafer E M (1995) Inhibition of Clathrin Assembly by High-Affinity Binding of Specific Inositol Polyphosphates to the Synapse-Specific Clathrin Assembly Protein AP-3. *J Biol Chem* **270**:1564-1568.

Yin W, Mu J and Birnbaum M J (2003) Role of AMP-Activated Protein Kinase in Cyclic AMP-Dependent Lipolysis In 3T3-L1 Adipocytes. *J Biol Chem* **278**:43074-43080.

York JD (2006) Regulation of Nuclear Processes by Inositol Polyphosphates. *Biochim Biophys Acta* **1761**:552-559.

York SJ, Armbruster B N, Greenwell P, Petes T D and York J D (2005) Inositol Diphosphate Signaling Regulates Telomere Length. *J Biol Chem* **280**:4264-4269.

Zhang T, Caffrey J J and Shears S B (2001) The Transcriptional Regulator, Arg82, Is a Hybrid Kinase With Both Monophosphoinositol- and Diphosphoinositol- Polyphosphate Synthase Activity. *FEBS Lett* **494**:208-212.

MOL #55897

Zhuo ML, Huang Y, Liu D P and Liang C C (2005) KATP Channel: Relation With Cell Metabolism and Role in the Cardiovascular System. *Int J Biochem Cell Biol* **37**:751-764.

Zwerschke W, Mazurek S, Stockl P, Hutter E, Eigenbrodt E and Jansen-Durr P (2003) Metabolic Analysis of Senescent Human Fibroblasts Reveals a Role for AMP in Cellular Senescence. *Biochem J* **376**:403-411.

MOL #55897

Footnote: My laboratory's research is funded by the Intramural Research Program of the NIH / National Institute of Environmental Health Sciences.

MOL #55897

Figure Legends:

Fig. 1. The axis of symmetry across the inositol ring.

The apparent simplicity of the *myo*-inositol molecule is deceptive. Here it is depicted both in its thermodynamically-preferred “chair” form (left graphic) and in a Haworth projection (right graphic). The latter is not so technically-discriminating, but it does have the advantage of more obviously illustrating the plane of symmetry that runs across the 2/5 axis (the broken line), thereby dividing the molecule into two chiral halves. The hydrogen atoms are omitted, for clarity. Agranoff’s turtle (Agranoff, 1978) provides a popular *aide-mémoire* for the carbon numbering (see also (Irvine and Schell, 2001)); the logic underlying the numbering system is described elsewhere (IUPAC-IUB Commission on Biochemical Nomenclature, 1969).

MOL #55897

Fig. 2. Synthesis of Ins(1,3,4,5,6)P₅ from Ins(1,4,5)P₃ in Mammals.

In the abbreviations of the chemical structures, “Ins” indicates the *myo*-inositol skeleton. The number of monophosphates around the inositol ring is denoted as a suffix after the ‘P’. The figure shows two pathways (ITPK1-dependent and ITPK1-independent) by which mammalian cells convert Ins(1,4,5)P₃ to Ins(1,3,4,5,6)P₅. It is unclear which of the two routes of Ins(1,3,4,5,6)P₅ synthesis are quantitatively the more important. IP3K = Ins(1,4,5)P₃-kinase, IPMK = inositol polyphosphate multikinase (also known as Ipk2 in yeast), 5-PASE = Ins(1,4,5)P₃ / Ins(1,3,4,5)P₄ 5-phosphatase, ITPK1 = inositol trisphosphate kinase. There is phylogenetic variability in the isomeric nature of the InsP₄ produced by Ins(1,4,5)P₃ phosphorylation by IPMK: it is mainly Ins(1,3,4,5)P₄ in mammals, mainly Ins(1,4,5,6)P₄ in yeasts, and IPK2 from *Drosophila* synthesizes almost equal amounts of both InsP₄ isomers ((Seeds et al., 2004)). Note that yeasts and *Drosophila* do not encode ITPK1 in their genomes (Seeds et al., 2004). Finally, plants (Brearley and Hanke, 1996) and slime molds (Stephens and Irvine, 1990) can synthesize Ins(1,3,4,5,6)P₅ by alternate metabolic pathways that do not utilize Ins(1,4,5)P₃.

MOL #55897

Fig. 3 Synthesis of PP-InsP₄, PP-InsP₅ and (PP)₂-InsP₄ in Yeasts and Mammals.

This figure focuses on the best-characterized diphosphoinositol polyphosphates. Other members of this class of compounds do exist, but are less well understood (see text for details). The prefixes denote the number of diphosphates (PP). The positions of the diphosphate groups have been determined in the following publications: (Albert et al., 1997; Lin et al., 2009; Draskovic et al., 2008). The "1/3" designation indicates that it is not yet known if the diphosphate is present at either (or both!) of positions 1 and 3; the choice of position 1 in the figure is arbitrary. The 5- and 1/3-isomers of PP-InsP₅ are sometimes referred to as 5-IP₇ and 1/3-IP₇, respectively. The 1/3,5-(PP)₂-InsP₄ is often termed "IP₈" in the literature (see text). In addition to 5-PP-InsP₄, some 1/3-PP-InsP₄ is also formed from Ins(1,3,4,5,6)P₅ by IP₆K (Draskovic et al., 2008).

MOL #55897

Table 1. Diphosphoinositol Polyphosphate Metabolism: Enzyme Nomenclature.

HGNC-Approved Gene Name	Catalytic Function	Other names	<i>S. cerevisiae</i> homologue
<i>HISPPD2A</i>	Kinase (and phosphatase?*)	VIP1 PIP5K1	Vip1
<i>HISPPD1</i>	Kinase (and phosphatase?*)	VIP2 PIP5K2	
<i>IP6K#</i>	Kinase	IHPK#	Kcs1
<i>NUDT#</i>	Phosphatase	DIPP#	Ddp1

There are three IP6K genes: 1, 2 and 3.

There are four DIPP genes: 1, 2 α , 2 β , 3.

* The protein has an “acid phosphatase”-like domain, but to date no catalytic activity has

MOL #55897

been reported.

Fig. 1




Fig. 2




Fig. 3

