

MOL #86322

Arginine Vasopressin Enhances Cell Survival Via a GRK2- β arrestin1-ERK1/2-Dependent Pathway in H9c2 Cells

Weizhong Zhu, Douglas G. Tilley, Valerie D. Myers, Ryan C. Coleman and Arthur M. Feldman

Author Affiliations:

Cardiovascular Research Center (W.Z., V.D.M., R.C.C.), Departments of Physiology (W.Z., V.D.M., R.C.C., A.M.F.), Pharmacology (D.G.T.) and Medicine (A.M.F.), and Center for Translational Medicine (D.G.T.), Temple University School of Medicine

MOL #86322

Running title: V_{1A}R-mediated protection against hypoxic stress

Address correspondence to:

Weizhong Zhu, MD, PhD
Cardiovascular Research Center
Department of Physiology
Tel: 215-707-3152
Fax: 215-707-5737
Email: weizhong.zhu@temple.edu

Douglas G. Tilley, PhD
Center for Translational Medicine
Department of Pharmacology
Tel: 215-707-9758
Fax: 215-707-9890
Email: douglas.tilley@temple.edu

Number of pages: 17
Number of tables: 0
Number of figures: 5
Number of supplemental figures: 3
Number of references: 35

Number of words in abstract: 245
Number of words in introduction: 433
Number of words in discussion: 1145

Nonstandard abbreviations: V_{1A}R (arginine vasopressin type 1A receptor); H/R (hypoxia/reoxygenation); GRK (G-protein coupled receptor kinase).

MOL #86322

Abstract

Circulating levels of arginine vasopressin (AVP) are elevated during hypovolemia and during cardiac stress. AVP activates V_{1A} - $G\alpha_q$ coupled receptors in the heart and vasculature and V_2 - $G\alpha_s$ coupled receptors in the kidney. However, little is known regarding the signaling pathways that influence the effects of V_{1A} receptor ($V_{1A}R$) activation during cellular injury. Using hypoxia-reoxygenation (H/R) as a cell injury model, we evaluated cell survival and caspase 3/7 activity in H9c2 myoblasts after treatment with AVP. Pretreatment of H9c2 cells with AVP significantly reduced H/R-induced cell death and caspase 3/7 activity, effects that were blocked via both selective $V_{1A}R$ inhibition and MEK1/2 inhibition. AVP increased ERK1/2 phosphorylation in a concentration-dependent manner that was sensitive to MEK1/2 inhibition and $V_{1A}R$ inhibition, but not $V_{1B}R$ or V_2R inhibition. Discrete elements of the V_{1A} - $G\alpha_q$ -protein kinase C (PKC) and V_{1A} -G protein-coupled receptor kinase (GRK)- β -arrestin signaling cascades were inhibited in order to dissect the pathways responsible for the protective effects of $V_{1A}R$ signaling: $G\alpha_q$ (over-expression of GqI), PKC (administration of Ro 31-82425), GRK2 (β ARKct overexpression and siRNA knockdown), GRK5 (siRNA knockdown) and β -arrestin1 (siRNA knockdown). These studies demonstrated that both $G\alpha_q$ /PKC- and GRK2/ β -arrestin1-dependent $V_{1A}R$ signaling were capable of inducing ERK1/2 phosphorylation in response to AVP stimulation. However, AVP-mediated protection against H/R was elicited only via GRK2- and β -arrestin1-dependent signaling. These results suggest that activation of the $V_{1A}R$ in H9c2 cells mediates protective signaling via a GRK2- β -arrestin1-ERK1/2-dependent mechanism that leads to decreased caspase 3/7 activity and enhanced survival under conditions of ischemic stress.

MOL #86322

Introduction

Circulating levels of arginine vasopressin (AVP), which are elevated during hypovolemia and during cardiac stress, mediate important physiological functions such as osmotic regulation, vasoconstriction, and release of adrenocorticotrophic hormone (ACTH). These physiological effects are mediated through the interaction of AVP to specific membrane G protein-coupled receptors (GPCR) in different target tissues: $V_{1A}R$ ($G\alpha_q$ -coupled; heart and vasculature), V_2R ($G\alpha_s$ -coupled; renal collecting tubule) and $V_{1B}R$ ($G\alpha_q$ -coupled; anterior pituitary), also termed V_3R (de Keyzer et al., 1994; Lolait et al., 1992; Morel et al., 1992). While administration of AVP to neonatal mouse cardiomyocytes has been reported to increase cell hypertrophy (Hiroshima et al., 2007), AVP has variable effects on the heart during injury associated with ischemia and reperfusion (Indrabarya et al., 2009; Nazari et al., 2011). Recently, we found that both constitutive and controlled cardiac-specific overexpression of $V_{1A}R$ induced the development of left ventricular hypertrophy, dilatation, diminished contractile performance and reprogramming of the heart failure (HF) gene program in transgenic mice, effects that were found to be mediated via $G\alpha_q$ protein-dependent signaling (Li et al., 2011).

In the heart, $G\alpha_q$ protein-coupled receptors induce the activation of protein kinase C (PKC) leading to cellular hypertrophy and activation of the HF gene program (Dorn and Force, 2005). $G\alpha_q$ protein-coupled receptors, including $V_{1A}R$, also undergo agonist-induced phosphorylation by PKC and G-protein coupled receptor kinase isoforms (GRK2 and GRK5) leading to internalization and desensitization of the $V_{1A}R$ (Berrada et al., 2000; Innamorati et al., 1998; Innamorati et al., 1999). The GRKs can also alter cellular physiology independent of PKC through phosphorylation-dependent recruitment of β -arrestins and subsequent activation of downstream pathways including extracellular-regulated kinase 1/2 (ERK1/2) signaling, which can act to inhibit the ubiquitous responses to cell stress including apoptosis (Dorn, 2009; Huang et al., 2012; Tilley, 2011). Indeed, in the heart, G protein-independent signaling has been shown to promote cardioprotection downstream of angiotensin II- and β -adrenergic receptor activation during conditions of myocardial stress (Kim et al., 2012; Noma et al., 2007). These findings led to the development of biased ligands; GPCR antagonists that activate unique G protein-independent pathways. However, the role of GRK/ β -arrestin-mediated $V_{1A}R$ signaling in promoting cell survival is unknown. Thus, in the present study we sought to determine the effects of $V_{1A}R$ -G protein-independent signaling on cell survival and caspase 3/7 activity in H9c2 cells exposed to the stress of hypoxia and re-oxygenation. This has become of more than an academic interest because of the recent development of V_{1A}/V_2 receptor antagonists for the treatment of patients with hyponatremia secondary to elevated levels of AVP and HF (Gheorghide et al., 2012; Rosner and Ronco, 2010; Zmily et al., 2013).

MOL #86322

Materials and Methods

Materials

Penicillin and streptomycin were purchased from Invitrogen (Gaithersburg, MD). AVP (V9879) was from Sigma-Aldrich. Ro 31-8425 (Ro-31, Cat#557514) and PD 98059 (Cat#513000) were from Calbiochem (San Diego, CA). V_{1A} receptor selective antagonist, SR45059 was from Tocris Bioscience (Minneapolis, MN, Cat#2310). V_{1B} receptor selective antagonist SSR149415 (Cat#1114) and V_2 receptor selective antagonist OPC41061 (Cat#41061) were purchased from Axon Medchem (Groningen, Netherlands). Cell viability assay kit and Caspase 3/7 assay kit were purchased from Promega (Madison, WI, USA). The hypoxia chamber was purchased from Billups-Rothenberg, Inc. (CA, USA). M199 and Opti-MEM I reduced serum medium were purchased from Invitrogen. Silencer FAM-labeled negative control siRNA (Ambion, AM4620), GRK 2 siRNA (Ambion, AM16708), GRK5 siRNA (Thermo Scientific, L-080156), and β arrestin1 siRNA (Thermo Scientific, L-080156) were from Ambion. Adenovirus containing β -galactosidase (β -gal) and GFP were gifted by Dr. Yibin Wang (University of California, Los Angeles). Gq-I-IRES-GFP and β ARK_{ct} were obtained from Dr. Satoru Eguchi and Dr. Walter Koch (Temple University), respectively. Anti-P-ERK1/2, anti-total-ERK1/2, anti- β -arrestin1/2 and anti-GAPDH antibodies were purchased from Cell Signaling. Anti-GFP, Anti-GRK2 and anti-GRK5 were from Santa Cruz.

Cell culture and adenovirus

H9c2 cells were purchased from ATCC (Rockville, MD), maintained in M199 supplemented with 10% Fetal calf serum (FCS), 100 U/ml penicillin, and 10 μ l/ml streptomycin, and grown in an atmosphere of 5% CO₂/95% humidified air at 37°C on 6-well plates, 100,000 cells seeded per well. Adenoviral infection with β ARK_{ct} versus β -galactosidase control (Zhu et al., 2001) or Gq-I-ires-GFP versus GFP control was performed at 100 MOI when cell confluence reached 60-70%. 24-36 h after infection cells were treated with the reagents under normal conditions or following hypoxia/re-oxygenation (H/R).

Transfection of siRNA

Cells (1×10^5 and 5000 cells per well in 6 well-plate and 96-wellplate, respectively) were seeded in plates incubated in M199 with 10% FBS. When cell confluent reached to 50-60%, the medium was replaced by 1800 μ l (6-well plate) or 60 μ l (96-well plate) of Opti-MEM I reduced serum medium. The negative control siRNA FAM (2.4 μ g in 100 μ l of Opti-MEM I per well in 6-well plate and 0.083 μ g in 3.2 μ l of Opti-MEM I per well in 96-well plate) and same amount of scrambled, GRK2, GRK5 and β arrestin1 siRNA were mixed with the same volume of transfection reagent diluted with Opti-MEM I. The mixture (200 μ l and 20 μ l per well in 6-well and 96-well plates, respectively) were added into the cells according to the instructions (at a final concentration

MOL #86322

of 100 nM). The protein level of GRK2, GRK5 and β arrestin1 were determined by western blot at 48-96 hours after transfection, as indicated.

Hypoxia/re-oxygenation

5000 cells per well were seeded in 96-well plates. When the cell confluence reached 80-90%, cells were starved with FCS-free medium overnight. Prior to H/R, the cells were pretreated with specific reagents and the culture cell plates were placed in the hypoxia-chamber. The chamber was sealed and aerated with 5% CO₂ and 95% Nitrogen at 3 liters per minute for 30 minutes. The cells were maintained in the hypoxic environment for 24 hrs at 37°C and then the cells were re-oxygenated for 24 hrs with 5% CO₂/95% humidified air.

Cell viability and caspase 3/7 assay

Sub-confluent cultures grown in 96-well plates with no prior interventions, or infection with adenovirus or transfection with siRNA, as indicated, were starved by withdrawing serum at least 6 h. After serum starvation, cells were pretreated with 10 pM to 0.1 μ M of vasopressin for 30 minutes in the presence or absence of pharmacological agents as indicated. Cells were then incubated in the hypoxia chamber for 24 h followed by re-oxygenation for 24 hours. The relative number of live cells and caspase 3/7 activity were measured following the manufacturer's instructions. Within each assay, triplicates were performed for each condition to attain a single n value.

Immunoblot analysis of phosphorylated ERK1/2

After cells were treated for 2-60 min with 1 nM to 10 μ M AVP, they were rapidly washed 2 times with ice-cold PBS and lysed with 250 μ l of ice-cold lysis buffer (50 mM Tris-HCl, 1 mM EGTA, 150 mM NaCl, 1% Triton X-100, 1 mM phenylmethylsulfonylflouride, 5 mg/ml leupeptin, 20 mg/ml aprotinin, 1 mM NaF, and 1 mM Na₃VO₄). After centrifugation at 13,000 x g for 10 min, equal amounts of total cell lysate (20 μ g protein) were subjected to 4-12% SDS-PAGE, followed by immunoblotting for phosphorylated ERK1/2 (1:1000), total ERK1/2 (1:1000) or GFP (1:200). T-ERK1/2 was used to normalize P-ERK1/2 responses.

Statistical Analysis

A commercial software package was used for statistical analysis (Graph Pad Software Inc). Comparison of means \pm SE was analyzed by unpaired t-test, one- or two-way ANOVA followed by Tukey's post hoc analysis, as indicated in figure legends. A value of P<0.05 was considered as statistically significant. Number of independent experiments performed in each assay is indicated in the figure legends.

MOL #86322

Results

AVP protects H9c2 cells against hypoxia/reoxygenation-induced cell death via V_{1A}R- and ERK1/2-dependent signaling

To test the ability of AVP to mediate protective signaling, we established an in vitro model of cellular stress wherein H9c2 myoblasts, derived from embryonic rat ventricle, underwent hypoxia for 24 hr followed by re-oxygenation for 24 hr (H/R). Under normoxic and serum-free conditions H9c2 cell survival decreased and caspase 3/7 activity increased, responses that were significantly enhanced when the cells underwent H/R (Figs. 1A and 1B). The addition of 10% fetal calf serum was sufficient to prevent these responses under either normoxic or H/R conditions. Since H/R performed in the absence of serum induced the most significant alteration in H9c2 cell survival and caspase 3/7 activity, we used these conditions for all subsequent experiments. As seen in Figure 1C, pretreatment of the H9c2 cells with AVP significantly attenuated the H/R-induced changes in both cell survival and caspase 3/7 activity in a concentration-dependent manner.

To confirm the role for the cardiac-expressed AVP-receptor (V_{1A}R) in mediating AVP-dependent protection against cell death, H9c2 cells underwent H/R in the presence or absence of AVP (10 nM) and the V_{1A}R selective antagonist, SR49059 (0.1 μM). Antagonism of V_{1A}R with SR49059 abolished the ability of AVP to reduce both caspase 3/7 activity and cell death (Figs. 1D and 1E). Neither V_{1B}R- nor V₂R-selective antagonists (SSR14941 and OPC41061, respectively) had an effect on AVP-dependent changes in caspase 3/7 activity or cell survival during H/R (Supplemental Figures 1A and 1B). Additionally, to assess the potential role for MEK1/2-ERK1/2 signaling in mediating the protective effect of AVP, H9c2 cells were treated with the MEK1/2 inhibitor PD98059 (10 μM) prior to H/R induction. Inhibition of MEK1/2 completely blocked the AVP-mediated effects on cell survival and caspase 3/7 activity (Figs. 1D and 1E).

AVP-mediated ERK1/2 phosphorylation dynamics

Since MEK1/2 inhibition abolished the ability of AVP to enhance cell survival during H/R, we sought to define the ERK1/2 signaling response to AVP. AVP stimulation of H9c2 cells for 5 min elicited ERK1/2 phosphorylation (P-ERK1/2) in a concentration-dependent manner with an EC₅₀ of 9.0 ± 2.0 nM (Figs. 2A and 2B). This response could only be blocked by V_{1A}R-selective antagonism, but not by V_{1B}R- or V₂R-selective antagonism. The IC₅₀ of SR49059 (V_{1A}R), SSR14941 (V_{1B}R), and OPC41061 (V₂R) for inhibition of AVP (10 nM)-induced P-ERK1/2 were 0.38 ± 0.08 , 104.4 ± 10.2 , and 374.9 ± 23.4 nM, respectively (Figs. 2C and 2D). ERK1/2 signaling often persists long after receptor stimulation, therefore we assessed the P-ERK1/2 response to AVP at both acute (2-10 min) and prolonged (30-60 min) timepoints. AVP-induced P-ERK1/2 peaked at 5 min

MOL #86322

and was sustained at ~50% peak activation even to 60 min post-stimulation (Figs. 2E and 2F). As expected, pretreatment of H9c2 cells with the MEK1/2 inhibitor PD98059 (10 μ M) significantly attenuated both the acute and prolonged P-ERK1/2 responses to AVP stimulation. Thus far, these results demonstrate that $V_{1A}R$ stimulation enhances H9c2 cell survival via MEK1/2-ERK1/2 signaling.

Inhibition of $G\alpha_q$ protein/PKC signaling blunts the AVP-induced acute P-ERK1/2 response but has no effect on the AVP-induced cell survival response

GPCR stimulation induces the simultaneous activation of both G protein-dependent and -independent signaling pathways that can each act to modulate ERK1/2 activity (Tilley, 2011). Since $V_{1A}R$ - $G\alpha_q$ protein coupling induces the activation of protein kinase C (PKC), regulating the hypertrophic gene program (Li et al., 2011), we tested whether $G\alpha_q$ protein/PKC-mediated signaling is involved in AVP-induced P-ERK1/2 and cell survival responses. Overexpression of the peptide inhibitor of $G\alpha_q$ (Gq-I-ires-GFP [GqI]) inhibited the acute P-ERK1/2 response to AVP (2 and 5 min, by $90\pm 3\%$ and $52\pm 6\%$, respectively, versus GFP control), but had no effect on the AVP-induced prolonged responses at 30 and 60 min (Figs. 3A and 3C). Further, suppression of $G\alpha_q$ protein by GqI overexpression did not impact AVP-induced effects on caspase3/7 activity or cell survival (Figs. 3D and 3E). Similarly, pretreatment of H9c2 cells with the PKC inhibitor Ro-31 (1 μ M) inhibited acute phosphorylation of ERK1/2 ($90\pm 6\%$ and $62\pm 3\%$ inhibition at 2 and 5 min, respectively), but had no effect on persistent ERK1/2 phosphorylation (30-60 min) (Fig. 3B and 3C). Pretreatment of H9c2 cells with Ro-31 prior to H/R led to a significant enhancement of H/R-induced cell death and caspase 3/7 activity (Supplemental Figure 1C). Interestingly, despite the enhanced negative effect on survival mediated by chronic inhibition of PKC by Ro-31, AVP was still capable of inducing a concentration-dependent reduction in cell death and caspase 3/7 activity (Fig. 3F). These results suggest a $G\alpha_q$ protein/PKC-independent component to $V_{1A}R$ -mediated cell survival signaling.

Overexpression of β ARKct blunts persistent AVP-induced P-ERK1/2 and abrogates AVP-mediated H9c2 cell survival

To begin to assess the contribution of G protein-independent signaling to $V_{1A}R$ -mediated P-ERK1/2 and survival responses in H9c2 cells, we overexpressed β ARKct, the c-terminal peptide of GRK2 that prevents its recruitment and activation upon GPCR stimulation, or β -galactosidase (β -gal) as a control (Koch et al., 1994; Zhu et al., 2001). Overexpression of β ARKct predominantly blunted the persistent P-ERK1/2 responses at 30 and 60min by $45\pm 2\%$ and $39\pm 4\%$, respectively, versus β -gal control (Figs. 4A and 4C). Additionally, AVP-elicited cell survival was completely abolished by overexpression of β ARKct (Figs. 4D and 4E). Comparison

MOL #86322

of the effects of the combined inhibition of AVP signaling by simultaneous treatment of H9c2 cells with Ro-31 and β ARKct were consistent with these results, as the combination of both Ro-31 and β ARKct did not further impact the cell survival responses to AVP (Supplemental Figs. 1D and 1E). These results suggest a role for GRK2-dependent $V_{1A}R$ signaling in the promotion of ERK1/2 phosphorylation and cell survival under conditions of cellular stress.

siRNA-mediated deletion of GRK2 and β -arrestin1, but not GRK5, blunts persistent AVP-induced P-ERK1/2 and cell survival responses in H9c2 cells

To confirm the role of GRK2 in mediating the effects of AVP on persistent ERK1/2 phosphorylation and survival in H9c2 cells, siRNA-mediated knockdown of GRK2 was performed versus a scrambled siRNA control. After a 48hr knockdown, GRK2 protein expression was significantly reduced by $68.5 \pm 6.5\%$ (Figs. 5A and 5D), at which timepoint the P-ERK1/2 response to AVP was assessed. Consistent with the effects of β ARKct on $V_{1A}R$ -mediated signaling, deletion of GRK2 significantly inhibited the persistent P-ERK1/2 response to AVP (Figs. 5A and 5E). After 96hr, including the H/R assay, GRK2 deletion was shown to impair the ability of AVP to promote H9c2 survival during H/R (Figs. 5F and 5G). Of note, the H/R procedure did not alter siRNA-mediated silencing of GRK2 in the presence or absence of AVP (Supplemental Figures 2A and 2B). In addition to GRK2, GRK5 is the other predominant GRK cardiac isoform that has been shown to modulate ERK1/2 activity and survival signaling (Huang et al., 2012). To determine if GRK5 contributes to G protein-independent effects of AVP on ERK1/2 phosphorylation and cell survival, we next performed GRK5 siRNA-mediated knockdown experiments. Although knockdown of GRK5 protein expression was attained to a similar extent as that of GRK2 (Figs. 5B and 5D), neither the P-ERK1/2 response (Figs. 5B and 5E) nor the caspase 3/7 activity and survival responses (Figs. 5F and 5G) to AVP were altered in comparison to the scrambled siRNA controls.

β -arrestins are recruited to GRK-phosphorylated GPCR to induce a number of cellular processes including persistent ERK1/2 activation, and have been shown to promote cardiac protection under conditions of stress (Kim et al., 2012; Luttrell et al., 2001; Noma et al., 2007). Thus, we sought to determine whether β -arrestins are required for the $V_{1A}R$ -GRK2-dependent effects on ERK1/2 and survival signaling. We began by transfecting the cells with with β -arrestin1-selective versus scrambled siRNA. 48 hours later, β -arrestin1 protein expression was significantly reduced by $79.5 \pm 3.5\%$ (Figs. 5C and 5D). As observed with GRK2 knockdown, deletion of β -arrestin1 inhibited the AVP-induced P-ERK1/2 response, again primarily at the persistent (30-60 min) timepoints (Figs 5C and 5E). Further, β -arrestin1 knockdown significantly impaired the ability of AVP to promote H9c2 cell survival and decrease caspase 3/7 activity during H/R (Figs. 5F and 5G).

MOL #86322

Via immunoblotting analysis, β -arrestin1 was shown to be the predominant β -arrestin isoform expressed in H9c2 cells (Supplemental Fig. 3A). To determine if there was a low level of β -arrestin2 expression that mediated V_{1A}R effects on ERK1/2 phosphorylation and cell survival, we also transfected H9c2 cells with siRNA directed against β -arrestin2. Perhaps due to the low/non-detectable expression of β -arrestin 2 in these cells, β -arrestin2 siRNA induced a substantial, though variable, knockdown of β -arrestin1, which was sufficient to significantly reduce AVP-mediated P-ERK1/2 and survival responses (Supplemental Fig. 3B-E), essentially recapitulating the data attained using β -arrestin1 siRNA. Altogether, these results reveal that under conditions of hypoxic stress, engagement of a V_{1A}R-GRK2- β -arrestin1-ERK1/2-mediated signaling pathway in H9c2 cells promotes survival.

Discussion

G protein-independent signaling through GRKs and β -arrestins has been shown to mediate a number of beneficial effects on cardiac function and survival during heart failure (Tilley, 2011). GRKs were initially identified as desensitizing regulators of GPCR, terminating their acute signaling responses to agonist stimulation via phosphorylation of the C-terminal tail of the receptor (Benovic et al., 1986; Hausdorff et al., 1990). It has since been clearly demonstrated that beyond desensitization, GRK also initiate G protein-independent signaling events that act to regulate ERK1/2 activity, for instance via the recruitment of the scaffolding proteins β -arrestins through differential phosphorylation of GPCRs (Heitzler et al., 2012). These events commonly promote more persistent ERK1/2 activity than that mediated by G proteins, and have been demonstrated to impact numerous cellular processes, including regulation of apoptosis and cell survival in numerous cells and, importantly, in the heart (Dorn, 2009; Huang et al., 2012; Tilley, 2011). Although relatively little is known about the pathophysiologic effects of cardiac-expressed V_{1A}R, our recent study utilizing transgenic mice with inducible cardiac-restricted overexpression of V_{1A}R was the first to conclusively demonstrate that enhanced V_{1A}R signaling in the heart itself directly leads to the development of left ventricular hypertrophy, dilatation, diminished contractile performance and reprogramming of the HF gene profile in a G α_q protein-dependent manner (Li et al., 2011), though a role for G α_q protein-independent signaling in the control of cardiac function and/or survival was not investigated.

In the present study, using H9c2 myoblasts as an *in vitro* model of hypoxia/reoxygenation, H/R, we demonstrate that AVP enhances cell survival during stress in a V_{1A}R-GRK2- β -arrestin1-ERK1/2-dependent manner. H9c2 myoblasts, derived from embryonic rat ventricle, were an excellent model system in which to directly test the effects of V_{1A}R signaling during *in vitro* stress as they express cardiac and skeletal isoforms of L-type Ca²⁺ channels, sarcolemmal ATPase splice variants characteristic of a normal heart and endogenous V_{1A}R, which respond to AVP with typical G α_q protein-coupled receptor responses (Chen and Chen, 1999; Guo

MOL #86322

et al., 2012; Hammes et al., 1994; Hescheler et al., 1991; Reilly et al., 1998; Sipido and Marban, 1991; Tran et al., 1995). During the development of the H/R model, we found initially that H/R increased caspase 3/7 activity and decreased cell viability with enhanced robustness over normoxic conditions only under serum-free conditions, whereas the presence of FBS protected the cells. This would be predicted as the presence of growth factors in the serum could act to constitutively promote cell survival and proliferation through many pathways, including activation of ERK1/2. A high level of ERK1/2 phosphorylation would also mask or blunt AVP-specific effects on ERK1/2 regulation. Since ischemia *in vivo* promotes a reduction in local oxygen and nutrient delivery, we feel the serum-free condition in our H/R model better replicates this condition. Additionally, it facilitates a controlled environment to specifically add AVP to the media alone to determine its effects on survival, without the confounding presence of other growth- and survival-promoting factors, and the underlying mechanisms, in this case through G protein-independent regulation of ERK1/2.

Our finding that V_{1A}R-mediated cell survival is mediated via G protein-independent regulation of ERK1/2 activity is consistent with other reports describing the molecular mechanisms of GPCR-induced survival signaling, including the cardiac-expressed β -adrenergic receptors (β AR) and the angiotensin type 1A receptor (AT1R) (Kim et al., 2012; Noma et al., 2007). As with the V_{1A}R, the AT1R is a G α_q -coupled receptor, and studies have shown the survival signaling response to AT1R stimulation depends not on rapid G α_q protein-mediated ERK1/2 signaling, but on prolonged GRK/ β -arrestin-mediated ERK1/2 signaling (Ahn et al., 2009; Kim et al., 2012). Our data demonstrates that this pattern of ERK1/2 activation is similar for V_{1A}R in H9c2 myoblasts, wherein G α_q protein/PKC-dependent signaling mediates the acute P-ERK1/2 response and GRK2- β -arrestin1-dependent signaling primarily regulates the persistent phase of the P-ERK1/2 response. This could account for our previous observation in the V_{1A}R transgenic mice that although inhibition of G α_q protein blocked the development of the heart failure phenotype, it only partially inhibited activation of ERK1/2 (Li et al., 2011). Thus, although ERK1/2 signaling is required for the AVP-mediated survival response, the G protein-dependent cohort did not contribute to this effect, implying the need for selective subcellular targeting of V_{1A}R-GRK2- β -arrestin1-ERK1/2 signaling in promoting survival.

A distinctive finding in our study is that GRK2, not GRK5, is responsible for driving the β -arrestin1-ERK1/2 signaling response to AVP, whereas previous studies have found that GRK2 mainly induces GPCR desensitization while other GRK isoforms, including GRK5, are responsible for mediating β -arrestin-dependent ERK1/2 signaling responses (Heitzler et al., 2012; Noma et al., 2007). This is an important finding as inhibition of GRK2 has been shown to be cardioprotective in models of heart failure (Lymperopoulos et al., 2012), but could limit the ability of V_{1A}R to exert survival signaling. In fact, in light of our findings, inhibition of GRK2 could potentially exacerbate negative V_{1A}R-mediated effects by preventing receptor desensitization and augmenting deleterious G α_q protein-dependent signaling during heart failure in which there are elevated levels

MOL #86322

of circulating AVP. Additionally, β -arrestins 1 and 2 are usually ubiquitously expressed and can have redundant or opposing effects on receptor internalization or signaling processes. Thus, our finding that H9c2 cells predominantly express β -arrestin1 with no detectable β -arrestin2 expression is unique, and although in this system β -arrestin1 clearly plays an essential role in mediating the protective effects of $V_{1A}R$ signaling, the contribution of β -arrestin2 to these effects in other cell types, such as fully differentiated adult cardiomyocytes, should be examined in a case-by-case basis.

The importance of understanding the relative contributions of both G protein-dependent and -independent pathways in the regulation of cell survival mechanisms has been recently highlighted in a number of studies investigating biased GPCR ligands, that is, ligands that have agonist-like effects but that are biased toward the activation of one particular pathway over another. A number of AT1R biased agonists that lack the ability to activate $G\alpha_q$ protein-dependent signaling but do promote GRK/ β -arrestin-dependent ERK1/2 signaling have been shown capable of regulating apoptotic pathways *in vitro* and enhancing cardiac contractility and improved cardiac performance *in vivo*, one of which is undergoing Phase II evaluation for the treatment of heart failure (Boerrigter et al., 2011; Violin et al., 2010). Late-stage heart failure is often associated with high levels of AVP that has been associated with symptomatic hyponatremia and increased mortality (Francis et al., 1990; Gheorghide et al., 2012). However, the V_2 -selective antagonist tolvaptan had no effect on survival and actually increased circulating levels of AVP (Lanfear et al., 2013) while the V_2 -selective antagonist lixivaptan was associated with increased mortality in patients hospitalized with acute heart failure and hyponatremia (Xu, 2012). The present data suggests that the use of a biased ligand that blocks both V_2R and $V_{1A}R$ - $G\alpha_q$ protein-dependent signaling, while at the same time activating $V_{1A}R$ -mediated GRK2/ β -arrestin signaling would provide a novel new therapeutic for the treatment of patients with chronic heart failure and elevated levels of AVP.

Acknowledgements

We thank Dr. Satoru Eguchi (Temple University School of Medicine) for kindly providing us the Gq-I-ires-GFP adenovirus.

Author contributions

Participated in research design: Zhu, Tilley and Myers.

Conducted experiments: Zhu, Myers and Coleman.

Performed data analysis: Zhu, Tilley, Myers and Feldman.

Wrote or contributed to the writing of the manuscript: Zhu, Tilley and Feldman.

MOL #86322

References

- Ahn S, Kim J, Hara MR, Ren XR and Lefkowitz RJ (2009) β -Arrestin-2 Mediates Anti-apoptotic Signaling through Regulation of BAD Phosphorylation. *J Biol Chem* **284**(13): 8855-8865.
- Benovic JL, Strasser RH, Caron MG and Lefkowitz RJ (1986) Beta-adrenergic receptor kinase: identification of a novel protein kinase that phosphorylates the agonist-occupied form of the receptor. *Proceedings of the National Academy of Sciences of the United States of America* **83**(9): 2797-2801.
- Berrada K, Plesnicher CL, Luo X and Thibonnier M (2000) Dynamic interaction of human vasopressin/oxytocin receptor subtypes with G protein-coupled receptor kinases and protein kinase C after agonist stimulation. *J Biol Chem* **275**(35): 27229-27237.
- Boerrigter G, Lark MW, Whalen EJ, Soergel DG, Violin JD and Burnett JC, Jr. (2011) Cardiorenal actions of TRV120027, a novel ss-arrestin-biased ligand at the angiotensin II type I receptor, in healthy and heart failure canines: a novel therapeutic strategy for acute heart failure. *Circ Heart Fail* **4**(6): 770-778.
- Chen WC and Chen CC (1999) Signal transduction of arginine vasopressin-induced arachidonic acid release in H9c2 cardiac myoblasts: role of Ca²⁺ and the protein kinase C-dependent activation of p42 mitogen-activated protein kinase. *Endocrinology* **140**(4): 1639-1648.
- de Keyser Y, Auzan C, Lenne F, Beldjord C, Thibonnier M, Bertagna X and Clauser E (1994) Cloning and characterization of the human V3 pituitary vasopressin receptor. *FEBS Lett* **356**(2-3): 215-220.
- Dorn GW, 2nd (2009) GRK mythology: G-protein receptor kinases in cardiovascular disease. *J Mol Med (Berl)* **87**(5): 455-463.
- Dorn GW, 2nd and Force T (2005) Protein kinase cascades in the regulation of cardiac hypertrophy. *J Clin Invest* **115**(3): 527-537.
- Francis GS, Benedict C, Johnstone DE, Kirlin PC, Nicklas J, Liang CS, Kubo SH, Rudin-Toretzky E and Yusuf S (1990) Comparison of neuroendocrine activation in patients with left ventricular dysfunction with and without congestive heart failure. A substudy of the Studies of Left Ventricular Dysfunction (SOLVD). *Circulation* **82**(5): 1724-1729.
- Gheorghide M, Pang PS, Ambrosy AP, Lan G, Schmidt P, Filippatos G, Konstam M, Swedberg K, Cook T, Traver B, Maggioni A, Burnett J, Grinfeld L, Udelson J and Zannad F (2012) A comprehensive, longitudinal description of the in-hospital and post-discharge clinical, laboratory, and neurohormonal course of patients with heart failure who die or are re-hospitalized within 90 days: analysis from the EVEREST trial. *Heart failure reviews* **17**(3): 485-509.
- Guo S, Olm-Shipman A, Walters A, Urciuoli WR, Devito S, Nadtochiy SM, Wojtovich AP and Brookes PS (2012) A cell-based phenotypic assay to identify cardioprotective agents. *Circ Res* **110**(7): 948-957.
- Hammes A, Oberdorf S, Strehler EE, Stauffer T, Carafoli E, Vetter H and Neyses L (1994) Differentiation-specific isoform mRNA expression of the calmodulin-dependent plasma membrane Ca(2+)-ATPase. *FASEB J* **8**(6): 428-435.
- Hausdorff WP, Caron MG and Lefkowitz RJ (1990) Turning off the signal: desensitization of beta-adrenergic receptor function. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* **4**(11): 2881-2889.
- Heitzler D, Durand G, Gallay N, Rizk A, Ahn S, Kim J, Violin JD, Dupuy L, Gauthier C, Piketty V, Crepieux P, Poupon A, Clement F, Fages F, Lefkowitz RJ and Reiter E (2012) Competing G protein-coupled receptor kinases balance G protein and beta-arrestin signaling. *Molecular systems biology* **8**: 590.
- Hescheler J, Meyer R, Plant S, Krautwurst D, Rosenthal W and Schultz G (1991) Morphological, biochemical, and electrophysiological characterization of a clonal cell (H9c2) line from rat heart. *Circ Res* **69**(6): 1476-1486.
- Hiroshima M, Wang S, Aoyagi T, Oikawa R, Sanbe A, Takeo S and Tanoue A (2007) Vasopressin promotes cardiomyocyte hypertrophy via the vasopressin V1A receptor in neonatal mice. *Eur J Pharmacol* **559**(2-3): 89-97.
- Huang ZM, Gold JJ and Koch WJ (2012) G protein-coupled receptor kinases in normal and failing myocardium. *Front Biosci* **17**: 3047-3060.

MOL #86322

- Indrambarya T, Boyd JH, Wang Y, McConechy M and Walley KR (2009) Low-dose vasopressin infusion results in increased mortality and cardiac dysfunction following ischemia-reperfusion injury in mice. *Crit Care* **13**(3): R98.
- Innamorati G, Sadeghi H and Birnbaumer M (1998) Transient phosphorylation of the V1a vasopressin receptor. *J Biol Chem* **273**(12): 7155-7161.
- Innamorati G, Sadeghi H and Birnbaumer M (1999) Phosphorylation and recycling kinetics of G protein-coupled receptors. *Journal of receptor and signal transduction research* **19**(1-4): 315-326.
- Kim KS, Abraham D, Williams B, Violin JD, Mao L and Rockman HA (2012) beta-Arrestin-biased AT1R stimulation promotes cell survival during acute cardiac injury. *American journal of physiology Heart and circulatory physiology* **303**(8): H1001-1010.
- Koch WJ, Hawes BE, Inglese J, Luttrell LM and Lefkowitz RJ (1994) Cellular expression of the carboxyl terminus of a G protein-coupled receptor kinase attenuates G beta gamma-mediated signaling. *J Biol Chem* **269**(8): 6193-6197.
- Lanfear DE, Sabbah HN, Goldsmith SR, Greene SJ, Ambrosy AP, Fought AJ, Kwasny MJ, Swedberg K, Yancy CW, Konstam MA, Maggioni AP, Zannad F, Gheorghiade M and on behalf of the Eti (2013) Association of Arginine Vasopressin Levels With Outcomes and the Effect of V2 Blockade in Patients Hospitalized for Heart Failure With Reduced Ejection Fraction: Insights From the EVEREST Trial. *Circulation Heart failure* **6**(1): 47-52.
- Li X, Chan TO, Myers V, Chowdhury I, Zhang XQ, Song J, Zhang J, Andrel J, Funakoshi H, Robbins J, Koch WJ, Hyslop T, Cheung JY and Feldman AM (2011) Controlled and cardiac-restricted overexpression of the arginine vasopressin V1A receptor causes reversible left ventricular dysfunction through Galphaq-mediated cell signaling. *Circulation* **124**(5): 572-581.
- Lolait SJ, O'Carroll AM, McBride OW, Konig M, Morel A and Brownstein MJ (1992) Cloning and characterization of a vasopressin V2 receptor and possible link to nephrogenic diabetes insipidus. *Nature* **357**(6376): 336-339.
- Luttrell LM, Roudabush FL, Choy EW, Miller WE, Field ME, Pierce KL and Lefkowitz RJ (2001) Activation and targeting of extracellular signal-regulated kinases by beta-arrestin scaffolds. *Proc Natl Acad Sci U S A* **98**(5): 2449-2454.
- Lymperopoulos A, Rengo G and Koch WJ (2012) GRK2 inhibition in heart failure: something old, something new. *Current pharmaceutical design* **18**(2): 186-191.
- Morel A, O'Carroll AM, Brownstein MJ and Lolait SJ (1992) Molecular cloning and expression of a rat V1a arginine vasopressin receptor. *Nature* **356**(6369): 523-526.
- Nazari A, Sadr SS, Faghihi M, Imani A and Moghimian M (2011) The cardioprotective effect of different doses of vasopressin (AVP) against ischemia-reperfusion injuries in the anesthetized rat heart. *Peptides* **32**(12): 2459-2466.
- Noma T, Lemaire A, Naga Prasad SV, Barki-Harrington L, Tilley DG, Chen J, Le Corvoisier P, Violin JD, Wei H, Lefkowitz RJ and Rockman HA (2007) Beta-arrestin-mediated beta1-adrenergic receptor transactivation of the EGFR confers cardioprotection. *J Clin Invest* **117**(9): 2445-2458.
- Reilly BA, Brostrom MA and Brostrom CO (1998) Regulation of protein synthesis in ventricular myocytes by vasopressin. The role of sarcoplasmic/endoplasmic reticulum Ca²⁺ stores. *J Biol Chem* **273**(6): 3747-3755.
- Rosner MH and Ronco C (2010) Hyponatremia in heart failure: the role of arginine vasopressin and its antagonism. *Congestive heart failure* **16 Suppl 1**: S7-14.
- Sipido KR and Marban E (1991) L-type calcium channels, potassium channels, and novel nonspecific cation channels in a clonal muscle cell line derived from embryonic rat ventricle. *Circ Res* **69**(6): 1487-1499.
- Tilley DG (2011) G protein-dependent and G protein-independent signaling pathways and their impact on cardiac function. *Circ Res* **109**(2): 217-230.
- Tran K, Zha X, Chan M and Choy PC (1995) Enhancement of phospholipid hydrolysis in vasopressin-stimulated BHK-21 and H9c2 cells. *Mol Cell Biochem* **151**(1): 69-76.

MOL #86322

Violin JD, DeWire SM, Yamashita D, Rominger DH, Nguyen L, Schiller K, Whalen EJ, Gowen M and Lark MW (2010) Selectively engaging beta-arrestins at the angiotensin II type 1 receptor reduces blood pressure and increases cardiac performance. *J Pharmacol Exp Ther* **335**(3): 572-579.

Xu NN (2012) Clinical Review: Lixivaptan (<http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/CardiovascularandRenalDrugsAdvisoryCommittee/UCM318867.pdf>), (Committee CaRDA ed), FDA, Silver Spring MD.

Zhu WZ, Zheng M, Koch WJ, Lefkowitz RJ, Kobilka BK and Xiao RP (2001) Dual modulation of cell survival and cell death by beta(2)-adrenergic signaling in adult mouse cardiac myocytes. *Proc Natl Acad Sci U S A* **98**(4): 1607-1612.

Zmily HD, Alani A and Ghali JK (2013) Evaluation of lixivaptan in euvoletic and hypervolemic hyponatremia and heart failure treatment. *Expert opinion on drug metabolism & toxicology* **9**(5): 645-655.

MOL #86322

Footnotes

This work was supported by the National Institutes of Health National Heart, Lung and Blood Institute [Grants HL105414 and HL091799].

MOL #86322

Figure Legends

Fig. 1. AVP protects H9c2 cells against hypoxia/reoxygenation-induced cell death via V_{1A}R- and ERK1/2-dependent signaling. (A) Caspase 3/7 activity in response to serum starvation under conditions of H/R or normoxia, **P<0.01 vs starvation + normoxia, ††P<0.01 vs FBS in same group, one-way ANOVA. (B) Cell survival in response to serum starvation under conditions of H/R or normoxia, *P<0.05 vs starvation + normoxia, †P<0.05 vs FBS in same group, one-way ANOVA. (C) AVP concentration-dependently increased cell survival and decreased caspase 3/7 activity under conditions of H/R, *P<0.05 for cell survival, **P<0.01 for caspase 3/7 activity, repeated two-way ANOVA. Pretreatment of cells with SR49059 (0.1 μM) or PD98059 (10 μM) for 30 min blunted effect of AVP on caspase 3/7 activity (D) and cell survival (E), *P<0.05 vs vehicle (PBS)-stimulated control, one-way ANOVA. All data are presented as the mean ± SEM from 3 independent experiments.

Fig. 2. V_{1A}R, not V_{1B}R or V₂R, mediates ERK1/2 phosphorylation in H9c2 cells. (A, B) An AVP concentration-P-ERK1/2 response analysis was performed in H9c2 cells, stimulation for 5 min, as determined via immunoblotting. (C, D) Inhibitory concentration-responses for V_{1A}R-, V_{1B}R- or V₂R-selective antagonists (30 min pretreatment at 1nM-10μM) against AVP (10nM, 5 min treatment)-induced P-ERK1/2 were generated in H9c2 cells. (E, F) MEK1/2 inhibition with PD98059 (10 μM) blunted the AVP-dependent phosphorylation of ERK1/2 at all timepoints analyzed, **P<0.01 vs control at the same timepoints, unpaired t-test. All data are presented as the mean ± SEM from 3-4 independent experiments.

Fig 3. Inhibition of Gα_q protein/PKC signaling blunts the AVP-induced acute P-ERK1/2 response but has no effect on the AVP-induced cell survival response. (A) Overexpression of GqI-GFP (vs GFP alone) with adenovirus or (B) inhibition of PKC with Ro-31 (1 μM, vs 0.1% DMSO vehicle) significantly blunted the acute (2-10 min), but not prolonged (30-60 min) phosphorylation of ERK1/2 evoked by AVP (10 nM) in H9c2 cells. (C) The averaged data of A and B, *P<0.05, **P<0.01 GqI vs GFP at the same timepoints, and †P<0.05, ††P<0.01 Ro-31 vs vehicle at the same timepoints, unpaired t-test. Overexpression of GqI failed to block the AVP-mediated decrease in caspase 3/7 activity (D) and increase in cell survival (E), **p<0.01 vs appropriate vehicle (PBS)-stimulated control, one-way ANOVA. (F) Pretreatment of H9c2 cells with Ro-31 (1 μM) did not impact the ability of AVP to protect cells from H/R, as AVP concentration-dependently decreased caspase 3/7 activity and increased cell survival. *P<0.05 for cell survival, **P<0.01 for caspase 3/7 activity, repeated two-way ANOVA. All data are presented as the mean ± SEM from at least 3 independent experiments.

MOL #86322

Fig. 4 Overexpression of β ARKct blunts persistent AVP-induced P-ERK1/2 and abrogates AVP-mediated H9c2 cell survival. (A) Overexpression of β ARKct (versus β -gal) significantly blunted the ERK1/2 phosphorylation response (from 10-60 min) following AVP stimulation. (B) The averaged data of A, *P<0.05, **P<0.01 vs the same timepoint, unpaired t-test. Overexpression of β ARKct significantly blocked the AVP-dependent decrease in caspase 3/7 activity (D) and increase in cell survival (E), *P<0.05 vs vehicle (PBS)-stimulated β -gal control, one-way ANOVA. All data are presented as the mean \pm SEM from 3 independent experiments.

Fig. 5 siRNA-mediated deletion of GRK2 and β -arrestin1, but not GRK5, blunts persistent AVP-induced P-ERK1/2 and cell survival responses in H9c2 cells. Representative immunoblots showing GRK2 (A), GRK5 (B) and β -arrestin1 (C) protein expression after siRNA transfection and the corresponding ERK1/2 phosphorylation responses to AVP stimulation (10nM, 0-60 min) in H9c2 cells. (D) The averaged data of GRK2, GRK5, or β -arrestin1 knockdown, compared to scrambled siRNA, **P<0.01 vs scrambled siRNA control, one-way ANOVA. (E) Summary of the effects of siRNA-mediated deletion of GRK2, GRK5 and β arr1 on the P-ERK1/2 response to AVP (10nM, 0-60 min), **P<0.01 (GRK2 siRNA), ††P<0.01 (β arr1 siRNA) vs scrambled siRNA at the same timepoint, unpaired t-test. The effects of siRNA-mediated knockdown of GRK2, GRK5 or β arr1 on AVP-mediated caspase 3/7 activity (F) and cell survival (G) during H/R in H9c2 cells are shown. *P<0.05 GRK2 siRNA vs Scr siRNA (cell survival, caspase 3/7 activity), ††P<0.01 β arr1 siRNA vs Scr siRNA (caspase 3/7 activity) and †P<0.05 β arr1 siRNA vs Scr siRNA (cell survival), repeated two-way ANOVA. All data are presented as mean \pm SEM of 3 independent experiments.




Fig. 1









A**B****C****D****E****F**

Fig. 2


A




B




C




D







E




F




A**B****D****E**


A




B




C




D



E



F



G

