

Accelerated Communication

## 5-Iodo-A-85380, an $\alpha 4\beta 2$ Subtype-Selective Ligand for Nicotinic Acetylcholine Receptors<sup>1</sup>

ALEXEY G. MUKHIN, DANIELA GÜNDISCH, ANDREW G. HORTI, ANDREI O. KOREN, GILLES TAMAGNAN, ALANE S. KIMES, JOANN CHAMBERS, D. BRUCE VAUPEL, SARAH L. KING, MARINA R. PICCIOTTO, ROBERT B. INNIS, and EDYTHE D. LONDON

Brain Imaging Center, Intramural Research Program, National Institute on Drug Abuse, Baltimore, Maryland (A.G.M., D.G., A.G.H., A.O.K., A.S.K., J.C., D.B.V., E.D.L.) and Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut (G.T., S.L.K., M.R.P., R.B.I.)

Received September 29, 1999; accepted December 10, 1999

This paper is available online at <http://www.molpharm.org>

### ABSTRACT

In an effort to develop selective radioligands for in vivo imaging of neuronal nicotinic acetylcholine receptors (nAChRs), we synthesized 5-iodo-3-(2(S)-azetidylmethoxy)pyridine (5-iodo-A-85380) and labeled it with <sup>125</sup>I and <sup>123</sup>I. Here we present the results of experiments characterizing this radioiodinated ligand in vitro. The affinity of 5-[<sup>125</sup>I]iodo-A-85380 for  $\alpha 4\beta 2$  nAChRs in rat and human brain is defined by  $K_d$  values of 10 and 12 pM, respectively, similar to that of epibatidine (8 pM). In contrast to epibatidine, however, 5-iodo-A-85380 is more selective in binding to the  $\alpha 4\beta 2$  subtype than to other nAChR subtypes. In rat adrenal glands, 5-iodo-A-85380 binds to nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits with 1/1000th the affinity of epibatidine, and exhibits 1/60th and 1/190th the affinity of epibatidine at  $\alpha 7$  and

muscle-type nAChRs, respectively. Moreover, unlike epibatidine and cytisine, 5-[<sup>125</sup>I]iodo-A-85380 shows no binding in any brain regions in mice homozygous for a mutation in the  $\beta 2$  subunit of nAChRs. Binding of 5-[<sup>125</sup>I]iodo-A-85380 in rat brain is reversible, and is characterized by high specificity and a slow rate of dissociation of the receptor-ligand complex ( $t_{1/2}$  for dissociation  $\sim 2$  h). These properties, along with other features observed previously in in vivo experiments (low toxicity, rapid penetration of the blood-brain barrier, and a high ratio of specific to nonspecific binding), suggest that this compound, labeled with <sup>125</sup>I or <sup>123</sup>I, is superior to other radioligands available for in vitro and in vivo studies of  $\alpha 4\beta 2$  nAChRs, respectively.

Nicotinic acetylcholine receptors (nAChRs) are excitatory ligand-gated cation channels that are widely distributed in mammalian organisms, appearing in the central and peripheral nervous systems, neuromuscular junctions, and adrenal glands. The nAChR channel complex is composed of five protein subunits, which form a pore that is permeable to Na<sup>+</sup>, K<sup>+</sup>, and Ca<sup>2+</sup> (Lindstrom, 1995; Holladay et al., 1997).

To date,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ , and  $\epsilon$  subunits have been isolated and cloned from mammalian and avian tissues, with nine varieties of  $\alpha$  and four varieties of  $\beta$  subunits identified. The  $\alpha 1$ ,  $\beta 1$ ,  $\gamma$ ,  $\delta$ , and  $\epsilon$  subunits form the neuromuscular junction receptor, the very first nAChR to be characterized. The other subunits ( $\alpha 2$ - $\alpha 9$  and  $\beta 2$ - $\beta 4$ ) are found predominantly throughout the nervous system (Lindstrom, 1995; Holladay et al., 1997). This subunit diversity affords a large potential for a variety of nAChR subtypes, exhibiting distinct cation-conducting properties and pharmacological heterogeneity.

The research described in this publication was supported by the Intramural Research Program of the National Institute on Drug Abuse, funding from the Office of the National Drug Control Policy, and funding from Department of Veterans Affairs (Connecticut/Massachusetts VA Mental Illness Research, Education, and Clinical Center). Samples of postmortem brain tissue were obtained from the Brain and Tissue Bank for Developmental Disorders under National Institutes of Health Contract no. NO1-HD-1-3138.

<sup>1</sup> A preliminary report of this work has been presented previously (Mukhin AG, Gündisch D, Horti AG, Koren AO, Kimes AS, Vaupel DB and London ED (1998) 5-Iodo-A-85380—a novel highly selective ligand for the  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors. *Soc Neurosci Abstr* 24:85).

Based on binding properties and pharmacological sensitivity, major nAChR subtypes in mammalian brain can be categorized as  $\alpha$ -bungarotoxin-sensitive ( $\alpha 7$ ) and  $\alpha$ -bungarotoxin-insensitive (e.g.,  $\alpha 4\beta 2$ ) (Lindstrom, 1995; Holladay et al., 1997). Accordingly, <sup>125</sup>I- $\alpha$ -bungarotoxin has been the radioligand of choice for in vitro characterization of the  $\alpha 7$  subtype of nAChR, whereas tritiated agonists, such as nico-

**ABBREVIATIONS:** nAChR, nicotinic acetylcholine receptor; IPH, ( $\pm$ )-*exo*-2-(2-iodo-5-pyridyl-7-azabicyclo[2.2.1]heptane; A-85380, 3-(2(S)-azetidylmethoxy)pyridine; HSS, HEPES-salt solution;  $\alpha$ -BTX,  $\alpha$ -bungarotoxin; DFP, diisopropyl fluorophosphate.

tine, acetylcholine, *N*-methylcarbamylcholine, cytisine, and epibatidine, have been used to study nAChRs of the latter group in vitro (Holladay et al., 1997). Of these ligands, epibatidine has the highest known affinity for  $\alpha 4\beta 2$  nAChRs, and outstanding in vitro binding characteristics (high specific-to-nonspecific binding ratio and slow kinetics of dissociation) (Dukat et al., 1993; Houghtling et al., 1995; Flores et al., 1996; Holladay et al., 1997; Stauderman et al., 1998; Xiao et al., 1998).

Radioligands developed for noninvasive in vivo imaging of  $\alpha$ -bungarotoxin-insensitive nAChRs have exhibited shortcomings, such as poor subtype selectivity and high levels of nonspecific binding (Nybäck et al., 1994). The radioligands [ $^{123}\text{I}$ ]IPH and [ $^{125}\text{I}$ ]IPH (( $\pm$ )-*exo*-2-(2-[ $^{123/125}\text{I}$ ]iodo-5-pyridyl)-7-azabicyclo[2.2.1]heptane), recently developed iodinated analogs of epibatidine, do not distinguish well between the  $\alpha 4\beta 2$  subtype and nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits (Dávila-García et al., 1997), much like epibatidine itself (Flores et al., 1996; Xiao et al., 1998). In contrast to  $\alpha 4\beta 2$  nAChRs, nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits, possibly in combination with  $\alpha 5$  subunits, are distributed mostly in the peripheral nervous system and adrenal glands (Holladay et al., 1997). Therefore, high affinity for the latter receptors could contribute to the untoward cardiovascular effects of epibatidine and its analogs (Molina et al., 1997; Horti et al., 1998) and might limit the use of epibatidine-based compounds for imaging nAChRs in human subjects.

Recently, 3-(2(*S*)-azetidinylmethoxy)pyridine (A-85380, Fig. 1) has been identified as a high-affinity nAChR ligand (Abreo et al., 1996). Subsequently, a chloro analog of A-85380, ABT-594 (Fig. 1), has been developed as a promising nonopioid analgesic having affinity for  $\alpha 4\beta 2$  nAChRs comparable to that of epibatidine, but lacking its toxicity (Bannon et al., 1998). In a search for improved radioligands suitable for noninvasive in vivo imaging of nAChRs, chemists in our group synthesized several halogenated analogs of A-85380 (Koren et al., 1998). Some of these compounds, particularly 5-iodo-A-85380 (Fig. 1), exhibited extremely high affinity for nAChRs in rat brain (Koren et al., 1998).

Initial evaluation of 5-[ $^{125}\text{I}$ ]iodo- and 5-[ $^{123}\text{I}$ ]iodo-A-85380 in vivo in mice (Musachio et al., 1998; Vaupel et al., 1998) and 5-[ $^{123}\text{I}$ ]iodo-A-85380 in rhesus monkey (Chefer et al., 1998) and baboon (Musachio et al., 1999) demonstrated that these radioligands readily crossed the blood-brain barrier, bound to cerebral nAChRs with high specificity, and had low toxicity. Here, we present an in vitro characterization of 5-[ $^{125}\text{I}$ ]iodo-A-85380, indicating that this ligand possesses excellent properties as a probe for studying the  $\alpha 4\beta 2$  nAChR subtype.

5-Iodo-A-85380 features high affinity for nAChRs, low non-specific binding, slow dissociation from the receptor, and exceptionally high selectivity for the  $\alpha 4\beta 2$  subtype among the major mammalian nAChR subtypes. These properties, together with the results of in vivo studies with this ligand (Chefer et al., 1998; Musachio et al., 1998, 1999; Vaupel et al., 1998), suggest that 5-[ $^{123}\text{I}$ ]iodo-A-85380 may have exceptional potential as a radioligand for in vivo imaging of  $\alpha 4\beta 2$  nAChRs with single photon emission computed tomography.

## Experimental Procedures

**Materials.** 5-[ $^{123}\text{I}$ ]iodo-A-85380 and 5-[ $^{125}\text{I}$ ]iodo-A-85380 were prepared according to the literature procedures (Musachio et al., 1998; Horti et al., 1999). Specific activity of 5-[ $^{125}\text{I}$ ]iodo-A-85380 was determined as described previously (Horti et al., 1999). On the day of each synthesis, the specific activities of the three batches of 5-[ $^{125}\text{I}$ ]iodo-A-85380 used in these studies were 1550, 1980, and 2200 Ci/mmol, respectively. [ $^{125}\text{I}$ ]- $\alpha$ -Bungarotoxin ( $^{125}\text{I}$ - $\alpha$ -BTX, 100 Ci/mmol), [ $^3\text{H}$ ]cytisine (32 Ci/mmol), and ( $\pm$ )-[ $^3\text{H}$ ]epibatidine (48 Ci/mmol) were obtained from New England Nuclear Corp. (Boston, MA). 5-Iodo-A-85380 was prepared by a published method (Koren et al., 1998). ( $\pm$ )-*exo*-2-(2-Iodo-5-pyridyl)-7-azabicyclo[2.2.1]heptane (IPH), a gift from Dr. Kellar, was synthesized by Dr. J. L. Musachio at the Johns Hopkins University as described previously (Musachio et al., 1997). 3-(2(*S*)-Azetidinylmethoxy)pyridine dihydrochloride (A-85380),  $\alpha$ -bungarotoxin ( $\alpha$ -BTX), ( $-$ )- and ( $\pm$ )-epibatidine, ( $-$ )-cytisine, (+)- and ( $-$ )-stereoisomers of nicotine, acetylcholine, carbachol, dihydro- $\beta$ -erythroidine, curare, mecamylamine, atropine, naloxone, (*R*)-( $-$ )-apomorphine, and haloperidol were purchased from Research Biochemicals International (Natick, MA). Physostigmine, diisopropyl fluorophosphate (DFP), and all other chemicals used were purchased from Sigma Chemical Co. (St. Louis, MO). Male Fischer-344 and Sprague-Dawley rats were obtained from Charles River Breeding Laboratories (Wilmington, MA). Rats, shipped at the age of 12 weeks, were housed in a temperature- and light-controlled vivarium for at least 2 weeks before being used for this study. Mice were generated by mating parents heterozygous for a mutation in the  $\beta 2$  nAChR subunit (Picciotto et al., 1995). Frozen *Torpedo californica* electric organ tissue was purchased from Marinus Inc. (Long Beach, CA). Frozen samples of postmortem tissue of human cerebral cortex (four subjects, 38 to 49 years of age, death from arteriosclerotic cardiovascular disease) were obtained from the Brain and Tissue Bank for Developmental Disorders (Baltimore, MD).

All animal procedures performed at the National Institute on Drug Abuse Brain Imaging Center were approved by the National Institute on Drug Abuse Institutional Animal Care and Use Committee and were in accordance with the Guide for the Care and Use of Laboratory Animals, as endorsed by the National Institutes of Health. All animal procedures performed at the Yale University School of Medicine were approved by the Yale Animal Care and Use Committee.

**Membrane Preparation.** After  $\text{CO}_2$  euthanasia and decapitation of the rats, brains were removed and prepared as follows. Brain tissue used for binding studies was obtained by a single cut just behind the inferior colliculi to exclude the cerebellum and medulla. In some experiments, specific brain regions and the adrenal glands were isolated. Frozen samples of *Torpedo californica* electric organ and postmortem human cerebral cortical tissue were thawed at room temperature for 30 to 60 min before membrane preparation. Total membrane fractions from all tissues were isolated by homogenization of the respective tissue with a Brinkmann Polytron homogenizer in 10 to 20 volumes of a HEPES-salt solution (HSS), containing HEPES (pH 7.4, 15 mM), 120 mM NaCl, 5.4 mM KCl, 0.8 mM  $\text{MgCl}_2$ , and 1.8 mM  $\text{CaCl}_2$ , followed by centrifugation at 40,000g for 10 min. The pellets were washed twice with HSS through rehomogenization and centrifugation at the same settings. Three additional washings




Fig. 1. Chemical structures of A-85380 and two of its analogs.

were performed in the case of the total membrane fraction from rat adrenal glands and *Torpedo californica* electric organ. Crude membrane fractions (P2) were isolated as described previously (Koren et al., 1998), and were stored in aliquots at  $-70^{\circ}\text{C}$  for at least 16 h but not more than 4 weeks before use. On the day of assay, pellets were thawed, homogenized in 30 volumes of HSS, and centrifuged at 40,000g for 10 min. The resultant pellets were resuspended in a freshly prepared HSS and used for binding assays.

**Binding Assays.** Assays were carried out in HSS at  $22^{\circ}\text{C}$  unless otherwise specified. Incubations were performed in polystyrene tubes except for the assays with  $^{125}\text{I}$ - $\alpha$ -BTX, for which borosilicate glass tubes were used. The HSS for the studies of membranes from *T. californica* electric organ contained 0.1% of BSA. Nonspecific binding was determined in the presence of  $300\ \mu\text{M}$  (-)-nicotine except for the assays with  $^{125}\text{I}$ - $\alpha$ -BTX, for which  $1\ \mu\text{M}$   $\alpha$ -BTX was used instead (see the figure legends for other specific conditions for particular binding assays). Incubation was terminated by filtration through Whatman GF/B glass fiber filters, presoaked in 1% polyethyleneimine, using a Brandel 48-channel cell harvester. Filters were washed three times with 3-ml aliquots of a rinse buffer (50 mM Tris  $\cdot$  HCl, pH 7.4). In  $^{125}\text{I}$ - $\alpha$ -BTX assays, the rinse buffer also contained 1% of nonfat dry milk to reduce nonspecific binding to filter material. Radioactivity was measured using a Beckman LS 3801 liquid scintillation counting system (efficiency 43%) or a LKB Wallac 1277 gamma counter (efficiency 68%).

**In Vitro Autoradiography.** Sagittal slices (20  $\mu\text{m}$  thick) at 0.4, 0.9, 1.4, 1.9, 3.9, and 4.2 mm lateral to midline from eight frozen Fischer-344 rat brains were obtained by sectioning in a cryostat at  $-20^{\circ}\text{C}$ , and were thaw-mounted onto gelatin-coated slides. Sections were refrozen and kept frozen at  $-80^{\circ}\text{C}$  until the day of the assay. Slices were preincubated for 20 min in 50 mM Tris  $\cdot$  HCl buffer (pH 7.0) containing 120 mM NaCl, 5 mM KCl, 2.5 mM  $\text{CaCl}_2$ , and 1 mM  $\text{MgCl}_2$ . Thereafter, they were incubated with 210 pM 5- $^{125}\text{I}$ iodo-A-85380 (specific activity 435 Ci/mmol) in the same buffer for 2 h at  $25^{\circ}\text{C}$ , then rinsed twice in ice-cold buffer for 5 min each and once in distilled water for 1 min. The slides were dried overnight in a vacuum desiccator. Nonspecific binding was assessed in adjacent slices incubated in 210 pM 5- $^{125}\text{I}$ iodo-A-85380 containing 10  $\mu\text{M}$  nicotine bitartrate. Slices and appropriate  $^{125}\text{I}$ -standards were apposed to  $^3\text{H}$ -Ultrafilm for 2 days at room temperature. The resulting autoradiograms were digitized using a video camera-based system. The digitized images were analyzed using a computer program (IN-QUIRY, Loats Associates, Inc., Westminster, MD).

Brain slices (12  $\mu\text{m}$ ) from 2- to 4-month-old mice were prepared and processed using a procedure similar to that used for rat slices, but the incubation with radioligands was carried for 30 min in 50 mM Tris  $\cdot$  HCl (pH 7.4). Details of the procedure were described previously (Perry and Kellar, 1995; Zoli et al., 1998). In each experiment, sections from three  $\beta 2^{+/+}$  and three  $\beta 2^{-/-}$  mice were run in

parallel. The brain sections were incubated with 200 pM [ $^{125}\text{I}$ ]IPH (2200 Ci/mmol) or with 200 pM 5- $^{125}\text{I}$ iodo-A-85380 (200 Ci/mmol). Slides were apposed to  $^3\text{H}$ -Hyperfilm for 2 days ([ $^{125}\text{I}$ ]IPH) or for 5 days (5- $^{125}\text{I}$ iodo-A-85380) at room temperature. In addition, slides from  $\beta 2^{-/-}$  mice, labeled with 5- $^{125}\text{I}$ iodo-A-85380, were apposed to  $^3\text{H}$ -Hyperfilm for 5 weeks.


**Protein Assay.** Protein measurements were performed using a dye reagent kit (Bio-Rad, Richmond, CA) and BSA as a standard.

**Data Analysis.** Saturation binding data were subjected to Scatchard and linear regression analyses. Competition binding data were analyzed using nonlinear regression methods. Values of  $K_i$  were derived from the measured  $\text{IC}_{50}$  and  $K_d$  values for radioactive ligands using the Cheng-Prusoff equation  $K_i = \text{IC}_{50}/(1 + F/K_d)$ , where F is the concentration of unbound radioligand. The  $K_d$  values were obtained from three to six independent experiments performed on the same membrane preparations that were used for the competition assays. Results of the kinetic experiments were analyzed using semilogarithmic plots and linear regression analysis. The values of equilibrium constant of dissociation,  $K_d$ , obtained from the kinetic studies, were calculated by the equation  $K_d = k_{\text{diss}}/k_{\text{ass}}$ , where  $k_{\text{diss}}$  and  $k_{\text{ass}}$  are the dissociation and association rate constants, respectively.

## Results and Discussion

**Kinetic and Equilibrium Binding Characteristics.** The specific binding of 5- $^{125}\text{I}$ iodo-A-85380, determined in rat brain membranes at  $22^{\circ}\text{C}$  and at a ligand concentration of 10 pM, reached one half the maximal (equilibrium) binding level in  $67 \pm 9$  min (Fig. 2a). The binding was completely reversible and was characterized by a very slow dissociation ( $t_{1/2} = 132 \pm 9$  min) (Fig. 2b). The rate constants of association ( $k_{\text{ass}}$ ) and dissociation ( $k_{\text{diss}}$ ) were  $(5.6 \pm 1.4) \cdot 10^{-4}$  pM/min and  $(54 \pm 4) \cdot 10^{-4}$ /min, respectively. The  $K_d$  value, calculated as the ratio of  $k_{\text{diss}}$  to  $k_{\text{ass}}$ , was  $9.7 \pm 1.8$  pM.

Based on these data, subsequent equilibrium binding studies with 5- $^{125}\text{I}$ iodo-A-85380 were performed using a 4-h incubation at  $22^{\circ}\text{C}$ . At the lowest concentration of 5- $^{125}\text{I}$ iodo-A-85380 (ca. 1 pM) used in saturation studies, radioligand depletion of up to 30% was observed. To account for this depletion, the concentrations of free radioligand at equilibrium were calculated by reducing the concentration of total added radioactivity by the concentration of total bound radioactivity. The specific binding of 5- $^{125}\text{I}$ iodo-A-85380 in rat brain was saturable and was represented by a single population of binding sites over a radioligand concentration range of 1 to 500 pM (Fig. 3). The binding parameters ( $K_d$  and  $B_{\text{max}}$ ) were  $10.6 \pm 0.3$  pM and



**Fig. 2.** Kinetics of ligand-receptor association (A) and dissociation (B) for 5- $^{125}\text{I}$ iodo-A-85380 in rat brain. In association experiments, rat brain P2 membrane fractions (4–5  $\mu\text{g}$  of protein) were incubated in a total volume of 0.4 ml with 10 pM 5- $^{125}\text{I}$ iodo-A-85380 at  $22^{\circ}\text{C}$ . In dissociation experiments, (-)-nicotine at final concentration of 1 mM was added to rat brain P2 membrane fractions preincubated with 10 pM 5- $^{125}\text{I}$ iodo-A-85380 for 4 h at  $22^{\circ}\text{C}$ . Each graph represents results of a single experiment performed in triplicate (S.E.M. < 10%). Similar results were obtained in three additional experiments for both association and dissociation studies. The mean  $\pm$  S.E. values from the four corresponding experiments were:  $t_{1/2\text{ass}}$   $67 \pm 9$  min [ $k_{\text{ass}}$   $(5.6 \pm 1.4) \cdot 10^{-4}$  /min/pM], and  $t_{1/2\text{diss}}$   $132 \pm 9$  min [ $k_{\text{diss}}$   $(5.4 \pm 0.4) \cdot 10^{-3}$  /min].

$3.8 \pm 0.6$  pmol/g tissue ( $160 \pm 25$  fmol/mg protein) in Fischer-344 rat forebrain (Fig. 3a), and  $10.0 \pm 0.2$  pM and  $3.9 \pm 0.2$  pmol/g tissue ( $178 \pm 6$  fmol/mg protein) in Sprague-Dawley rat forebrain (Fig. 3b), respectively.

The observed  $K_d$  values in saturation studies with 5- $^{125}$ I]iodo-A-85380 agreed with both the  $K_d$  value of 9.7 pM derived from the kinetic experiments (Fig. 2) and the  $K_i$  value of 11 pM obtained in our previous competition assays with ( $\pm$ )- $^3$ H]epibatidine (Koren et al., 1998). The density of 5- $^{125}$ I]iodo-A-85380 binding sites in rat forebrain was comparable to densities obtained using ( $-$ )- $^3$ H]cytisine and ( $-$ )- $^3$ H]nicotine (Lippiello and Fernandes, 1986; Pabreza et al., 1991; Flores et al., 1992), ligands that primarily label the  $\alpha 4\beta 2$  nAChR subtype in rat brain. Figure 3c depicts binding of 5- $^{125}$ I]iodo-A-85380 in a postmortem sample of human brain cortex. As in rat brain, a single population of binding sites with a  $K_d$  value of  $11.6 \pm 0.7$  pM was observed. The density of binding sites in the human cortex was characterized by  $B_{max} = 0.98 \pm 0.04$  pmol/g tissue, ( $53 \pm 6$  fmol/mg protein). This density was close to values obtained in studies of postmortem human cortex using ( $\pm$ )- $^3$ H]epibatidine, ( $-$ )- $^3$ H]nicotine, and ( $-$ )- $^3$ H]cytisine (Sihver et al., 1998).

Nonspecific binding of 5- $^{125}$ I]iodo-A-85380 was proportional to the concentration of the radioligand (data not shown) and, at a concentration of 100 pM (approximately 10 times the  $K_d$  value), constituted ca. 10% of total binding. Much of this value was attributable to binding of the radioligand to filter material. True nonspecific binding to tissue typically did not exceed 5% of total binding.


As seen in Table 1, the binding affinity of 5- $^{125}$ I]iodo-A-85380 in rat brain membranes was moderately sensitive to variations in temperature during the incubation period. Thus, increasing the incubation temperature from 4–37°C resulted in a modest increase in the  $K_d$  value (from  $9.9 \pm 0.8$  to  $20 \pm 2$  pM, respectively). It should be emphasized that incubation at 4°C required an extended incubation time (24 h) to reach equilibrium. The 4-h incubation time routinely used seemed to be insufficient to reach equilibrium at this temperature and resulted in an inaccurate  $K_d$  value of  $15.5 \pm$

0.9 pM ( $n = 2$ ). On the other hand, when incubating at 22°C, increasing the duration beyond 4 h (up to 18 h) did not produce significant changes in the observed  $K_d$  or  $B_{max}$  values (data not shown). This observation suggests that the 4-h incubation time was sufficient to reach equilibrium at 22°C and that neither the radioligand nor the receptor protein underwent degradation under the assay conditions used.

**Competition Studies.** In competition assays with 5- $^{125}$ I]iodo-A-85380, affinities for nAChRs in rat brain for six well-characterized nicotinic agonists and four nicotinic antagonists (Table 2) fell into an order that was consistent with that previously observed in assays using other radioligands for  $\alpha$ -bungarotoxin-insensitive nAChRs (Pabreza et al., 1991; Decker et al., 1995; Houghtling et al., 1995). Compounds that did not effectively inhibit binding of 5- $^{125}$ I]iodo-A-85380 ( $K_i > 25 \mu\text{M}$ ) included a noncompetitive nicotinic antagonist (mecamylamine), an antagonist at muscarinic acetylcholine receptors (atropine), cholinesterase inhibitors (physostigmine and DFP), an antagonist at opiate receptors (naloxone), an agonist at dopamine receptors (apomorphine), and an antagonist at  $D_2$ -like dopamine receptors (haloperidol).

A  $K_i$  value of  $10 \pm 1$  pM, measured for the unlabeled 5-iodo-A-85380, was very close to that obtained for ( $-$ )-epibatidine (Table 2) and nearly identical with the  $K_d$  values for 5- $^{125}$ I]iodo-A-85380 measured in direct binding assays (Figs. 2 and 3). In addition, the  $K_i$  values for ( $-$ )-epibatidine, 5-iodo-A-85380, and A-85380 from Table 2 agreed well with those derived previously from competition assays with ( $\pm$ )- $^3$ H]epibatidine (Koren et al., 1998).

In our competition studies, the concentration of 5- $^{125}$ I]iodo-A-85380 (130 pM) was more than 10-fold higher than its  $K_d$  value. Because more than 90% of the binding sites were occupied by the radioligand at this concentration, the results obtained effectively characterized the entire population of 5- $^{125}$ I]iodo-A-85380 binding sites in rat forebrain. In all cases where  $K_i$  values were determined, the pseudo-Hill coefficient values were close to 1. These findings support our previous conclusion that in the rat forebrain, over the concentration range used, 5- $^{125}$ I]iodo-A-85380 labels a homog-



**Fig. 3.** Scatchard plots of 5- $^{125}$ I]iodo-A-85380 binding data obtained from saturation studies in P2 membrane fraction of Fischer-344 rat brain (A), in total membrane fraction of Sprague-Dawley rat brain (B), and in total membrane fraction of postmortem tissue of human cerebral cortex (C). Rat brain membranes ( $18$ – $25 \mu\text{g}$  of protein) or human cortical membranes ( $26$ – $30 \mu\text{g}$  of protein) were incubated in a total volume of, respectively, 1 or 0.5 ml with 1 to 500 pM 5- $^{125}$ I]iodo-A-85380 for 4 h at 22°C. Data were analyzed as described in *Experimental Procedures*. Each point represents the mean from three or four replicates (S.E.M. < 7%). A, pooled data from two independent experiments performed on the same membrane preparation. Four additional experiments were performed on membranes from two separate membrane preparations. The  $K_d$  and  $B_{max}$  values (mean  $\pm$  S.E.) obtained from these six experiments were  $10.6 \pm 0.3$  pM and  $160 \pm 25$  fmol/mg protein ( $3.8 \pm 0.6$  pmol/g tissue), respectively. B, results of a single experiment. Similar binding characteristics were observed in four additional experiments performed on membranes from four independent preparations. The  $K_d$  and  $B_{max}$  values (mean  $\pm$  S.E.) obtained from these five experiments were  $10.0 \pm 0.2$  pM and  $178 \pm 6$  fmol/mg protein ( $3.9 \pm 0.2$  pmol/g tissue), respectively. C, results of a single experiment. Similar results were obtained in three additional experiments performed on membranes of human postmortem cortical tissue obtained from different subjects. The  $K_d$  and  $B_{max}$  values (mean  $\pm$  S.E.) obtained from all four experiments were  $11.6 \pm 0.7$  pM and  $53 \pm 6$  fmol/mg protein ( $0.98 \pm 0.04$  pmol/g tissue).

enous population of agonist binding sites associated with  $\alpha$ -bungarotoxin-insensitive nAChRs, presumably the  $\alpha 4\beta 2$  subtype.

**Regional Distribution in Brain.** To test our hypothesis that 5-[<sup>125</sup>I]iodo-A-85380 labels  $\alpha 4\beta 2$  nAChRs, we investigated distribution of the radioligand binding in the rat brain using both in vitro binding assays and autoradiography. In all brain regions studied, 5-[<sup>125</sup>I]iodo-A-85380 binding was characterized by interaction with a single population of homogenous binding sites (Scatchard plots not shown) with  $K_d$  values close to 11 pM (Fig. 4). The average of  $K_d$  values from all regions studied was  $11.0 \pm 0.2$  pM. These constants closely agreed with the  $K_d$  and  $K_i$  values observed for the whole rat forebrain (Fig. 3, Table 2). The regional distribution of binding sites (Fig. 4) closely matched that of the  $\alpha 4\beta 2$  nAChR subtype measured previously in rat brain using (-)-[<sup>3</sup>H]cytisine and ( $\pm$ )-[<sup>3</sup>H]epibatidine (Pabreza et al., 1991; Houghtling et al., 1995); with the highest densities being observed in thalamic nuclei and superior colliculi; intermediate densities in the striatum, cortex, and hippocampus; and the lowest concentrations in the hypothalamus and cerebellum.

Results of the autoradiographic study with 5-[<sup>125</sup>I]iodo-A-85380 (Fig. 5) corroborated the pattern of distribution determined in the in vitro binding experiments. The study was carried out using 210 pM 5-[<sup>125</sup>I]iodo-A-85380, a concentration that theoretically should have saturated nearly 95% of the high-affinity binding sites. In the presence of 10  $\mu$ M (-)-nicotine, the binding was blocked almost completely (Fig. 5a), indicating that, at the concentration used, the

TABLE 1

Effect of temperature on binding parameters of 5-[<sup>125</sup>I]iodo-A-85380

Rat brain P2 membrane fractions (5  $\mu$ g of protein) were incubated in a total volume of 0.3 ml with 1 to 500 pM 5-[<sup>125</sup>I]iodo-A-85380 for 4 h at 22°C or 37°C, or for 24 h at 4°C. Results were analyzed as described in *Experimental Procedures*. Data represent mean  $\pm$  S.E. obtained from three to four experiments per incubation temperature. Experiments were performed in quadruplicate.

Binding Parameter	Incubation Temperature		
	4°C	22°C	37°C
$K_d$ , pM	$9.9 \pm 0.8$	$10.6 \pm 0.3$	$20 \pm 2$
$B_{max}$ , fmol/mg protein	$162 \pm 5$	$167 \pm 5$	$166 \pm 6$

TABLE 2

Inhibition of 5-[<sup>125</sup>I]iodo-A-85380 binding by nAChR and non-nAChR ligands

Rat brain P2 membrane fractions (10–11  $\mu$ g of protein) were incubated in a total volume of 0.2 ml with 130 pM 5-[<sup>125</sup>I]iodo-A-85380 and 9 to 11 concentrations of competitors for 4 h at 22°C and analyzed as described in *Experimental Procedures*. The inhibition constants ( $K_i$  values) were calculated by the Cheng-Prusoff equation from measured  $IC_{50}$  values using a  $K_d$  value of 10 pM for 5-[<sup>125</sup>I]iodo-A-85380 binding. In all assays, the pseudo-Hill coefficients ( $n_H$ ) did not differ significantly from 1. Data represent mean  $\pm$  S.E. obtained from four to six experiments per compound. Experiments were performed in duplicate.


nAChR Ligands	$K_i$ , nM	Non-nAChR Ligands	$K_i$ , nM
<b>Agonists</b>		Atropine	>25,000
(-)-Epibatidine	$0.009 \pm 0.001$	Physostigmine	>25,000
5-Iodo-A-85380	$0.010 \pm 0.001$	DFP	>1,000,000
A-85380	$0.015 \pm 0.002$	Naloxone	>25,000
(-)-Cytisine	$0.16 \pm 0.02$	Apomorphine	>25,000
(-)-Nicotine	$0.96 \pm 0.06$	Haloperidol	>25,000
Acetylcholine	$4.0 \pm 0.1$		
(+)-Nicotine	$10 \pm 1$		
Carbachol	$73 \pm 6$		
<b>Antagonists</b>			
Dihydro- $\beta$ -erythroidine	$7.4 \pm 0.8$		
Curare	$730 \pm 70$		
$\alpha$ -Bungarotoxin	>2,500		
Mecamylamine	>25,000		

binding of 5-[<sup>125</sup>I]iodo-A-85380 was limited to interactions with nAChRs in the rat brain.

Taken together, the regional densities of binding sites, labeled with 5-[<sup>125</sup>I]iodo-A-85380, and the results of the competition studies suggest that the labeled sites correspond to agonist binding sites on the  $\alpha 4\beta 2$  nAChR subtype. Still, these findings do not rule out the possibility that 5-iodo-A-85380 binds to other major mammalian nAChR subtypes, such as the  $\alpha 7$ , muscle-type, or subtypes containing  $\alpha 3$  and  $\beta 4$  subunits.

**Subtype Selectivity.** To address the issue of binding selectivity among nAChR subtypes, we measured affinities of unlabeled 5-iodo-A-85380 in four different competition assays. To determine the affinity of 5-iodo-A-85380 for the  $\alpha 4\beta 2$  subtype, we used previously described competition assays with ( $\pm$ )-[<sup>3</sup>H]epibatidine or (-)-[<sup>3</sup>H]cytisine and P2 membrane fractions of Fischer-344 rat forebrain. Consistent with the previous observations (Koren et al., 1998), Scatchard plots of ( $\pm$ )-[<sup>3</sup>H]epibatidine binding to these rat brain membranes, at ligand concentrations of 1 to 800 pM, revealed a single population of binding sites (data not shown) with a  $K_d$  of  $9.5 \pm 0.5$  pM ( $n = 4$ ). Similar high-affinity binding of ( $\pm$ )-[<sup>3</sup>H]epibatidine was observed in the rat (Houghtling et al., 1995), mouse (Marks et al., 1998), and human brain (Houghtling et al., 1995; Sihver et al., 1998), as well as in transfected cells stably expressing the  $\alpha 4\beta 2$  nAChR subtype (Houghtling et al., 1995; Whiteaker et al., 1998). In addition, the affinity of 5-iodo-A-85380 for the  $\alpha 4\beta 2$  nAChR subtype was measured in competition assays with (-)-[<sup>3</sup>H]cytisine, which is the most widely used ligand for characterization of this subtype. In our saturation experiments, (-)-[<sup>3</sup>H]cytisine revealed binding sites with a  $K_d$  value of  $300 \pm 50$  pM ( $n = 3$ ) in the rat brain membranes, consistent with published data (Pabreza et al., 1991).

To characterize binding of 5-iodo-A-85380 to the  $\alpha 7$  and muscular nAChR subtypes, we modified (see *Experimental*




**Fig. 4.** Distribution of 5-[<sup>125</sup>I]iodo-A-85380 in Fischer-344 rat brain. The individual brain structures obtained from ten animals were pooled and total membrane fractions from each region were isolated as described in *Experimental Procedures*. Membrane samples from each brain region (correspond to 0.1–0.5 mg of wet tissue) were incubated in a total volume of 0.25 ml with 1 to 500 pM 5-[<sup>125</sup>I]iodo-A-85380 for 4 h at 22°C. For all brain regions, Scatchard analyses produced data consistent with homogeneous populations of binding sites with similar  $K_d$  values. Gray ( $B_{max}$  values) and open ( $K_d$  values) columns represent means obtained from three to four saturation assays performed on membranes from two separate preparations. For all regions studied, S.E.M. values were less than 10% except for the hypothalamus, where S.E.M. = 25%. Cb, cerebellum; Cx, frontal cortex; F-Br, forebrain; Hipp, hippocampus; Hyp, hypothalamus; SC, superior colliculus; Str, striatum; Th, thalamus.

Procedures) previously described techniques (Bougis et al., 1986; Arneric et al., 1994). These techniques use  $^{125}\text{I}$ - $\alpha$ -bungarotoxin and membrane fractions isolated either from the rat brain (for the  $\alpha 7$  subtype) or from *T. californica* electroplax (for muscle-type nAChRs). In our experiments,  $^{125}\text{I}$ - $\alpha$ -bungarotoxin bound to a single population of binding sites in each of the two membrane fractions, exhibiting  $K_d$  values of  $1.5 \pm 0.2$  nM ( $n = 3$ ) in rat brain and  $2.3 \pm 0.3$  nM ( $n = 3$ ) in electroplax, consistent with published data (Zeghloul et al., 1988; Quik et al., 1996).

To complete the study on nAChR subtype selectivity, we developed an assay using  $(\pm)$ - $^3\text{H}$ ]epibatidine and a membrane

fraction from rat adrenal glands to estimate the affinity of 5-iodo-A-85380 for nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits. This assay was based on a previous study, which showed that  $(\pm)$ - $^3\text{H}$ ]epibatidine, in addition to its high affinity for  $\alpha 4\beta 2$  nAChRs in rat brain (Houghtling et al., 1995), bound to cells stably expressing receptors of the  $\alpha 3\beta 4$  subtype (Stauderman et al., 1998; Xiao et al., 1998) and to membranes from rat adrenal glands (Houghtling et al., 1995; Flores et al., 1997). Results of studies with bovine adrenals (Criado et al., 1997; Wenger et al., 1997) and cultured rat pheochromocytoma cells (PC12) (Rogers et al., 1992; Henderson et al., 1994) suggested that the adrenal glands were rich in nAChR subtype(s) containing  $\alpha 3$  and  $\beta 4$



**Fig. 5.** In vitro autoradiography of rat brain with 5- $^{125}\text{I}$ ]iodo-A-85380. Slices (20- $\mu\text{m}$  thick; taken approximately 0.4, 1.4, and 1.9 mm lateral to the midline) were incubated with 210 pM 5- $^{125}\text{I}$ ]iodo-A-85380 for 2 h. Nonspecific binding was assessed in the presence of 10  $\mu\text{M}$  (-)-nicotine. Pseudocolor-transformed autoradiograms were expressed in femtomoles per milligram of tissue. Acb, nucleus accumbens; Cb, cerebellum; CPU, caudate putamen; DTg, dorsal tegmental area; fr, fasciculus retroflexus; Hipp, hippocampus; IP, interpeduncular nucleus; MHb, medial habenula; Pn, pontine nucleus; RS, retrosplenial cortex; S, subiculum; SuG, superior colliculus, superficial gray; Th, thalamus; VTA, ventral tegmental area.

TABLE 3

Binding selectivity of 5-iodo-A-85380 and other nAChR ligands at four major mammalian nAChR subtypes

In assays of the  $\alpha 4\beta 2$  subtype, rat brain P2 membrane fractions (200  $\mu\text{g}$  of protein) were incubated in a total volume of 0.5 ml with 0.5 nM  $(\pm)$ - $^3\text{H}$ ]epibatidine and 9 to 11 concentrations of competitors for 1.5 h at 22°C. In assays of the nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits ( $\alpha 3\beta 4x$ ), total membrane fractions of rat adrenal glands (250  $\mu\text{g}$  of protein) were incubated in a total volume of 1 ml with 0.4 nM  $(\pm)$ - $^3\text{H}$ ]epibatidine and five to nine concentrations of competitors for 1.5 h at 22°C. In assays of the  $\alpha 7$  subtype, rat brain P2 membrane fractions (100  $\mu\text{g}$  of protein) were incubated in a total volume of 0.1 ml with 2 nM  $^{125}\text{I}$ - $\alpha$ -bungarotoxin and 9 to 11 concentrations of competitors for 3 h at 22°C. In assays of the muscle subtype, total membrane fractions from *T. californica* electric organ (0.1  $\mu\text{g}$  of protein) were incubated in a total volume of 0.1 ml with 2 nM  $^{125}\text{I}$ - $\alpha$ -bungarotoxin and 9 to 11 concentrations of competitors for 1.5 h at 22°C. The inhibition constants ( $K_i$  values) were calculated by the Cheng-Prusoff equation from measured  $\text{IC}_{50}$  values using the following  $K_d$  values: 10 and 55 pM for  $(\pm)$ - $^3\text{H}$ ]epibatidine binding in rat forebrain ( $\alpha 4\beta 2$ ) and in rat adrenal glands ( $\alpha 3\beta 4x$ ), respectively; and 1.5 and 2.3 nM for  $^{125}\text{I}$ - $\alpha$ -bungarotoxin binding in rat forebrain ( $\alpha 7$ ) and in *T. californica* electric organ ( $\alpha 1\beta 1\delta\epsilon$ ), respectively. These  $K_d$  values were obtained from three to six separate saturation assays per constant. Data represent means  $\pm$  S.E. obtained from three to seven competition assays per constant.

Ligand	$K_i$ at nAChR Subtype, nM (Ratio to $K_i$ at $\alpha 4\beta 2$ )			
	$\alpha 4\beta 2$	$\alpha 3\beta 4x$	$\alpha 7$	Muscle
$(\pm)$ -IPH	$0.027 \pm 0.004$ (1)	$0.11 \pm 0.04$ (4)	$30 \pm 3$ (1,100)	$6.5 \pm 0.1$ (240)
$(\pm)$ -Epibatidine	$0.008 \pm 0.001$ (1)	$0.049 \pm 0.02$ (6)	$4.0 \pm 0.5$ (500)	$7.5 \pm 0.5$ (900)
(-)-Nicotine	$0.84 \pm 0.13$ (1)	$100 \pm 20$ (120)	$130 \pm 10$ (150)	$1000 \pm 100$ (1,200)
(-)-Cytisine	$0.18 \pm 0.02$ (1)	$54 \pm 9$ (300)	$260 \pm 20$ (1,400)	$190 \pm 20$ (1,100)
A-85380	$0.017 \pm 0.002$ (1)	$14 \pm 2$ (800)	$17 \pm 2$ (1,000)	$320 \pm 20$ (19,000)
5-Iodo-A-85380	$0.010 \pm 0.001$ (1)	$51 \pm 5$ (5,000)	$250 \pm 20$ (25,000)	$1400 \pm 200$ (140,000)

subunits as well as  $\alpha 7$  subtype, but expressed few, if any, receptors of the  $\alpha 4\beta 2$  subtype. Binding assays with ( $\pm$ )-[ $^3\text{H}$ ]epibatidine using rat adrenal gland membranes demonstrated a single population of binding sites (data not shown) with a  $K_d$  value of  $55 \pm 5$  pM ( $n = 3$ ). ( $\pm$ )-[ $^3\text{H}$ ]Epibatidine binding to the rat adrenal gland membranes at a radioligand concentration of 0.5 nM was not blocked (data not shown) by  $\alpha$ -bungarotoxin at concentrations as high as 10,000 times its affinity ( $K_i = 1$  nM) at  $\alpha 7$  nAChRs (Quik et al., 1996). This observation suggests that, at conditions used for the competition assays, the binding of ( $\pm$ )-[ $^3\text{H}$ ]epibatidine in rat adrenal glands does not reflect interactions with the  $\alpha 7$  subtype. In light of the above-cited reports, the present data are consistent with the view that [ $^3\text{H}$ ]epibatidine binds to nAChRs containing  $\alpha 3$  and  $\beta 4$  subunits. Nonetheless, we cannot exclude the possibility that some portion of binding could reflect interactions with nAChRs including some other subunits (e.g.,  $\alpha 5$ ).


Results of the competition assays for different nAChR ligands/subtypes are summarized in Table 3. It is notable that the affinity of 5-iodo-A-85380 for the  $\alpha 4\beta 2$  receptor exceeded its affinities for other major mammalian nAChR subtypes by three to five orders of magnitude. In this regard, 5-iodo-A-85380 is vastly superior to all  $\alpha 4\beta 2$ -specific nAChR ligands known to date, including (-)-cytisine, which has long been the ligand of choice for characterizing the  $\alpha 4\beta 2$  subtype. The high affinity of 5-iodo-A-85380 for the  $\alpha 4\beta 2$  nAChR subtype measured in competition assays with ( $\pm$ )-[ $^3\text{H}$ ]epibatidine was confirmed in additional assays with (-)-[ $^3\text{H}$ ]cytisine, which yielded a nearly identical  $K_i$  value of  $10.5 \pm 0.7$  pM ( $n = 3$ ), and was consistent with results from binding assays with radiolabeled 5-iodo-A-85380 (Figs. 2 and 3), which provided  $K_d$  values of 9.7 to 10.6 pM. It should be noted that the exceptionally high  $\alpha 4\beta 2$ -subtype selectivity of 5-iodo-A-85380 is consistent with previous studies on interactions of the structurally related compounds, A-85380 (Sullivan et al., 1996) and ABT-594 (Bannon et al., 1998), with the  $\alpha 4\beta 2$ ,  $\alpha 7$ , and muscle-type nAChRs.

The high selectivity of 5-iodo-A-85380 for the  $\alpha 4\beta 2$  nAChR subtype was additionally confirmed by in vitro autoradiographic studies of the  $\beta 2$ -knockout mouse brain. In brain from the wild-type mouse, distribution of 5-[ $^{125}\text{I}$ ]iodo-A-85380 binding resembled that of [ $^{125}\text{I}$ ]IPH (Fig. 6), and the known pattern of distribution of  $\alpha 4\beta 2$  nAChRs. Unlike the case of the wild-type mouse, 5-[ $^{125}\text{I}$ ]iodo-A-85380 did not exhibit binding in any brain region of mice homozygous for a mutation in the  $\beta 2$  subunit of nAChRs. Unlabeled were the medial habenula and interpeduncular nucleus (Fig. 6), which were labeled with [ $^{125}\text{I}$ ]IPH (Fig. 6), and which were labeled previously with [ $^3\text{H}$ ]epibatidine and [ $^3\text{H}$ ]cytisine in mice that had a mutation in the  $\beta 2$  subunit (Zoli et al., 1998). As shown previously (Perry and Kellar, 1995), the medial habenula and interpeduncular nucleus contain substantially higher densities of binding sites for [ $^3\text{H}$ ]epibatidine than for [ $^3\text{H}$ ]cytisine. Taken together, the results of studies of nAChRs in the medial habenula and interpeduncular nucleus suggest the presence of at least two distinct types of nAChRs in these regions. Of these types, only one, namely, that containing  $\beta 2$  subunit (presumably, the  $\alpha 4\beta 2$  nAChR subtype), can be labeled with 5-[ $^{125}\text{I}$ ]iodo-A-85380. Thus, 5-[ $^{125}\text{I}$ ]iodo-A-85380 appears to be more selective than either epibatidine or, more importantly, cytisine, which has been accepted heretofore as the most selective high-affinity ligand for the  $\alpha 4\beta 2$  nAChR subtype.

Available data do not rule out the possibility that 5-iodo-

A-85380 has high affinity for subtypes other than  $\alpha 4\beta 2$ , which are far less abundant in mammalian brain. The investigation of such a possibility is a subject for future studies. Nonetheless, taking into consideration recent unpublished findings (K. J. Kellar, personal communication) that the parent compound, A-85380, has picomolar affinities toward several  $\beta 2$ -containing nAChR subtypes (including the  $\alpha 3\beta 2$  subtype) but only nanomolar affinities toward  $\beta 4$ -containing nAChRs (including  $\alpha 4\beta 4$ ), it is reasonable to assume that 5-iodo-A-85380 would follow the same pattern. Additionally, taken together with the above-cited results from Dr. Kellar's laboratory, the present observation of low affinity of A-85380 toward nAChRs in rat adrenal glands suggests that these receptors do not contain  $\beta 2$  subunits and are represented by  $\alpha 3\beta 4x$  nAChRs, where  $x$  may or may not represent another subunit, e.g.,  $\alpha 5$ . In this regard, it is noteworthy that the ratios of affinity for the  $\alpha 4\beta 2$  subtype to affinity for  $\alpha 3\beta 4x$  nAChRs, derived in the present work for ( $\pm$ )-epibatidine, cytisine, (-)-nicotine, and A-85380 (Table 3), closely matched recently published data obtained for the same compounds using rat brain membranes ( $\alpha 4\beta 2$  nAChRs) and a cell line stably expressing  $\alpha 3\beta 4$  nAChRs (Xiao et al., 1998).

In summary, the present results demonstrate that 5-iodo-A-85380 is an excellent ligand for studying nAChRs. It features extremely high affinity, slow dissociation from the receptor-ligand complex, high specific-to-nonspecific binding



**Fig. 6.** In vitro autoradiography of wild-type mouse brain and of brains from mice homozygous for a mutation in the  $\beta 2$  subunit of nAChRs performed with [ $^{125}\text{I}$ ]IPH and with 5-[ $^{125}\text{I}$ ]iodo-A-85380. Slices (12- $\mu\text{m}$  thick) at bregma -1.5 mm (four top panels) and at bregma -3.4 mm (four bottom panels) were incubated with 200 pM [ $^{125}\text{I}$ ]IPH (2200 Ci/mmol) and with 200 pM 5-[ $^{125}\text{I}$ ]iodo-A-85380 (420 Ci/mmol) for 0.5 h. Processed slides were exposed to Hyperfilm for 2 days ([ $^{125}\text{I}$ ]IPH) or 5 days (5-[ $^{125}\text{I}$ ]iodo-A-85380). Autoradiograms from mice homozygous for a mutation in the  $\beta 2$  subunit showed an absence of 5-[ $^{125}\text{I}$ ]iodo-A-85380 binding. Similar results were observed in all slices analyzed. Arrow and double arrow indicate, respectively, the medial habenula and interpeduncular nucleus, where binding of [ $^{125}\text{I}$ ]IPH, but not of 5-[ $^{125}\text{I}$ ]iodo-A-85380, was seen in brains of  $\beta 2$ -subunit knockout mice. Exposure for 5 weeks of 5-[ $^{125}\text{I}$ ]iodo-A-85380-labeled slices from mice homozygous for a mutation in the  $\beta 2$  subunit of nAChRs to Hyperfilm did not result in the detection of any additional specific signal. Results shown were reproduced in additional experiments with two control and two mutant animals as well as in experiments with 5-[ $^{125}\text{I}$ ]iodo-A-85380 in three control and three mutant animals.

ratio, and exceptionally high selectivity for the  $\alpha 4\beta 2$  nAChR subtype. Furthermore, the ability to produce 5- $^{125}\text{I}$ iodo-A-85380 with a specific activity of up to 2200 Ci/mmol makes it possible to detect nAChRs in the femtomolar range.

Recent *in vivo* studies with 5- $^{123/125}\text{I}$ iodo-A-85380 in the mouse (Musachio et al., 1998; Vaupel et al., 1998) and rhesus monkey (Chefer et al., 1998) and baboon (Musachio et al., 1999) demonstrated that this radioligand readily crosses the blood-brain barrier, specifically accumulates in the brain regions enriched with the  $\alpha 4\beta 2$  nAChRs, and exhibits low toxicity. These data, together with the results of the present *in vitro* characterization of 5-iodo-A-85380, suggest that radiolabeled with  $^{123}\text{I}$ , this compound would be particularly promising for noninvasive imaging of nAChRs with single photon emission computed tomography in both animals and humans.

#### Acknowledgments

We thank Cindy Ambriz (National Institute on Drug Abuse Brain Imaging Center) for preparing this manuscript and excellent administrative support, and Louis Amici (Yale University) for technical assistance.

#### References

- Abreo MA, Lin NH, Garvey DS, Gunn DE, Hettinger AM, Wasicak JT, Pavlik PA, Martin YC, Donnelly-Robert DL, Anderson DJ, Sullivan JP, Williams M, Arneric SP and Holladay MW (1996) Novel 3-pyridyl ethers with subnanomolar affinity for central neuronal nicotinic acetylcholine receptors. *J Med Chem* **39**:817–825.
- Arneric SP, Sullivan JP, Briggs CA, Donnelly-Roberts D, Anderson DJ, Raszkievicz JL, Hughes ML, Cadman ED, Adams P and Garvey DS (1994) (*S*)-3-Methyl-5-(1-methyl-2-pyrrolidinyl)isoxazole (ABT 418): A novel cholinergic ligand with cognition-enhancing and anxiolytic activities: I. *In vitro* characterization. *J Pharmacol Exp Ther* **270**:310–318.
- Bannon AW, Decker MW, Holladay MW, Curzon P, Donnelly-Robert DL, Putt-farcken PS, Bitner RS, Diaz A, Dickenson AH, Porsolt RD, Williams M and Arneric SP (1998) Broad-spectrum, non-opioid analgesic activity by selective modulation of neuronal nicotinic acetylcholine receptors. *Science (Wash DC)* **279**:77–81.
- Bougis PE, Marchot P and Rochat H (1986) Characterization of elapidae snake venom components using optimized reverse-phase high-performance liquid chromatographic conditions and screening assays for alpha-neurotoxin and phospholipase A2 activities. *Biochemistry* **25**:7235–7243.
- Chefer SI, Horti AG, Lee KS, Koren AO, Jones DW, Gorey JG, Links JM, Mukhin AG, Weinberger DR and London ED (1998) *In vivo* imaging of brain nicotinic acetylcholine receptors with 5- $^{123}\text{I}$ iodo-A-85380 using single photon emission computed tomography. *Life Sci* **63**:PL355–PL360.
- Criado M, Domínguez del Toro E, Carrasco-Serrano C, Smillie FI, Juiz JM, Viniestra S and Ballesta JJ (1997) Differential expression of  $\alpha$ -bungarotoxin-sensitive neuronal nicotinic receptors in adrenergic chromaffin cells: A role for transcription factor Egr-1. *J Neurosci* **17**:6554–6564.
- Dávila-García MI, Musachio JL, Perry DC, Xiao Y, Horti A, London ED, Dannals RF and Kellar KJ (1997)  $^{125}\text{I}$ IPH, an epibatidine analog, binds with high affinity to neuronal nicotinic cholinergic receptors. *J Pharmacol Exp Ther* **282**:445–451.
- Decker MW, Brioni JD, Bannon AW and Arneric SP (1995) Diversity of neuronal nicotinic acetylcholine receptors: Lessons from behavior and implications for CNS therapeutics. *Life Sci* **56**:545–570.
- Dukat M, Damaj MI, Glassco W, Duman D, May EL, Martin BR and Glennon RA (1993) Epibatidine: A very high affinity nicotine-receptor ligand. *Med Chem Res* **4**:131–139.
- Flores CM, Dávila-García MI, Ulrich YM and Kellar KJ (1997) Differential regulation of neuronal nicotinic receptor binding sites following chronic nicotine administration. *J Neurochem* **69**:2216–2219.
- Flores CM, DeCamp RM, Kilo S, Rogers SW and Hargreaves KM (1996) Neuronal nicotinic receptor expression in sensory neurons of the rat trigeminal ganglion: Demonstration of  $\alpha 3\beta 4$ , a novel subtype in the mammalian nervous system. *J Neurosci* **16**:7892–7901.
- Flores CM, Rogers SW, Pabreza LA, Wolfe BB and Kellar KJ (1992) A subtype of nicotinic cholinergic receptor in rat brain is composed of  $\alpha 4$  and  $\beta 2$  subunits and is up-regulated by chronic nicotine treatment. *Mol Pharmacol* **41**:31–37.
- Henderson LP, Gdovin MJ, Liu CL, Gardner PD and Maue RA (1994) Nerve growth factor increases nicotinic ACh receptor gene expression and current density in wild-type and protein kinase A-deficient PC12 Cells. *J Neurosci* **14**:1153–1163.
- Holladay MW, Dart MJ and Lynch JK (1997) Neuronal nicotinic acetylcholine receptors as targets for drug discovery. *J Med Chem* **40**:4169–4194.
- Horti AG, Koren AO, Lee KS, Mukhin AG, Vaupel DB, Kimes AS, Stratton M and London ED (1999) Radiosynthesis and preliminary evaluation of 5- $^{123/125}\text{I}$ iodo-3-(2*S*)-azetidinylmethoxy)pyridine: A radioligand for nicotinic acetylcholine receptors. *Nucl Med Biol* **26**:175–182.
- Horti AG, Scheffel U, Kimes AS, Musachio JL, Ravert HT, Mathews WB, Zhan Y, Finley PA, London ED and Dannals RF (1998) Synthesis and evaluation of

- N*- $^{11}\text{C}$ methylated analogues of epibatidine as tracers for positron emission tomographic studies of nicotinic acetylcholine receptors. *J Med Chem* **41**:4199–4206.
- Houghtling RA, Dávila-García MI and Kellar KJ (1995) Characterization of ( $\pm$ )- $^3\text{H}$ epibatidine binding to nicotinic cholinergic receptors in rat and human brain. *Mol Pharmacol* **48**:280–287.
- Koren AO, Horti AG, Mukhin AG, Gündisch D, Kimes AS, Dannals RF and London ED (1998) 2-, 5-, and 6-Halo-3-(2*S*)-azetidinylmethoxy)pyridines: Synthesis, affinity for nicotinic acetylcholine receptors, and molecular modeling. *J Med Chem* **41**:3690–3698.
- Lindstrom JM (1995) Nicotinic acetylcholine receptors, in *Handbook of Receptors and Channels: Ligand- and Voltage-Gated Ion Channels* (North RA ed) pp 153–175, CRC Press, Boca Raton.
- Lippiello PM and Fernandes KG (1986) The binding of  $l$ - $^3\text{H}$ nicotine to a single class of high affinity sites in rat brain membranes. *Mol Pharmacol* **29**:448–454.
- Marks MJ, Smith KW and Collins AC (1998) Differential agonist inhibition identifies multiple epibatidine binding sites in mouse brain. *J Pharmacol Exp Ther* **285**:377–386.
- Molina PE, Ding YS, Carroll I, Liang F, Volkow ND, Pappas N, Kuhar M, Abumrad N, Gatley SJ and Fowler JS (1997) Fluoro-norchlorepipatidine: Preclinical assessment of acute toxicity. *Nucl Med Biol* **24**:743–747.
- Musachio JL, Horti A, London ED and Dannals RF (1997) Synthesis of a radioiodinated analog of epibatidine: ( $\pm$ )-*exo*-2-(2-iodo-5-pyridyl)-7-azabicyclo[2.2.1]heptane for *in vitro* and *in vivo* studies of nicotinic acetylcholine receptors. *J Labelled Cmp Radiopharm* **39**:39–48.
- Musachio JL, Scheffel U, Finley PA, Zhan Y, Mochizuki T, Wagner HN and Dannals RF (1998) 5- $^{123/125}\text{I}$ iodo-3-(2*S*)-azetidinylmethoxy)pyridine, a radioiodinated analog of A-85380 for *in vivo* studies of central nicotinic acetylcholine receptors. *Life Sci* **62**:PL351–PL357.
- Musachio JL, Villemagne VL, Scheffel UA, Dannals RF, Dogan AS, Yokoi F and Wong DF (1999) Synthesis of an I-123 analog of A-85380 and preliminary SPECT imaging of nicotinic receptors in baboon. *Nucl Med Biol* **26**:201–207.
- Nybäck H, Halldin C, Åhlin A, Curvall M and Eriksson L (1994) PET studies of the uptake of (*S*)- and (*R*)- $^{11}\text{C}$ nicotine in the human brain: Difficulties in visualizing specific receptor binding *in vivo*. *Psychopharmacology (Berl)* **115**:31–36.
- Pabreza LA, Dhawan S and Kellar KJ (1991)  $^3\text{H}$ Cytisine binding to nicotinic cholinergic receptors in brain. *Mol Pharmacol* **39**:9–12.
- Perry DC and Kellar KJ (1995)  $^3\text{H}$ Epibatidine labels nicotinic receptors in rat brain: An autoradiographic study. *J Pharmacol Exp Ther* **275**:1030–1034.
- Piccio MR, Zoli M, Léna C, Bessis A, Lallemand Y, LeNovère N, Vincent P, Pich EM, Brulet P and Changeux J-P (1995) Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature (Lond)* **374**:65–67.
- Quik M, Choremis J, Komourian J, Lukas RJ and Puchacz E (1996) Similarity between rat brain nicotinic  $\alpha$ -bungarotoxin receptors and stably expressed  $\alpha$ -bungarotoxin binding sites. *J Neurochem* **67**:145–154.
- Rogers SW, Mandelzys A, Deneris ES, Cooper E and Heinemann S (1992) The expression of nicotinic acetylcholine receptors by PC12 cells treated with NGF. *J Neurosci* **12**:4611–4623.
- Silver W, Gillberg PG and Nordberg A (1998) Laminar distribution of nicotinic receptor subtypes in human cerebral cortex as determined by  $^3\text{H}$ (-)-nicotine,  $^3\text{H}$ cytisine and  $^3\text{H}$ epibatidine *in vitro* autoradiography. *Neuroscience* **85**:1121–1133.
- Stauderman KA, Mahaffy LS, Akong M, Veliceli G, Chavez-Noriega LE, Crona JH, Johnson EC, Elliott KJ, Gillespie A, Reid RT, Adams P, Harpold MM and Corey-Naeve J (1998) Characterization of human recombinant neuronal nicotinic acetylcholine receptor subunit combinations  $\alpha 2\beta 4$ ,  $\alpha 3\beta 4$  and  $\alpha 4\beta 4$  stably expressed in HEK293 cells. *J Pharmacol Exp Ther* **284**:777–789.
- Sullivan JP, Donnelly-Robert DL, Briggs CA, Anderson DJ, Gopalakrishnan M, Piattoni-Kaplan M, Campbell JE, McKenna DG, Molinari E, Hettinger AM, Garvey DS, Wasicak JT, Holladay MW, Williams M and Arneric SP (1996) A-85380 [3-(2*S*)-Azetidinylmethoxy)pyridine]: *In vitro* pharmacological properties of a novel, high affinity  $\alpha 4\beta 2$  nicotinic acetylcholine receptor ligand. *Neuropharmacology* **35**:725–734.
- Vaupel DB, Mukhin AG, Kimes AS, Horti AG, Koren AO and London ED (1998) *In vivo* studies with  $^{125}\text{I}$ 5-I-A-85380, a nicotinic acetylcholine receptor radioligand. *NeuroReport* **9**:2311–2317.
- Wenger BW, Bryant DL, Boyd RT and McKay DB (1997) Evidence for spare nicotinic acetylcholine receptors and a  $\beta 4$  subunit in bovine adrenal chromaffin cells: Studies using bromoacetylcholine, epibatidine, cytosine and mAb35. *J Pharmacol Exp Ther* **281**:905–913.
- Whiteaker P, Sharples CG and Wonnacott S (1998) Agonist-induced up-regulation of  $\alpha 4\beta 2$  nicotinic acetylcholine receptors in M10 cells: Pharmacological and spatial definition. *Mol Pharmacol* **53**:950–962.
- Xiao Y, Meyer EL, Thompson JM, Surin A, Wroblewski J and Kellar KJ (1998) Rat  $\alpha 3\beta 4$  subtype of neuronal nicotinic acetylcholine receptor stably expressed in a transfected cell line: Pharmacology of ligand binding and function. *Mol Pharmacol* **54**:322–333.
- Zegloul S, Marchot P, Bougis PE and Ronin C (1988) Selective loss of binding sites for the iodinated  $\alpha$ -neurotoxin I from *Naja mossaibica mossaibica* venom upon enzymatic deglycosylation of Torpedo electric organ membranes. *Eur J Biochem* **174**:543–550.
- Zoli M, Léna C, Piccioletto MR and Changeux J-P (1998) Identification of four classes of brain nicotinic receptors using  $\beta 2$  mutant mice. *J Neurosci* **18**:4461–4472.

**Send reprint requests to:** Edythe D. London, Brain Imaging Center, National Institute on Drug Abuse, 5500 Nathan Shock Dr., Baltimore, MD 21224. E-mail: elondon@tracer.org