

β_2 -Adrenergic Receptor Lacking the Cyclic AMP-Dependent Protein Kinase Consensus Sites Fully Activates Extracellular Signal-Regulated Kinase 1/2 in Human Embryonic Kidney 293 Cells: Lack of Evidence for G_s/G_i Switching.

JACQUELINE FRIEDMAN, BONITA BABU, and RICHARD B. CLARK

Department of Integrative Biology and Pharmacology, the University of Texas Health Science Center at Houston Medical School, Houston, Texas

Received February 26, 2002; accepted July 5, 2002

This article is available online at <http://molpharm.aspetjournals.org>

ABSTRACT

Stimulation of the β_2 -adrenergic receptor (β_2 AR) in human embryonic kidney (HEK) 293 cells causes a transient activation of Extracellular Signal-Regulated Kinase (ERK) 1/2. One of the mechanisms proposed for this activation is a PKA-mediated phosphorylation of the β_2 AR that switches receptor coupling from G_s to G_i and triggers internalization of the receptor. To examine these phenomena, we characterized agonist activation of ERK1/2 in HEK293 cells by the endogenous β_2 AR and in HEK293 cells stably overexpressing either the wild-type β_2 AR or a substitution mutant β_2 AR (PKA⁻) that lacks the cyclic AMP-dependent protein kinase (PKA) consensus phosphorylation sites (S261A, S262A and S345A, S346A). As the baseline, we established that epinephrine stimulation of the endogenous β_2 AR in HEK293 cells (20–30 fmol/mg) caused a rapid and transient activation of ERK1/2 with an EC₅₀ of 5 to 6 nM. In contrast, the potency of epinephrine stimulation of ERK1/2 in cells stably overexpressing WT β_2 AR and PKA⁻ (2–4 pmol of β_2 AR/mg) was increased by over 100-fold relative to HEK293

cells, the EC₅₀ values being 20 to 60 pM. The nearly identical 100-fold shift in EC₅₀ for ERK1/2 activation in the PKA⁻ and WT β_2 AR relative to that in the HEK293 showed that the PKA⁻ are fully capable of activating ERK1/2. We also found maximal activation of ERK1/2 in the overexpressing cell lines at concentrations of epinephrine that cause no internalization (i.e., the EC₅₀ for internalization was 75 nM). Pertussis toxin pretreatment caused only a weak inhibition of epinephrine activation of ERK1/2 in the HEK293 (7–16%) and no inhibition in the PKA⁻ cells. Finally we found that the Src family kinase inhibitor 4-amino-5-(4-chlorophenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine (10 μ M) caused a >90% inhibition of epinephrine or forskolin activation of ERK1/2 in both cell lines. Our results indicate that the dominant mechanism of β_2 AR activation of ERK1/2 does not require PKA phosphorylation of the β_2 AR, receptor internalization or switching from activation of G_s to G_i , but clearly requires activation of a Src family member that may be downstream of PKA.

Early reports that purified β_2 AR or a synthetic peptide corresponding to the third intracellular loop of the β_2 AR could activate pure G_i in reconstituted preparations in vitro raised the possibility that the β_2 AR could activate G_i in vivo (Cerione et al., 1985; Rubenstein et al., 1991), although with reduced efficiency relative to G_s . It was then demonstrated that PKA phosphorylation of a III₁ loop peptide of the β_2 AR increased activation of G_i in reconstituted preparations and slightly reduced its activation of G_s (Okamoto et al., 1991), leading to the proposal that β_2 AR activation could switch from G_s to G_i after PKA phosphorylation of the III₁ loop.

Recent support for this scheme was derived from studies showing that isoproterenol activation of ERK1/2 in HEK293 cells was blocked more than 85% by pretreatment with pertussis toxin, that isoproterenol activation of ERK1/2 was inhibited by H89, and that transient expression of a β_2 AR lacking the PKA consensus sites (S261A, S262A, S345A, S346A) inhibited isoproterenol activation of ERK1/2 by 40% (Daaka et al., 1997). Other studies are not consistent with the switching hypothesis. First, it has been shown that forskolin activates ERK1/2 consistent with a PKA-mediated pathway of activation downstream of the receptor (Daaka et al., 1997; Schmitt and Stork, 2000). Second, a recent study found that isoproterenol activation of ERK1/2 in HEK293 cells was not inhibited by pertussis toxin (Schmitt and Stork,

This work was supported by National Institutes of Health grant GM31208 (to R.B.C.).

ABBREVIATIONS: β_2 AR, β_2 -adrenergic receptor; HEK, human embryonic kidney; ERK, extracellular signal-regulated kinase; PKA, cyclic AMP-dependent protein kinase; WT, wild-type; CGP-12177, 4-[3-[(1,1-dimethylethyl)amino]2-hydroxypropoxy]-1,3-dihydro-2H-benzimidazol-2-one; PP2, 4-amino-5-(4-chlorophenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine; DMEM, Dulbecco's modified Eagle's medium; HA, hemagglutinin; AT, ascorbate/thiourea; HE, HEPES/EDTA; PAGE, polyacrylamide gel electrophoresis; ICI-118551, (\pm)-1-[2,3-(dihydro-5,7-methyl-1H-inden-4-yl)oxy]-3-[(methyl)amino]-2-butanol; EGF, epidermal growth factor.

2000). Third, we have shown that pertussis toxin pretreatment does not alter epinephrine-induced desensitization of HEK293 cells and S49 lymphoma cells, as would be expected if a G_s to G_i switch occurred (Clark et al., 1986; Seibold et al., 2000). Fourth, isoproterenol activation of ERK1/2 does not occur in the kin^- and cyc^- mutants of S49 mouse lymphoma cells (Wan and Huang, 1998).

Reports on the role of internalization of the β_2 AR in ERK1/2 activation, studied through the use of transient expression of dominant negative dynamin (K44A) and mutant arrestins, have also been inconsistent. Several studies suggested that internalization was required for ERK1/2 activation by the β_2 AR in HEK293 cells (Daaka et al., 1998; Luttrell et al., 1999b) and COS-7 cells (Pierce et al., 2000), whereas others found that internalization was not required for β_2 AR activation of ERK1/2 in COS-1 cells (DeGraff et al., 1999).

Several lines of evidence suggest a role for Src in activation of ERK1/2 by the β_2 AR: i) isoproterenol activation caused the formation of a multiprotein complex of receptor, β -arrestin, and Src, and dominant negative mutants of Src reduced ERK1/2 activation (Daaka et al., 1997; Luttrell et al., 1999b; Zou et al., 1999; Miller et al., 2000); ii) Src was shown to be activated by $G_s\alpha$ and $G_i\alpha$ in reconstituted *in vitro* preparations suggesting direct activation of Src by G proteins (Ma et al., 2000); and iii) it was suggested that Src binds to tyrosine 350 of the β_2 AR after phosphorylation of this residue by an unidentified tyrosine kinase (Fan et al., 2001).

Obviously there seems to be considerable complexity in β_2 AR activation of ERK1/2, and many issues are unresolved. In this article, we examine the role of receptor switching, G_i -internalization, and Src family kinases in β_2 AR activation of ERK1/2 in HEK293 cells. We had previously generated a mutant β_2 AR in which the two PKA consensus sites were eliminated by substitution of serines 261, 262, 345, and 346 with alanine, and this mutant receptor, termed PKA⁻, was stably overexpressed in the HEK293 cell line. We reported previously that the coupling efficiency of this mutant and the EC₅₀ for activation of adenylyl cyclase were nearly identical to those of WT β_2 AR overexpressed at comparable levels (Seibold et al., 1998, 2000). Stable overexpression of the PKA⁻ and the WT β_2 AR (~100-fold relative to endogenous receptor) caused a dramatic left-shift in the EC₅₀ for epinephrine activation of adenylyl cyclase. This property of the β_2 AR allows the analysis of mutant receptors overexpressed in HEK293 cells because the response of the endogenous receptor is overwhelmed if expression is high enough. Because the EC₅₀ for epinephrine activation of adenylyl cyclase in cells expressing either the PKA⁻ or WT β_2 AR is left-shifted relative to the endogenous receptor, it follows that activation of ERK1/2 should show a similar left-shift if the mechanism involves receptor activation of G_s . The prediction of the switching model in which the PKA-phosphorylated β_2 AR is proposed to activate G_i is that there should be no left shift in ERK1/2 activation relative to the endogenous receptor; rather, there should be an inhibition of the activation by endogenous β_2 AR (Daaka et al., 1997). The potency of agonist activation of ERK1/2 in HEK293 cells by transiently expressed wild-type or PKA⁻ mutant β_2 ARs was not examined in previous studies (Daaka et al., 1997; Schmitt and Stork, 2000).

In this article, we demonstrate that the stably overexpressed PKA⁻ mutant of the β_2 AR shows the predicted enormous left shift in the potency of activation of ERK1/2 by

epinephrine relative to that of the endogenous low level of expression in the HEK293 and that the shift is nearly identical to that found with overexpressed wild-type β_2 AR. This demonstrated that the PKA consensus sites were not required for full activation of ERK1/2. Our data also show only a minor role for G_i in epinephrine activation of ERK1/2, no requirement for internalization (because the EC₅₀ for epinephrine-induced internalization in the PKA⁻ was about 1000-fold higher than that for activation of ERK1/2), and a requirement for Src family kinase activation.

Materials and Methods

Materials. Antibodies to ERK1/2 and phospho-ERK1/2 were purchased from Cell Signaling Technology Inc. (Beverly, MA). [³²P]NAD and [³H]CGP-12177 were obtained from PerkinElmer Life Sciences (Boston, MA). Pertussis toxin was from List Biological Laboratories Inc. (Campbell, CA). Enhanced chemiluminescence reagents and Hyperfilm were from Amersham Biosciences (Piscataway, NJ). Agonists and antagonists were purchased from Sigma-Aldrich. BA85 nitrocellulose was from Schleicher & Schuell (Keene, NH). Cell culture reagents were purchased from Mediatech (Herndon, VA). The Src family kinase inhibitor PP2 was from Alexis Corporation (Läufelfingen, Switzerland).

Cell Culture. HEK293 cells purchased from the American Type Culture Collection (Manassas, VA) were grown in 5% CO₂ at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum. The stably transfected HEK293 cell lines used in this study expressing either WT or mutant receptors have been described previously (January et al., 1997; Seibold et al., 1998; Seibold et al., 2000), and were as follows: double epitope-tagged wild-type receptor (HA- β_2 AR-His₆) and the double epitope-tagged PKA⁻ (S261A, S262A, S345A, S346A), both with the hemagglutinin (HA) tag on the N terminus and His₆ tag on the C terminus; and HA-tagged wild-type (HA- β_2 AR). The stably transfected cell lines were cultured in medium containing 400 μ g/ml G418. The levels of β_2 AR in these transfect lines were 2 to 4 pmol/mg membrane protein, whereas the level of endogenous β_2 AR was 30 to 40 fmol/mg.

Cell Treatment and Preparation of Solubilized Extract. Cells were grown to confluence in growth medium in 12-well plates coated with poly(L-lysine). The medium was removed 18 h before treatment, cells were incubated for 30 min with serum-free DMEM, and that medium was replaced with 2 ml of serum-free DMEM with or without 100 ng/ml of pertussis toxin. Cells were treated with hormones or carrier as indicated at 37°C with continuous rocking; pretreatments with β_2 AR antagonists were for 2 min before agonist treatment, and PP2 was added 1 h before agonist treatment. Epinephrine was stored in 10 mM ascorbate/100 mM thiourea pH 7 (AT). Stock solutions were diluted 100-fold when additions were made to cells such that the final concentration of AT in all incubations (control and treated) was 0.1 mM ascorbate/1.0 mM thiourea. PP2 was dissolved and stored in DMSO and was diluted 100-fold for cell treatments. To terminate treatments, the medium was removed and 0.5 ml of solubilization buffer (20 mM HEPES, pH 7.4, 150 mM NaCl, 0.8% dodecyl maltoside, 1 mM EDTA, pH 7, 20 mM tetrasodium pyrophosphate, 10 mM NaF, 10 μ g/ml benzamidine, 10 μ g/ml trypsin inhibitor, 10 μ g/ml leupeptin, and 0.1 μ M okadaic acid) was added to each well. The plates were placed on ice, and the solubilized cells were pipetted into 1.5-ml microcentrifuge tubes. The tubes were rocked at 4°C for 30 min, and then centrifuged at 12,500 rpm for 15 min. The supernatants were removed and frozen at -80°.

Measurement of ERK1/2 and Phospho-ERK1/2. To measure activation of ERK1/2 by Western blotting, 20 μ l of the solubilized extracts were resolved by SDS-PAGE (10% gels) and blotted onto BA-85 nitrocellulose. The blots were blocked for 1 h in 5% nonfat dried milk in wash buffer (150 mM NaCl, 50 mM Tris pH 7.5, and 0.1% Tween 20). After two 10-min washes, the blots were incubated

overnight at 4°C on a rocker with a 1:1000 dilution of anti-phospho-ERK, washed twice for 10 min each, and incubated for 1 h at room temperature with a 1:5000 dilution of secondary antibody (goat anti-rabbit HRP-conjugate, Bio-Rad, Hercules, CA) After 2 washes, the blots were exposed to ECL reagents for 1 min, dried, and exposed to Hyperfilm (Amersham Biosciences) for 15 sec to 2 min. The blots were then stripped and reprobed identically with anti-ERK. Western blots were quantitated using Scion Image software (Scion Corp., Frederick, MD). All results with the anti-phospho-ERK were normalized to the corresponding anti-ERK.

Membrane Preparation and Adenylyl Cyclase Assay. Cells were plated into 100-mm dishes that had been precoated with poly(L-lysine). Treatments were administered at 37°C and were stopped by removal of media followed by six washes with 5 ml of ice-cold HE buffer (20 mM HEPES, pH 8.0, 1 mM EDTA, pH 7). The cells were scraped into HE containing 10 µg/ml leupeptin and 0.1 µM okadaic acid and homogenized with seven strokes in a type B Dounce homogenizer (Bellco Glass, Vineland, NJ). The homogenates were layered onto sucrose step gradients (23 and 43% prepared in HE buffer) and centrifuged at 25,000 rpm in a Beckman SW28.1 rotor for 35 min. The fraction at the 23/43% sucrose interface was removed, frozen in liquid nitrogen, and stored at -80°C. Adenylyl cyclase activity was measured as described previously (Seibold et al., 2000).

ADP Ribosylation. Membranes from control or pertussis toxin-treated cells were incubated for 30 min at 30°C with 10 µM NAD, 0.5 mM ATP, 0.2 mM GDP, 5 mM MgCl₂, 1 mM EDTA, pH 7, 20 mM Tris, pH 7.5, 5 mM dithiothreitol, 5 mM thymidine, 8 mM creatine phosphate, 8 U/ml creatine phosphokinase, and 5 µCi/tube [³²P]NAD. The incubation mix was diluted in 5 ml of 20 mM Tris, pH 7.5, and centrifuged for 15 min at 30,000 rpm in a Beckman 50Ti rotor (Beckman Coulter, Fullerton, CA). The pellets were each dissolved in SDS sample buffer and resolved on 12% SDS-PAGE gels. The proteins were transferred to nitrocellulose and exposed to a Storm Phosphorimager screen (Molecular Dynamics, Sunnyvale, CA). The phosphorylated bands were quantitated using ImageQuant (Molecular Dynamics).

Internalization. The procedure for measuring internalization of the β₂AR has been described in detail previously (Seibold et al., 2000). Cells in 12-well dishes were treated with epinephrine or AT for 5 min. Cells were then washed five times with ice-cold, serum-free DMEM, placed on ice, then incubated with 5 to 10 nM [³H]CGP-12177 with or without 1 µM alprenolol in serum-free DMEM. Dishes were then incubated for 1 h on ice, washed twice with ice-cold PBS to remove [³H]CGP-12177, and cells released by trypsin were transferred to scintillation vials for counting.

Results

Characterization of β₂AR Agonist Activation of ERK1/2 in HEK293 Cells.

To examine the activation of ERK1/2 by the endogenous β₂AR (≈20–30 fmol/mg membrane protein), HEK293 cells grown to confluence were placed in serum-free medium for 18 h before treatment with epinephrine. As shown in Fig. 1A, activation of ERK1/2 by 100 nM epinephrine showed a slight lag after which levels peaked at 5 min and then sharply declined, returning to near control levels by 20 min. To determine whether the activation of ERK1/2 by epinephrine was mediated by its binding to the β₂AR, several approaches were taken. We found that 5 µM ICI-118551, a selective β₂AR antagonist, added 2 min before a 5-min incubation with various concentrations of epinephrine, caused a nearly complete inhibition of ERK1/2 activation (Fig. 1B), consistent with the competitive action of ICI-118551. We also made the following observations (data not shown): i) the activation of ERK1/2 by 100 nM isoproterenol was equivalent to epineph-




Fig. 1. Characterization of epinephrine activation of ERK1/2 in HEK293 cells. **A**, time course of epinephrine activation of ERK1/2. HEK293 cells grown to confluence in 12-well plates were treated with 100 nM epinephrine or carrier (AT) for the indicated times. **B**, ICI-118551 inhibition of epinephrine-activated ERK1/2. HEK293 cells were pretreated for 2 min ± 5 µM ICI-118551, followed by 5-min treatments with the indicated concentrations of epinephrine or carrier (AT). After treatment, cells were solubilized and extracts were subjected to SDS-PAGE, transferred onto BA85 nitrocellulose membranes, and Western blotted with anti-phospho-ERK1/2 as described under *Materials and Methods*. Gels were then stripped and reprobed with anti-ERK1/2. Results with anti-phospho-ERK were normalized to the corresponding anti-ERK data and shown as arbitrary units (AU) on the y-axis.

rine and was also blocked by ICI-118551; ii) 20 nM salmeterol, a partial agonist and one of the most selective β_2 AR agonists, showed $\approx 80\%$ of the activation of ERK1/2 induced by epinephrine and was also blocked by ICI-118551; iii) 100 nM prazosin (an antagonist of both α_1 - and α_2 -adrenergic receptors) had no effect on epinephrine activation of ERK1/2. These data demonstrated that epinephrine activation of ERK1/2 was specific for the β_2 AR.

To determine the EC_{50} for epinephrine activation of ERK1/2, HEK293 cells were incubated for 5 min with a range of epinephrine concentrations. Immunoblots from one representative experiment are shown in Fig. 2, as well as a data summary from five identical experiments. The EC_{50} for epinephrine activation of ERK1/2 was 5.7 nM. It should be noted that this EC_{50} is about 100- to 200-fold lower than that for epinephrine activation of adenylyl cyclase in membranes prepared from these cells (600–800 nM, data not shown).

Epinephrine Stimulation of ERK1/2 in HEK293 Cells Overexpressing Either PKA⁻, HA- β_2 AR-His₆, or HA- β_2 AR. As discussed above, overexpression of β_2 AR in a variety of cell types causes a progressive decrease in the EC_{50} (increased potency) for β_2 AR activation of adenylyl cyclase (Whaley et al., 1994; Seibold et al., 1998; Clark et al., 1999). We have shown previously that the EC_{50} values for epinephrine activation of adenylyl cyclase in cells overexpressing either the WT β_2 AR or the PKA⁻ were left-shifted 50- to 100-fold relative to the endogenous receptor in HEK293 cells. It follows that activation of ERK1/2 should show a similar left shift if the mechanism involved for both processes was

similar at the level of receptor coupling to G_s/adenylyl cyclase.

To determine whether the EC_{50} for epinephrine activation of ERK1/2 in the PKA⁻ overexpression line was left-shifted relative to the endogenous receptor and whether the shift in the potency of the PKA⁻ was similar to that of two lines overexpressing the WT β_2 AR to similar levels, the experiments shown in Fig. 3 were performed. HEK293 cells stably overexpressing either the PKA⁻, HA- β_2 AR-His₆, or HA- β_2 AR (2–4 pmol/mg) were incubated with a range of epinephrine concentrations for 5 min and the EC_{50} values for ERK1/2 activation determined. The EC_{50} for epinephrine activation of ERK1/2 in the PKA⁻ cell line was ≈ 36 pM, more than 100-fold left-shifted relative to epinephrine activation of ERK1/2 by the endogenous receptor in HEK293 cells. Concentrations as low as 10 pM gave a significant increase in ERK1/2 phosphorylation. The EC_{50} values for epinephrine activation of ERK1/2 in the HA- β_2 AR-His₆ and HA- β_2 AR were in the range of 20 to 40 pM. These data demonstrated that the PKA⁻ β_2 AR was unimpaired in its activation of ERK1/2 because the potency of epinephrine activation was similar in the two lines overexpressing the wild-type receptors. Furthermore, as with the endogenous receptor, the left shift in the EC_{50} for ERK1/2 activation by the three overexpressing clones relative to their EC_{50} for adenylyl cyclase activation (January et al., 1998; Seibold et al., 2000) was more than 100-fold.

Because there was the possibility that the time course of ERK1/2 activation by epinephrine was altered at these very low concentrations, we followed the time course of ERK1/2 activation in the PKA⁻ by 1.0 and 10.0 nM epinephrine as shown in Fig. 4. The data show that the time course of activation by 1.0 and 10.0 nM epinephrine in the PKA⁻ cells was similar to that of the endogenous receptor in HEK293 cells. With lower concentrations of epinephrine, there is some indication that the lag is extended; however, the peak level of ERK1/2 remained at 5 min. A similar time course of ERK1/2 activation was found in the cells expressing the two wild-type receptors.

Effect of Pertussis Toxin Pretreatment of HEK293 Cells on the β_2 -Adrenergic Activation of ERK1/2. To determine the role of G_i in epinephrine activation of ERK1/2, we examined the effect of pertussis toxin pretreatment. HEK293 cells expressing only endogenous levels of receptor and the PKA⁻ cell line were pretreated with or without 100 ng/ml of pertussis toxin for 18 h in serum-free medium. Cells were then incubated with various concentrations of epinephrine for 5 min. As shown in Fig. 5A (summary of seven independent experiments with a typical Western blot above), pertussis toxin pretreatment of HEK293 cells expressing the endogenous receptor caused only a 7 to 16% inhibition of ERK1/2 activation by the various concentrations of epinephrine. We also found only marginal pertussis toxin inhibition of either isoproterenol or salmeterol activation of ERK1/2 in the HEK293 cell line (data not shown). No significant pertussis toxin inhibition of epinephrine activation of ERK1/2 in the PKA⁻-expressing cells was observed (Fig. 5B).

Because pertussis toxin treatment caused only a weak inhibition of epinephrine activation of ERK1/2 by the endogenous receptor, and no inhibition of ERK1/2 activation by the PKA⁻, we performed several controls to determine the extent of toxin activity. First, cells were pretreated overnight with




Fig. 2. Dose response of epinephrine activation of ERK1/2 in HEK293 cells. HEK293 cells were treated for 5 min with the indicated concentrations of epinephrine. The cells were solubilized and ERK1/2 phosphorylation measured and quantitated as described under *Materials and Methods*. The data on the graph are the mean values \pm S.E.M. from five experiments, each performed in duplicate. The data from each experiment were converted to percentage of the maximum stimulation before determination of the means. The Western blot is from one representative experiment.




Fig. 3. Dose response of epinephrine activation of ERK1/2 in HEK293 cells stably expressing either the PKA⁻, HA-β₂AR-His₆, or HA-β₂AR. Cells expressing the various receptors were treated for 5 min with the indicated concentrations of epinephrine. The cells were solubilized and ERK1/2 phosphorylation was measured and quantitated as described under *Materials and Methods*. A, Western blots are from one representative experiment for each cell line. B, the data are the mean ± S.E.M. of seven (PKA⁻) or four (HA-β₂AR-His₆ and HA-β₂AR) experiments, each performed in duplicate. The data from each experiment were converted to percentage of maximum epinephrine stimulation of ERK1/2 phosphorylation before calculating the means.

pertussis toxin, and membranes derived from control and pertussis toxin-treated HEK293 or PKA⁻ cells were ADP-ribosylated in the presence of [³²P]NAD. Overnight treatment with pertussis toxin resulted in a >90% inhibition of ADP-ribosylation of G_i/G_o in membrane preparations (Fig. 6A). As a second control, we examined pertussis toxin's effect on epinephrine stimulation of adenylyl cyclase. We found that pertussis toxin treatment of HEK293 cells caused an approximate doubling of epinephrine-stimulated adenylyl cyclase activity with no change in the EC₅₀ (Fig. 6B). This result is consistent with our prior study of this toxin's effects on epinephrine stimulation of adenylyl cyclase in the HEK293 cells overexpressing the wild-type and mutant β_2 AR (Seibold et al., 2000) (i.e., a doubling of the V_{max} and no change in the EC₅₀).

Internalization of the β_2 AR in PKA⁻. As discussed, there have been some inconsistencies concerning the role of β_2 AR internalization in activation of ERK1/2. It occurred to us that we could determine how closely these two processes were related by comparing their EC₅₀ values. To measure the EC₅₀ for epinephrine-induced internalization of the PKA⁻ β_2 AR, cells were treated for 5 min with or without various epinephrine concentrations as indicated in Fig. 7 (■). Internalization was measured using [³H]CGP-12177 as described under *Materials and Methods*. Epinephrine (10 nM) produced a barely detectable level of internalization, and the EC₅₀ for epinephrine-induced internalization was 75 nM. To compare the EC₅₀ for epinephrine-induced internalization with that for activation of adenylyl cyclase and ERK1/2, the PKA⁻ data from Fig. 3 and the results from a typical adenylyl cyclase assay are also plotted in Fig. 7. It can be seen that the EC₅₀ for epinephrine activation of internalization is approximately 1000-fold higher than that for ERK1/2 and 10-fold higher relative to activation of adenylyl cyclase.

Inhibition of ERK1/2 Activation by the Src Family Kinase Inhibitor PP2. To assess the role of the Src family in ERK1/2 activation, PKA⁻ cells were pretreated with the Src family inhibitor PP2 for 1 h before stimulation with




Fig. 4. Time course of epinephrine activation of ERK1/2 in PKA⁻ cells. PKA⁻ cells were treated with either 1.0 nM epinephrine (■) or 10.0 nM epinephrine (▲) for the times indicated on the graph. The value at zero time is the mean of controls determined at each of the time points. The cells were solubilized and ERK1/2 phosphorylation measured and quantitated as described under *Materials and Methods*. The data are from one representative experiment.

either epinephrine, forskolin, or EGF (Fig. 8). PP2 blocked 90 to 100% of epinephrine and forskolin activation of ERK1/2, but produced only a 33% inhibition of EGF stimulation. Similar results were obtained with the HEK293 cells expressing only endogenous β_2 AR (data not shown). These data strongly suggest that Src family kinases play an essential role in β_2 AR activation of ERK1/2. Furthermore, the inhibition of forskolin activation of ERK1/2 shows that it is unlikely that a direct receptor or G protein interaction with Src is required.




Fig. 5. Effect of pertussis toxin on epinephrine activation of ERK1/2 in HEK293 cells. HEK293 cells expressing either the endogenous β_2 AR only (A) or the PKA⁻ β_2 AR (B) growing in 12-well plates were placed in serum-free medium and then treated with or without 100 ng/ml pertussis toxin (PTOX) for 18 h as described under *Materials and Methods*. Cells were incubated with increasing concentrations of epinephrine for 5 min as indicated. Phosphorylation of ERK1/2 was measured as described under *Materials and Methods*. The data are the means \pm S.E.M. of seven (HEK293) or three (PKA⁻) experiments. One representative Western blot is shown for each cell line.

Discussion

In this article, we focused on the proposal that PKA phosphorylation of the β_2 AR switches receptor activation from G_s to G_i and that this switch of receptor activation of G proteins was required for ERK1/2 activation, as was internalization of the receptor. Two approaches were used to assess the capacity of the PKA⁻ receptor to activate ERK1/2, both of which were based on our prior studies demonstrating that β_2 AR overexpression predictably left-shifts the EC_{50} for agonist activation of adenylyl cyclase relative to that for the endogenous β_2 AR. First, we found that the potencies (EC_{50} values) for epinephrine activation of ERK1/2 in the PKA⁻ and the clones overexpressing WT β_2 AR were similar (20–60 pM) and about 100- to 200-fold lower than the EC_{50} for epinephrine activation of endogenous receptor in HEK293 cells (5–6 nM). Second, we determined whether the amplification of epinephrine activation of ERK1/2 relative to its activation of adenylyl cyclase for the overexpressed β_2 ARs was similar to that for the endogenous receptor. For the endogenous receptor, the EC_{50} for epinephrine activation of adenylyl cyclase in cell-free membrane preparations (\approx 600–800 nM) was about 100-fold greater than the EC_{50} for ERK1/2 activation (5–6 nM), whereas the comparable ratio for the PKA⁻ was about 200- to 300-fold (15 nM for adenylyl cyclase activation, and 30–60

pM for ERK1/2 activation). The shift in sensitivity of the PKA⁻ for ERK1/2 activation relative to adenylyl cyclase was actually somewhat greater than that for the HEK293 cell line expressing only endogenous β_2 AR.

Our work also addressed the role of G_i in β_2 AR activation of ERK1/2. Previous studies had found either a major role for G_i for agonist activation of ERK1/2 after stimulation of the endogenous β_2 AR in HEK293 cells (Daaka et al., 1997) or none (Schmitt and Stork, 2000). In the latter study, activa-




Fig. 6. Evidence for the efficacy of pertussis toxin treatments. A, ADP-ribosylation of G_i/G_o in membranes from HEK293 and PKA⁻ cells pretreated for 18 h \pm pertussis toxin. HEK293 or PKA⁻ cells plated into 100-mm dishes were treated with or without pertussis toxin (100 ng/ml) and membranes were harvested and frozen as described under *Materials and Methods*. Thawed membranes were then incubated for 30 min in the presence or absence of pertussis toxin in the ADP-ribosylation mix containing [³²P]NAD. Membranes were washed and aliquots resolved on 12% SDS-PAGE. B, epinephrine-stimulated adenylyl cyclase activity in membranes from HEK293 cells treated with pertussis toxin or not. HEK293 cells in 100-mm dishes were treated overnight with or without 100 ng/ml of pertussis toxin, membranes were prepared, and adenylyl cyclase stimulation by the indicated concentrations of epinephrine was measured in the membranes. The data are the means \pm S.E.M. of triplicate determinations from one representative experiment.




Fig. 7. Dose response of epinephrine-induced internalization in the PKA⁻ and comparison with that for activation of ERK1/2 and adenylyl cyclase. Cells in 12-well dishes were treated with various concentrations of epinephrine for 5 min, and internalization of the β_2 AR was measured by surface binding of [³H]CGP-12177 as described under *Materials and Methods*. The data shown are the average values of two identical experiments, each performed in triplicate at the various concentrations of epinephrine (variation in the values was less than 5% at each time point). Results are expressed as the percentage of the total surface binding that was internalized. The data for epinephrine activation of ERK1/2 for the PKA⁻ expressing cells were taken from Fig. 3. The dose response for epinephrine stimulation of adenylyl cyclase in PKA⁻ membrane preparations is from one representative assay.




Fig. 8. Inhibition of epinephrine-, EGF-, and forskolin-activated ERK1/2 by PP2. To assess a possible role for Src in ERK1/2 activation, PKA⁻ cells were pretreated for 1 h at 37°C with 1% DMSO or 10 μ M PP2. The cells were then incubated for 5 min with 100 pM or 10 nM epinephrine, 100 ng/ml EGF, or 20 μ M forskolin. Phosphorylation of ERK1/2 was measured as described under *Materials and Methods*. The data from each experiment were normalized as percentage of maximum ERK1/2 stimulation. The data are the means \pm S.E.M. from four experiments.

tion of ERK1/2 in HEK293 cells by isoproterenol stimulation of the endogenous β_2 AR involved G_s /PKA activation of Rap1 and B raf with no role for G_i . In agreement with this study, we found no pertussis toxin inhibition of epinephrine activation of ERK1/2 in HEK293 cells expressing the PKA⁻ and only a slight inhibition of activation in HEK293 cells expressing only the endogenous β_2 AR, suggesting that G_i played little if any role. In further support of our conclusion concerning the effects of pertussis toxin, we previously reported that conditions (low concentrations of epinephrine) that provoke a PKA-mediated desensitization of the β_2 AR result in only a modest 2- to 3-fold increase in the EC_{50} for epinephrine stimulation of adenylyl cyclase in L cells (Yuan et al., 1994). If the β_2 AR switched to activation of G_i , a considerably larger loss of epinephrine-stimulated adenylyl cyclase activity would be expected. Also, we have not found that pertussis toxin treatment alters the extent of epinephrine-induced desensitization of HEK293 cells (Clark et al., 1996; Seibold et al., 2000) as would be predicted if PKA-mediated switching occurred.

There are a number of reasons that might account for the differences between our work and that of others (Daaka et al., 1997) that were also performed on clones of the HEK293 cells. Clonal cell lines are adapted for fast growth and it is possible that there are important differences in the expression of factors involved in the complex activation of ERK1/2 that shunt activation to alternate pathways in different clones, as has been suggested previously (Liebmann, 2001). That is, it is entirely possible that for unknown but potentially very interesting reasons, the clonal lines of HEK293 cells used by other groups (Daaka et al., 1998; Luttrell et al., 1999a; Maudsley et al., 2000) express such factors as reduced levels of G_i -specific RGS proteins and different levels of receptor binding proteins that alter localization to microdomains in the plasma membrane thus diminishing the coupling efficiency of β_2 AR activation of G_i (Hall et al., 1998; Ostrom et al., 2001). For reasons such as these (and one could imagine many other scenarios) the role of G_i may vary significantly from cell line to cell line. Important experimental differences in the present work were that we used only stable transfection of the β_2 ARs and determined EC_{50} values for epinephrine activation of ERK1/2 with overexpression of the PKA⁻ and wild-type β_2 ARs. EC_{50} values were not determined in the previous study of the PKA⁻ mutant (Daaka et al., 1997). It is possible that the use of transient overexpression paradigms in previous studies actually resulted in much higher levels of receptor expression in a fraction of the cells and/or differential localization, and these factors could affect specificity and/or coupling efficiency of β_2 AR activation of G_s versus G_i . Our studies with transiently expressed β_2 ARs in HEK293 indicate that they couple poorly to G_s /adenylyl cyclase (i.e., they show almost no left shift in EC_{50} for epinephrine activation of adenylyl cyclase with high expression). Finally, it is important not to overinterpret results after an 18-h pretreatment with pertussis toxin (Piiper et al., 2000), because cAMP levels are elevated for prolonged periods.

With regard to a role of internalization in ERK1/2 activation, previous studies based on the use of transient expression of dominant negative blockers of internalization have given somewhat ambiguous results. Taking a different approach that circumvents the need for expression of dominant-negative dynamin or arrestins, which may have nonspecific

effects, we found that activation of ERK1/2 in the PKA⁻ overexpressing cells occurred at approximately 1000-fold lower concentrations ($EC_{50} = 30$ – 60 pM) of epinephrine than those required for internalization ($EC_{50} = 75$ nM), demonstrating a huge amplification of ERK1/2 activation relative to internalization. Our result contrasts with the prior study showing a correlation of the two processes (Daaka et al., 1998). Because we found full ERK1/2 activation with extremely low levels of β_2 AR occupancy that do not cause internalization, there should be no GRK phosphorylation of the β_2 AR and arrestin binding, because it is generally accepted that they are correlated with and required for internalization. The very small amplification of internalization ($EC_{50} = 75$ nM) we observed relative to occupation of the β_2 AR (K_d for epinephrine is ~ 500 – 600 nM) is entirely consistent with previous work that GRK-mediated phosphorylation and binding of β -arrestin occurs at much higher concentrations of epinephrine than those required for PKA activation.

A number of findings suggest that Src activation plays a role in β_2 AR stimulation of ERK1/2 (Daaka et al., 1997; Ma et al., 2000; Maudsley et al., 2000; Fan et al., 2001), and it has been suggested that Src is activated by either formation of a β_2 AR/ β -arrestin/Src complex or by direct G_s or G_i activation of Src. Consistent with a role for Src, we found that the Src family kinase inhibitor PP2 (10 μ M) caused a $>90\%$ inhibition of epinephrine and forskolin activation of ERK1/2 in the HEK293 cells. However, that we see full activation of ERK1/2 by very low concentrations of both epinephrine (which are most unlikely to cause formation of a β_2 AR/ β -arrestin/Src complex) and forskolin (which bypasses the need for receptor or G proteins by direct binding to adenylyl cyclase), raises the question of the mechanism of Src activation. In this regard Schmitt and Stork (2002) recently reported that PKA activates Src by phosphorylation of serine 17 in NIH3T3 cells, as well as in HEK293 cells (P. Stork, personal communication). Whereas this phosphorylation site on Src has been known for some time (Collett et al., 1979), the physiological relevance has not been understood. These studies provide evidence that PKA phosphorylates and activates Src directly and, combined with our studies showing full activation of ERK1/2 by extremely low receptor occupancy in the PKA⁻ cells, leads us to propose that Src is activated by the β_2 AR by the classic G_s /adenylyl cyclase(AC)/PKA pathway and that the dominant pathway for β_2 AR activation of ERK1/2 in HEK293 cells is as follows: β_2 AR $\rightarrow G_s \rightarrow AC \rightarrow PKA \rightarrow Src \rightarrow ERK$. The similar amplification in the potency of ERK1/2 activation relative to adenylyl cyclase activation by the endogenous and overexpressed β_2 ARs is further indication that G_s and PKA are probably major upstream players in ERK1/2 activation.

Acknowledgments

We are grateful for the technical contributions of Bruce Williams and to Dr. Anita Seibold for providing the stably transfected cell lines.

References

- Cerione RA, Staniszewski C, Benovic JL, Lefkowitz RJ, Caron MG, Gierschik P, Somers R, Spiegel AM, Codina J, and Birnbaumer L (1985) Specificity of the functional interactions of the β -adrenergic receptor and rhodopsin with guanine nucleotide regulatory proteins reconstituted in phospholipid vesicles. *J Biol Chem* 260:1493–1500.
- Clark RB, Allal C, Friedman J, Johnson M, and Barber R (1996) Stable activation and desensitization of beta 2-adrenergic receptor stimulation of adenylyl cyclase

- by salmeterol: evidence for quasi-irreversible binding to an exosite. *Mol Pharmacol* **49**:182–189.
- Clark RB, Goka TJ, Proll MA, and Friedman J (1986) Homologous desensitization of beta-adrenergic receptors in lymphoma cells is not altered by the inactivation of Ni (Gi), the inhibitory guanine nucleotide regulatory protein. *Biochem J* **235**:399–405.
- Clark RB, Knoll BJ, and Barber R (1999) Partial agonists and G protein-coupled receptor desensitization. *Trends Pharmacol Sci* **20**:279–286.
- Collett MS, Erikson E, and Erikson RL (1979) Structural analysis of the avian sarcoma virus transforming protein: sites of phosphorylation. *J Virol* **29**:770–781.
- Daaka Y, Luttrell LM, Ahn S, Della Rocca GJ, Ferguson SS, Caron MG, and Lefkowitz RJ (1998) Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. *J Biol Chem* **273**:685–688.
- Daaka Y, Luttrell LM, and Lefkowitz RJ (1997) Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. *Nature (Lond)* **390**:88–91.
- DeGraff JL, Gagnon AW, Benovic JL, and Orsini MJ (1999) Role of arrestins in endocytosis and signaling of alpha2-adrenergic receptor subtypes. *J Biol Chem* **274**:11253–11259.
- Fan G, Shumay E, Malbon CC, and Wang H (2001) c-Src tyrosine kinase binds the beta2-adrenergic receptor via phospho-Tyr-350, phosphorylates G-protein-linked receptor kinase 2 and mediates agonist-induced receptor desensitization. *J Biol Chem* **276**:13240–13247.
- Hall RA, Premont RT, Chow CW, Blitzer JT, Pitcher JA, Claing A, Stoffel RH, Barak LS, Shenolikar S, Weinman EJ, et al. (1998) The beta2-adrenergic receptor interacts with the Na⁺/H⁺-exchanger regulatory factor to control Na⁺/H⁺ exchange. *Nature (Lond)* **392**:626–630.
- January B, Seibold A, Allal C, Whaley BS, Knoll BJ, Moore RH, Dickey BF, Barber R, and Clark RB (1998) Salmeterol-induced desensitization, internalization and phosphorylation of the human beta2-adrenoceptor. *Br J Pharmacol* **123**:701–711.
- January B, Seibold A, Whaley B, Hipkin RW, Lin D, Schonbrunn A, Barber R, and Clark RB (1997) beta2-Adrenergic receptor desensitization, internalization and phosphorylation in response to full and partial agonists. *J Biol Chem* **272**:23871–23879.
- Liebmann C (2001) Regulation of MAP kinase activity by peptide receptor signalling pathway: Paradigms of multiplicity. *Cell Signal* **13**:777–785.
- Luttrell LM, Daaka Y, and Lefkowitz RJ (1999a) Regulation of tyrosine kinase cascades by G-protein-coupled receptors. *Curr Opin Cell Biol* **11**:177–183.
- Luttrell LM, Ferguson SS, Daaka Y, Miller WE, Maudsley S, Della Rocca GJ, Lin F, Kawakatsu H, Owada K, Luttrell DK, et al. (1999b) Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. *Science (Wash DC)* **283**:655–661.
- Ma YC, Huang J, Ali S, Lowry W, and Huang XY (2000) Src tyrosine kinase is a novel direct effector of G proteins. *Cell* **102**:635–646.
- Maudsley S, Pierce KL, Zamah AM, Miller WE, Ahn S, Daaka Y, Lefkowitz RJ, and Luttrell LM (2000) The beta2-adrenergic receptor mediates extracellular signal-regulated kinase activation via assembly of a multi-receptor complex with the epidermal growth factor receptor. *J Biol Chem* **275**:9572–9580.
- Miller WE, Maudsley S, Ahn S, Khan KD, Luttrell LM, and Lefkowitz RJ (2000) beta-arrestin1 interacts with the catalytic domain of the tyrosine kinase c-SRC. Role of beta-arrestin1-dependent targeting of c-SRC in receptor endocytosis. *J Biol Chem* **275**:11312–11319.
- Okamoto T, Murayama Y, Hayashi Y, Inagaki M, Ogata E, and Nishimoto I (1991) Identification of a Gs activator region of the beta 2-adrenergic receptor that is autoregulated via protein kinase A-dependent phosphorylation. *Cell* **67**:723–730.
- Ostrom RS, Gregorian C, Drenan RM, Xiang Y, Regan JW, and Insel PA (2001) Receptor number and caveolar co-localization determine receptor coupling efficiency to adenylyl cyclase. *J Biol Chem* **276**:42063–42069.
- Pierce KL, Maudsley S, Daaka Y, Luttrell LM, and Lefkowitz RJ (2000) Role of endocytosis in the activation of the extracellular signal-regulated kinase cascade by sequestering and nonsequestering G protein-coupled receptors. *Proc Natl Acad Sci USA* **97**:1489–1494.
- Piiper A, Gebhardt R, Kronenberger B, Giannini CD, Elez R, and Zeuzem S (2000) Pertussis toxin inhibits cholecystokinin- and epidermal growth factor-induced mitogen-activated protein kinase activation by disinhibition of the cAMP signaling pathway and inhibition of c-Raf-1. *Mol Pharmacol* **58**:608–613.
- Rubenstein RC, Linder ME, and Ross EM (1991) Selectivity of the beta-adrenergic receptor among Gs, Gi's and Go: assay using recombinant alpha subunits in reconstituted phospholipid vesicles. *Biochemistry* **30**:10769–10777.
- Schmitt JM and Stork PJ (2000) beta2-Adrenergic receptor activates extracellular signal-regulated kinases (ERKs) via the small G protein rap1 and the serine/threonine kinase B-Raf. *J Biol Chem* **275**:25342–25350.
- Schmitt JM and Stork PJ (2002) PKA phosphorylation of Src mediates cAMP's inhibition of cell growth via Rap1. *Mol Cell* **9**:85–94.
- Seibold A, January BG, Friedman J, Hipkin RW, and Clark RB (1998) Desensitization of beta2-adrenergic receptors with mutations of the proposed G protein-coupled receptor kinase phosphorylation sites. *J Biol Chem* **273**:7637–7642.
- Seibold A, Williams B, Huang ZF, Friedman J, Moore RH, Knoll BJ, and Clark RB (2000) Localization of the sites mediating desensitization of the beta2-adrenergic receptor by the GRK pathway. *Mol Pharmacol* **58**:1162–1173.
- Wan Y and Huang XY (1998) Analysis of the G_i/mitogen-activated protein kinase pathway in mutant S49 cells. *J Biol Chem* **273**:14533–14537.
- Whaley BS, Yuan N, Birnbaumer L, Clark RB, and Barber R (1994) Differential expression of the beta-adrenergic receptor modifies agonist stimulation of adenylyl cyclase: a quantitative evaluation. *Mol Pharmacol* **45**:481–489.
- Yuan N, Friedman J, Whaley BS, and Clark RB (1994) cAMP-dependent protein kinase and protein kinase C consensus site mutations of the beta-adrenergic receptor. Effect on desensitization and stimulation of adenylylcyclase. *J Biol Chem* **269**:23032–23038.
- Zou Y, Komuro I, Yamazaki T, Kudoh S, Uozumi H, Kadowaki T, and Yazaki Y (1999) Both Gs and Gi proteins are critically involved in isoproterenol-induced cardiomyocyte hypertrophy. *J Biol Chem* **274**:9760–9770.

Address correspondence to: Richard B. Clark, Department of Integrative Biology and Pharmacology, the University of Texas Health Science Center at Houston Medical School, P.O. Box 20708, Houston, TX 77225. E-mail: richard.b.clark@uth.tmc.edu
