

PERSPECTIVE

Brain-Derived Neurotrophic Factor: Linking Fear Learning to Memory Consolidation

Marie-H. Monfils, Kiriana K. Cowansage, and Joseph E. LeDoux

Center for Neural Science, New York University, New York, New York

Received May 17, 2007; accepted May 23, 2007

ABSTRACT

Brain-derived neurotrophic factor (BDNF), a member of the neurotrophin family, plays an important role in synaptic plasticity. In this issue of *Molecular Pharmacology*, Ou and Gean (p. 350) thoroughly describe the molecular cascade by which fear learning leads to an increase in BDNF expression in the lateral amygdala (LA). Calcium influx through *N*-methyl-D-aspartate receptors and L-type voltage-dependent calcium channels,

which occurs in the LA during fear conditioning, activates protein kinase A and Ca²⁺/calmodulin-dependent protein kinase IV. Each induces phosphorylation of cAMP response element-binding protein, which binds to the BDNF promoter, leading to BDNF expression in the LA, and contributes to fear memory consolidation.

The activity-dependent modification of synapses, a process known as synaptic plasticity, permits the brain to generate efficient neural networks that facilitate advantageous behavioral adaptations. Although an extensive body of research has demonstrated the importance of synaptic strengthening in the beneficial functions of learning and memory, recent evidence also suggests that a host of pathologic conditions, including mood disorders and drug addiction, engage overlapping mechanisms. Thus, the elucidation of molecular pathways involved in driving these physiological changes has broad clinical implications (Malenka and Bear, 2004; MacKinnon and Zamoiski, 2006).

The neurotrophins, a family of structurally related proteins known for their role in promoting neuronal differentiation and survival during development (Levi-Montalcini, 1987; Leibrock et al., 1989; Barde, 1994), have recently surfaced as playing an important role in mediating synaptic plasticity (Schinder and Poo, 2000; Lu, 2003, 2004; Bramham and Messaoudi, 2005; Arancio and Chao, 2007). In this issue

of *Molecular Pharmacology*, Ou and Gean (2007) report on their investigation of the mechanism by which one member of this family, brain-derived neurotrophic factor (BDNF), mediates fear memory consolidation in the amygdala. BDNF and its main receptor, TrkB, are fast emerging as major regulators of synaptic transmission and plasticity in the adult brain. In mammals, BDNF is synthesized, stored, and released from excitatory neurons containing glutamate (Lessmann et al., 2003) and, in some populations, dopamine (Berton et al., 2006).

Past investigations into the role of BDNF in plasticity have predominantly focused on the hippocampus, a structure traditionally associated with learning and memory. In that structure, TrkB receptors have been specifically localized to the pre- and postsynaptic elements of glutamatergic synapses (Drake et al., 1999) and coimmunoprecipitate with the transmembrane NMDA receptor protein, which allows calcium entry into the cell in response to detection of coincidental rises in both presynaptic glutamate and postsynaptic voltage (Aoki et al., 2000). Because this calcium influx is necessary for the initiation of cellular changes, the colocalization of NMDA receptors with BDNF and its receptor, TrkB, at synaptic junctions sets the stage to synchronize bidirectional synaptic optimization.

M.-H.M. and K.K.C. contributed equally to this work
Article, publication date, and citation information can be found at
<http://molpharm.aspetjournals.org>
doi:10.1124/mol.107.038232.
Please see the related article on page 350.

ABBREVIATIONS: BDNF, brain-derived neurotrophic factor; NMDA, *N*-methyl-D-aspartate; LA, lateral amygdala; LTP, long-term potentiation; MAPK, mitogen-activated protein kinase; MEK, mitogen-activated protein kinase kinase; L-VDCC, L-type voltage-dependent calcium channel; PKA, protein kinase A; CaMKIV, Ca²⁺/calmodulin-dependent protein kinase IV; CREB, cAMP response element-binding protein.

Much evidence indicates that learning initiates alterations in glutamate-dependent excitatory synaptic transmission that subsequently stabilize through structural changes at postsynaptic sites on dendritic spines (for review, see Lamprecht and LeDoux, 2004). Direct infusion of BDNF into the hippocampus enhances synaptic strength, both *in vitro* (Kang and Schuman, 1995; Levine et al., 1995) and *in vivo* (Messaoudi et al., 1998), as well as modulates the induction of long-term potentiation (Patterson et al., 1996; Messaoudi et al., 2002) and structural changes in dendritic spines (Alonso et al., 2004). Recent work by Tyler et al. (2002) also suggests that BDNF activation is necessary for the learning-induced modification of hippocampal spines, which may also involve the activation of TrkB (von Bohlen und Halbach, 2006). Furthermore, Bekinschtein et al. (2007) recently demonstrated that BDNF-dependent storage of long-term memories occurs within hours after acquisition of an associative learning task, suggesting that BDNF is likely to be involved in memory stabilization.

Despite the traditional popularity of the hippocampus in all matters of learning and memory, there is increasing empirical support for the role of another structure—the amygdala—in the types of synaptic changes facilitated by BDNF. In particular, mounting evidence now indicates a role for BDNF signaling in the basal and lateral nuclei of the amygdala (Rattiner et al., 2004a; Ou and Gean, 2006), areas known to be necessary for the formation of learned fear associations (LeDoux, 2000). Because the amygdala has been implicated in many pathologic conditions, including post-traumatic stress (Garakani et al., 2006), anxiety (Rauch et al., 2006), and autism spectrum disorders (Baron-Cohen et al., 2000; Bachevalier and Loveland, 2006), considerable efforts have been devoted to the characterization of this circuitry as a central site for emotion-induced neuronal plasticity (LeDoux, 2000; Maren, 2001; Paré et al., 2004; Wilensky et al., 2006). A number of studies have shown, using Pav-

lovian fear conditioning paradigms, that the physiological basis for such changes begins with the relay of sensory information from the medial geniculate nucleus of the thalamus to the lateral amygdala (LA), where the initial association is made via an LTP-like mechanism, followed by the intra-amygdala transfer of signals to the central nucleus of the amygdala, which facilitates the expression of a fear response by way of projections to brainstem and hypothalamic targets (Davis, 1997; LeDoux, 2000; Paré et al., 2004). Together, these findings support the notion that changes in synaptic strength are required for the acquisition of emotional memories. At a more profound level, however, our knowledge of these larger-scale anatomical modifications remains bound by our more limited comprehension of the molecular machinery governing those changes.

A recent study found temporally specific increases in BDNF gene expression to occur in the basal/lateral portion of the amygdala (BLA) after paired stimuli that supported learning, but not after exposure to neutral or aversive stimuli alone (Rattiner et al., 2004a). BDNF signaling through TrkB receptors was also found to be necessary for the consolidation of fear memories (Rattiner et al., 2004a). In agreement with the findings of Rattiner et al. (2004a), Ou and Gean (2006) reported increases in BDNF protein expression and activation of TrkB receptors in the amygdala. Their study further revealed that intra-amygdala infusion of a TrkB ligand scavenger or the inhibition of Trk receptors impaired fear memory assessed 24 h after training. In addition, they showed that BDNF phosphorylates mitogen-activated protein kinase (MAPK), and this is blocked by the Trk receptor inhibitor K252a (Ou and Gean, 2006). The BDNF-induced phosphorylation of MAPK occurs via Shc binding to the TrkB receptor, which leads to the activation of Ras, Raf, MEK, and MAPK. BDNF also phosphorylates MAPK via activation of PI-3 kinase (Ou and Gean, 2006).




Fig. 1. BDNF signaling cascade involved during fear conditioning. Calcium influx through NMDA receptors and L-VDCC channels, which occurs in the LA during fear conditioning, activates adenylyl cyclase (AC) and PKA. Activated PKA translocates to the nucleus and induces CREB phosphorylation. Increase in intracellular calcium also activates CaMKIV and leads to phosphorylation of CREB at Ser-133. Activated CREB binds to the BDNF promoter, leading to BDNF expression in the LA, and contributes to fear memory consolidation (from Ou and Gean, 2007). Fear conditioning is also associated with binding of BDNF to TrkB receptors. This results in the association of Shc and TrkB receptor [structure of ligand binding domain after the work of Ultsch et al. (1999)], and leads to the activation of Ras, Raf, MEK, and MAPK (not shown). BDNF also phosphorylates MAPK via activation of PI3 kinase (Ou and Gean, 2006) (not shown).

In this issue of *Molecular Pharmacology*, Ou and Gean (2007) extend their earlier work to a characterization of the molecular cascades underlying fear conditioning that exert transcriptional and translational control over BDNF expression in the amygdala. Consistent with the previous work of Rattiner et al. (2004b), they show a significant increase in BDNF exon I- and III-containing mRNA in the amygdala of fear-conditioned rats. Inhibition of protein synthesis and translation, using intra-LA anisomycin or actinomycin D, respectively, attenuates this increase in fear-conditioning-induced BDNF expression. Furthermore, they demonstrate that the increase in BDNF depends on the activation of NMDA receptors as well as L-type voltage-dependent calcium channels (L-VDCC), the blockade of which significantly attenuates BDNF expression. A similar reduction was also apparent after the pharmacological inhibition of PKA and CaMKIV activity. In addition, through the use of DNA affinity precipitation and chromatin immunoprecipitation assays, Ou and Gean (2007) demonstrate a specific increase in the binding of phosphorylated cAMP response element binding protein (CREB) to exon I and III promoters after fear conditioning. They found that sequestration of endogenous BDNF during fear conditioning by infusion of a TrkB IgG did not affect the BDNF protein level increases typically observed 1 h after conditioning. This suggests that whereas BDNF signaling through TrkB receptors in the amygdala is required for long-term memory (Rattiner et al., 2004a; Ou and Gean, 2006), it is not necessary to regulate the increase in BDNF protein levels induced by fear conditioning (Ou and Gean, 2007).

Many of the most common psychiatric disorders that afflict humans are emotional disorders, a number of which involve the activation of fear circuitry in the brain. To develop suitable treatments for anxiety-related disorders, it is necessary to develop a better understanding of the molecular mechanisms that underlie their development and manifestation. Ou and Gean's (2007) findings elegantly illustrate that calcium influx through NMDA receptors and L-VDCC channels, known to occur during fear conditioning, activates PKA and CaMKIV, each inducing CREB phosphorylation. In turn, phosphorylated CREB binds to the BDNF promoter, leading to an increase in BDNF expression in the amygdala, and probably contributes to fear memory consolidation (see Fig. 1). Ou and Gean (2007) describe a tight molecular cascade linking the initial physiological events that take place during fear conditioning to the expression of BDNF—a potent modulator of synaptic plasticity that could lead to the restructuring of synapses in the LA. Taken together, these findings implicate the BDNF signaling cascade in the amygdala as a potential target for novel pharmacological interventions.

References

- Aoki C, Wu K, Elste A, Len G, Lin S, McAuliffe G, and Black IB (2000) Localization of brain-derived neurotrophic factor and TrkB receptors to postsynaptic densities of adult rat cerebral cortex. *J Neurosci Res* **59**:454–463.
- Alonso M, Medina JH, Pozzo-Miller L (2004) ERK1/2 activation is necessary for BDNF to increase dendritic spine density in hippocampal CA1 pyramidal neurons. *Learn Mem* **11**:172–178.
- Arancio O and Chao MV (2007) Neurotrophins, synaptic plasticity and dementia. *Curr Opin Neurobiol*, in press.
- Bachevalier J and Loveland KA (2006) The orbitofrontal-amygdala circuit and self-regulation of social-emotional behavior in autism. *Neurosci Biobehav Rev* **30**:97–117.
- Baron-Cohen S, Ring HA, Bullmore ET, Wheelwright S, Ashwin C, and Williams SC (2000) The amygdala theory of autism. *Neurosci Biobehav Rev* **24**:355–364.
- Barde (1994) Neurotrophins: a family of proteins supporting the survival of neurons. *Prog Clin Biol Res* **390**:45–56.
- Beckinschtein P, Cammarota M, Igaz LM, Bevilacqua LR, Izquierdo I, and Medina JH (2007) Persistence of long-term memory storage requires a late protein synthesis- and BDNF- dependent phase in the hippocampus. *Neuron* **53**:261–277.
- Berton O, McClung CA, Dileone RJ, Krishnan V, Renthal W, Russo SJ, Graham D, Tsankova NM, Bolanos CA, Rios M, et al. (2006) Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* **311**:864–868.
- Bramham CR and Messaoudi E (2005) BDNF function in adult synaptic plasticity: the synaptic consolidation hypothesis. *Prog Neurobiol* **76**:99–125.
- Davis M (1997) Neurobiology of fear responses: the role of the amygdala. *J Neuro-psychiatry Clin Neurosci* **9**:382–402.
- Drake CT, Milner TA, and Patterson SL (1999) Ultrastructural localization of full-length trkB immunoreactivity in rat hippocampus suggests multiple roles in modulating activity-dependent synaptic plasticity. *J Neurosci* **19**:8009–8026.
- Garakani A, Mathew SJ, and Charney DS (2006) Neurobiology of anxiety disorders and implications for treatment. *Mt Sinai J Med* **73**:941–949.
- Kang H and Schuman EM (1995) Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. *Science* **267**:1658–1662.
- Lamprecht and LeDoux (2004) Structural plasticity and memory. *Nat Rev Neurosci* **5**:45–54.
- LeDoux JE (2000) Emotion circuits in the brain. *Annu Rev Neurosci* **23**:155–184.
- Leibrock J, Lottspeich F, Hohn A, Hofer M, Hengerer B, Masiakowski P, Thoenen H, and Barde YA (1989) Molecular cloning and expression of brain-derived neurotrophic factor. *Nature* **341**:149–152.
- Lessmann V, Gottmann K, and Malcangio M (2003) Neurotrophin secretion: current facts and future prospects. *Prog Neurobiol* **69**:341–374.
- Levi-Montalcini R (1987) The nerve growth factor 35 years later. *Science* **237**:1154–1162.
- Levine ES, Dreyfus CF, Black IB, and Plummer MR (1995) Brain-derived neurotrophic factor rapidly enhances synaptic transmission in hippocampal neurons via postsynaptic tyrosine kinase receptors. *Proc Natl Acad Sci U S A* **92**:8074–8077.
- Lu B (2003) BDNF and activity-dependent synaptic modulation. *Learn Mem* **10**:86–98.
- Lu B (2004) Acute and long-term synaptic modulation by neurotrophins. *Prog Brain Res* **146**:137–150.
- MacKinnon and Zamoiski (2006) Panic comorbidity with bipolar disorder: what is the manic-panic connection? *Bipolar Disord* **8**:648–664.
- Malenka RC and Bear MF (2004) LTP and LTD: an embarrassment of riches. *Neuron* **44**:5–21.
- Maren S (2001) Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* **24**:897–931.
- Messaoudi E, Bardsen K, Srebro B, and Bramham CR (1998) Acute intrahippocampal infusion of BDNF induces lasting potentiation of synaptic transmission in the rat dentate gyrus. *J Neurophysiol* **79**:496–499.
- Messaoudi E, Ying SW, Kanhema T, Croll SD, and Bramham CR (2002) Brain-derived neurotrophic factor triggers transcription-dependent, late phase long-term potentiation in vivo. *J Neurosci* **22**:7453–7461.
- Ou LC and Gean PW (2006) Regulation of amygdala-dependent learning by brain-derived neurotrophic factor is mediated by extracellular signal-regulated kinase and phosphatidylinositol-3-kinase. *Neuropsychopharmacology* **31**:287–296.
- Ou LC and Gean PW (2007) Transcriptional regulation of brain-derived neurotrophic factor in the amygdala during consolidation of fear memory. *Mol Pharmacol* **72**:350–358.
- Paré D, Quirk GJ, and LeDoux JE (2004) New vistas on amygdala networks in conditioned fear. *J Neurophysiol* **92**:1–9.
- Patterson SL, Abel T, Deuel TA, Martin KC, Rose JC, and Kandel ER (1996) Recombinant BDNF rescues deficits in basal synaptic transmission and hippocampal LTP in BDNF knockout mice. *Neuron* **16**:1137–1145.
- Rattiner LM, Davis M, French CT, and Ressler KJ (2004a) Brain-derived neurotrophic factor and tyrosine kinase receptor B involvement in amygdala-dependent fear conditioning. *J Neurosci* **24**:4796–4806.
- Rattiner LM, Davis M, and Ressler KJ (2004b) Differential regulation of brain-derived neurotrophic factor transcripts during the consolidation of fear learning. *Learn Mem* **11**:727–731.
- Rauch SL, Shin LM, and Phelps EA (2006) Neurocircuitry models of posttraumatic stress disorder and extinction: human neuroimaging research—past, present, and future. *Biol Psychiatry* **60**:376–382.
- Schinder AF and Poo M (2000) The neurotrophin hypothesis for synaptic plasticity. *Trends Neurosci* **23**:639–645.
- Tyler WJ, Alonso M, Bramham CR, and Pozzo-Miller LD (2002) From acquisition to consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-dependent learning. *Learn Mem* **9**:224–237.
- Ullsch MH, Wiesmann C, Simmons LC, Henrich J, Yang M, Reilly D, Bass SH, and de Vos AM (1999) Crystal structures of the neurotrophin-binding domain of TrkA, TrkB, and TrkC. *J Mol Biol* **290**:149–159.
- von Bohlen and Halbach O, Krause S, Medina D, Sciarretta C, Minichiello L, and Unsicker K (2006) Regional- and age-dependent reduction in trkB receptor expression in the hippocampus is associated with altered spine morphologies. *Biol Psychiatry* **59**:793–800.
- Wilensky AE, Schafe GE, Kristensen MP, and LeDoux (2006) Rethinking the fear circuit: the central nucleus of the amygdala is required for the acquisition, consolidation, and expression of Pavlovian fear conditioning. *J Neurosci* **26**:12387–12396.

Address correspondence to: Marie-H. Monfils, Center for Neural Science, 4 Washington Place, Room 809, New York University, New York, NY 10003-6621. E-mail: monfils@cns.nyu.edu