

MINIREVIEW

# Tristetraprolin and Its Role in Regulation of Airway Inflammation

Pavan Prabhala and Alaina J. Ammit

Faculty of Pharmacy, University of Sydney, Sydney, New South Wales, Australia

Received September 17, 2014; accepted November 26, 2014

## ABSTRACT

Chronic inflammatory diseases, such as asthma and chronic obstructive pulmonary disease (COPD), are clinically and socioeconomically important diseases globally. Currently the mainstay of anti-inflammatory therapy in respiratory diseases is corticosteroids. Although corticosteroids have proven clinical efficacy in asthma, many asthmatic inflammatory conditions (e.g., infection, exacerbation, and severe asthma) are not responsive to corticosteroids. Moreover, despite an understanding that COPD progression is driven by inflammation, we currently do not have effective anti-inflammatory strategies to combat this disease. Hence, alternative anti-inflammatory strategies are required. p38 mitogen-activated protein kinase (MAPK) has emerged as an important signaling molecule driving airway inflammation, and pharmacological inhibitors against p38 MAPK may provide potential

therapies for chronic respiratory disease. In this review, we discuss some of the recent *in vitro* and *in vivo* studies targeting p38 MAPK, but suggest that p38 MAPK inhibitors may prove less effective than originally considered because they may block anti-inflammatory molecules along with proinflammatory responses. We propose that an alternative strategy may be to target an anti-inflammatory molecule farther downstream of p38 MAPK, i.e., tristetraprolin (TTP). TTP is an mRNA-destabilizing, RNA-binding protein that enhances the decay of mRNAs, including those encoding proteins implicated in chronic respiratory diseases. We suggest that understanding the molecular mechanism of TTP expression and its temporal regulation will guide future development of novel anti-inflammatory pharmacotherapeutic approaches to combat respiratory disease.

## Introduction

Many chronic inflammatory diseases are a consequence of overactive inflammatory signaling pathways. Hence, repressing these pathways as potential drug targets represents a way in which to re-establish control and attenuate the severity of various chronic inflammatory diseases. However, these pathways are not exclusively responsible for proinflammatory signaling, and targeting these pathways may cause the ablation of other vital, anti-inflammatory signals necessary for restoring normal physiology. We propose that this may be the case with targeting one of the members of the mitogen-activated protein kinase (MAPK) superfamily, p38 MAPK, in respiratory disease. That is, while p38 MAPK inhibitors may repress proinflammatory cytokines, they may also inhibit the expression of crucial anti-inflammatory proteins, such as tristetraprolin (TTP). TTP is an mRNA-destabilizing, RNA-binding protein that promotes

the decay of the mRNAs of many proteins, including those implicated as playing a role in chronic respiratory diseases. In this review, we focus on the role of the p38 MAPK signaling pathway in respiratory disease and highlight some *in vitro* and *in vivo* evidence with p38 MAPK inhibitors. We outline the role of TTP in the regulation of inflammation in other disease settings and highlight the potential of understanding the molecular mechanism of TTP expression and its temporal regulation for future development of novel anti-inflammatory pharmacotherapeutic approaches in respiratory disease.

## MAPK

Members of the MAPK superfamily play important roles in the regulation of airway inflammation in respiratory disease pathogenesis. Three well defined phosphoproteins, extracellular signal-related kinase (ERK), c-Jun N-terminal kinase, and p38 MAPK, are the MAPKs central to this signaling cascade. The signaling cascade that results in the activation of MAPKs is derived from a relatively linear framework (Fig. 1). Stimuli such as cellular stressors or mitogens interact in a predominantly receptor-mediated manner to trigger the phosphorylation of

This research was supported by the National Health and Medical Research Council of Australia [Grant APP1025637] and philanthropic funding from Maurice Renshaw (President, Faculty of Pharmacy Foundation). P.P. received scholarship support from Renshaw.  
dx.doi.org/10.1124/mol.114.095984.


**ABBREVIATIONS:** ARE, adenosine/uridine-rich element; BIRB 796, 1-(3-tert-butyl-1-*p*-tolyl-1*H*-pyrazol-5-yl)-3-(4-(2-morpholinoethoxy)naphthalen-1-yl)urea; COPD, chronic obstructive pulmonary disease; ERK, extracellular signal-related kinase; IL, interleukin; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; miR, microRNA; NF- $\kappa$ B, nuclear factor  $\kappa$ B; PH-797804, 3-(3-bromo-4-((2,4-difluorobenzyl)oxy)-6-methyl-2-oxopyridin-1(2*H*)-yl)-*N*,4-dimethylbenzamide; PKC, protein kinase C; SB202190, 4-(4-fluorophenyl)-2-(4-hydroxyphenyl)-5-(4-pyridyl)-1*H*-imidazole; SB203580, 4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)imidazole; SB239063, *trans*-4-[4-(4-fluorophenyl)-5-(2-methoxy-4-pyrimidinyl)-1*H*-imidazol-1-yl]cyclohexanol; TNF, tumor necrosis factor; TTP, tristetraprolin; UTR, untranslated region.

a specific MAPK kinase kinase, which then causes the phosphorylation of its specific downstream MAPK kinase. Once the MAPK kinase is phosphorylated, it is subsequently able to phosphorylate its own substrate MAPK. Activated MAPKs then represent a point of divergence leading to the phosphorylation of numerous downstream transcription factors, protein kinases, and proto-oncogenes (Ashwell, 2006; Hu et al., 2007; Balakathiresan et al., 2009). In this review, we focus on p38 MAPK and its downstream signaling effectors. The p38 MAPK-mediated pathway regulates a plethora of cellular processes that drive chronic airway inflammatory disease (Chung, 2011). Extensive investigation of the p38 MAPK-driven molecular mechanisms responsible for respiratory disease pathogenesis has led to drug discovery programs that target this pathway as an approach to potential anti-inflammatory therapy. Moreover, p38 MAPK has also been significantly implicated in corticosteroid insensitivity (Ammit, 2013).

### p38 MAPK: Role in Airway Inflammation and Corticosteroid Insensitivity

There are four isoforms of p38 MAPK— $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ —transcribed by four separate genes and characterized by

different expression patterns. p38 $\alpha$  is expressed ubiquitously and, along with p38 $\beta$ , is considered to represent the major isoform responsible for inflammatory responses. Inflammation is a central component of many respiratory diseases, especially asthma and chronic obstructive pulmonary disease (COPD), and the use of selective pharmacological inhibitors has positioned p38 MAPK as playing a major role in disease severity and progression. The majority of the *in vitro* studies have used the inhibitor SB203580 [4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)imidazole], a pyridinyl imidazole that binds competitively at the ATP-binding site of p38 MAPK. It is widely considered to predominantly target p38 $\alpha$  and p38 $\beta$ , but not the  $\delta$  and  $\gamma$  isoforms of p38 MAPK, although questions regarding selectivity have been raised, especially at higher doses (Bain et al., 2007). Nevertheless, extensive investigations utilizing this first-generation inhibitor have yielded much valuable information regarding the role of p38 MAPK in chronic respiratory disease. p38 MAPK regulates the immunomodulatory function of various airway cells through the control of chemokine and cytokine release (Amrani et al., 2001; Hallsworth et al., 2001; Henness et al., 2006; Quante et al., 2008). p38 MAPK is also responsible for neutrophil and eosinophil migration (Rousseau et al., 1997);



**Fig. 1.** Current and potential inhibitory targets to control p38 MAPK-dependent signaling in respiratory disease. The three members of the MAPK superfamily—ERK, c-Jun N-terminal kinase (JNK), and p38 MAPK (p38)—are stimulated via a series of steps involving MAPK kinase kinases (MAPKKKs) phosphorylating downstream MAPK kinases (MAPKKs), then ultimately MAPKs. This figure focuses on downstream effectors of the p38 MAPK pathway, and the numbers indicate inhibitory strategies along this pathway. While (1) and (2) will repress p38 MAPK, we propose that blocking p38 MAPK may have a negative impact on anti-inflammatory proteins, as TTP is regulated downstream of p38 MAPK. Thus, we propose (3) to control the phosphorylation status and activity of TTP. PP2A is an endogenous protein able to activate TTP. Once active, TTP is able to elicit anti-inflammatory actions by destabilizing cytokine mRNA after recruiting CAF1 and CNOT1.

altering the airway smooth muscle contractility, causing obstruction of airflow (Lakser et al., 2002; Bhavsar et al., 2008); releasing degradative enzymes, like matrix metalloproteinases (Woo et al., 2004); and causing the remodeling of airway architecture by increasing smooth muscle proliferation (Nath et al., 2006).

p38 MAPK has also been significantly implicated in corticosteroid insensitivity (Bhavsar et al., 2008; Chang et al., 2012). Inhaled corticosteroids are the mainstay of chronic asthma treatment and are used effectively at low doses for the majority of asthmatics. However, there is a subset of the asthmatic population with severe asthma who need maximal doses of inhaled corticosteroids and another small subset of the asthmatic population who are resistant to corticosteroids. Details of the molecular mechanisms thought to be responsible for corticosteroid insensitivity and resistance have been outlined in recent reviews (Ammit, 2013; Barnes, 2013). Interestingly, p38 MAPK isoforms, p38 $\gamma$  (Bhattacharyya et al., 2011) and p38 $\alpha/\beta$  (Mercado et al., 2012) in particular, may serve as legitimate targets for restoring corticosteroid responsiveness. Thus, targeting p38 MAPK, perhaps in an isoform-specific manner, might prove to have clinical efficacy as an anti-inflammatory treatment in chronic respiratory disease, especially in patients who respond very poorly to inhaled corticosteroids.

### Use of p38 MAPK Inhibitors To Repress Inflammatory Disease: What Have We Learned?

First-generation inhibitors of p38 MAPK include pharmacological inhibitors like SB203580, BIRB 796 [1-(3-tert-butyl-1-p-tolyl-1H-pyrazol-5-yl)-3-(4-(2-morpholinoethoxy)naphthalen-1-yl)urea], SB202190 [4-(4-fluorophenyl)-2-(4-hydroxyphenyl)-5-(4-pyridyl)-1H-imidazole], and SB239063 [*trans*-4-[4-(4-fluorophenyl)-5-(2-methoxy-4-pyrimidinyl)-1H-imidazol-1-yl]cyclohexanol]. These pyridinyl imidazole compounds competitively antagonize the ATP-binding site, thereby blocking the activity of p38 MAPK (Lee et al., 1994). Recent studies of p38 MAPK inhibitors have used these compounds as a template to create a new set of slightly remodeled and more selective p38 MAPK inhibitors. These inhibitors offer a broad range of anti-inflammatory effects because p38 MAPK is responsible for a large portion of inflammatory responses. Inhibiting the p38 MAPK pathway in animal models showed a dramatic reduction in levels of proinflammatory cytokines and thereby reasonably attenuated chronic inflammatory responses (Young et al., 1993; Ronkina et al., 2010). These compounds have not yet been used in the treatment of asthma, but they have been used marginally in clinical trials for rheumatoid arthritis (Haddad, 2001) and Crohn's disease (Schreiber et al., 2006).

More recently, three molecules have been trialed as potential anti-inflammatory molecules in COPD. Losmapimod, an oral inhibitor of p38 MAPK, was shown to be effective in improving plasma fibrinogen levels and levels of hyperinflation in patients with COPD (Lomas et al., 2012). However, it was also revealed to be ineffective in reducing interleukin (IL)-6, IL-8, and C-reactive protein levels. Similarly, another p38 MAPK inhibitor, dilmapiomod, was administered as a single oral dose in a 4-week clinical trial and demonstrated reduction in phosphorylated heat shock protein 27 and tumor necrosis factor (TNF)  $\alpha$  production (Singh et al., 2010). Similarly to losmapimod, dilmapiomod was also able to reduce serum fibrinogen and forced vital capacity but unable to alter levels of IL-6 or

IL-8. Another 6-week clinical study has suggested that a small-molecule p38 MAPK inhibitor, PH-797804 [3-(3-bromo-4-((2,4-difluorobenzyl)oxy)-6-methyl-2-oxopyridin-1(2H)-yl)-N,4-dimethylbenzamide] (Xing et al., 2012), was effective in increasing some measurable airway criteria like forced expiratory volume in 1 second. This study also reported that PH-797804 reduced systemic fibrinogen and circulating cytokines in patients with moderate to severe COPD in comparison with patients given placebo. A randomized clinical trial assessing safety and efficacy of PH-797804 in COPD has been conducted (MacNee et al., 2013), demonstrating improvements over placebo, albeit with a limited sample size. This has led to "cautious optimism" (Singh, 2013), although there is the possibility of unwanted effects and potential for transient efficacy of p38 MAPK inhibition, as observed with rheumatoid arthritis.

Efficacy in these trials can be used as a yardstick to determine if any generation of p38 MAPK inhibitors can be used as monotherapy or in combination with corticosteroids. But there are important lessons to be learned from trials of first-generation p38 MAPK inhibitors in rheumatoid arthritis. The trials have been largely unsuccessful in rheumatoid arthritis due to the unexpected inefficacy and side effects. The side effects associated with the use of p38 MAPK inhibitors, such as hepatotoxicity, dizziness, and skin rash, may occur as a result of potential nonspecific inhibition (Heinrichsdorff et al., 2008). The unexpected inefficacy could also be attributed to the activation of alternate pathways to induce proinflammatory molecules, including nuclear factor  $\kappa$ B (NF- $\kappa$ B) (Hammaker and Firestein, 2010). Notably, Clark and Dean (2012) have raised the important issue of p38 MAPK-driven anti-inflammatory proteins to explain the apparent failure of p38 MAPK inhibitors in the rheumatology clinic. They state, "It is likely that the failure of p38 inhibitors is connected to the important role of the p38 pathway in the negative feedback control of inflammatory responses" (Clark and Dean, 2012). That is, inhibiting p38 MAPK also blocks the endogenous anti-inflammatory proteins present that naturally curtail the inflammatory response. Important anti-inflammatory molecules, including IL-10, MAPK phosphatase-1, and TTP, are upregulated in response to increased p38 MAPK activation. Hence, p38 MAPK inhibitors represent a double-edged sword. Going farther upstream in the MAPK signaling pathway (see Fig. 1) could be an alternative option as the therapeutic efficacy could be higher, but there is a greater risk of incurring further nonspecific side effects. Another possible option could be to target the anti-inflammatory molecules that regulate the MAPK pathway. Preventing the suppression or prolonging the activation of the transiently activated anti-inflammatory molecules could be of therapeutic benefit. This is the promise of TTP.

### Alternative Anti-Inflammatory Molecule: TTP

TTP is a critical anti-inflammatory protein that functions to promote mRNA decay. Many clinically important proteins, including cytokines, are controlled by post-transcriptional mRNA-stabilizing mechanisms (Brooks and Blackshear, 2013). Notably, many of these proteins have been implicated in asthma (Table 1). We assert that an important consideration that will likely affect the success of p38 MAPK inhibitors in respiratory disease is that the expression and function of this critical anti-inflammatory protein is controlled by p38 MAPK.

Because of this, inhibiting p38 MAPK to repress proinflammatory proteins will also repress important anti-inflammatory molecules such as TTP.

### TTP Is an mRNA-Destabilizing, RNA-Binding Protein That Enhances mRNA Decay

A role for TTP as a *trans*-acting anti-inflammatory protein was first elucidated when the TTP knockout mouse developed a proinflammatory phenotype due to the overexpression of the cytokine TNF- $\alpha$  in macrophages, resulting in cachexia, myeloid hyperplasia, and a host of other inflammatory conditions (Taylor et al., 1996a). Based on evidence from earlier work (Caput et al., 1986), it was concluded that TTP confers mRNA instability and degradation by binding the conserved adenosine/uridine-rich element (ARE) present within the 3' untranslated region (UTR) of mRNA transcripts containing the nonameric sequence UUAUUUAUU (Carballo et al., 1998; Lai et al., 1999). This promotes the poly(A) tail shortening shown in transcripts of granulocyte-macrophage colony-stimulating factor and TNF- $\alpha$  (Carballo et al., 2000; Marchese et al., 2010). The importance of this *cis*-element is highlighted by its presence in ~8% of the human transcriptome (Bakheet et al., 2006).

Regulation of inflammatory mediators can occur at the transcriptional level, controlling gene expression, or it can occur at the post-transcriptional level, whereby mRNA stability determines whether a particular transcript is translated to protein or is rapidly degraded. Regulation can also occur at the post-translational level by modulating the stability and

longevity of translated proteins. This typically involves either preventing the degradation of the protein (via the proteasome) or modifying it to increase its stability. Although all three levels of regulatory control can be targeted to modulate an inflammatory response, post-transcriptional modification presents a feasible opportunity to exert fine control over the proinflammatory process.

The mRNA transcripts of many inflammatory mediators are unstable and are regulated post-transcriptionally; this ensures that low protein levels are maintained under homeostatic conditions. However, they are rapidly induced and then degraded depending on the nature of the extracellular cues. In this way, immunomodulatory cells are able to react with the appropriate physiologic responses to match the stimuli. Inflammatory conditions arise when inflammatory proteins are induced at a time they are not needed or in an amount that is excessive. Hence, mRNA stability plays a crucial role in the pathogenesis of inflammatory diseases. This is evident in the pathogenesis of asthma. Many of the characteristics associated with asthma, such as basement membrane thickening, increased interstitial matrix deposition, airway smooth muscle mass increase, increased airway fibrosis, and neovascularization in the subepithelial mucosa, can be attributed to elevated levels of the inflammatory mediators that give rise to these characteristics (Ammit, 2005). As outlined in Table 1, many of the TTP target mRNAs encode proteins responsible for these disease characteristics (Tang and Fiscus, 2001; Mattos et al., 2002; Hirst, 2003; Batra et al., 2007; Kato and Schleimer, 2007; Barnes, 2008; Ishmael, 2011; Kang et al., 2011; Rael and

TABLE 1  
TTP targets implicated in asthma: potential role of TTP in airway inflammation  
Adapted from Brooks and Blackshear (2013).

TTP Target Implicated in Asthma	Function in Asthma Pathogenesis	Cell Lines/Cell Types in which Target Is Shown To Be Regulated by TTP
IL-17A	Inducing and mediating proinflammatory responses	Human T-cell lines (Lee et al., 2012)
IL-6	Proinflammatory (neutrophil production)	Various (Neininger et al., 2002; Jalonen et al., 2006; Patil et al., 2008; Tudor et al., 2009; Van Tubergen et al., 2011; Zhao et al., 2011; Brahma et al., 2012; Mercado et al., 2012; Chafin et al., 2013; Banerjee et al., 2014)
IL-1 $\beta$	Many proinflammatory functions	Mouse RAW264.7 cells (Chen et al., 2006)
TNF- $\alpha$	Many proinflammatory functions	Mouse macrophages (Taylor et al., 1996a; Carballo et al., 1998)
CCL2	Recruitment of inflammatory mediators	Mouse macrophages, fibroblasts (Sauer et al., 2006)
CXCL8	Recruitment of neutrophils	Various cell lines (Suswam et al., 2008; Balakathiresan et al., 2009; Bhattacharyya et al., 2011; Al Ghoulh and Magder, 2012; Galbiati et al., 2012; Mercado et al., 2012; Suswam et al., 2013; Colin et al., 2014)
GM-CSF	Increasing levels of neutrophils and eosinophils	Mouse bone marrow-derived stromal cells (Carballo et al., 2000)
VEGF	Promoting angiogenesis	Various (Essafi-Benkhadir et al., 2007, 2010; Suswam et al., 2008; Lee et al., 2010; Bhattacharyya et al., 2011; Banerjee et al., 2014)
IL-13	Involved in airway remodeling	Mast cells (Barnstein et al., 2006)
IL-23	Neutrophil infiltration	Mouse dendritic cells, macrophages (Qian et al., 2011; Molle et al., 2013)
CCL5	Chemotaxis of chemokines and cytokines	Mouse dendritic cells, peripheral blood mononuclear cells (Jin et al., 2012; Rosenberger et al., 2012)
ICAM-1	Recruitment of circulating proinflammatory mediators	Human pulmonary microvascular endothelial cells (Mercado et al., 2012; Colin et al., 2014)
COX-2	Regulation of prostaglandin D <sub>2</sub>	Various (Phillips et al., 2004; Sully et al., 2004; Tudor et al., 2009; Hammaker and Firestein, 2010; Bhattacharyya et al., 2011)
iNOS	Modulation of airway and vascular smooth muscle tone	DLD-1 cells, rat cortical astrocytes (Linker et al., 2005; Lisi et al., 2011)
IL-2	Inducing and mediating proinflammatory responses	Mouse splenocytes (Ogilvie et al., 2005)
CXCL1	Neutrophil chemoattractant	Mouse macrophages (Datta et al., 2008)
MMP-9	Remodeling of the extracellular matrix	Three-dimensional cancer model, chick chorioallantoic membrane (Van Tubergen et al., 2013)
CCL3	Chemotaxis of chemokines and cytokines, inflammatory mediator release	Mouse macrophages, fibroblasts (Sauer et al., 2006; Kang et al., 2011)

CCL, C-C ligand; COX, cyclooxygenase; CXCL, C-X-C ligand; GM-CSF, granulocyte-macrophage colony-stimulating factor; ICAM, intercellular adhesion molecule; iNOS, inducible nitric oxide synthase; MMP, matrix metalloproteinase; VEGF, vascular endothelial growth factor.

Lockey, 2011; Boonpiyathad et al., 2013; Banerjee et al., 2014; Masaki et al., 2014).

## TTP in Respiratory Disease: What Is Known?

The important anti-inflammatory role of TTP has been extensively studied in mouse models and cell lines, such as peripheral blood monocytes, bone marrow-derived macrophages (Carballo et al., 2001), HeLa cells and embryonic fibroblasts (Chen et al., 2012), and a plethora of other cell lines, to elucidate the molecular mechanisms that drive these anti-inflammatory functions (Brooks and Blackshear, 2013) (Table 1). To date, the potential of TTP as an anti-inflammatory therapy has primarily focused on its role in rheumatoid arthritis, leaving investigations into the anti-inflammatory function of TTP in airway disease relatively unexplored. Lessons may also be learned from studies examining tumors, as many inflammatory tumors have features that parallel chronic inflammatory diseases like asthma. Among the similarities are structural remodeling, inflammatory cell migration, cell proliferation, the induction of proteinases, and dysregulation of endogenous anti-inflammatory proteins. The latter characteristic is perhaps the most important similarity, as there is a prolific induction of inflammatory mediators in both inflammatory tumors and chronic inflammatory diseases. This suggests that the endogenous feedback machinery present within the cell is being dysregulated in cancer. TTP acted as a tumor suppressor in a model of a specific mast cell tumor by reducing the pivotal cytokine IL-3 that propagated that tumor (Stoecklin et al., 2003). Similarly, IL-6, a proinflammatory mediator shown to maintain a pro-cancerous inflammatory state, was reduced by TTP when cells were stimulated with lipopolysaccharide (LPS) (Sauer et al., 2006). This, along with studies that showed that TTP was attenuated in many cancerous cell lines relative to noncancerous lines (Carrick and Blackshear, 2007; Brennan et al., 2009), presents compelling evidence to suggest that TTP could be a viable therapeutic target. This is, however, just a guide, and caution should be exercised when using tumor models, as they have many limitations. Among the main limitations, we see that tumors under resistance mechanisms can induce a host of neoplastic characteristics not mentioned here. It is not just in cancer models, however, that TTP has been shown to be effective; TTP was shown to downregulate cytokines in acute respiratory disease (Colin et al., 2014), arthritis (Carrick et al., 2004), and psoriasis (Colin et al., 2014), to highlight a few. It was also shown to be effective in reducing the immune cell-driven cytokine IL-17A in the presence of TTP overexpression that was increased when TTP was silenced (Lee et al., 2012). TTP was also a key anti-inflammatory protein upregulated in response to antiasthma drugs in cellular models utilizing human bronchial epithelial BEAS-2B cells and airway smooth muscle cells (Kaur et al., 2008).

## TTP Function: Multiple Levels of Regulatory Control

To assess the potential of TTP as an anti-inflammatory molecule, we have to first gain an understanding of the mechanisms and molecules that control its regulation. TTP, also known as Nup475 (DuBois et al., 1990), Tis 11 (Varnum et al., 1991), G0S24 (Heximer and Forsdyke, 1993), or Zfp36 (Taylor et al., 1991), is the prototypical member of the family

of Cys-Cys-Cys-His class tandem zinc-finger proteins. It is an immediate early response gene with low levels expressed within quiescent cells. It can be rapidly expressed and translated into protein to exert regulatory control upon mRNA decay. Various studies have shown that TTP is regulated at different stages of its life cycle. As with many critical molecules, TTP regulation is possible at the transcriptional level, the post-transcriptional level, and the post-translational level.

**Transcriptional Regulation.** At the transcriptional level, it was shown that the deletion of a single intron from the promoter region in the Zfp36 5' UTR caused an 85% drop in serum-induced TTP mRNA (Lai et al., 1995). Structural analysis of the 5' UTR revealed promoter element binding sites identified as transcription factor binding sites (Lai et al., 1995, 1998), including early growth response protein 1, TTP promoter element 1, activating protein 2-like, and specificity protein 1. These binding sites in the Zfp36 promoter region were shown to be partially responsible for TTP transcriptional activation through mutational analysis of splice variants in the presence of serum (DuBois et al., 1990; Lai et al., 1995, 1998; Kaneda et al., 2000). The same studies also showed that NF- $\kappa$ B had a binding site on the intron, suggesting that an interaction could exist between transcription factors to induce TTP mRNA. A strong interaction was not proven, only that the intron had to remain in close proximity to the promoter elements. These data, taken in concert with studies showing the ability of TTP mRNA to be upregulated by molecules such as TNF- $\alpha$ , LPS, IL-1 $\beta$ , IL-4, transforming growth factor- $\beta$ , interferon  $\gamma$ , mitogenic serum stimuli, cAMP, and phorbol esters, have allowed us to gain insight into the molecular pathways that drive TTP mRNA upregulation. Inhibitor studies involving inhibition of NF- $\kappa$ B, in the presence of IL-1 $\beta$  and LPS, have shown that TTP mRNA is a direct substrate of the NF- $\kappa$ B signaling pathway (King et al., 2009; Chen et al., 2013). Other signaling pathways include the protein kinase C (PKC) pathway, via transcription factor activating protein 2, which was shown to regulate TTP mRNA in the presence of a phorbol ester (Chen et al., 1998; Tan and Parker, 2003; Jalonen et al., 2007; Leppänen et al., 2008, 2010) and the epidermal growth factor pathway, via the transcription factor ETS domain-containing protein ELK-1 (Florowska et al., 2012).

The signaling molecule most extensively studied in relation to TTP mRNA regulation is p38 MAPK. The p38 MAPK-mediated pathway exerts most of its control over TTP via the downstream protein kinase MK2 (Mahtani et al., 2001; Chrestensen et al., 2004; Stoecklin et al., 2004; Hitti et al., 2006; Tudor et al., 2009; Marchese et al., 2010); this is discussed in greater detail below. TTP mRNA is expressed with an immediate early peak followed by a decline back to near baseline levels after ~4 hours (Tchen et al., 2004; Hitti et al., 2006). In the presence of a p38 MAPK inhibitor, this peak at the early time point was reduced, suggesting p38 MAPK-dependent regulation of TTP mRNA expression. It has also been suggested that because TTP mRNA is increased via the p38 MAPK pathway along with other pathways, it may be increased in a temporally specific manner, with NF- $\kappa$ B responsible for the early mRNA production and p38 MAPK responsible for later time points (Chen et al., 2013).

Taken together, these studies demonstrate that targeting TTP at the transcriptional level represents an opportunity to inhibit multiple inhibitory pathways at a downstream level, because there are a number of signaling molecules and pathways that regulate TTP mRNA expression.

**Post-Transcriptional Regulation.** It has been widely reported that the p38 MAPK pathway is responsible for post-transcriptional regulation of TTP by influencing mRNA stability. Actinomycin D chase experiments performed in the presence of a p38 MAPK inhibitor demonstrated direct reliance on the p38 MAPK for TTP mRNA stability (Carballo et al., 2001; Tchen et al., 2004). TTP mRNA transcripts were more stable in the presence of the p38 MAPK inhibitor. The ERK pathway has also been synergistically linked with p38 MAPK to induce TTP mRNA stability (Brook et al., 2006; Deleault et al., 2008; Essafi-Benkhadir et al., 2010). Indirectly p38 MAPK is able to destabilize TTP mRNA through TTP self-regulated mRNA decay. Examination of TTP 3' UTR constructs in the presence of TTP protein revealed that TTP has a high affinity for its own mRNA, and as such, it is able to negatively regulate itself (Brooks et al., 2004; Tchen et al., 2004). This differential regulation is representative of the complex interaction between these molecules. Moreover, actinomycin D analysis of TTP mRNA stability in the presence of a PKC pathway inhibitor showed that PKC had a similar effect on TTP mRNA stability as p38 MAPK, thereby suggesting that part of its impact on TTP may be due to an increase in TTP mRNA stability (Leppänen et al., 2010).

Another interesting area of focus on the regulation and function of TTP has been with microRNAs (miRs) and their ability to impact TTP and bind 3' UTRs of specific mRNA transcripts. MiRs are small, ~20 nucleotides in length; they are regulators of mRNA decay. Accumulating evidence demonstrates that miRs can regulate TTP; miR-346, for example, was shown to be important in the regulation of TNF- $\alpha$  in rheumatoid arthritis, by controlling release of TNF- $\alpha$  directly via TTP (Semaan et al., 2011), while miR29 was observed to bind TTP mRNA in a cancer model (Gebeshuber et al., 2009; Al-Ahmadi et al., 2013). Another study proposed that TTP does not directly bind to miR-16 but interacts through association with Ago/eiF2C family members (proteins required for miR activity) in a complex with miR16 to assist with the targeting of AREs (Jing et al., 2005). Moreover, it was reported that miR-466 bound competitively to IL-10 and conferred anti-inflammatory properties by antagonizing TTP in Toll-like receptor-triggered macrophages (Ma et al., 2010). These reports reveal that miRs have an impact on TTP both directly or indirectly. So far, whether miRs present within the airways and implicated in asthma are linked to TTP remains an open question; however, data from rheumatoid arthritis and cancer models provide us with an inflammatory model in which miRs have been shown to alter TTP binding. Further studies exploring the regulation of TTP by miRs in the context of asthmatic inflammation are warranted.

**Post-Translational Control.** Once the stable mRNA transcripts are translated into protein, there is a level of control that is exercised over the translated protein. Being an immediate early response gene, TTP protein is rapidly expressed after gene transcription; hence, TTP protein expression bears a similar temporal profile to mRNA expression (Mahtani et al., 2001; Brook et al., 2006). Post-translational regulation of TTP protein occurs predominantly in the form of phosphorylation. TTP protein appears in two distinct forms, a phosphorylated form and an unphosphorylated form. Phosphorylation of TTP is regulated post-translationally via the ERK and p38 MAPK pathways (Brook et al., 2006; Deleault et al., 2008; Essafi-Benkhadir et al., 2010). Other kinase pathways, such as the

PKC pathway, were shown not to have an impact on TTP protein expression. Post-translational regulation of TTP is also able to alter its localization to the cytoplasm with the aid of 14-3-3 proteins (Johnson et al., 2002; Benjamin et al., 2006; Brook et al., 2006; Sun et al., 2007). Post-translational control of TTP function represents a promising means to harness the anti-inflammatory capacity of TTP, because its phosphorylation status dictates its activity. Hence, this is explored in greater depth in the following sections.

## TTP Is a Substrate for MK2

TTP's function as an mRNA-destabilizing factor is mainly controlled by the p38 MAPK–MK2 axis. Critically, p38 MAPK also activates the downstream kinase MK2, which phosphorylates murine TTP protein on two key serine residues (Ser52 and Ser178) (Chrestensen et al., 2004). MK2 is a substrate of the p38 MAPK pathway and has been implicated in post-translational control of cytokines. This was demonstrated by using MK2 knockout mice, which had attenuated production of TNF- $\alpha$  protein when compared with wild-type equivalents. However, the mRNA levels in wild-type mice were very similar to those in knockout mice, suggesting that the control exerted was at the translational level. This also suggested that control might be imparted via a downstream substrate of MK2. TTP was shown to be a substrate for MK2 in primary cells by the use of MK2/TTP double-knockout mice, p38 MAPK inhibitors, and a reconstituted p38 MAPK transduction pathway using recombinant kinases to detect TTP phosphorylation (Mahtani et al., 2001; Hitti et al., 2006; Tudor et al., 2009). Production of cytokines was also impaired compared with wild type, with IL-6 mRNA half-life, in particular, markedly attenuated (Kotlyarov et al., 1999; Neiningner et al., 2002). In vitro studies revealed that TTP is phosphorylated by MK2 (Mahtani et al., 2001; Chrestensen et al., 2004) and phosphorylation of MK2 is able to modulate TTP function (Stoecklin et al., 2004).

## Phosphorylation of TTP Protein

As detailed above, the main function of TTP protein is to decay target mRNAs; however, its activity is also profoundly affected by post-translational modification (Cao et al., 2007). Post-translational phosphorylation appears to be the main factor responsible for determining its stability, activity, and ability to form protein complexes. Phosphorylation of TTP has been extensively studied and mapped, revealing that TTP is a hyperphosphorylated protein in vivo, with its mRNA-destabilizing activity depending on its phosphorylation status. When TTP protein is phosphorylated, it is unable to affect the stability of target mRNA; however, when unphosphorylated, it is active and therefore able to confer its mRNA-destabilizing ability. TTP protein is more stable when phosphorylated and unstable when unphosphorylated. This allows TTP protein to be rapidly degraded by the proteasome when unphosphorylated (Brook et al., 2006; Hitti et al., 2006). It has also been shown that Ser220 on mouse TTP is phosphorylated via the ERK MAPK pathway (Taylor et al., 1995), suggesting that dual MAPK involvement may be required to confer post-translational modifications to TTP (Brooks et al., 2004). The functions of all the various TTP phosphorylation sites are not completely understood; however, the sites for MK2 protein binding have been characterized at Ser52 and Ser178 on murine TTP (Chrestensen et al., 2004;

Stoecklin et al., 2004). Phosphorylation at these sites enables TTP to form complexes with multifunction adaptor 14-3-3 proteins that ablate its function as an mRNA-destabilizing protein (Chrestensen et al., 2004; Sun et al., 2007; Marchese et al., 2010). This complex also serves to protect TTP from dephosphorylation in a protein phosphatase 2A (PP2A)-dependent manner (Sun et al., 2007) and prevent degradation of TTP via the proteasome (Deleault et al., 2008; Pfeiffer and Brooks, 2012). There are, however, some data to suggest that the role of 14-3-3 proteins might be limited to chaperoning TTP and only preventing its dephosphorylation (Marchese et al., 2010; Ngoc et al., 2014). The additional role as a block on proteasomal degradation is challenged, as recombinant 14-3-3 in the presence of TTP did not affect TTP stability and degradation by the proteasome (Ngoc et al., 2014). This is in line with data that showed that TTP levels increased when cells were in the presence of a proteasome inhibitor. This complex that forms alters the functionality of TTP without altering its ability to bind mRNA (Cao et al., 2003; Rigby et al., 2005; Marchese et al., 2010; Clement et al., 2011), suggesting that phosphorylated TTP remains inactive while being bound to target mRNA until it is dephosphorylated. It is then able to recruit the Ccr4-CAF1 deadenylase components via Not1, which are then able to rapidly deadenylate the target mRNA, resulting in their destabilization (Sandler et al., 2011; Colin et al., 2014). Phosphorylation of TTP also has an impact on its subcellular localization, with studies showing that upon stimulation with a mitogen, TTP is rapidly translocated from the nucleus to the cytoplasm (Taylor et al., 1996b; Johnson et al., 2002; Brook et al., 2006). This is in line with data suggesting that degradation of ARE-containing mRNAs occurs at different locations within the cytoplasm (Franks and Lykke-Andersen, 2007). Taken together, these studies demonstrate the importance of the phosphorylation status of TTP in controlling TTP activity and thus represent a novel area to target in future pharmacotherapeutic strategies.

### Regulation of the Temporal Kinetics of TTP by p38 MAPK

Knowing the temporal kinetics of TTP may allow us to tailor pharmacological intervention to increase the anti-inflammatory efficiency of the process. This would be beneficial as it might allow us to target specific events within the TTP temporal regulation rather than the molecule as a whole. The kinetics of TTP originate with the p38 MAPK pathway or other such inflammatory pathways. When p38 MAPK is phosphorylated, it becomes active, at which point it then phosphorylates downstream targets like MK2. This p38 MAPK upregulation occurs very soon after the cell is stimulated by some extracellular trigger. Our own studies in human airway smooth muscle cells indicate that activation of p38 MAPK occurs as early as 10–15 minutes after stimulation (Quante et al., 2008; Moutzouris et al., 2010) and possibly even earlier. TTP is a known substrate for p38 MAPK and MK2, both of which are required for the full induction of TTP expression. As an immediate early gene, TTP is rapidly expressed, with our preliminary studies indicating an increase in mRNA as early as 30 minutes and with one study showing upregulation as early as 10 minutes (Taylor et al., 1995). This causes the initial accumulation of TTP mRNA within the nucleus followed by efficient translation (Taylor et al., 1996b; Johnson et al., 2002; Brook et al., 2006). During

this period when TTP is inactive, the cytokine mRNA is stabilized and results in a wave of cytokine upregulation, peaking at 1 hour (Balakathiresan et al., 2009). This is derived from the fact that active p38 MAPK has a peak at 15–30 minutes, with activation returning to basal levels after about 1 hour (Quante et al., 2008; Moutzouris et al., 2010). Once p38 MAPK activation has subsided, MK2 is no longer phosphorylated and hence is unable to phosphorylate TTP. To determine the phosphorylation status of TTP using Western blotting can be challenging; however, earlier studies have used phosphatases to identify and confirm the phosphorylated immunoreactive band from the unphosphorylated band (Carballo et al., 2001; Mahtani et al., 2001; Balakathiresan et al., 2009). Hence, at the 1-hour time point, the unphosphorylated immunoreactive band of about 40–45 kDa appears, suggesting that TTP is active (King et al., 2009). The literature suggests that this may be in part due to the action of PP2A dephosphorylating phosphorylated TTP, which leads to the rapid deadenylation and degradation of cytokine mRNAs, thereby potentially attenuating the degree of inflammation. MAPK phosphatase-1 regulation of p38 MAPK phosphorylation also plays an important role. Because there appears to be a sequential order to the control of TTP activity, it can be refined and modified to aid in the fine-tuning of a potential anti-inflammatory agent.

### Regulation of TTP by PP2A

To fully explore TTP and its potential as an anti-inflammatory agent, it is important to highlight the key molecules that may positively affect TTP function. We know that TTP is a hyperphosphorylated protein, and any protein that could reverse its inactivation would be of therapeutic value. This is where PP2A may play a role. PP2A is a heterotrimeric serine-threonine protein phosphatase consisting of a scaffold subunit, a catalytic subunit for its phosphatase activity, and a regulatory subunit to confer substrate specificity (Janssens et al., 2008; Shi, 2009). PP2A is a master controller of cell signaling and regulates many cellular functions, including cell growth, inflammation, and apoptosis (Santoro et al., 1998; Yang et al., 2001). Many of these functions have been attributed to PP2A-mediated regulation of MAPKs (Junttila et al., 2008). PP2A is able to act on p38 MAPK and its upstream kinase MKK3 (Prickett and Brautigan, 2007). Inhibitors of PP2A, like okadaic acid, calyculin A, and simian viral protein SV40, were shown to induce phosphorylation and activate upstream kinases of p38 MAPK and ERK in vitro (Junttila et al., 2008). It was also shown that inhibition of PP2A increases stability of cytokine mRNA by increasing the activity of p38 MAPK and its downstream kinase MK2 (Sun et al., 2007; Clement et al., 2011). These studies are evidence of indirect regulation of TTP by PP2A. More directly, Brook et al. (2006) showed that the PP2A inhibitor calyculin A caused accumulation of hyperphosphorylated and stable TTP protein and highlighted an MK2-Ser52/Ser178-PP2A-dependent mechanism for controlling TTP stability. Because PP2A is able to regulate p38 MAPK as well as dephosphorylate TTP, TTP is both an indirect and direct target of PP2A. Thus, PP2A activation offers promise as a means to control the activity of TTP, although the fine temporal control needs to be delineated. Most of the work on PP2A regulation has been performed in cancer cells (Janssens et al., 2008; Perrotti and Neviani, 2008; Tay et al., 2012), with some work in mast cells (Kranias et al., 2010) and respiratory epithelial cells (Cornell et al., 2009).

Notably, PP2A has emerged as playing an important role in vivo models of asthma (Collison et al., 2013; Hatchwell et al., 2014), and PP2A inhibition (with okadaic acid) resulted in corticosteroid insensitivity in vitro (Kobayashi et al., 2011). Strategies targeting PP2A could become therapeutically important as an anti-inflammatory approach in asthma.

## Summary and Future Directions

The substantial in vitro evidence implicating the p38 MAPK signaling pathway in the pathogenesis of respiratory diseases driven by inflammation has propelled studies targeting p38 MAPK as a novel anti-inflammatory strategy in vivo. However, as discussed in this review, p38 MAPK activation also results in repression of critical anti-inflammatory proteins, in addition to inhibition of proinflammatory molecules. The RNA-destabilizing molecule TTP is one such key anti-inflammatory protein whose function is significantly repressed by p38 MAPK inhibition. We propose that a greater understanding of TTP upregulation and its activation by phosphatases such as PP2A in airway inflammation represents an alternative anti-inflammatory strategy in chronic respiratory diseases.

### Authorship Contributions

Wrote or contributed to the writing of the manuscript: Prabhala, Ammit.

### References

Al-Ahmadi W, Al-Ghamdi M, Al-Souhibani N, and Khabar KS (2013) miR-29a inhibition normalizes HuR over-expression and aberrant AU-rich mRNA stability in invasive cancer. *J Pathol* **230**:28–38.

Al Ghouleh I and Magder S (2012) NADPH oxidase-derived superoxide destabilizes lipopolysaccharide-induced interleukin 8 mRNA via p38, extracellular signal-regulated kinase mitogen-activated protein kinase, and the destabilizing factor tristetraprolin. *Shock* **37**:433–440.

Ammit AJ (2005) The role of mRNA stability in airway remodelling. *Pulm Pharmacol Ther* **18**:405–415.

Ammit AJ (2013) Glucocorticoid insensitivity as a source of drug targets for respiratory disease. *Curr Opin Pharmacol* **13**:370–376.

Amrani Y, Ammit AJ, and Panettieri RA, Jr (2001) Tumor necrosis factor receptor (TNFR) 1, but not TNFR2, mediates tumor necrosis factor- $\alpha$ -induced interleukin-6 and RANTES in human airway smooth muscle cells: role of p38 and p42/44 mitogen-activated protein kinases. *Mol Pharmacol* **60**:646–655.

Ashwell JD (2006) The many paths to p38 mitogen-activated protein kinase activation in the immune system. *Nat Rev Immunol* **6**:532–540.

Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevvernic I, Arthur JSC, Alessi DR, and Cohen P (2007) The selectivity of protein kinase inhibitors: a further update. *Biochem J* **408**:297–315.

Bakheet T, Williams BR, and Khabar KS (2006) ARED 3.0: the large and diverse AU-rich transcriptome. *Nucleic Acids Res* **34**:D111–D114.

Balakathiresan NS, Bhattacharyya S, Gutti U, Long RP, Jozwik C, Huang W, Srivastava M, Pollard HB, and Biswas R (2009) Tristetraprolin regulates IL-8 mRNA stability in cystic fibrosis lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **296**:L1012–L1018.

Banerjee R, Van Tubergen EA, Scanlon CS, Vander Broek R, Lints JP, Liu M, Russo N, Inglehart RC, Wang Y, and Polverini PJ et al. (2014) The G protein-coupled receptor GALR2 promotes angiogenesis in head and neck cancer. *Mol Cancer Ther* **13**:1323–1333.

Barnes PJ (2008) The cytokine network in asthma and chronic obstructive pulmonary disease. *J Clin Invest* **118**:3546–3556.

Barnes PJ (2013) Corticosteroid resistance in patients with asthma and chronic obstructive pulmonary disease. *J Allergy Clin Immunol* **131**:636–645.

Barnstein BO, Li G, Wang Z, Kennedy S, Chalfant C, Nakajima H, Bunting KD, and Ryan JJ (2006) Stat5 expression is required for IgE-mediated mast cell function. *J Immunol* **177**:3421–3426.

Batra J, Chatterjee R, and Ghosh B (2007) Inducible nitric oxide synthase (iNOS): role in asthma pathogenesis. *Indian J Biochem Biophys* **44**:303–309.

Benjamin D, Schmidlin M, Min L, Gross B, and Moroni C (2006) BRP1 protein turnover and mRNA decay activity are regulated by protein kinase B at the same phosphorylation sites. *Mol Cell Biol* **26**:9497–9507.

Bhattacharyya S, Gutti U, Mercado J, Moore C, Pollard HB, and Biswas R (2011) MAPK signaling pathways regulate IL-8 mRNA stability and IL-8 protein expression in cystic fibrosis lung epithelial cell lines. *Am J Physiol Lung Cell Mol Physiol* **300**:L81–L87.

Bhavsar P, Hew M, Khorasani N, Torrego A, Barnes PJ, Adcock I, and Chung KF (2008) Relative corticosteroid insensitivity of alveolar macrophages in severe asthma compared with non-severe asthma. *Thorax* **63**:784–790.

Boonpiyathad S, Pornsuriyasak P, Buranapraditkun S, and Klaewsongkram J (2013) Interleukin-2 levels in exhaled breath condensates, asthma severity, and asthma control in nonallergic asthma. *Allergy Asthma Proc* **34**:e35–e41.

Brahma PK, Zhang H, Murray BS, Shu FJ, Sidell N, Seli E, and Kallen CB (2012) The mRNA-binding protein Zfp36 is upregulated by  $\beta$ -adrenergic stimulation and represses IL-6 production in 3T3-L1 adipocytes. *Obesity (Silver Spring)* **20**:40–47.

Brennan SE, Kuwano Y, Alkharouf N, Blackshear PJ, Gorospe M, and Wilson GM (2009) The mRNA-destabilizing protein tristetraprolin is suppressed in many cancers, altering tumorigenic phenotypes and patient prognosis. *Cancer Res* **69**:5168–5176.

Brook M, Tchen CR, Santalucia T, McIlrath J, Arthur JS, Saklatvala J, and Clark AR (2006) Posttranslational regulation of tristetraprolin subcellular localization and protein stability by p38 mitogen-activated protein kinase and extracellular signal-regulated kinase pathways. *Mol Cell Biol* **26**:2408–2418.

Brooks SA and Blackshear PJ (2013) Tristetraprolin (TTP): interactions with mRNA and proteins, and current thoughts on mechanisms of action. *Biochim Biophys Acta* **1829**:666–679.

Brooks SA, Connolly JE, and Rigby WF (2004) The role of mRNA turnover in the regulation of tristetraprolin expression: evidence for an extracellular signal-regulated kinase-specific, AU-rich element-dependent, autoregulatory pathway. *J Immunol* **172**:7263–7271.

Cao H, Deterding LJ, and Blackshear PJ (2007) Phosphorylation site analysis of the anti-inflammatory and mRNA-destabilizing protein tristetraprolin. *Expert Rev Proteomics* **4**:711–726.

Cao H, Dzineku F, and Blackshear PJ (2003) Expression and purification of recombinant tristetraprolin that can bind to tumor necrosis factor- $\alpha$  mRNA and serve as a substrate for mitogen-activated protein kinases. *Arch Biochem Biophys* **412**:106–120.

Caput D, Beutler B, Hartog K, Thayer R, Brown-Shimer S, and Cerami A (1986) Identification of a common nucleotide sequence in the 3'-untranslated region of mRNA molecules specifying inflammatory mediators. *Proc Natl Acad Sci USA* **83**:1670–1674.

Carballo E, Cao H, Lai WS, Kennington EA, Campbell D, and Blackshear PJ (2001) Decreased sensitivity of tristetraprolin-deficient cells to p38 inhibitors suggests the involvement of tristetraprolin in the p38 signaling pathway. *J Biol Chem* **276**:42580–42587.

Carballo E, Lai WS, and Blackshear PJ (1998) Feedback inhibition of macrophage tumor necrosis factor- $\alpha$  production by tristetraprolin. *Science* **281**:1001–1005.

Carballo E, Lai WS, and Blackshear PJ (2000) Evidence that tristetraprolin is a physiological regulator of granulocyte-macrophage colony-stimulating factor messenger RNA deadenylation and stability. *Blood* **95**:1891–1899.

Carrick DM and Blackshear PJ (2007) Comparative expression of tristetraprolin (TTP) family member transcripts in normal human tissues and cancer cell lines. *Arch Biochem Biophys* **462**:278–285.

Carrick DM, Lai WS, and Blackshear PJ (2004) The tandem CCCH zinc finger protein tristetraprolin and its relevance to cytokine mRNA turnover and arthritis. *Arthritis Res Ther* **6**:248–264.

Chafin CB, Regna NL, Dai R, Caudell DL, and Reilly CM (2013) MicroRNA-let-7a expression is increased in the mesangial cells of NZB/W mice and increases IL-6 production in vitro. *Autoimmunity* **46**:351–362.

Chang PJ, Bhavsar PK, Michaeloudes C, Khorasani N, and Chung KF (2012) Corticosteroid insensitivity of chemokine expression in airway smooth muscle of patients with severe asthma. *J Allergy Clin Immunol* **130**:877–885.

Chen CC, Wang JK, Chen WC, and Lin SB (1998) Protein kinase C  $\epsilon$  mediates lipopolysaccharide-induced nitric-oxide synthase expression in primary astrocytes. *J Biol Chem* **273**:19424–19430.

Chen X, Wei Z, Wang W, Yan R, Xu X, and Cai Q (2012) Role of RNA-binding protein tristetraprolin in tumor necrosis factor- $\alpha$  mediated gene expression. *Biochem Biophys Res Commun* **428**:327–332.

Chen YL, Huang YL, Lin NY, Chen HC, Chiu WC, and Chang CJ (2006) Differential regulation of ARE-mediated TNF $\alpha$  and IL-1 $\beta$  mRNA stability by lipopolysaccharide in RAW264.7 cells. *Biochem Biophys Res Commun* **346**:160–168.

Chen YL, Jiang YW, Su YL, Lee SC, Chang MS, and Chang CJ (2013) Transcriptional regulation of tristetraprolin by NF- $\kappa$ B signaling in LPS-stimulated macrophages. *Mol Biol Rep* **40**:2867–2877.

Chrestensen CA, Schroeder MJ, Shabanowitz J, Hunt DF, Peló JW, Worthington MT, and Sturgill TW (2004) MAPKAP kinase 2 phosphorylates tristetraprolin in vivo sites including Ser178, a site required for 14-3-3 binding. *J Biol Chem* **279**:10176–10184.

Chung KF (2011) p38 mitogen-activated protein kinase pathways in asthma and COPD. *Chest* **139**:1470–1479.

Clark AR and Dean JL (2012) The p38 MAPK pathway in rheumatoid arthritis: a sideways look. *Open Rheumatol J* **6**:209–219.

Clement SL, Scheckel C, Stoecklin G, and Lykke-Andersen J (2011) Phosphorylation of tristetraprolin by MK2 impairs AU-rich element mRNA decay by preventing deadenylase recruitment. *Mol Cell Biol* **31**:256–266.

Colin S, Darné B, Kadi A, Favier M, Lesaffre C, Conduzorgues JP, Al-Mahmoud S, and Doss N (2014) The antiangiogenic insulin receptor substrate-1 antisense oligonucleotide aganirsin impairs AU-rich mRNA stability by reducing 14-3-3 $\beta$ -tristetraprolin protein complex, reducing inflammation and psoriatic lesion size in patients. *J Pharmacol Exp Ther* **349**:107–117.

Collison A, Hatchwell L, Verrills N, Wark PA, de Siqueira AP, Toozé M, Carpenter H, Don AS, Morris JC, and Zimmermann N et al. (2013) The E3 ubiquitin ligase midline 1 promotes allergen and rhinovirus-induced asthma by inhibiting protein phosphatase 2A activity. *Nat Med* **19**:232–237.

Cornell TT, Hinkovska-Galcheva V, Sun L, Cai Q, Hershenson MB, Vanway S, and Shanley TP (2009) Ceramide-dependent PP2A regulation of TNF $\alpha$ -induced IL-8 production in respiratory epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **296**:L849–L856.

- Datta S, Biswas R, Novotny M, Pavicic PG, Jr, Herjan T, Mandal P, and Hamilton TA (2008) Tristetraprolin regulates CXCL1 (KC) mRNA stability. *J Immunol* **180**: 2545–2552.
- Deleault KM, Skinner SJ, and Brooks SA (2008) Tristetraprolin regulates TNF TNF- $\alpha$  mRNA stability via a proteasome dependent mechanism involving the combined action of the ERK and p38 pathways. *Mol Immunol* **45**:13–24.
- DuBois RN, McLane MW, Ryder K, Lau LF, and Nathans D (1990) A growth factor-inducible nuclear protein with a novel cysteine/histidine repetitive sequence. *J Biol Chem* **265**:19185–19191.
- Essafi-Benkhadir K, Onesto C, Stebe E, Moroni C, and Pagès G (2007) Tristetraprolin inhibits Ras-dependent tumor vascularization by inducing vascular endothelial growth factor mRNA degradation. *Mol Biol Cell* **18**:4648–4658.
- Essafi-Benkhadir K, Pouyssegur J, and Pagès G (2010) Implication of the ERK pathway on the post-transcriptional regulation of VEGF mRNA stability. *Methods Mol Biol* **661**:451–469.
- Florkowska M, Tymoszek P, Balwierz A, Skucha A, Kochan J, Wawro M, Stalinska K, and Kasza A (2012) EGF activates TTP expression by activation of ELK-1 and EGR-1 transcription factors. *BMC Mol Biol* **13**:8.
- Franks TM and Lykke-Andersen J (2007) TTP and BRF proteins nucleate processing body formation to silence mRNAs with AU-rich elements. *Genes Dev* **21**: 719–735.
- Galbiati V, Carne A, Mitjans M, Galli CL, Marinovich M, and Corsini E (2012) Isoeugenol destabilizes IL-8 mRNA expression in THP-1 cells through induction of the negative regulator of mRNA stability tristetraprolin. *Arch Toxicol* **86**:239–248.
- Gebelhuber CA, Zatloukal K, and Martinez J (2009) miR-29a suppresses tristetraprolin, which is a regulator of epithelial polarity and metastasis. *EMBO Rep* **10**:400–405.
- Haddad JJ (2001) VX-745. Vertex Pharmaceuticals. *Curr Opin Investig Drugs* **2**: 1070–1076.
- Hallsworth MP, Moir LM, Lai D, and Hirst SJ (2001) Inhibitors of mitogen-activated protein kinases differentially regulate eosinophil-activating cytokine release from human airway smooth muscle. *Am J Respir Crit Care Med* **164**:688–697.
- Hammaker D and Firestein GS (2010) “Go upstream, young man”: lessons learned from the p38 saga. *Ann Rheum Dis* **69** (Suppl 1):i77–i82.
- Hatchwell L, Girkin J, Dun MD, Morten M, Verrills N, Toop HD, Morris JC, Johnston SL, Foster PS, and Collison A et al. (2014) Salmeterol attenuates chemotactic responses in rhinovirus-induced exacerbation of allergic airways disease by modulating protein phosphatase 2A. *J Allergy Clin Immunol* **133**:1720–1727.
- Heinrichsdorff J, Luedde T, Perdiguerio E, Nebreda AR, and Pasparakis M (2008) p38  $\alpha$  MAPK inhibits JNK activation and collaborates with IkappaB kinase 2 to prevent endotoxin-induced liver failure. *EMBO Rep* **9**:1048–1054.
- Henness S, van Thoor E, Ge Q, Armour CL, Hughes JM, and Ammit AJ (2006) IL-17A acts via p38 MAPK to increase stability of TNF- $\alpha$ -induced IL-8 mRNA in human ASM. *Am J Physiol Lung Cell Mol Physiol* **290**:L1283–L1290.
- Heximer SP and Forsdyke DR (1993) A human putative lymphocyte G0/G1 switch gene homologous to a rodent gene encoding a zinc-binding potential transcription factor. *DNA Cell Biol* **12**:73–88.
- Hirst SJ (2003) Regulation of airway smooth muscle cell immunomodulatory function: role in asthma. *Respir Physiol Neurobiol* **137**:309–326.
- Hitti E, Iakovleva T, Brook M, Deppenmeier S, Gruber AD, Radzioch D, Clark AR, Blackshear PJ, Kotlyarov A, and Gaestel M (2006) Mitogen-activated protein kinase-activated protein kinase 2 regulates tumor necrosis factor mRNA stability and translation mainly by altering tristetraprolin expression, stability, and binding to adenine/uridine-rich element. *Mol Cell Biol* **26**:2399–2407.
- Hu JH, Chen T, Zhuang ZH, Kong L, Yu MC, Liu Y, Zang JW, and Ge BX (2007) Feedback control of MKP-1 expression by p38. *Cell Signal* **19**:393–400.
- Ishmael FT (2011) The inflammatory response in the pathogenesis of asthma. *J Am Osteopath Assoc* **111**(11, Suppl 7):S11–S17.
- Jalonen U, Leppänen T, Kankaanranta H, and Moilanen E (2007) Salbutamol increases tristetraprolin expression in macrophages. *Life Sci* **81**:1651–1658.
- Jalonen U, Nieminen R, Vuolteenaho K, Kankaanranta H, and Moilanen E (2006) Down-regulation of tristetraprolin expression results in enhanced IL-12 and MIP-2 production and reduced MIP-3 $\alpha$  synthesis in activated macrophages. *Mediators Inflamm* **2006**:40691.
- Janssens V, Longin S, and Goris J (2008) PP2A holoenzyme assembly: in cauda venenum (the sting is in the tail). *Trends Biochem Sci* **33**:113–121.
- Jin WJ, Chen CF, Liao HY, Gong LL, Yuan XH, Zhao BB, Zhang D, Feng X, Liu JJ, and Wang Y et al. (2012) Downregulation of the AU-rich RNA-binding protein ZFP36 in chronic HBV patients: implications for anti-inflammatory therapy. *PLoS One* **7**:e33356.
- Jing Q, Huang S, Guth S, Zarubin T, Motoyama A, Chen J, Di Padova F, Lin SC, Gram H, and Han J (2005) Involvement of microRNA in AU-rich element-mediated mRNA instability. *Cell* **120**:623–634.
- Johnson BA, Stehn JR, Yaffe MB, and Blackwell TK (2002) Cytoplasmic localization of tristetraprolin involves 14-3-3-dependent and -independent mechanisms. *J Biol Chem* **277**:18029–18036.
- Junttila MR, Li S-P, and Westermarck J (2008) Phosphatase-mediated crosstalk between MAPK signaling pathways in the regulation of cell survival. *FASEB J* **22**: 954–965.
- Kaneda N, Murata T, Niimi Y, and Minamiyama M (2000) Cloning and characterization of the rat TIS11 gene. *Mol Cell Biochem* **213**:119–126.
- Kang JG, Amar MJ, Remaley AT, Kwon J, Blackshear PJ, Wang PY, and Hwang PM (2011) Zinc finger protein tristetraprolin interacts with CCL3 mRNA and regulates tissue inflammation. *J Immunol* **187**:2696–2701.
- Kato A and Schleimer RP (2007) Beyond inflammation: airway epithelial cells are at the interface of innate and adaptive immunity. *Curr Opin Immunol* **19**: 711–720.
- Kaur M, Holden NS, Wilson SM, Sukkar MB, Chung KF, Barnes PJ, Newton R, and Giembycz MA (2008) Effect of beta2-adrenoceptor agonists and other cAMP-elevating agents on inflammatory gene expression in human ASM cells: a role for protein kinase A. *Am J Physiol Lung Cell Mol Physiol* **295**:L505–L514.
- King EM, Kaur M, Gong W, Rider CF, Holden NS, and Newton R (2009) Regulation of tristetraprolin expression by interleukin-1 beta and dexamethasone in human pulmonary epithelial cells: roles for nuclear factor-kappa B and p38 mitogen-activated protein kinase. *J Pharmacol Exp Ther* **330**:575–585.
- Kobayashi Y, Mercado N, Barnes PJ, and Ito K (2011) Defects of protein phosphatase 2A causes corticosteroid insensitivity in severe asthma. *PLoS One* **6**:e27627.
- Kotlyarov A, Neining A, Schubert C, Eckert R, Birchmeier C, Volk HD, and Gaestel M (1999) MAPK kinase 2 is essential for LPS-induced TNF- $\alpha$  biosynthesis. *Nat Cell Biol* **1**:94–97.
- Kranias G, Watt LF, Carpenter H, Holst J, Ludowyke R, Strack S, Sim AT, and Verrills NM (2010) Protein phosphatase 2A carboxymethylation and regulatory B subunits differentially regulate mast cell degranulation. *Cell Signal* **22**: 1882–1890.
- Lai WS, Carballo E, Strum JR, Kennington EA, Phillips RS, and Blackshear PJ (1999) Evidence that tristetraprolin binds to AU-rich elements and promotes the deadenylation and destabilization of tumor necrosis factor alpha mRNA. *Mol Cell Biol* **19**:4311–4323.
- Lai WS, Thompson MJ, and Blackshear PJ (1998) Characteristics of the intron involvement in the mitogen-induced expression of Zfp-36. *J Biol Chem* **273**: 506–517.
- Lai WS, Thompson MJ, Taylor GA, Liu Y, and Blackshear PJ (1995) Promoter analysis of Zfp-36, the mitogen-inducible gene encoding the zinc finger protein tristetraprolin. *J Biol Chem* **270**:25266–25272.
- Lakser OJ, Lindeman RP, and Fredberg JJ (2002) Inhibition of the p38 MAP kinase pathway destabilizes smooth muscle length during physiological loading. *Am J Physiol Lung Cell Mol Physiol* **282**:L1117–L1121.
- Lee HH, Son YJ, Lee WH, Park YW, Chae SW, Cho WJ, Kim YM, Choi HJ, Choi DH, and Jung SW, et al. (2010) Tristetraprolin regulates expression of VEGF and tumorigenesis in human colon cancer. *Int J Cancer* **126**:1817–1827.
- Lee HH, Yoon NA, Vo MT, Kim CW, Woo JM, Cha HJ, Cho YW, Lee BJ, Cho WJ, and Park JW (2012) Tristetraprolin down-regulates IL-17 through mRNA destabilization. *FEBS Lett* **586**:41–46.
- Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, and Landvatter SW et al. (1994) A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* **372**: 739–746.
- Leppänen T, Jalonen U, Kankaanranta H, Tuominen R, and Moilanen E (2008) Inhibition of protein kinase C beta II downregulates tristetraprolin expression in activated macrophages. *Inflamm Res* **57**:230–240.
- Leppänen T, Jalonen U, Korhonen R, Tuominen RK, and Moilanen E (2010) Inhibition of protein kinase Cdelta reduces tristetraprolin expression by destabilizing its mRNA in activated macrophages. *Eur J Pharmacol* **628**:220–225.
- Linker K, Pautz A, Fehrich M, Hubrich T, Greeve J, and Kleiner H (2005) Involvement of KSRP in the post-transcriptional regulation of human iNOS expression-complex interplay of KSRP with TTP and HuR. *Nucleic Acids Res* **33**: 4813–4827.
- Lisi L, Navarra P, Feinstein DL, and Dello Russo C (2011) The mTOR kinase inhibitor rapamycin decreases iNOS mRNA stability in astrocytes. *J Neuroinflammation* **8**:1.
- Lomas DA, Lipson DA, Miller BE, Willits L, Keene O, Barnacle H, Barnes NC, and Tal-Singer R; Losmapimod Study Investigators (2012) An oral inhibitor of p38 MAP kinase reduces plasma fibrinogen in patients with chronic obstructive pulmonary disease. *J Clin Pharmacol* **52**:416–424.
- Ma F, Liu X, Li D, Wang P, Li N, Lu L, and Cao X (2010) MicroRNA-4661 upregulates IL-10 expression in TLR-triggered macrophages by antagonizing RNA-binding protein tristetraprolin-mediated IL-10 mRNA degradation. *J Immunol* **184**: 6053–6059.
- MacNee W, Allan RJ, Jones I, De Salvo MC, and Tan LF (2013) Efficacy and safety of the oral p38 inhibitor PH-797804 in chronic obstructive pulmonary disease: a randomised clinical trial. *Thorax* **68**:738–745.
- Mahtani KR, Brook M, Dean JL, Sully G, Saklatvala J, and Clark AR (2001) Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor alpha mRNA stability. *Mol Cell Biol* **21**:6461–6469.
- Marchese FP, Aubareda A, Tudor C, Saklatvala J, Clark AR, and Dean JL (2010) MAPK kinase 2 blocks tristetraprolin-directed mRNA decay by inhibiting CAF1 deadenylation recruitment. *J Biol Chem* **285**:27590–27600.
- Masaki K, Suzuki Y, Kagawa S, Kodama M, Kabata H, Miyata J, Tanaka K, Fukunaga K, Sayama K, and Oguma T, et al. (2014) Dual role of interleukin-23 in epicutaneously-sensitized asthma in mice. *Allergol Int* **63**:13–22.
- Mattos W, Lim S, Russell R, Jatakanon A, Chung KF, and Barnes PJ (2002) Matrix metalloproteinase-9 expression in asthma: effect of asthma severity, allergen challenge, and inhaled corticosteroids. *Chest* **122**:1543–1552.
- Mercado N, Hakim A, Kobayashi Y, Meah S, Usmani OS, Chung KF, Barnes PJ, and Ito K (2012) Restoration of corticosteroid sensitivity by p38 mitogen activated protein kinase inhibition in peripheral blood mononuclear cells from severe asthma. *PLoS One* **7**:e41582.
- Molle C, Zhang T, Ysebrant de Lendonck L, Gueydan C, Andrianne M, Sherer F, Van Sinaeys G, Blackshear PJ, Leo O, and Goriely S (2013) Tristetraprolin regulation of interleukin 23 mRNA stability prevents a spontaneous inflammatory disease. *J Exp Med* **210**:1675–1684.
- Moutzouris JP, Che W, Ramsay EE, Manetsch M, Alkhoury H, Bjorkman AM, Schuster F, Ge Q, and Ammit AJ (2010) Proteasomal inhibition upregulates the endogenous MAPK deactivator MKP-1 in human airway smooth muscle: mechanism of action and effect on cytokine secretion. *Biochim Biophys Acta* **1803**: 414–423.
- Nath P, Leung SY, Williams A, Noble A, Chakravarty SD, Luedtke GR, Medicherla S, Higgins LS, Protter A, and Chung KF (2006) Importance of p38 mitogen-activated protein kinase pathway in allergic airway remodelling and bronchial hyperresponsiveness. *Eur J Pharmacol* **544**:160–167.

- Neininger A, Kontoyiannis D, Kotlyarov A, Winzen R, Eckert R, Volk HD, Holtmann H, Kollias G, and Gaestel M (2002) MK2 targets AU-rich elements and regulates biosynthesis of tumor necrosis factor and interleukin-6 independently at different post-transcriptional levels. *J Biol Chem* **277**:3065–3068.
- Ngoc LV, Wauquier C, Soin R, Bousbata S, Twyffels L, Kruijs V, and Gueydan C (2014) Rapid proteasomal degradation of posttranscriptional regulators of the TIS11/tristetraprolin family is induced by an intrinsically unstructured region independently of ubiquitination. *Mol Cell Biol* **34**:4315–4328.
- Ogilvie RL, Abelson M, Hau HH, Vlasova I, Blackshear PJ, and Bohjanen PR (2005) Tristetraprolin down-regulates IL-2 gene expression through AU-rich element-mediated mRNA decay. *J Immunol* **174**:953–961.
- Patil CS, Liu M, Zhao W, Coatney DD, Li F, VanTubergen EA, D'Silva NJ, and Kirkwood KL (2008) Targeting mRNA stability arrests inflammatory bone loss. *Mol Ther* **16**:1657–1664.
- Perrotti D and Neviani P (2008) Protein phosphatase 2A (PP2A), a drugable tumor suppressor in Ph1(+) leukemias. *Cancer Metastasis Rev* **27**:159–168.
- Pfeiffer JR and Brooks SA (2012) Cullin 4B is recruited to tristetraprolin-containing messenger ribonucleoproteins and regulates TNF- $\alpha$  mRNA polysome loading. *J Immunol* **188**:1828–1839.
- Phillips K, Kedersha N, Shen L, Blackshear PJ, and Anderson P (2004) Arthritis suppressor genes TIA-1 and TTP dampen the expression of tumor necrosis factor alpha, cyclooxygenase 2, and inflammatory arthritis. *Proc Natl Acad Sci USA* **101**:2011–2016.
- Prickett TD and Brautigan DL (2007) Cytokine activation of p38 mitogen-activated protein kinase and apoptosis is opposed by alpha-4 targeting of protein phosphatase 2A for site-specific dephosphorylation of MEK3. *Mol Cell Biol* **27**:4217–4227.
- Qian X, Ning H, Zhang J, Hoft DF, Stumpo DJ, Blackshear PJ, and Liu J (2011) Posttranscriptional regulation of IL-23 expression by IFN-gamma through tristetraprolin. *J Immunol* **186**:6454–6464.
- Quante T, Ng YC, Ramsay EE, Henness S, Allen JC, Parmentier J, Ge Q, and Ammit AJ (2008) Corticosteroids reduce IL-6 in ASM cells via up-regulation of MKP-1. *Am J Respir Cell Mol Biol* **39**:208–217.
- Rael EL and Lockey RF (2011) Interleukin-13 signaling and its role in asthma. *World Allergy Organ J* **4**:54–64.
- Rigby WF, Roy K, Collins J, Rigby S, Connolly JE, Bloch DB, and Brooks SA (2005) Structure/function analysis of tristetraprolin (TTP): p38 stress-activated protein kinase and lipopolysaccharide stimulation do not alter TTP function. *J Immunol* **174**:7883–7893.
- Ronkina N, Menon MB, Schwermann J, Tiedje C, Hitti E, Kotlyarov A, and Gaestel M (2010) MAPKAP kinases MK2 and MK3 in inflammation: complex regulation of TNF biosynthesis via expression and phosphorylation of tristetraprolin. *Biochem Pharmacol* **80**:1915–1920.
- Rosenberger CM, Podymnagin RL, Navarro G, Zhao GW, Askovich PS, Weiss MJ, and Aderem A (2012) miR-451 regulates dendritic cell cytokine responses to influenza infection. *J Immunol* **189**:5965–5975.
- Rousseau S, Houle F, Landry J, and Huot J (1997) p38 MAP kinase activation by vascular endothelial growth factor mediates actin reorganization and cell migration in human endothelial cells. *Oncogene* **15**:2169–2177.
- Sandler H, Kreth J, Timmers HT, and Stoecklin G (2011) Not1 mediates recruitment of the deadenylase Caf1 to mRNAs targeted for degradation by tristetraprolin. *Nucleic Acids Res* **39**:4373–4386.
- Santoro MF, Annand RR, Robertson MM, Peng YW, Brady MJ, Mankovich JA, Hackett MC, Ghayur T, Walter G, and Wong WW et al. (1998) Regulation of protein phosphatase 2A activity by caspase-3 during apoptosis. *J Biol Chem* **273**:13119–13128.
- Sauer I, Schaljo B, Vogl C, Gattermeier I, Kolbe T, Müller M, Blackshear PJ, and Kovarik P (2006) Interferons limit inflammatory responses by induction of tristetraprolin. *Blood* **107**:4790–4797.
- Schreiber S, Feagan B, D'Haens G, Colombel JF, Geboes K, Yurcov M, Isakov V, Golovenko O, Bernstein CN, and Ludwig D, et al BIRB 796 Study Group (2006) Oral p38 mitogen-activated protein kinase inhibition with BIRB 796 for active Crohn's disease: a randomized, double-blind, placebo-controlled trial. *Clin Gastroenterol Hepatol* **4**:325–334.
- Semaan N, Frenzel L, Alsaleh G, Suffert G, Gottenberg JE, Sibilia J, Pfeffer S, and Wachsmann D (2011) miR-346 controls release of TNF- $\alpha$  protein and stability of its mRNA in rheumatoid arthritis via tristetraprolin stabilization. *PLoS One* **6**:e19827.
- Shi Y (2009) Assembly and structure of protein phosphatase 2A. *Sci China C Life Sci* **52**:135–146.
- Singh D (2013) P38 inhibition in COPD; cautious optimism. *Thorax* **68**:705–706.
- Singh D, Smyth L, Borrill Z, Sweeney L, and Tal-Singer R (2010) A randomized, placebo-controlled study of the effects of the p38 MAPK inhibitor SB-681323 on blood biomarkers of inflammation in COPD patients. *J Clin Pharmacol* **50**:94–100.
- Stoecklin G, Gross B, Ming XF, and Moroni C (2003) A novel mechanism of tumor suppression by destabilizing AU-rich growth factor mRNA. *Oncogene* **22**:3554–3561.
- Stoecklin G, Stubbs T, Kedersha N, Wax S, Rigby WF, Blackwell TK, and Anderson P (2004) MK2-induced tristetraprolin:14-3-3 complexes prevent stress granule association and ARE-mRNA decay. *EMBO J* **23**:1313–1324.
- Sully G, Dean JL, Wait R, Rawlinson L, Santalucia T, Saklatvala J, and Clark AR (2004) Structural and functional dissection of a conserved destabilizing element of cyclo-oxygenase-2 mRNA: evidence against the involvement of AUF-1 [AU-rich element/poly(U)-binding/degradation factor-1], AUF-2, tristetraprolin, HuR (Hu antigen R) or FBP1 (far-upstream-sequence-element-binding protein 1). *Biochem J* **377**:629–639.
- Sun L, Stoecklin G, Van Way S, Hinkovska-Galcheva V, Guo RF, Anderson P, and Shanley TP (2007) Tristetraprolin (TTP)-14-3-3 complex formation protects TTP from dephosphorylation by protein phosphatase 2a and stabilizes tumor necrosis factor-alpha mRNA. *J Biol Chem* **282**:3766–3777.
- Suswam E, Li Y, Zhang X, Gillespie GY, Li X, Shacka JJ, Lu L, Zheng L, and King PH (2008) Tristetraprolin down-regulates interleukin-8 and vascular endothelial growth factor in malignant glioma cells. *Cancer Res* **68**:674–682.
- Suswam EA, Shacka JJ, Walker K, Lu L, Li X, Si Y, Zhang X, Zheng L, Nabors LB, and Cao H et al. (2013) Mutant tristetraprolin: a potent inhibitor of malignant glioma cell growth. *J Neurooncol* **113**:195–205.
- Tan SL and Parker PJ (2003) Emerging and diverse roles of protein kinase C in immune cell signalling. *Biochem J* **376**:545–552.
- Tang ML and Fiscus LC (2001) Important roles for L-selectin and ICAM-1 in the development of allergic airway inflammation in asthma. *Pulm Pharmacol Ther* **14**:203–210.
- Tay KH, Jin L, Tseng HY, Jiang CC, Ye Y, Thorne RF, Liu T, Guo ST, Verrills NM, and Hersey P et al. (2012) Suppression of PP2A is critical for protection of melanoma cells upon endoplasmic reticulum stress. *Cell Death Dis* **3**:e337.
- Taylor GA, Carballo E, Lee DM, Lai WS, Thompson MJ, Patel DD, Schenkman DI, Gilkeson GS, Broxmeyer HE, and Haynes BF et al. (1996a) A pathogenetic role for TNF alpha in the syndrome of cachexia, arthritis, and autoimmunity resulting from tristetraprolin (TTP) deficiency. *Immunity* **4**:445–454.
- Taylor GA, Lai WS, Oakey RJ, Seldin MF, Shows TB, Eddy RL, Jr, and Blackshear PJ (1991) The human TTP protein: sequence, alignment with related proteins, and chromosomal localization of the mouse and human genes. *Nucleic Acids Res* **19**:3454.
- Taylor GA, Thompson MJ, Lai WS, and Blackshear PJ (1995) Phosphorylation of tristetraprolin, a potential zinc finger transcription factor, by mitogen stimulation in intact cells and by mitogen-activated protein kinase in vitro. *J Biol Chem* **270**:13341–13347.
- Taylor GA, Thompson MJ, Lai WS, and Blackshear PJ (1996b) Mitogens stimulate the rapid nuclear to cytosolic translocation of tristetraprolin, a potential zinc-finger transcription factor. *Mol Endocrinol* **10**:140–146.
- Tchen CR, Brook M, Saklatvala J, and Clark AR (2004) The stability of tristetraprolin mRNA is regulated by mitogen-activated protein kinase p38 and by tristetraprolin itself. *J Biol Chem* **279**:32393–32400.
- Tudor C, Marchese FP, Hitti E, Aubareda A, Rawlinson L, Gaestel M, Blackshear PJ, Clark AR, Saklatvala J, and Dean JL (2009) The p38 MAPK pathway inhibits tristetraprolin-directed decay of interleukin-10 and pro-inflammatory mediator mRNAs in murine macrophages. *FEBS Lett* **583**:1933–1938.
- Van Tubergen E, Vander Broek R, Lee J, Wolf G, Carey T, Bradford C, Prince M, Kirkwood KL, and D'Silva NJ (2011) Tristetraprolin regulates interleukin-6, which is correlated with tumor progression in patients with head and neck squamous cell carcinoma. *Cancer* **117**:2677–2689.
- Van Tubergen EA, Banerjee R, Liu M, Vander Broek R, Light E, Kuo S, Feinberg SE, Willis AL, Wolf G, and Carey T, et al. (2013) Inactivation or loss of TTP promotes invasion in head and neck cancer via transcript stabilization and secretion of MMP9, MMP2, and IL-6. *Clin Cancer Res* **19**:1169–1179.
- Varnum BC, Ma QF, Chi TH, Fletcher B, and Herschman HR (1991) The TIS11 primary response gene is a member of a gene family that encodes proteins with a highly conserved sequence containing an unusual Cys-His repeat. *Mol Cell Biol* **11**:1754–1758.
- Woo CH, Lim JH, and Kim JH (2004) Lipopolysaccharide induces matrix metalloproteinase-9 expression via a mitochondrial reactive oxygen species-p38 kinase-activator protein-1 pathway in Rat 264.7 cells. *J Immunol* **173**:6973–6980.
- Xing L, Devadas B, Devraj RV, Selness SR, Shieh H, Walker JK, Mao M, Messing D, Samas B, and Yang JZ et al. (2012) Discovery and characterization of atropisomer PH-797804, a p38 MAP kinase inhibitor, as a clinical drug candidate. *Chem-MedChem* **7**:273–280.
- Yang J, Fan GH, Wadzinski BE, Sakurai H, and Richmond A (2001) Protein phosphatase 2A interacts with and directly dephosphorylates RelA. *J Biol Chem* **276**:47828–47833.
- Young P, McDonnell P, Dunnington D, Hand A, Laydon J, and Lee J (1993) Pyridinyl imidazoles inhibit IL-1 and TNF production at the protein level. *Agents Actions* **39**:C67–C69.
- Zhao W, Liu M, D'Silva NJ, and Kirkwood KL (2011) Tristetraprolin regulates interleukin-6 expression through p38 MAPK-dependent affinity changes with mRNA 3' untranslated region. *J Interferon Cytokine Res* **31**:629–637.

**Address correspondence to:** Dr. Alaina J. Ammit, Faculty of Pharmacy, University of Sydney, Sydney, New South Wales 2006, Australia. E-mail: alaina.ammit@sydney.edu.au