

MINIREVIEW

Arrestin-Dependent and -Independent Internalization of G Protein–Coupled Receptors: Methods, Mechanisms, and Implications on Cell Signaling

Ee Von Moo, Jeffrey R. van Senten, Hans Bräuner-Osborne, and  Thor C. Møller

Department of Drug Design and Pharmacology, University of Copenhagen, Copenhagen, Denmark

Received October 27, 2020; accepted January 7, 2021

ABSTRACT

Agonist-induced endocytosis is a key regulatory mechanism for controlling the responsiveness of the cell by changing the density of cell surface receptors. In addition to the role of endocytosis in signal termination, endocytosed G protein–coupled receptors (GPCRs) have been found to signal from intracellular compartments of the cell. Arrestins are generally believed to be the master regulators of GPCR endocytosis by binding to both phosphorylated receptors and adaptor protein 2 (AP-2) or clathrin, thus recruiting receptors to clathrin-coated pits to facilitate the internalization process. However, many other functions have been described for arrestins that do not relate to their role in terminating signaling. Additionally, there are now more than 30 examples of GPCRs that internalize independently of arrestins. Here we review the methods, pharmacological tools, and cellular backgrounds used to determine the role of arrestins in receptor internalization, highlighting their advantages and caveats. We also summarize key examples of arrestin-independent GPCR endocytosis in the literature

and their suggested alternative endocytosis pathway (e.g., the caveolae-dependent and fast endophilin-mediated endocytosis pathways). Finally, we consider the possible function of arrestins recruited to GPCRs that are endocytosed independently of arrestins, including the catalytic arrestin activation paradigm. Technological improvements in recent years have advanced the field further, and, combined with the important implications of endocytosis on drug responses, this makes endocytosis an obvious parameter to include in molecular pharmacological characterization of ligand-GPCR interactions.

SIGNIFICANCE STATEMENT

G protein–coupled receptor (GPCR) endocytosis is an important means to terminate receptor signaling, and arrestins play a central role in the widely accepted classical paradigm of GPCR endocytosis. In contrast to the canonical arrestin-mediated internalization, an increasing number of GPCRs are found to be endocytosed via alternate pathways, and the process appears more diverse than the previously defined “one pathway fits all.”

Introduction

G protein–coupled receptors (GPCRs) form the largest class of transmembrane cell surface receptors, and they regulate intracellular signaling in response to a diverse range of extracellular stimuli (Lefkowitz, 2013). Initiation, processing,

and termination of these signals are tightly regulated in a spatiotemporal manner to ensure homeostasis. To date, approximately one-third of the marketed drugs target GPCRs, illustrating their importance in human pathologies (Hauser et al., 2017).

Agonist-stimulated GPCRs catalyze the activation of heterotrimeric G proteins and thereby modulate downstream effector proteins, including adenylyl cyclase, phospholipase C, and Rho guanine nucleotide exchange factor (Hilger et al., 2018). Stimulation of these pathways can evoke receptor phosphorylation on serine and threonine residues in the third

This project received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreements no. 797497 (T.C.M.) and no. 846827 (E.V.M.). The authors declare no conflicts of interest.

<https://doi.org/10.1124/molpharm.120.000192>

ABBREVIATIONS: AP-2, adaptor protein 2; AT, angiotensin; BAR, Bin/amphiphysin/Rvs; BRET, bioluminescence resonance energy transfer; CaS, calcium-sensing; Cdc42, cell division control protein 42 homolog; C-domain, C-terminal domain; CIP4, Cdc42-interacting protein 4; C-tail, carboxy-terminal tail; ERK, extracellular signal-regulated kinase; FEME, fast endophilin-mediated endocytosis; FIAsh fluorescein arsenical hairpin FPR1, formylpeptide receptor 1; GLP, glucagon-like peptide; GPCR, G protein–coupled receptor; GRK, GPCR kinase; HEK293, human embryonic kidney 293; ICL, intracellular loop; MEF, mouse embryonic fibroblast; MHC-I, major histocompatibility complex class I; mGlu, metabotropic glutamate; N-domain, N-terminal domain; PAR, proteinase-activated receptor; PI(4,5)P₂, phosphatidylinositol 4,5-bisphosphate; PKC, protein kinase C; PTH, parathyroid hormone; RBL, rat basophilic leukemia; SH3, Src homology 3; siRNA, small interfering RNA; 7TM, seven transmembrane; TR-FRET, time-resolved Förster resonance energy transfer.

intracellular loop (ICL) and the carboxy-terminal tail (C-tail), mediated by GPCR kinases (GRKs) and second messenger-dependent kinases from the protein kinase A and protein kinase C (PKC) families (Ferguson, 2001; Gurevich and Gurevich, 2019). Recruitment of arrestin proteins to phosphorylated active-state receptors can lead to desensitization of second messenger signaling by sterically precluding G protein coupling to receptors. Furthermore, arrestins can facilitate and regulate receptor signaling by scaffolding a wide range of proteins involved in signaling pathways, for instance, components of mitogen-activated protein kinase cascades, Src family tyrosine kinases, and E3 ubiquitin ligases (Peterson and Luttrell, 2017).

To protect cells from overstimulation upon prolonged or repeated exposure to agonists, activated GPCRs are removed from the cell surface by means of endocytosis. In this way, the majority of the cell surface population of a GPCR may be internalized within minutes of agonist stimulation (January et al., 1997). Internalized receptors are processed and sorted in the endosomal network for recycling to the plasma membrane (resensitization) or degradation via the lysosomal pathway (downregulation) (Pavlos and Friedman, 2017). Moreover, GPCRs can signal from intracellular compartments as well. This may result in different signaling consequences compared with cell surface signaling due to distinct location and timing (Lobingier and von Zastrow, 2019).

In this mini review we summarize the endocytosis pathways that have been described for GPCRs—arrestin-dependent and -independent—and the receptors that have been shown to be endocytosed independently of arrestins, including the tools and methods that have been used to assess the arrestin dependence. Finally, we discuss the possible functional implications of arrestin recruitment to receptors that internalize independently of arrestins.

Clathrin-Mediated Endocytosis

Several mechanisms of GPCR endocytosis have been described. The best characterized endocytic route and the predominant pathway of endocytosis in mammalian cells is clathrin-mediated endocytosis (Pearse, 1976) (Fig. 1A). Ligand-activated GPCRs are in many cases targeted to clathrin-coated pits by binding to arrestins (Traub, 2009). There are four subtypes of arrestin: arrestin-1 and -4 that are mainly expressed in the visual system and arrestin-2 and -3 (also called β -arrestin-1 and -2, respectively) that are ubiquitously expressed (Mundell et al., 2002). Activated arrestins are recruited to clathrin-coated pits through interaction with the $\beta 2$ adaptin subunit of adaptor protein-2 (AP-2), via an RXR motif located in the autoinhibitory C-tail of all arrestin subtypes (Laporte et al., 2000; Schmid et al., 2006). In addition, arrestins can interact with clathrin through an LIELD or LIEFE motif located in the autoinhibitory segment of arrestin-2 and -3 and/or through a loop in the C-terminal domain (C-domain) of arrestin that is present in the long splice isoform of arrestin-2 and in arrestin-1 and -4 (Goodman et al., 1996; Kang et al., 2009). In the inactive form, the C-tail of arrestin functions as an autoinhibitory segment that is bound to a groove in the N-terminal domain (N-domain) of arrestin, thus masking the AP-2 and clathrin motifs (Hirsch et al., 1999). Binding of a phosphorylated receptor C-tail displaces the autoinhibitory segment and enables arrestin to interact

with AP-2 and clathrin (Xiao et al., 2004; Nobles et al., 2007). Alternatively, the $\mu 2$ adaptin subunit of AP-2 has been shown to interact directly with GPCRs, through which it might facilitate clathrin-mediated endocytosis independently of arrestins (Diviani et al., 2003; Paing et al., 2006).

AP-2 and other clathrin adaptor proteins initiate clathrin-mediated endocytosis by binding to plasma membrane domains enriched for the phospholipid phosphatidylinositol 4,5-bisphosphate [PI(4,5)P₂]. AP-2 is a tetrameric protein complex comprising α , $\beta 2$, $\mu 2$, and $\sigma 2$ adaptin subunits. Interaction with PI(4,5)P₂ induces a conformational change in AP-2, exposing binding sites for clathrin, receptors, and other cargo proteins and additional PI(4,5)P₂ interaction motifs (Kelly et al., 2014) (Fig. 1A, step 2a and b). Clathrin molecules interact with AP-2 and form polymeric lattices, enveloping the plasma membrane into clathrin-coated pits (Fotin et al., 2004; Dannhauser and Ungewickell, 2012) (Fig. 1A, step 3). Bin/amphiphysin/Rvs (BAR) domain-containing proteins, such as endophilins, induce and stabilize membrane curvature and thereby mediate constriction of the invagination neck, whereas membrane scission is powered by oligomers of dynamin, which is a GTPase (Sundborger et al., 2011). Dephosphorylation of PI(4,5)P₂ (Cremona et al., 1999) and chaperone-mediated disruption of clathrin-clathrin interactions (Schlossman et al., 1984) drive the uncoating of clathrin-coated vesicles, followed by vesicle fusion to early endosomes.

Another class of clathrin adaptors is constituted by epsins, which are monomeric proteins that can interact simultaneously with PI(4,5)P₂, clathrin, and AP-2 (Traub, 2009). Through their ubiquitin-interacting motifs, epsins can directly recruit polyubiquitinated proteins to clathrin-coated pits, including GPCRs (Chen et al., 2011) (Fig. 1A, step 2b). Many other proteins contribute to the formation of clathrin-coated pits (Traub, 2011), potentially including yet unidentified GPCR adaptor proteins.

Clathrin-Independent Endocytosis

Although the majority of endocytic vesicles arise from clathrin-mediated endocytosis (Pearse, 1976; Anderson et al., 1977), several other endocytic mechanisms have been shown to operate at the plasma membrane of mammalian cells. Some of these have been implicated in the internalization of GPCRs. Characterization of clathrin-independent endocytosis pathways, however, is hampered by a lack of knowledge about specific cargo and cellular machinery and an absence of pathway-specific manipulation tools (Sandvig et al., 2018). Hence, clathrin-independent endocytosis routes are generally less well delineated than clathrin-mediated endocytosis.

Fast Endophilin-Mediated Endocytosis. In addition to their contribution to clathrin-mediated endocytosis, endophilins (endophilin A1–3, but not B1 and B2) are essential for a rapid, clathrin-independent internalization pathway, called fast endophilin-mediated endocytosis (FEME) (Boucrot et al., 2015) (Fig. 1B). This rapid response to receptor activation (few seconds) relies on the presence of endophilin-enriched domains in the plasma membrane, which are dynamically assembled and disassembled in the absence of activated receptors or other cargo proteins.

FEME priming starts with the recruitment of the FES/CIP4 homology–BAR domain proteins formin-binding protein 17




Fig. 1. Endocytosis pathways involved in GPCR internalization. (A) Clathrin-mediated endocytosis of GPCRs. (1) Ligand-mediated receptor activation leads to conformational changes, for instance, exposing AP-2 binding motifs in the receptor C-tail, and/or post-translational modification of the receptor, such as phosphorylation or ubiquitination. (2a) Most GPCRs are recruited to clathrin-coated pits through arrestins. Binding of arrestin to the phosphorylated receptor C-tail enables interaction of the arrestin C-tail with AP-2 and clathrin. (2b) Additionally, receptors can be recruited to clathrin-coated pits independently of arrestins, through direct interaction with AP-2 or binding of their polyubiquitinated C-tails to epsins. (3) Clathrin is

and Cdc42-interacting protein 4 (CIP4) by active, membrane-bound cell division control protein 42 homolog (Cdc42). In turn, these proteins recruit the phosphatidylinositol 3,4,5-trisphosphate 5' phosphatase 1 and 2, thus increasing the local concentration of phosphatidylinositol-3,4-bisphosphate (Chan Wah Hak et al., 2018). Through its phosphatidylinositol-3,4-bisphosphate-binding pleckstrin homology domain and multiple endophilin-binding motifs, lamellipodin facilitates the concentration of endophilins in patches at the cell surface (Vehlow et al., 2013; Boucrot et al., 2015) (Fig. 1B, step 1). Endophilins containing BAR domains with an additional N-terminal amphipathic helix induce plasma membrane curvature by insertion of amphipathic helices in the inner membrane leaflet as well as membrane scaffolding through their BAR domains (Boucrot et al., 2012), whereas protein scaffolding is mediated via their Src homology 3 (SH3) domains (Boucrot et al., 2012, 2015; Vehlow et al., 2013). FEME vesicle formation requires membrane scaffolding by endophilin, actin polymerization, and dynamin-mediated scission (Boucrot et al., 2015; Renard et al., 2015) (Fig. 1B, step 3). Multiple ligand-stimulated GPCRs can internalize via the FEME route through direct interaction between proline-rich motifs in their ICL3 and the SH3 domain of endophilins (Tang et al., 1999; Boucrot et al., 2015), but it is not yet understood how receptor stimulation triggers FEME (Fig. 1B, step 2).

Endocytosis via Caveolae. Caveolae are invaginations of the plasma membrane, enriched for cholesterol and sphingomyelin (Ortegren et al., 2004). Their formation relies on the coordinated membrane-deforming activity of caveolin, cavin, and pacsin (also named syndapin) proteins (Ludwig et al., 2013) (Fig. 1C). Caveolins are cholesterol-binding proteins integrated in the inner leaflet of the plasma membrane that form 12- to 16-meric complexes (Ariotti et al., 2015). Mature caveolae are formed upon phosphatidylserine- and PI(4,5)P₂-mediated association of coat-forming trimeric cavin complexes (Kovtun et al., 2014), FES/CIP4 homology-BAR domain-containing pacsin proteins (Hansen et al., 2011) and the EH domain-containing 2 protein (Yeow et al., 2017) with these caveolin-rich domains (Fig. 1C, step 2). Pacsin furthermore interacts with dynamin via its SH3 domain, facilitating caveolae vesicle scission (Koch et al., 2011).

Several GPCRs have been reported to localize in caveolae, interact with caveolin, and/or internalize via caveolae-mediated endocytosis (Chini and Parenti, 2004). GPCRs can directly bind to caveolin through $\phi X\phi XXXX\phi$ or $\phi XXXX\phi XX\phi$ motifs (X = any amino acid; ϕ = Phe, Trp, or Tyr) (Couet et al., 1997) (Fig. 1C, step 1a). Interaction with caveolin, however, does not necessarily dictate receptor endocytosis via caveolae, as caveolin can also function as chaperone during receptor transport to the cell surface or facilitate caveolae localization without triggering receptor internalization (Chini and Parenti, 2004). Also, G α_q subunits, but not G $\beta\gamma$ or other

G α proteins, can interact with caveolin, which facilitates association between receptors and caveolae (Sengupta et al., 2008; Calizo and Scarlata, 2012) (Fig. 1C, step 1b).

In contrast to clathrin-coated pits, the density of caveolae differs between tissues and cell types: whereas caveolae can be undetectable in some cell types, they can occupy up to 50% of the plasma membrane surface in others (Thorn et al., 2003; Zhuang et al., 2011). It is unclear whether differences in caveolae abundance affect endocytosis of receptors via this pathway. Furthermore, even though caveolae can bud from the plasma membrane and translocate to early endosomes (Hayer et al., 2010), caveolae are now primarily viewed as membrane structures with functions different from endocytosis, including membrane tension buffer and specialized lipid rafts important for signaling (Sinha et al., 2011; Shvets et al., 2015). Altogether, the molecular mechanisms driving GPCR internalization via caveolae are still poorly understood, and it is unknown how agonist stimulation triggers caveolae-mediated endocytosis of GPCRs.

Arrestin-Independent Agonist-Induced Endocytosis

Despite the well characterized role of arrestins in GPCR endocytosis, including inhibiting GPCR/G protein coupling and initiating internalization (Kang et al., 2013; Tian et al., 2014), there is now a considerable number of examples of arrestin-independent GPCR internalization upon agonist stimulation (Table 1). In several cases the mechanism has been investigated further, and alternative mediators of endocytosis have been identified, such as caveolae, endophilin, GRKs, clathrin, and clathrin adaptors.

Clathrin- and AP-2-Dependent Pathway. In addition to the role of AP-2 in arrestin-mediated endocytosis, the $\mu 2$ adaptin subunit of AP-2 can directly interact with polyarginine motifs (Diviani et al., 2003), dileucine motifs ([D/E]XXXL [L/I], X = any amino acid) (Pandey, 2010), or tyrosine motifs (YXX Φ , Φ = bulky hydrophobic amino acid) (Ohno et al., 1995) in the intracellular loops or C-tail of GPCRs, through which it might facilitate clathrin-mediated endocytosis independently of arrestins.

The C-tail of proteinase-activated receptor (PAR) 1 contains a YXXL $\mu 2$ adaptin binding motif (Paing et al., 2004). Similar to the role of arrestins for other receptors, AP-2 interacts directly with PAR1 and is required for its constitutive and agonist-induced internalization through a clathrin- and dynamin-dependent pathway (Paing et al., 2006). Arrestins play a critical role in PAR1 desensitization to uncouple G protein signaling but are not essential for PAR1 internalization (Paing et al., 2002). Native receptors are found in two distinct pools: one in the cell membrane and one in an intracellular compartment (Shapiro et al., 1996).

recruited to the plasma membrane by adaptor proteins, such as AP-2 and epsins, to form clathrin carriers upon scission of clathrin-coated pits from the plasma membrane by dynamin. (B) Fast endophilin-mediated endocytosis. (1) Endophilin is concentrated at the plasma membrane through interaction with lamellipodin (Lmpd), which depends on the sequential action of Cdc42, formin-binding protein 17 (FBP17), CIP4, and phosphatidylinositol 3,4,5-trisphosphate 5' phosphatase (SHIP) 1/2. (2) Interaction of a ligand-activated GPCR with endophilin triggers FEME. (3) FEME carriers are formed upon scission of endophilin stabilized invaginations by dynamin. (C) Endocytosis via caveolae. GPCRs can be recruited to caveolae through direct interaction with caveolin (1a) or via G α_q -mediated interaction with caveolin (1b). (2) Caveolae membrane deformations consist of oligomeric caveolin complexes stabilized by coat-forming cavins and membrane curvature inducing pacsins. Caveolae carrier formation relies on dynamin for membrane scission. For clarity, only a selection of the components involved in the various endocytosis pathways are represented in the schematic. PI(3,4,5)P₃, phosphatidylinositol 3,4-bisphosphate; PI(3,4,5)P₂, phosphatidylinositol 3,4,5-trisphosphate; PS, phosphatidylserine.

TABLE 1
GPCRs that have been shown to undergo arrestin-independent agonist-induced or constitutive endocytosis

Receptor	Mechanism of internalization	Cell line	Method used to show arrestin independence	Reference
Arrestin-independent agonist-induced endocytosis				
β_1 - and α_9 -adrenoceptors and D ₄ adrenoceptor and AT _{1A}	FEME	BS-C-1	Indirect evidence (FEME is clathrin- and arrestin-independent)	Boucrot et al., 2015
β_2 -adrenoceptor and AT _{1A}	Caveolae (partial arrestin-dependence)	HEK293, COS-7	Dominant negative arrestin (arrestin-2-V53D)	Guo et al., 2015
5-HT _{2A}	Dynamamin-dependent	HEK293, C6	Dominant negative arrestins (arrestin-2-V53D, arrestin-2 ³¹⁹⁻⁴¹⁸ , arrestin-3 ²⁸⁴⁻⁴⁰⁹)	Bhatnagar et al., 2001; Gray et al., 2001
Apelin	Clathrin-dependent	HEK293	Dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Pope et al., 2016
BLT ₁	Dynamamin- and GRK2 phosphorylation-dependent	RBL-2H3	Dominant negative arrestin (arrestin-2-V53D)	Chen et al., 2004b
D ₃	Clathrin- and dynamamin-dependent, FEME	CHO, HEK293, BS-C-1	Dominant negative arrestins (arrestin-2-V53D, arrestin-3-V54D)	Boucrot et al., 2015; Xu et al., 2019
ET _A	Caveolae (partial arrestin-dependence)	HEK293	Arrestin-AP-2 BRET, arrestin-AP-2 complex inhibitor (barbadin), arrestin-2/3 siRNA	Okamoto et al., 2000; Hamdan et al., 2007; Beautrait et al., 2017
ET _B	Unknown (partial arrestin-dependence)	HEK293	Arrestin-AP-2 BRET, arrestin-2/3 siRNA	Hamdan et al., 2007
FPR1	GRK phosphorylation-dependent	MEFs, U937	Arrestin-2/3 ^{-/-} MEFs, dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Hsu et al., 1997; Gilbert et al., 2001; Vines et al., 2003
GLP-1	Dynamamin-dependent and caveolae	HEK293, MIN6	HEK293-Arrestin-2/3, arrestin-3 overexpression	Syme et al., 2006; Jones et al., 2018
GLP-2	Caveolae	BHK-21, DLD-1	Indirect evidence (dynamamin-independent and C-terminal truncated receptor mutant)	Estall et al., 2004, 2005
IP	Clathrin- and dynamamin-dependent	HEK293	Dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Smyth et al., 2000
M ₁	Clathrin- and dynamamin-dependent	tsA201, MDCK, HEK293	Overexpression of arrestin-2 and -3, dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Lee et al., 1998; Vögler et al., 1998; Shmuel et al., 2007
M ₂	Phosphorylation- and Arf6 GTPase-dependent	tsA201, HeLa	Arrestin-2 and -3 overexpression	Pals-Rylaarsdam et al., 1997; Delaney et al., 2002
M ₃	Dynamamin-dependent	tsA201, HEK293	Overexpression of arrestin-2 and -3, dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Lee et al., 1998; Vögler et al., 1998
M ₄	Dynamamin-dependent, FEME	tsA201, BS-C-1	Overexpression of arrestin-2 and -3, dominant negative arrestin (arrestin-2 ³¹⁹⁻⁴¹⁸)	Lee et al., 1998; Vögler et al., 1998; Boucrot et al., 2015
PAR1	Dependent on clathrin, dynamamin, phosphorylation, epsin-1 and AP-2	MEFs, COS-7, HeLa, HEK293	Arrestin-2/3 ^{-/-} MEFs	Paing et al., 2002; Paing et al., 2006; Chen et al., 2011
PAR4	Dependent on clathrin, AP-2 and dynamamin	HeLa	Arrestin-2/3 ^{-/-} MEFs	Smith et al., 2016
PKR ₂	GRK2 phosphorylation- and clathrin-dependent	HEK293	Dominant negative arrestins (arrestin-2-V53D, arrestin-3-V54D), arrestin-2 and -3 siRNA	Yin et al., 2014
Secretin	PKA-dependent	HEK293	Dominant negative arrestin (arrestin-2-V53D)	Walker et al., 1999
UT	Unknown	HEK293, MEFs	Dominant negative arrestin (arrestin-3-V54D), arrestin-2/3 ^{-/-} MEFs	Giebing et al., 2005
Constitutive internalization				
CaS	Unknown (partial arrestin-dependence)	HEK293	HEK293-Arrestin-2/3	Mos et al., 2019
CXCR4	Dynamamin- and PKC-dependent	HEK293 COS-7	Indirect evidence (C-terminal truncated mutant)	Signoret et al., 1997, 1998
ADGRA3	Clathrin-dependent	HEK293	HEK293-Arrestin-2/3	Spieß et al., 2019
M ₃ and β_2 -adrenoceptor	MHC-1	HeLa, COS-7	Indirect evidence (clathrin-independent)	Scarselli and Donaldson, 2009
mGlu ₅	Dynamamin-dependent, caveolae	COS-7	Indirect evidence (clathrin-independent)	Fourgeaud et al., 2003
mGlu ₇	MHC-1	HeLa	Indirect evidence (clathrin-independent)	Lavezzari and Roche, 2007
PAR1	Clathrin- and AP-2-dependent	MEFs, COS-7, HeLa, HEK293	Arrestin-2/3 ^{-/-} MEFs	Paing et al., 2002, 2006
TP β	Clathrin- and dynamamin-dependent	HEK293, CHO	Dominant negative arrestin (arrestin-3 ²⁰¹⁻⁴⁰⁹)	Parent et al., 2001
US28	Clathrin-dependent	HeLa, MEFs	Arrestin-2/3 ^{-/-} MEFs	Fraile-Ramos et al., 2003
XCR1	Unknown	HEK293	HEK293-Arrestin-2/3	Bauer et al., 2019

Arf6, ADP-ribosylation factor 6; BHK-21, baby hamster kidney fibroblasts; MDCK, Madin-Darby canine kidney; PKA, protein kinase A.

In the absence of agonist, the native receptor cycles between the cell membrane and the intracellular pool. Instead, internalized, activated PAR1 is sorted from endosomes to lysosomes, where it is rapidly degraded (Trejo et al., 1998). Studies using a combination of microscopy, mutant receptors, and pharmacological and genetic inhibitors showed that the two processes are dependent on distinct mechanisms. Constitutive internalization is effected upon mutation of the PAR1 tyrosine motif (Y383A/L386A) and by depletion of AP-2 using small interfering RNA (siRNA), indicating the importance of AP-2 in the process (Paing et al., 2006). On the other hand, agonist-induced internalization is only partially inhibited in AP-2-depleted cells and is reliant on additional sequences in the C-tail, suggesting the involvement of other clathrin adaptors (Paing et al., 2004, 2006; Trejo et al., 2000). Trejo and colleagues proposed that epsin-1 is the other key clathrin adaptor protein for active PAR1 internalization (Chen et al., 2011). Epsin-1-mediated endocytosis requires ubiquitination of PAR1, and the ubiquitin-interacting motifs of epsin-1 are crucial for this pathway. In cells depleted of epsin-1 and/or AP-2 by siRNA, activated PAR1 internalization was impaired, further confirming that both adaptor proteins are required for agonist-induced internalization.

Several other receptors that internalize through a partially or completely arrestin-independent mechanism also contain tyrosine motifs. PAR4 has a YXXL motif in ICL3, and a Y263A/L268A double mutation in the AP-2 binding motif disrupted the ability of the receptor to internalize (Smith et al., 2016). Similarly, depletion of AP-2 and clathrin by siRNA inhibited receptor internalization, thus confirming an AP-2- and clathrin-dependent mechanism. The IP prostanoid receptor has a YXXL motif in ICL2 (Smyth et al., 2000). Microscopy studies showed that the active receptor was present in clathrin-coated vesicles. This process is repressed by dominant negative dynamin (K44A), but it is arrestin- and GRK-independent and is not affected by PKC-mediated receptor phosphorylation. In contrast, the α_{1B} -adrenoceptor contains several YXX Φ and dileucine motifs (Diviani et al., 2003), but AP-2 has not been shown to interact with these motifs. Instead, AP-2 binds to a stretch of eight arginine residues on the receptor C-tail. Arrestins contribute partially to α_{1B} -adrenoceptor endocytosis, demonstrating that multiple pathways can regulate internalization of an individual receptor.

GRK-Dependent Pathway. GRKs are serine/threonine kinases that phosphorylate active GPCRs, thus facilitating arrestin binding to the receptor and inhibiting G protein interactions (Benovic et al., 1986, (Bouvier et al., 1988); Komolov and Benovic, 2018). Seven mammalian GRKs (GRK1–7) that regulate GPCRs have been identified: GRK1 and GRK7 are expressed exclusively in the retina; GRK4 is only found in significant amounts in the testes, whereas GRK2, -3, -5 and -6 are universally expressed (Komolov and Benovic, 2018). In addition to their role in arrestin binding, there are several reports of an internalization pathway that is GRK phosphorylation-dependent but arrestin-independent.

Upon agonist stimulation, the BLT₁ leukotriene receptor internalizes through a GRK2- and dynamin-dependent mechanism without involving arrestins (Chen et al., 2004b). Agonist stimulation of the receptor does not cause arrestin redistribution within the cell, and the receptor does not associate with arrestins. Moreover, receptor internalization is not affected by overexpression of wild-type and dominant

negative arrestin-2 (V53D). The receptor undergoes endocytosis in rat basophilic leukemia (RBL)-2H3 cells that express high levels of endogenous GRK2, but not in human embryonic kidney 293 (HEK293) or COS-7 cells that express 5–10-fold less GRK2 than RBL-2H3 cells. Additionally, the process is blocked by coexpression of a catalytically inactive GRK2 mutant (K220R) and enhanced when wild-type GRK2 is overexpressed, which indicates that it is phosphorylation-dependent. A BLT₁ receptor mutant with truncated C-tail lost the ability to internalize and associate with GRK2 and dominant negative dynamin (K44A) inhibited BLT₁ endocytosis. However, the mechanism of how GRK2-mediated phosphorylation can trigger arrestin-independent endocytosis remains to be clarified.

Formylpeptide receptor 1 (FPR1) has also been suggested to internalize via a GRK phosphorylation-dependent and arrestin-independent pathway (Prossnitz et al., 1995; Hsu et al., 1997; Vines et al., 2003). A phosphorylation-deficient mutant of FPR1 where the serine and threonine residues in the C terminus are converted to alanine and glycine residues was unable to desensitize and internalize, thus indicating that phosphorylation is required for the processes (Hsu et al., 1997). GRK2 and to a lesser degree GRK3 were shown to be the kinases responsible for phosphorylating the FPR1 C-tail (Prossnitz et al., 1995). Although arrestins are colocalized with FPR1 in membranes and endosomes during receptor internalization (Bennett et al., 2000), they are not involved in internalization as determined by the use of dominant negative arrestin-2 (arrestin-2^{S319–418}) and mouse embryonic fibroblasts (MEFs) from arrestin-2/3 knockout mice (Gilbert et al., 2001; Vines et al., 2003). Nonetheless, arrestins have been suggested to play a role in FPR1 recycling to the plasma membrane because in arrestin-2/3 knockout MEFs the receptor accumulated in the perinuclear endosome compartment instead of recycling through an unknown mechanism (Vines et al., 2003). FPR1 internalization is furthermore insensitive to dominant negative mutants of dynamin (K44A) and clathrin (hub region, competes for binding to clathrin light chain) (Gilbert et al., 2001), which suggests that it is internalized through a different mechanism than the BLT₁ receptor.

FEME Pathway. β_1 - and α_{2A} -adrenoceptors, D₃ and D₄ dopamine receptors, and the M₄ muscarinic receptor have been demonstrated to internalize through the FEME pathway (Boucrot et al., 2015) (Fig. 1B). Agonist-induced internalization of these receptors was strongly reduced upon siRNA knockdown of endophilin-A1–3, but not with knockdown of clathrin or AP-2, thus indicating that it is an endocytic route that is independent of clathrin. Correspondingly, the process is not affected by overexpression or siRNA depletion of arrestin-2/3. Dynamin is identified as the main driver of the endophilin-mediated fission because dominant negative dynamin (K44A and K65A) and several small molecule dynamin inhibitors negatively affected the formation of endophilin buds. Using pharmacological inhibitors, this pathway was further shown to depend on cholesterol, actin, Rho GTPase, phosphatidylinositol 3-kinase, and the serine/threonine protein kinase PAK1 (Boucrot et al., 2015).

Caveolae Pathway. Using arrestin-2/3 knockout HEK293 cells, the glucagon-like peptide (GLP)-1 receptor has been shown to internalize independently of arrestins (Jones et al., 2018). Studies applying confocal microscopy showed that the GFP-tagged receptor localizes in membrane lipid rafts and

caveolae (Fig. 1C). As the receptor contains a classic caveolin-1 binding motif (EGVYLYTLLAFSVF) in ICL2, the receptor could interact directly with caveolin-1 and be endocytosed via caveolae. This is further evidenced using dominant negative mutants of caveolin-1 (P132L) and dynamin (K44A) that inhibit GLP-1 receptor endocytosis (Syme et al., 2006). The GLP-2 receptor within the same receptor family can most likely also be desensitized, internalized, and recycled independently of arrestins, since these processes were inert to receptor C-tail truncation (Estall et al., 2004, 2005). GLP-2 receptor endocytosis is inhibited by cholesterol sequestration with filipin or cholesterol depletion with methyl- β -cyclodextrin, thus suggesting that the receptor internalizes via a lipid-raft-mediated pathway. After endocytosis, the GLP-2 receptor colocalizes with caveolin-1 in early endosomes and perinuclear recycling compartments (Estall et al., 2004). However, unlike the GLP-1 receptor, the GLP-2 receptor lacks the classic caveolin-1 binding motif and is internalized independently of dynamin, thus suggesting different internalization pathways for the two receptors.

The ET_A and ET_B endothelin receptors are able to recruit arrestins but unable to promote association with AP-2 upon agonist stimulation, as shown using bioluminescence resonance energy transfer (BRET)-based assays for arrestin-2 and -3/ β 2 adaptin interactions (Hamdan et al., 2007). Furthermore, internalization of the ET_A receptor is not blocked by the arrestin/ β 2 adaptin inhibitor barbadin (Beautrait et al., 2017). However, knockdown of arrestin-2/3 using siRNA showed partial inhibition of ET_A and ET_B receptor endocytosis (Hamdan et al., 2007). Thus, this indicates that these receptors internalize partially through a mechanism that involves a direct interaction between arrestin and clathrin and partially through an arrestin- and AP-2-independent pathway. In fact, around a third of all clathrin-coated pits have been found not to contain AP-2 (Pascolutti et al., 2019), which suggests that AP-2-independent clathrin-mediated endocytosis could be more common than previously anticipated. The ET_A receptor undergoes caveolae-mediated internalization in HEK293 cells (Okamoto et al., 2000), which could account for the arrestin-independent component.

β_2 -adrenoceptors and angiotensin (AT) $_{1A}$ angiotensin receptors are shown to internalize via two distinct pathways, clathrin-mediated endocytosis and the caveolae endocytic route (Guo et al., 2015). Using specific inhibitors for clathrin (dominant negative epsin²⁰⁴⁻⁴⁵⁸) and caveolae (methyl- β -cyclodextrin) in cells depleted of clathrin and caveolin-1 by small hairpin RNA, the authors confirmed that these are different pathways. Moreover, clathrin-mediated endocytosis is mediated by GRKs, but caveolar endocytosis is not dependent on GRKs. The AT_{1A} receptor also showed cell type-specific internalization. In HEK293 and COS-7 cells, receptor internalization was not mediated by dynamin or arrestins, but in CHO cells, it was abolished by hypertonic sucrose, dominant negative arrestins (arrestin-2-V53D and arrestin-2¹⁻³⁴⁹) and dominant negative dynamin K44A, thereby indicating that arrestins are involved in clathrin-mediated AT_{1A} internalization in this cell line (Zhang et al., 1996; Oakley et al., 2000; Gáborik et al., 2001).

Arrestin-Independent Constitutive Endocytosis

GPCRs can also undergo endocytosis in the absence of agonist stimulation. In contrast to agonist-induced endocytosis, the

mechanisms and functions of constitutive endocytosis are less well understood. Similar to active receptor endocytosis, there are examples of GPCRs that are able to internalize constitutively without the need for arrestins (Table 1).

Major Histocompatibility Complex Class I Pathway. β_2 -adrenoceptor and M_3 muscarinic receptors internalize without agonist stimulation and colocalize with major histocompatibility complex class I (MHC-I) on peripheral endosomal structures. MHC-I marks a clathrin-independent endocytic pathway. Their constitutive internalization is also not inhibited by dominant negative dynamin (K44A) and only slightly affected by siRNA depletion of clathrin, suggesting that dynamin and clathrin are not required. Upon agonist stimulation, these receptors switch to a clathrin-dependent trafficking pathway (Scarselli and Donaldson, 2009). The clathrin-independent endocytosis pathway used by MHC-I is believed to be independent of arrestins, but it remains to be confirmed if arrestins play a role in the constitutive internalization of the β_2 -adrenoceptor and the M_3 receptor.

It is well established that the metabotropic glutamate (mGlu) receptors are unable to recruit arrestins (Pin and Bettler, 2016). In the absence of ligand, the mGlu₇ receptor colocalizes with internalized MHC-I in endosomes (Lavezzari and Roche, 2007). It was found to traffic there via an ADP-ribosylation factor 6-positive endosomal pathway that is not regulated by clathrin. The mGlu₅ receptor also internalizes through a clathrin-independent pathway in the absence of receptor activation (Fourgeaud et al., 2003); however, it contains caveolin-1 binding motifs in ICL1 and ICL3 and colocalizes with caveolin-1 in hippocampal neurons, and constitutive mGlu₅ receptor internalization is inhibited by nystatin-mediated sequestration of cholesterol (Francesconi et al., 2009), thus suggesting that mGlu₅ is constitutively internalized via caveolae.

Other Pathways. Other GPCRs also internalize constitutively without the need for arrestins, but their exact mechanisms are not defined. Using ELISA and confocal microscopy, the orphan adhesion receptor ADGRA3 (previously called GPR125) was found to undergo rapid constitutive internalization in an arrestin-independent, but clathrin-dependent manner (Spiess et al., 2019). The internalized receptor colocalized with transferrin receptor 1 in early endosomes. Chemokine receptors CXCR4 and XCR1 and the viral GPCR US28 also show constitutive activity and internalization in the absence of arrestins (Fraile-Ramos et al., 2003; Bauer et al., 2019; Spiess et al., 2019). Like the β_2 -adrenoceptor and the M_3 receptor, constitutive and agonist-induced internalization of CXCR4 occurs through distinct pathways. Constitutive internalization of the receptor is dependent on PKC and dynamin but seems to be independent of arrestins, as internalization was not affected by deleting potential arrestin-2/3 binding sites from the C-tail of CXCR4. However, activated CXCR4 internalizes through the arrestin-mediated pathway (Signoret et al., 1997, 1998). Similarly, the calcium-sensing (CaS) receptor undergoes constitutive internalization through a pathway that is partially arrestin-independent, whereas agonist-mediated internalization was found to be arrestin-2/3-dependent (Mos et al., 2019). Altogether, several GPCRs have been reported to internalize constitutively in an arrestin-independent manner. Although the molecular mechanisms at play during arrestin-independent internalization are often not investigated, these studies show that ligand-stimulated and

constitutive receptor endocytosis can differ in terms of arrestin dependency.

Arrestin Activation without Triggering Internalization. As described above, FPR1 and PAR1 recruit arrestins upon activation but internalize independently of arrestins. Instead, arrestins are proposed to regulate PAR1 desensitization (Paing et al., 2002) and FPR1 recycling (Vines et al., 2003). Most of the receptors that have been reported to internalize independently of arrestins (Table 1) have in fact been shown to recruit arrestin-2 and/or -3 to the plasma membrane in a recent systematic study using a receptor-independent enhanced bystander BRET assay (Avet C et al., preprint, DOI: <https://doi.org/10.1101/2020.04.20.052027>). These inconsistencies could be a result of different cellular backgrounds or experimental conditions, such as the need for overexpression of arrestin in most assays measuring arrestin recruitment. However, it is intriguing to consider the possibility that GPCRs can interact with arrestin without triggering internalization and the potential functional consequences of such an interaction.

Would it be possible for a receptor to interact with arrestin without exposing the AP-2 and clathrin binding motifs in the arrestin C-tail that is bound to the arrestin N-domain in inactive arrestin (Fig. 2A)? Several studies have used intramolecular biosensors to show that arrestins can adopt multiple active conformations with distinct functions (“active arrestin” is defined in the following as a conformation that involves a major conformational change from the inactive state, which exposes functionalities, such as protein and lipid binding sites that were inaccessible in the inactive state). This was first shown with arrestin-2 or -3 biosensors with BRET donors and acceptors fused to each end of arrestin (Shukla et al., 2008; Nobles et al., 2011; Zimmerman et al., 2012). Whereas endogenous agonists for AT_{1A} angiotensin and parathyroid (PTH) 1 parathyroid hormone receptors [angiotensin II and parathyroid hormone (PTH)-1-34, respectively] increased the intramolecular arrestin BRET signal, the arrestin-biased ligands [Sar¹, Ile⁴, Ile⁸]-angiotensin II and

[D-Trp¹², Tyr³⁴]-PTH-(7–34) decreased the BRET signal, thus indicating different arrestin conformations (Shukla et al., 2008). Similarly, when introducing point mutations in the transmembrane segment of the AT_{1A} receptor or the β₂-adrenoceptor that interfere with G protein coupling, thus generating an arrestin-biased receptor, the intramolecular BRET signal upon stimulation with angiotensin II or isoproterenol changed from increasing to decreasing. The intramolecular biosensor approach was later extended by inserting short (six amino acids) binding motifs for fluorescein arsenical hairpin (FlAsH) in different places in arrestin-3 and monitoring either Förster resonance energy transfer with a C-terminal cyan fluorescent protein (Nuber et al., 2016) or BRET with an N-terminal *Renilla* luciferase (Lee et al., 2016). These biosensors confirmed the existence of receptor and ligand specific arrestin conformations. The conformational signatures of arrestin furthermore correlated with receptor trafficking and arrestin-dependent extracellular signal-regulated kinase (ERK) 1/2 phosphorylation patterns (Lee et al., 2016). Although it is tempting to speculate that some of these biosensors could provide evidence for arrestin activation without arrestin C-tail displacement, such inferences are notoriously difficult to make for biosensors. Recently, the biosensor experiments were corroborated by a study using an intracellularly expressed antibody fragment (intrabody) specific for active arrestin-2 (Baidya et al., 2020). The binding of the intrabody to arrestin-2 was triggered by ligand stimulation of the V₂ vasopressin receptor, but not by the B₂ bradykinin receptor, although both recruited arrestin-2 to a similar extent. It is thus possible that some GPCRs can stabilize an arrestin conformation that does not lead to internalization but supports other arrestin functions, although such a conformation has not yet been demonstrated directly.

A possible mechanism for stabilizing different arrestin conformations could be by arrestin interactions with discrete receptor sites. The prevailing model for arrestin activation was for many years a multistep model where arrestin first




Fig. 2. Nonconventional receptor-arrestin interactions that do not lead to arrestin-dependent internalization. (A) Hypothetical interaction between arrestin and the 7TM core of a receptor that stabilizes an intermediate state of arrestin without displacement of the arrestin C-tail from the arrestin N-domain. The clathrin and AP-2 interaction motifs in the arrestin C-tail are still masked, but arrestin blocks G protein activation and could potentially potentiate signaling by scaffolding kinases. This model allows simultaneous interaction of the receptor C-tail with a hypothetical protein (depicted as interactor) that mediates receptor internalization. (B) Catalytic activation of arrestin. Arrestin is activated by a transient interaction with the receptor 7TM core and is stabilized at the membrane in this conformation by binding of PI(4,5)P₂. Activated arrestins accumulate in clathrin-coated structures where they can scaffold kinases to enhance signaling.

recognizes either the phosphorylated C-tail of the receptor or the seven transmembrane (7TM) core of an activated receptor, which constitutes a low-affinity precomplex. Arrestin can then proceed to engage the other binding site if it is present, thus forming the high-affinity complex that is required for arrestin activation (Gurevich and Benovic, 1993). Later studies have, however, revealed that engagement of only one of these sites is sufficient for activation of arrestin (Richardson et al., 2003; Kumari et al., 2016, 2017; Cahill et al., 2017; Latorraca et al., 2018). Several studies have looked at the correlation between the functional consequences of receptor-arrestin interactions and whether arrestin is binding to the receptor 7TM core, the phosphorylated C-tail, or both. For receptors with a C-tail that forms stable complexes with arrestins, such as the V_2 receptor, the core interaction was shown to be important for rapid desensitization of G protein signaling, but the interaction between the phosphorylated receptor C-tail and arrestin was sufficient to mediate internalization and arrestin-dependent ERK phosphorylation (Kumari et al., 2016, 2017; Cahill et al., 2017). This is consistent with PAR1 where G protein desensitization by arrestin is independent of phosphorylation (Chen et al., 2004a). Conversely, a truncation mutant of the substance P receptor that removes all serine and threonine residues from the C-tail desensitized and internalized like the full-length receptor (Richardson et al., 2003). Similarly, a phosphorylation-deficient mutant of the BLT₁ leukotriene receptor was still able to recruit arrestin and internalize, although with a delay compared with the wild-type receptor (Jala et al., 2005). Thus, only desensitization of G protein signaling seems to be specifically linked to interaction with the receptor 7TM core, whereas internalization and enhancement of signaling can be mediated by interaction with either of the two binding sites but seem to be linked as long as arrestin is associated with the receptor.

Displacement of the arrestin C-tail has been shown to enable arrestin to spontaneously undergo conformational changes to a presumably active conformation where the C-domain of arrestin is twisted 20° relative to the N-domain (Latorraca et al., 2018). This is supported by the fact that the naturally occurring p44 splice variant of arrestin-1 that lacks the C-tail has been crystallized in both active and inactive conformations (Granzin et al., 2012; Kim et al., 2013). Allosteric coupling between the two events would suggest that the converse is also true, i.e., that the conformational rearrangement of the N- and C-domains that normally occurs upon receptor binding would lead to C-tail displacement and subsequently receptor endocytosis. However, if multiple active arrestin conformations exist, as suggested by intramolecular biosensor experiments, we speculate that some of these conformations might not displace the arrestin C-tail. Indeed, differential phosphorylation of the receptor C-tail has been suggested to induce different arrestin conformations by selectively interacting with a distinct subset of the key elements that stabilize the inactive state of arrestin (Sente et al., 2018). Interestingly, proximal phosphorylation appears to release the finger loop that is important for interacting with the receptor 7TM core, but possibly not the three-element interaction between α -helix 1 and β -strand 1 of the arrestin N-domain and the arrestin C-tail.

None of the structures of receptor-arrestin complexes published so far has retained autoinhibitory binding of the arrestin C-tail. However, these structures used either phosphorylated

receptor C-tails that are known to bind strongly to arrestin (Shukla et al., 2014; Lee et al., 2020; Staus et al., 2020), truncated the arrestin C-tail (Huang et al., 2020; Staus et al., 2020), or destabilized the arrestin C-tail binding through mutations in the arrestin C-tail (Kang et al., 2015; Yin et al., 2019), thus making it unlikely or impossible to preserve autoinhibitory arrestin C-tail binding. Structures of receptors without a phosphorylated C-tail in complex with full-length arrestin are eagerly awaited to shed light on whether it is possible to retain binding of the arrestin C-tail when bound to a receptor.

An alternative model of arrestin activation termed catalytic activation has been described, where arrestin interacts transiently with the receptor 7TM core but remains bound to the membrane in an active conformation stabilized by phosphoinositide binding (Fig. 2B) (Eichel et al., 2016, 2018; Nuber et al., 2016). When arrestin dissociates from the receptor, it can no longer drive receptor internalization, but it can still traffic to clathrin-coated structures and mediate ERK1/2 phosphorylation from there. This catalytic activation mechanism was found for several receptors that are known to interact transiently with arrestin: the β_1 - and β_2 -adrenoceptors, the D₂ dopamine receptor, and the μ and κ opioid receptors (Eichel et al., 2016, 2018).

In conclusion, arrestins can block G protein activation without triggering endocytosis by binding to the 7TM core of stimulated receptors in a mechanism that most likely does not involve major conformational changes in arrestin. There is evidence from intramolecular arrestin biosensor experiments that a given receptor-ligand combination could stabilize a specific arrestin conformation, but the role of arrestins in FPR1 recycling after arrestin-independent endocytosis remains the only functional or structural evidence of a receptor-arrestin complex that mediates arrestin functions requiring arrestin activation without also driving receptor internalization (Vines et al., 2003). Catalytic activation of arrestin does, however, provide such a mechanism, and it would be interesting to determine if receptors that internalize in an arrestin-independent way accumulate active arrestins at clathrin-coated structures.

Methods for Measuring Endocytosis

It has for many years been technically challenging to measure receptor endocytosis with the same level of sensitivity and robustness as receptor signaling, which made detailed analysis of endocytic pathways challenging. However, there are now sensitive and robust internalization assays that can be combined with imaging for a complete picture of receptor endocytosis.

Imaging. The most direct method to track subcellular distribution of GPCRs is by fluorescence microscopy (Hislop and von Zastrow, 2011; Foster and Bräuner-Osborne, 2018). The receptor usually has to be genetically tagged with an epitope or fluorescent protein or undergo enzyme-directed covalent modification to allow for examination and localization using fluorescence microscopy (Daunt et al., 1997; Hislop and von Zastrow, 2011; Cahill et al., 2017). The method is flexible, widely available, and applicable to most GPCRs. Furthermore, it can be used *in vivo*, by knocking in the tagged receptor in animals, allowing for real-time imaging of receptor trafficking in e.g., neurons (Ehrlich et al., 2019). A potential

drawback is that recombinantly expressed and modified receptors may not exactly mimic the properties of native receptors.

BRET and Time-Resolved Förster Resonance Energy Transfer. BRET and time-resolved Förster resonance energy transfer (TR-FRET) internalization assays have higher throughput than imaging assays but lack the spatial resolution. BRET trafficking assays require heterologous coexpression of fusion proteins with a luciferase variant that can catalyze the generation of donor bioluminescence and a fluorescent protein acceptor to measure proximity between GPCRs and compartment markers in real time (Hamdan et al., 2005, 2006; Pflieger et al., 2007). The acceptor can be anchored to distinct cell compartments by genetically fusing it to targeting sequences for the plasma membrane (e.g., CAAX), the endoplasmic reticulum (e.g., PTP1B), or different stages of endosomes (e.g., Rab5, Rab7, or Rab11) (Pflieger and Eidne, 2006; Lan et al., 2012; (Szakadáti et al., 2015) Namkung et al., 2016; Cahill et al., 2017). When GPCRs relocate to these compartments, bystander BRET will be generated that allows for the study of the complete trafficking cycle (Balla et al., 2012; Cao et al., 2019). However, the specificity of the bystander BRET assay should be taken into consideration, as it relies on the assumption of a homogenous cell population with similar levels of fusion protein expression and requires fusion of a luciferase on the intracellular side of receptors, which could interfere with trafficking.

The TR-FRET internalization assay can also effectively assess the time course of receptor internalization. It requires fusion of an N-terminal SNAP-tag to receptors, which is less likely to interfere with trafficking than an intracellular tag and is applicable to a wide range of GPCRs (Roed et al., 2014; Jacobsen et al., 2017; Foster et al., 2019). The SNAP-tagged receptors are covalently labeled with a cell-impermeant terbium cryptate substrate (SNAP-Lumi4-Tb) that acts as energy donor and ensures that only receptors that are at the cell surface during labeling are tracked. Lanthanide complexes, such as terbium cryptate, have millisecond lifetimes, which makes it possible to introduce a delay between excitation and recording of emission, eliminating short-lived fluorescence background from, for example, cellular autofluorescence (Levoye et al., 2015). After washing to remove excess donor substrate, cells are incubated with a cell-impermeant energy acceptor (e.g., fluorescein-O'-acetic acid). For receptors at the cell surface, energy is transferred from the donor to the acceptor upon excitation of the donor, thus resulting in a low donor-acceptor ratio. Receptor internalization (constitutive or agonist-induced) causes an increased donor-acceptor distance that is incompatible with energy transfer, which leads to an increased donor/acceptor ratio (Foster and Bräuner-Osborne, 2018). The TR-FRET internalization assay uses a synthetic fluorophore in solution as the acceptor, which limits the assay to measuring receptor endocytosis.

ELISA, Flow Cytometry, Immunoblotting, and Radioligand Binding. Other high-throughput GPCR endocytosis assays include biochemical measurements of receptors with N-terminal epitope tags, such as hemagglutinin and Flag tags (Kang et al., 2013; Foster and Bräuner-Osborne, 2018). Examples include ELISA, flow cytometry, and immunoblotting to quantify loss of cell surface GPCR and determine the rate of internalization upon agonist stimulation (Okamoto et al., 2000). For receptors with readily available radioligands,

different protocols of binding experiments can be performed to compare cell surface with total receptor density (Heilker, 2007). However, this is limited to receptors with specific high-affinity radiolabeled ligands. Also, these assays are only able to measure net changes in receptor number, which is why it can be challenging to effectively distinguish between the effects of receptor biosynthesis and degradation, in contrast to the BRET assays that can track receptor quantity changes in different cellular compartments (Hislop and von Zastrow, 2011).

Tools for Testing Arrestin Involvement in Endocytosis. A variety of methods have been used to determine whether receptor endocytosis depends on arrestins, to further examine the molecular machinery involved, and to identify the distinct endocytic routes. Here we present a critical overview of some of the most commonly used tools, including their advantages and possible pitfalls, with focus on probing the involvement of arrestins.

Genetic Approaches. To determine the involvement of arrestins in GPCR desensitization, internalization, and down-regulation, a wide range of pharmacological and genetic tools have been developed. Arguably, the most effective method to study their impact on receptor endocytosis is to genetically deplete endogenous arrestins from the cellular background. Different methods have been used to successfully achieve this, including the use of RNA interference to knockdown arrestins (Ahn et al., 2003; Wei et al., 2003; Shenoy et al., 2006; O'Hayre et al., 2017; Luttrell et al., 2018). However, full knockdown of expression is rarely achieved with this method, which complicates analysis of the results. In contrast, MEFs prepared from arrestin-2/3 knockout mice can be used to perform experiments in the complete absence of arrestins. These MEFs have been used to study the contribution of arrestin to the endocytosis of several GPCRs, including the β_2 -adrenoceptor, the FPR1 formylpeptide receptor, the PAR1 and PAR4 proteinase-activated receptors, and the viral chemokine receptor US28 (Kohout et al., 2001; Paing et al., 2002; Fraile-Ramos et al., 2003; Vines et al., 2003; Smith et al., 2016). More recently, advancement of the CRISPR/CRISPR-associated protein genome editing technology has made specific gene targeting widely available. This led to the generation of a wide range of HEK293 cell lines where genes encoding different proteins have been knocked out, including GPCRs, G proteins, GRKs, and arrestins (Milligan and Inoue, 2018; Møller et al., 2020). The arrestin-2/3 knockout cells have been used to express and study GPCRs such as the β_2 -adrenoceptor and the CaS, gastric inhibitory polypeptide, μ -opioid and V_2 vasopressin receptors and to assess the role of arrestins in receptor endocytosis as well as their function in cell signaling (Gabe et al., 2018; Luttrell et al., 2018; Mos et al., 2019; Møller et al., 2020). The cell lines were validated to be completely deprived of arrestins, and, compared with knockout animals, they are less likely to have altered levels of other network components to compensate for the loss of the deleted genes, as arrestin-2/3 knockout cells did not have significantly altered expression of G proteins (Alvarez-Curto et al., 2016).

Dominant Negative Protein Mutants. Another commonly used group of tools to study GPCR endocytosis is dominant negative mutants. These mutants are able to compete with the wild-type protein for some of the same binding partners but are unable to perform certain key functions. They will effectively inhibit the action of the wild-type protein when

expressed in excess (Sheppard, 1994). For arrestins, there are several dominant negative mutants that effectively inhibit agonist-promoted endocytosis by interfering with the ability of arrestins to bind to GPCRs (arrestin-2-V53D, arrestin-3-V54D) (Ferguson et al., 1996), clathrin (arrestin-2- Δ LIELD), or β 2 adaptin (arrestin-2-F391A) (Kim and Benovic, 2002) or by competing with clathrin binding (arrestin-2^{319–418} and arrestin-3^{284–409}) (Krupnick et al., 1997; Orsini and Benovic, 1998).

Pharmacological Inhibitors. Finally, pharmacological inhibitors can be used to decipher the contribution of their target protein in the receptor endocytosis process. Barbadin is an inhibitor that selectively binds to AP-2 and disrupts the interaction between arrestins and the β 2 adaptin subunit of AP-2 (Beautrait et al., 2017). Other pharmacological inhibitors include pitstop2 that blocks both clathrin-dependent and -independent endocytosis (Dutta et al., 2012), dynasore and dyngo-4a that inhibit dynamin-dependent pathways (Macia et al., 2006; Hill et al., 2009; McCluskey et al., 2013), nystatin and filipin that disrupt caveolae-mediated internalization, and chlorpromazine that blocks internalization via clathrin-coated pits (Okamoto et al., 2000). However, experimental results based on these inhibitors need to be interpreted with care, as the selectivity of these compounds may not be as defined as previously thought (Ivanov, 2008; Park et al., 2013; Guo et al., 2015). Pitstop2, for instance, is still being advertised as a clathrin-selective inhibitor, even though equipotent inhibition of clathrin-independent endocytosis was reported long ago (Dutta et al., 2012).

Conclusions

GPCR endocytosis is often assumed to be an arrestin-mediated process. Here we have presented more than 30 examples of arrestin-independent agonist-induced or constitutive endocytosis (Table 1), which show that GPCR endocytosis is a more diverse process than initially expected. However, for most of the receptors that are shown to internalize independently of arrestins, there is little known about the alternative pathway. The absence of clear discriminators between the pathways has resulted in a lack of tools to specifically study them. Further complicating matters, agonists that selectively activate one endocytosis pathway have been described, for example, for the D₃ dopamine receptor (Xu et al., 2019). Moreover, receptors display cell type-specific endocytosis, indicating that cellular background and molecular makeup of the environment can drive distinct internalization pathways. It is thus necessary for the field to apply the recent technological developments that are outlined in this review, such as high-throughput internalization assays and CRISPR/CRISPR-associated protein genome editing, to delineate the arrestin-independent internalization pathways and thereby expand our understanding of GPCR regulation.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Moo, van Senten, Bräuner-Osborne, Möller.

References

Ahn S, Nelson CD, Garrison TR, Miller WE, and Lefkowitz RJ (2003) Desensitization, internalization, and signaling functions of β -arrestins demonstrated by RNA interference. *Proc Natl Acad Sci USA* **100**:1740–1744.
 Alvarez-Curto E, Inoue A, Jenkins L, Raihan SZ, Prihandoko R, Tobin AB, and Milligan G (2016) Targeted elimination of G proteins and arrestins defines

their specific contributions to both intensity and duration of G protein-coupled receptor signaling. *J Biol Chem* **291**:27147–27159.
 Anderson RGW, Brown MS, and Goldstein JL (1977) Role of the coated endocytic vesicle in the uptake of receptor-bound low density lipoprotein in human fibroblasts. *Cell* **10**:351–364.
 Ariotti N, Rae J, Leneva N, Ferguson C, Loo D, Okano S, Hill MM, Walser P, Collins BM, and Parton RG (2015) Molecular characterization of caveolin-induced membrane curvature. *J Biol Chem* **290**:24875–24890.
 Baidya M, Kumari P, Dwivedi-Agnihotri H, Pandey S, Chaturvedi M, Stepniwski TM, Kawakami K, Cao Y, Laporte SA, Selent J, et al. (2020) Key phosphorylation sites in GPCRs orchestrate the contribution of β -arrestin 1 in ERK1/2 activation. *EMBO Rep* **21**:e49886.
 Balla A, Tóth DJ, Soltész-Katona E, Szakadati G, Erdélyi LS, Várnai P, and Hunyady L (2012) Mapping of the localization of type 1 angiotensin receptor in membrane microdomains using bioluminescence resonance energy transfer-based sensors. *J Biol Chem* **287**:9090–9099.
 Bauer A, Madela J, Berg C, Daugvilaite V, Gurka S, Mages HW, Kroczek RA, Rosenkilde MM, and Voigt S (2019) Rat cytomegalovirus-encoded γ -chemokine vXCL1 is a highly adapted, species-specific agonist for rat XCR1-positive dendritic cells. *J Cell Sci* **133**:jcs236190.
 Beautrait A, Paradis JS, Zimmerman B, Giubilaro J, Nikolajev L, Armando S, Kobayashi H, Yamani L, Namkung Y, Heydenreich FM, et al. (2017) A new inhibitor of the β -arrestin/AP2 endocytic complex reveals interplay between GPCR internalization and signalling. *Nat Commun* **8**:15054.
 Bennett TA, Maestas DC, and Prossnitz ER (2000) Arrestin binding to the G protein-coupled N-formyl peptide receptor is regulated by the conserved “DRY” sequence. *J Biol Chem* **275**:24590–24594.
 Benovic JL, Strasser RH, Caron MG, and Lefkowitz RJ (1986) β -adrenergic receptor kinase: identification of a novel protein kinase that phosphorylates the agonist-occupied form of the receptor. *Proc Natl Acad Sci USA* **83**:2797–2801.
 Bhatnagar A, Willins DL, Gray JA, Woods J, Benovic JL, and Roth BL (2001) The dynamin-dependent, arrestin-independent internalization of 5-hydroxytryptamine 2A (5-HT_{2A}) serotonin receptors reveals differential sorting of arrestins and 5-HT_{2A} receptors during endocytosis. *J Biol Chem* **276**:8269–8277.
 Boucrot E, Ferreira AP, Almeida-Souza L, Debard S, Vallis Y, Howard G, Bertot L, Sauvonnnet N, and McMahon HT (2015) Endophilin marks and controls a clathrin-independent endocytic pathway. *Nature* **517**:460–465.
 Boucrot E, Pick A, Çamdere G, Liska N, Evergren E, McMahon HT, and Kozlov MM (2012) Membrane fission is promoted by insertion of amphipathic helices and is restricted by crescent BAR domains. *Cell* **149**:124–136.
 Bouvier M, Hausdorff WP, De Blasi A, O’Dowd BF, Kobilka BK, Caron MG, and Lefkowitz RJ (1988) Removal of phosphorylation sites from the β -adrenergic receptor delays onset of agonist-promoted desensitization. *Nature* **333**:370–373.
 Cahill TJ III, Thomsen AR, Tarrasch JT, Plouffe B, Nguyen AH, Yang F, Huang LY, Khsai AW, Bassoni DL, Gavino BJ, et al. (2017) Distinct conformations of GPCR- β -arrestin complexes mediate desensitization, signaling, and endocytosis. *Proc Natl Acad Sci USA* **114**:2562–2567.
 Calizo RC and Scarlata S (2012) A role for G-proteins in directing G-protein-coupled receptor-caveolae localization. *Biochemistry* **51**:9513–9523.
 Cao Y, Namkung Y, and Laporte SA (2019) Methods to monitor the trafficking of β -arrestin/G protein-coupled receptor complexes using enhanced bystander BRET. *Methods Mol Biol* **1957**:59–68.
 Chan Wah Hak L, Khan S, Di Meglio I, Law AL, Lucken-Ardjomande Häslar S, Quintaneiro LM, Ferreira APA, Krause M, McMahon HT, and Boucrot E (2018) FBP17 and CIP4 recruit SHIP2 and lamellipodin to prime the plasma membrane for fast endophilin-mediated endocytosis. *Nat Cell Biol* **20**:1023–1031.
 Chen B, Dores MR, Grimsey N, Canto I, Barker BL, and Trejo J (2011) Adaptor protein complex-2 (AP-2) and epsin-1 mediate protease-activated receptor-1 internalization via phosphorylation- and ubiquitination-dependent sorting signals. *J Biol Chem* **286**:40760–40770.
 Chen CH, Paing MM, and Trejo J (2004a) Termination of protease-activated receptor-1 signaling by β -arrestins is independent of receptor phosphorylation. *J Biol Chem* **279**:10020–10031.
 Chen Z, Gaudreau R, Le Gouill C, Rola-Pleszczynski M, and Stanková J (2004b) Agonist-induced internalization of leukotriene B₄ receptor 1 requires G-protein-coupled receptor kinase 2 but not arrestins. *Mol Pharmacol* **66**:377–386.
 Chini B and Parenti M (2004) G-protein coupled receptors in lipid rafts and caveolae: how, when and why do they go there? *J Mol Endocrinol* **32**:325–338.
 Couet J, Li S, Okamoto T, Ikezu T, and Lisanti MP (1997) Identification of peptide and protein ligands for the caveolin-scaffolding domain. Implications for the interaction of caveolin with caveolae-associated proteins. *J Biol Chem* **272**:6525–6533.
 Cremona O, Di Paolo G, Wenk MR, Lüthi A, Kim WT, Takei K, Daniell L, Nemoto Y, Shears SB, Flavell RA, et al. (1999) Essential role of phosphoinositide metabolism in synaptic vesicle recycling. *Cell* **99**:179–188.
 Dannhauser PN and Ungewickell EJ (2012) Reconstitution of clathrin-coated bud and vesicle formation with minimal components. *Nat Cell Biol* **14**:634–639.
 Daunt DA, Hurt C, Hein L, Kallio J, Feng F, and Kobilka BK (1997) Subtype-specific intracellular trafficking of α 2-adrenergic receptors. *Mol Pharmacol* **51**:711–720.
 Delaney KA, Murph MM, Brown LM, and Radhakrishna H (2002) Transfer of M2 muscarinic acetylcholine receptors to clathrin-derived early endosomes following clathrin-independent endocytosis. *J Biol Chem* **277**:33439–33446.
 Diviani D, Lattion AL, Abuin L, Staub O, and Cotecchia S (2003) The adaptor complex 2 directly interacts with the α 1b-adrenergic receptor and plays a role in receptor endocytosis. *J Biol Chem* **278**:19331–19340.
 Dutta D, Williamson CD, Cole NB, and Donaldson JG (2012) Pitstop 2 is a potent inhibitor of clathrin-independent endocytosis. *PLoS One* **7**:e45799.
 Ehrlich AT, Semache M, Gross F, Da Fonte DF, Runtz L, Colley C, Mezni A, Le Gouill C, Lukasheva V, Hogue M, et al. (2019) Biased signaling of the mu opioid receptor revealed in native neurons. *iScience* **14**:47–57.

- Eichel K, Jullié D, Barsi-Rhyne B, Latorraca NR, Masureel M, Sibarita JB, Dror RO, and von Zastrow M (2018) Catalytic activation of β -arrestin by GPCRs. *Nature* **557**: 381–386.
- Eichel K, Jullié D, and von Zastrow M (2016) β -Arrestin drives MAP kinase signaling from clathrin-coated structures after GPCR dissociation. *Nat Cell Biol* **18**: 303–310.
- Estall JL, Koehler JA, Yusta B, and Drucker DJ (2005) The glucagon-like peptide-2 receptor C terminus modulates β -arrestin-2 association but is dispensable for ligand-induced desensitization, endocytosis, and G-protein-dependent effector activation. *J Biol Chem* **280**:22124–22134.
- Estall JL, Yusta B, and Drucker DJ (2004) Lipid raft-dependent glucagon-like peptide-2 receptor trafficking occurs independently of agonist-induced desensitization. *Mol Biol Cell* **15**:3673–3687.
- Ferguson SS (2001) Evolving concepts in G protein-coupled receptor endocytosis: the role in receptor desensitization and signaling. *Pharmacol Rev* **53**: 1–24.
- Ferguson SSG, Downey WE III, Colapietro A-M, Barak LS, Ménard L, and Caron MG (1996) Role of β -arrestin in mediating agonist-promoted G protein-coupled receptor internalization. *Science* **271**:363–366.
- Foster SR and Bräuner-Osborne H (2018) Investigating internalization and intracellular trafficking of GPCRs: new techniques and real-time experimental approaches. *Handb Exp Pharmacol* **245**:41–61.
- Foster SR, Hauser AS, Vedel L, Strachan RT, Huang XP, Gavin AC, Shah SD, Nayak AP, Haugaard-Kedström LM, Penn RB, et al. (2019) Discovery of human signaling systems: pairing peptides to G protein-coupled receptors. *Cell* **179**:895–908.e21.
- Fotin A, Cheng Y, Sliz P, Grigorieff N, Harrison SC, Kirchhausen T, and Walz T (2004) Molecular model for a complete clathrin lattice from electron cryomicroscopy. *Nature* **432**:573–579.
- Fourgeaud L, Bessis AS, Rossignol F, Pin JP, Olivo-Marin JC, and Hémar A (2003) The metabotropic glutamate receptor mGluR5 is endocytosed by a clathrin-independent pathway. *J Biol Chem* **278**:12222–12230.
- Fraille-Ramos A, Kohout TA, Waldhoer M, and Marsh M (2003) Endocytosis of the viral chemokine receptor US28 does not require β -arrestins but is dependent on the clathrin-mediated pathway. *Traffic* **4**:243–253.
- Francesconi A, Kumari R, and Zukin RS (2009) Regulation of group I metabotropic glutamate receptor trafficking and signaling by the caveolar/lipid raft pathway. *J Neurosci* **29**:3590–3602.
- Gabe MBN, Sparre-Ulrich AH, Pedersen MF, Gasbjerg LS, Inoue A, Bräuner-Osborne H, Hartmann B, and Rosenkilde MM (2018) Human GIP(3–30)NH₂ inhibits G protein-dependent as well as G protein-independent signaling and is selective for the GIP receptor with high-affinity binding to primate but not rodent GIP receptors. *Biochem Pharmacol* **150**:97–107.
- Gáborik Z, Szaszák M, Szidonya L, Balla B, Paku S, Catt KJ, Clark AJL, and Hunyady L (2001) β -arrestin- and dynamin-dependent endocytosis of the AT1 angiotensin receptor. *Mol Pharmacol* **59**:239–247.
- Giebing G, Tölle M, Jürgensen J, Eichhorst J, Furkert J, Beyermann M, Neuschäfer-Rube F, Rosenthal W, Zidek W, van der Giet M, et al. (2005) Arrestin-independent internalization and recycling of the uterotensin receptor contribute to long-lasting uterotensin II-mediated vasoconstriction. *Circ Res* **97**:707–715.
- Gilbert TL, Bennett TA, Maestas DC, Cimino DF, and Prossnitz ER (2001) Internalization of the human N-formyl peptide and C5a chemoattractant receptors occurs via clathrin-independent mechanisms. *Biochemistry* **40**:3467–3475.
- Goodman OB Jr, Krupnick JG, Santini F, Gurevich VV, Penn RB, Gagnon AW, Keen JH, and Benovic JL (1996) β -arrestin acts as a clathrin adaptor in endocytosis of the β_2 -adrenergic receptor. *Nature* **383**:447–450.
- Granzin J, Cousin A, Weirauch M, Schlessinger R, Büldt G, and Batra-Safferling R (2012) Crystal structure of p44, a constitutively active splice variant of visual arrestin. *J Mol Biol* **416**:611–618.
- Gray JA, Sheffler DJ, Bhatnagar A, Woods JA, Hufeisen SJ, Benovic JL, and Roth BL (2001) Cell-type specific effects of endocytosis inhibitors on 5-hydroxytryptamine(2A) receptor desensitization and resensitization reveal an arrestin-, GRK2-, and GRK5-independent mode of regulation in human embryonic kidney 293 cells. *Mol Pharmacol* **60**:1020–1030.
- Guo S, Zhang X, Zheng M, Zhang X, Min C, Wang Z, Cheon SH, Oak MH, Nah SY, and Kim KM (2015) Selectivity of commonly used inhibitors of clathrin-mediated and caveolae-dependent endocytosis of G protein-coupled receptors. *Biochim Biophys Acta* **1848** (10 Pt A):2101–2110.
- Gurevich VV and Benovic JL (1993) Visual arrestin interaction with rhodopsin. Sequential multisite binding ensures strict selectivity toward light-activated phosphorylated rhodopsin. *J Biol Chem* **268**:11628–11638.
- Gurevich VV and Gurevich EV (2019) GPCR signaling regulation: the role of GRKs and arrestins. *Front Pharmacol* **10**:125.
- Hamdan FF, Audet M, Garneau P, Pelletier J, and Bouvier M (2005) High-throughput screening of G protein-coupled receptor antagonists using a bioluminescence resonance energy transfer 1-based β -arrestin2 recruitment assay. *J Biomol Screen* **10**:463–475.
- Hamdan FF, Percherancier Y, Breton B, and Bouvier M (2006) Monitoring protein-protein interactions in living cells by bioluminescence resonance energy transfer (BRET). *Curr Protoc Neurosci* **Chapter 5**:Unit 5.23.
- Hamdan FF, Rochdi MD, Breton B, Fessard D, Michaud DE, Charest PG, Laporte SA, and Bouvier M (2007) Unraveling G protein-coupled receptor endocytosis pathways using real-time monitoring of agonist-promoted interaction between β -arrestins and AP-2. *J Biol Chem* **282**:29089–29100.
- Hansen CG, Howard G, and Nichols BJ (2011) Paclins 2 is recruited to caveolae and functions in caveolar biogenesis. *J Cell Sci* **124**:2777–2785.
- Hauser AS, Attwood MM, Rask-Andersen M, Schiöth HB, and Gloriam DE (2017) Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov* **16**:829–842.
- Hayer A, Stoerber M, Ritz D, Engel S, Meyer HH, and Helenius A (2010) Caveolin-1 is ubiquitinated and targeted to intraluminal vesicles in endolysosomes for degradation. *J Cell Biol* **191**:615–629.
- Heilker R (2007) High content screening to monitor G protein-coupled receptor internalisation. *Ernst Schering Found Symp Proc* (2):229–247.
- Hilger D, Masureel M, and Kobilka BK (2018) Structure and dynamics of GPCR signaling complexes. *Nat Struct Mol Biol* **25**:4–12.
- Hill TA, Gordon CP, McGeachie AB, Venn-Brown B, Odell LR, Chau N, Quan A, Mariana A, Sakoff JA, Chircop M, et al. (2009) Inhibition of dynamin mediated endocytosis by the dynoles-synthesis and functional activity of a family of indoles. *J Med Chem* **52**:3762–3773.
- Hirsch JA, Schubert C, Gurevich VV, and Sigler PB (1999) The 2.8 Å crystal structure of visual arrestin: a model for arrestin's regulation. *Cell* **97**:257–269.
- Hislop JN and von Zastrow M (2011) Analysis of GPCR localization and trafficking. *Methods Mol Biol* **746**:425–440.
- Hsu MH, Chiang SC, Ye RD, and Prossnitz ER (1997) Phosphorylation of the N-formyl peptide receptor is required for receptor internalization but not chemotaxis. *J Biol Chem* **272**:29426–29429.
- Huang W, Masureel M, Qu Q, Janetzko J, Inoue A, Kato HE, Robertson MJ, Nguyen KC, Glenn JS, Skiniotis G, et al. (2020) Structure of the neurotensin receptor 1 in complex with β -arrestin 1. *Nature* **579**:303–308.
- Ivanov AI (2008) Pharmacological inhibition of endocytic pathways: is it specific enough to be useful? *Methods Mol Biol* **440**:15–33.
- Jacobsen SE, Ammendrup-Johnsen I, Jansen AM, Gether U, Madsen KL, and Bräuner-Osborne H (2017) The GPRC6A receptor displays constitutive internalization and sorting to the slow recycling pathway. *J Biol Chem* **292**: 6910–6926.
- Jala VR, Shao WH, and Haribabu B (2005) Phosphorylation-independent β -arrestin translocation and internalization of leukotriene B₄ receptors. *J Biol Chem* **280**: 4880–4887.
- January B, Seibold A, Whaley B, Hipkin RW, Lin D, Schonbrunn A, Barber R, and Clark RB (1997) β_2 -adrenergic receptor desensitization, internalization, and phosphorylation in response to full and partial agonists. *J Biol Chem* **272**: 23871–23879.
- Jones B, Buenaventura T, Kanda N, Chabosseau P, Owen BM, Scott R, Goldin R, Angkathunyakul N, Corréa IR Jr, Bosco D, et al. (2018) Targeting GLP-1 receptor trafficking to improve agonist efficacy. *Nat Commun* **9**:1602.
- Kang DS, Kern RC, Puthenveedu MA, von Zastrow M, Williams JC, and Benovic JL (2009) Structure of an arrestin2-clathrin complex reveals a novel clathrin binding domain that modulates receptor trafficking. *J Biol Chem* **284**:29860–29872.
- Kang DS, Tian X, and Benovic JL (2013) β -Arrestins and G protein-coupled receptor trafficking. *Methods Enzymol* **521**:91–108.
- Kang Y, Zhou XE, Gao X, He Y, Liu W, Ishchenko A, Barty A, White TA, Yefanov O, Han GW, et al. (2015) Crystal structure of rhodopsin bound to arrestin by femto-second X-ray laser. *Nature* **523**:561–567.
- Kelly BT, Graham SC, Liska N, Dannhauser PN, Höning S, Ungewickell EJ, and Owen JD (2014) Clathrin adaptors. AP2 controls clathrin polymerization with a membrane-activated switch. *Science* **345**:459–463.
- Kim YJ, Hofmann KP, Ernst OP, Scheerer P, Choe HW, and Sommer ME (2013) Crystal structure of pre-activated arrestin p44. *Nature* **497**:142–146.
- Kim YM and Benovic JL (2002) Differential roles of arrestin-2 interaction with clathrin and adaptor protein 2 in G protein-coupled receptor trafficking. *J Biol Chem* **277**:30760–30768.
- Koch D, Spiwoaks-Becker I, Sabanov V, Sinning A, Dugladze T, Stellmacher A, Ahuja R, Grimm J, Schüller S, Müller A, et al. (2011) Proper synaptic vesicle formation and neuronal network activity critically rely on synapdin I. *EMBO J* **30**: 4955–4969.
- Kohout TA, Lin FS, Perry SJ, Conner DA, and Lefkowitz RJ (2001) β -Arrestin 1 and 2 differentially regulate heptahelical receptor signaling and trafficking. *Proc Natl Acad Sci USA* **98**:1601–1606.
- Komolov KE and Benovic JL (2018) G protein-coupled receptor kinases: past, present and future. *Cell Signal* **41**:17–24.
- Kovtun O, Tilla VA, Jung W, Leneva N, Ariotti N, Chaudhary N, Mandyam RA, Ferguson C, Morgan GP, Johnston WA, et al. (2014) Structural insights into the organization of the cavin membrane coat complex. *Dev Cell* **31**:405–419.
- Krupnick JG, Santini F, Gagnon AW, Keen JH, and Benovic JL (1997) Modulation of the arrestin-clathrin interaction in cells. Characterization of β -arrestin dominant-negative mutants. *J Biol Chem* **272**:32507–32512.
- Kumari P, Srivastava A, Banerjee R, Ghosh E, Gupta P, Ranjan R, Chen X, Gupta B, Gupta C, Jaiman D, et al. (2016) Functional competence of a partially engaged GPCR- β -arrestin complex. *Nat Commun* **7**:13416.
- Kumari P, Srivastava A, Ghosh E, Ranjan R, Dogra S, Yadav PN, and Shukla AK (2017) Core engagement with β -arrestin is dispensable for agonist-induced vasopressin receptor endocytosis and ERK activation. *Mol Biol Cell* **28**: 1003–1010.
- Lan TH, Liu Q, Li C, Wu G, and Lambert NA (2012) Sensitive and high resolution localization and tracking of membrane proteins in live cells with BRET. *Traffic* **13**: 1450–1456.
- Laporte SA, Oakley RH, Holt JA, Barak LS, and Caron MG (2000) The interaction of β -arrestin with the AP-2 adaptor is required for the clustering of β_2 -adrenergic receptor into clathrin-coated pits. *J Biol Chem* **275**:23120–23126.
- Latorraca NR, Wang JK, Bauer B, Townshend RJL, Hollingsworth SA, Olivieri JE, Xu HE, Sommer ME, and Dror RO (2018) Molecular mechanism of GPCR-mediated arrestin activation. *Nature* **557**:452–456.
- Lavezzari G and Roche KW (2007) Constitutive endocytosis of the metabotropic glutamate receptor mGluR7 is clathrin-independent. *Neuropharmacology* **52**: 100–107.
- Lee KB, Pals-Rylaarsdam R, Benovic JL, and Hosey MM (1998) Arrestin-independent internalization of the m1, m3, and m4 subtypes of muscarinic cholinergic receptors. *J Biol Chem* **273**:12967–12972.
- Lee MH, Appleton KM, Strungs EG, Kwon JY, Morinelli TA, Peterson YK, Laporte SA, and Luttrell LM (2016) The conformational signature of β -arrestin2 predicts its trafficking and signalling functions. *Nature* **531**:665–668.

- Lee Y, Warne T, Nehmé R, Pandey S, Dwivedi-Agnihotri H, Chaturvedi M, Edwards PC, García-Nafria J, Leslie AGW, Shukla AK, et al. (2020) Molecular basis of β -arrestin coupling to formoterol-bound β_1 -adrenoceptor. *Nature* **583**:862–866.
- Lefkowitz RJ (2013) A brief history of G-protein coupled receptors (Nobel Lecture). *Angew Chem Int Ed Engl* **52**:6366–6378.
- Levoye A, Zwier JM, Jaracz-Ros A, Klipfel L, Cottet M, Maurel D, Bdioui S, Bala-banay K, Prézéau L, Trinquet E, et al. (2015) A broad G protein-coupled receptor internalization assay that combines SNAP-tag labeling, diffusion-enhanced resonance energy transfer, and a highly emissive terbium cryptate. *Front Endocrinol (Lausanne)* **6**:167.
- Lobingier BT and von Zastrow M (2019) When trafficking and signaling mix: how subcellular location shapes G protein-coupled receptor activation of heterotrimeric G proteins. *Traffic* **20**:130–136.
- Ludwig A, Howard G, Mendoza-Topaz C, Deerinck T, Mackey M, Sandin S, Ellisman MH, and Nichols BJ (2013) Molecular composition and ultrastructure of the caveolar coat complex. *PLoS Biol* **11**:e1001640.
- Luttrell LM, Wang J, Plouffe B, Smith JS, Yamani L, Kaur S, Jean-Charles P-Y, Gauthier C, Lee MH, Pani B, et al. (2018) Manifold roles of β -arrestins in GPCR signaling elucidated with siRNA and CRISPR/Cas9. *Sci Signal* **11**:eaat7650.
- Macia E, Ehrlich M, Massol R, Boucrot E, Brunner C, and Kirchhausen T (2006) Dynasore, a cell-permeable inhibitor of dynamin. *Dev Cell* **10**:839–850.
- McCluskey A, Daniel JA, Hadzic G, Chau N, Clayton EL, Mariana A, Whiting A, Gorgani NN, Lloyd J, Quan A, et al. (2013) Building a better dynasore: the dyngo compounds potentially inhibit dynamin and endocytosis. *Traffic* **14**:1272–1289.
- Milligan G and Inoue A (2018) Genome editing provides new insights into receptor-controlled signalling pathways. *Trends Pharmacol Sci* **39**:481–493.
- Möller TC, Pedersen MF, van Senten JR, Seiersen SD, Mathiesen JM, Bouvier M, and Bräuner-Osborne H (2020) Dissecting the roles of GRK2 and GRK3 in μ -opioid receptor internalization and β -arrestin2 recruitment using CRISPR/Cas9-edited HEK293 cells. *Sci Rep* **10**:17395.
- Mos I, Jacobsen SE, Foster SR, and Bräuner-Osborne H (2019) Calcium-sensing receptor internalization is β -arrestin-dependent and modulated by allosteric ligands. *Mol Pharmacol* **96**:463–474.
- Mundell SJ, Orsini MJ, and Benovic JL (2002) Characterization of arrestin expression and function. *Methods Enzymol* **343**:600–611.
- Namkung Y, Le Gouill C, Lukashova V, Kobayashi H, Hogue M, Khoury E, Song M, Bouvier M, and Laporte SA (2016) Monitoring G protein-coupled receptor and β -arrestin trafficking in live cells using enhanced bystander BRET. *Nat Commun* **7**:12178.
- Nobles KN, Guan Z, Xiao K, Oas TG, and Lefkowitz RJ (2007) The active conformation of β -arrestin1: direct evidence for the phosphate sensor in the N-domain and conformational differences in the active states of β -arrestins1 and -2. *J Biol Chem* **282**:21370–21381.
- Nobles KN, Xiao K, Ahn S, Shukla AK, Lam CM, Rajagopal S, Strachan RT, Huang TY, Bressler EA, Hara MR, et al. (2011) Distinct phosphorylation sites on the β_2 -adrenergic receptor establish a barcode that encodes differential functions of β -arrestin. *Sci Signal* **4**:ra51.
- Nuber S, Zabel U, Lorenz K, Nuber A, Milligan G, Tobin AB, Lohse MJ, and Hoffmann C (2016) β -Arrestin biosensors reveal a rapid, receptor-dependent activation/deactivation cycle. *Nature* **531**:661–664.
- Oakley RH, Laporte SA, Holt JA, Caron MG, and Barak LS (2000) Differential affinities of visual arrestin, β arrestin1, and β arrestin2 for G protein-coupled receptors delineate two major classes of receptors. *J Biol Chem* **275**:17201–17210.
- O'Hayre M, Eichel K, Avino S, Zhao X, Steffen DJ, Feng X, Kawakami K, Aoki J, Messer K, Sunahara R, et al. (2017) Genetic evidence that β -arrestins are dispensable for the initiation of β_2 -adrenergic receptor signaling to ERK. *Sci Signal* **10**:eaal3395.
- Ohno H, Stewart J, Fournier M-C, Bosshart H, Rhee I, Miyatake S, Saito T, Gallusser A, Kirchhausen T, and Bonifacino JS (1995) Interaction of tyrosine-based sorting signals with clathrin-associated proteins. *Science* **269**:1872–1875.
- Okamoto Y, Ninomiya H, Miwa S, and Masaki T (2000) Cholesterol oxidation switches the internalization pathway of endothelin receptor type A from caveolae to clathrin-coated pits in Chinese hamster ovary cells. *J Biol Chem* **275**:6439–6446.
- Orsini MJ and Benovic JL (1998) Characterization of dominant negative arrestins that inhibit β_2 -adrenergic receptor internalization by distinct mechanisms. *J Biol Chem* **273**:34616–34622.
- Ortengren U, Karlsson M, Blazic N, Blomqvist M, Nystrom FH, Gustavsson J, Fredman P, and Strålfors P (2004) Lipids and glycosphingolipids in caveolae and surrounding plasma membrane of primary rat adipocytes. *Eur J Biochem* **271**:2028–2036.
- Paing MM, Johnston CA, Siderovski DP, and Trejo J (2006) Clathrin adaptor AP2 regulates thrombin receptor constitutive internalization and endothelial cell resensitization. *Mol Cell Biol* **26**:3231–3242.
- Paing MM, Stutts AB, Kohout TA, Lefkowitz RJ, and Trejo J (2002) β -Arrestins regulate protease-activated receptor-1 desensitization but not internalization or down-regulation. *J Biol Chem* **277**:1292–1300.
- Paing MM, Temple BR, and Trejo J (2004) A tyrosine-based sorting signal regulates intracellular trafficking of protease-activated receptor-1: multiple regulatory mechanisms for agonist-induced G protein-coupled receptor internalization. *J Biol Chem* **279**:21938–21947.
- Pals-Rylaarsdam R, Gurevich VV, Lee KB, Ptasiński JA, Benovic JL, and Hosey MM (1997) Internalization of the m2 muscarinic acetylcholine receptor. Arrestin-independent and -dependent pathways. *J Biol Chem* **272**:23682–23689.
- Pandey KN (2010) Small peptide recognition sequence for intracellular sorting. *Curr Opin Biotechnol* **21**:611–620.
- Parent JL, Labrecque P, Driss Rochdi M, and Benovic JL (2001) Role of the differentially spliced carboxyl terminus in thromboxane A₂ receptor trafficking: identification of a distinct motif for tonic internalization. *J Biol Chem* **276**:7079–7085.
- Park RJ, Shen H, Liu L, Liu X, Ferguson SM, and De Camilli P (2013) Dynamin triple knockout cells reveal off target effects of commonly used dynamin inhibitors. *J Cell Sci* **126**:5305–5312.
- Pascolutti R, Algisì V, Conte A, Raimondi A, Pasham M, Upadhyayula S, Gaudin R, Maritzen T, Barbieri E, Caldieri G, et al. (2019) Molecularly distinct clathrin-coated pits differentially impact EGFR fate and signaling. *Cell Rep* **27**:3049–3061.e6.
- Pavlos NJ and Friedman PA (2017) GPCR signaling and trafficking: the long and short of it. *Trends Endocrinol Metab* **28**:213–226.
- Pearse BMF (1976) Clathrin: a unique protein associated with intracellular transfer of membrane by coated vesicles. *Proc Natl Acad Sci USA* **73**:1255–1259.
- Peterson YK and Luttrell LM (2017) The diverse roles of arrestin scaffolds in G protein-coupled receptor signaling. *Pharmacol Rev* **69**:256–297.
- Pfleger KD and Eidne KA (2006) Illuminating insights into protein-protein interactions using bioluminescence resonance energy transfer (BRET). *Nat Methods* **3**:165–174.
- Pfleger KD, Dalrymple MB, Dromey JR, and Eidne KA (2007) Monitoring interactions between G-protein-coupled receptors and β -arrestins. *Biochem Soc Trans* **35**:764–766.
- Pin JP and Bettler B (2016) Organization and functions of mGlu and GABA_B receptor complexes. *Nature* **540**:60–68.
- Pope GR, Tilve S, McArdle CA, Lolait SJ, and O'Carroll AM (2016) Agonist-induced internalization and desensitization of the apelin receptor. *Mol Cell Endocrinol* **437**:108–119.
- Prossnitz ER, Kim CM, Benovic JL, and Ye RD (1995) Phosphorylation of the N-formyl peptide receptor carboxyl terminus by the G protein-coupled receptor kinase, GRK2. *J Biol Chem* **270**:1130–1137.
- Renard HF, Simunovic M, Lemière J, Boucrot E, Garcia-Castillo MD, Arumugam S, Chambon V, Lamaze C, Wunder C, Kenworthy AK, et al. (2015) Endophilin-A2 functions in membrane scission in clathrin-independent endocytosis. *Nature* **517**:493–496.
- Richardson MD, Balius AM, Yamaguchi K, Freilich ER, Barak LS, and Kwatra MM (2003) Human substance P receptor lacking the C-terminal domain remains competent to desensitize and internalize. *J Neurochem* **84**:854–863.
- Roed SN, Wismann P, Underwood CR, Kulahin N, Iversen H, Cappelen KA, Schäffer L, Lehtonen J, Hecksher-Soerensen J, Secher A, et al. (2014) Real-time trafficking and signaling of the glucagon-like peptide-1 receptor. *Mol Cell Endocrinol* **382**:938–949.
- Sandvig K, Kavaliauskienė S, and Skotland T (2018) Clathrin-independent endocytosis: an increasing degree of complexity. *Histochem Cell Biol* **150**:107–118.
- Scarselli M and Donaldson JG (2009) Constitutive internalization of G protein-coupled receptors and G proteins via clathrin-independent endocytosis. *J Biol Chem* **284**:3577–3585.
- Schlossman DM, Schmid SL, Braell WA, and Rothman JE (1984) An enzyme that removes clathrin coats: purification of an uncoating ATPase. *J Cell Biol* **99**:723–733.
- Schmid EM, Ford MG, Burtay A, Praefcke GJ, Peak-Chew SY, Mills IG, Benmerah A, and McMahon HT (2006) Role of the AP2 β -appendage hub in recruiting partners for clathrin-coated vesicle assembly. *PLoS Biol* **4**:e262.
- Sengupta P, Philip F, and Scarlata S (2008) Caveolin-1 alters Ca²⁺ signal duration through specific interaction with the G α_q family of G proteins. *J Cell Sci* **121**:1363–1372.
- Sente A, Peer R, Srivastava A, Baidya M, Lesk AM, Balaji S, Shukla AK, Babu MM, and Flock T (2018) Molecular mechanism of modulating arrestin conformation by GPCR phosphorylation. *Nat Struct Mol Biol* **25**:538–545.
- Shapiro MJ, Trejo J, Zeng D, and Coughlin SR (1996) Role of the thrombin receptor's cytoplasmic tail in intracellular trafficking. Distinct determinants for agonist-triggered versus tonic internalization and intracellular localization. *J Biol Chem* **271**:32874–32880.
- Shenoy SK, Drake MT, Nelson CD, Houtz DA, Xiao K, Madabushi S, Reiter E, Pre-mont RT, Lichtarge O, and Lefkowitz RJ (2006) β -arrestin-dependent, G protein-independent ERK1/2 activation by the β_2 adrenergic receptor. *J Biol Chem* **281**:1261–1273.
- Sheppard D (1994) Dominant negative mutants: tools for the study of protein function in vitro and in vivo. *Am J Respir Cell Mol Biol* **11**:1–6.
- Shmuel M, Nodel-Berner E, Hyman T, Rouvinski A, and Altschuler Y (2007) Caveolin 2 regulates endocytosis and trafficking of the M1 muscarinic receptor in MDCK epithelial cells. *Mol Biol Cell* **18**:1570–1585.
- Shukla AK, Violin JD, Whalen EJ, Gesty-Palmer D, Shenoy SK, and Lefkowitz RJ (2008) Distinct conformational changes in β -arrestin report biased agonism at seven-transmembrane receptors. *Proc Natl Acad Sci USA* **105**:9988–9993.
- Shukla AK, Westfield GH, Xiao K, Reis RI, Huang LY, Tripathi-Shukla P, Qian J, Li S, Blanc A, Oleskie AN, et al. (2014) Visualization of arrestin recruitment by a G-protein-coupled receptor. *Nature* **512**:218–222.
- Shvets E, Bitsikas V, Howard G, Hansen CG, and Nichols BJ (2015) Dynamic caveolae exclude bulk membrane proteins and are required for sorting of excess glycosphingolipids. *Nat Commun* **6**:6867.
- Signoret N, Oldridge J, Pelchen-Matthews A, Klasse PJ, Tran T, Brass LF, Rosenkilde MM, Schwartz TW, Holmes W, Dallas W, et al. (1997) Phorbol esters and SDF-1 induce rapid endocytosis and down modulation of the chemokine receptor CXCR4. *J Cell Biol* **139**:651–664.
- Signoret N, Rosenkilde MM, Klasse PJ, Schwartz TW, Malim MH, Hoxie JA, and Marsh M (1998) Differential regulation of CXCR4 and CCR5 endocytosis. *J Cell Sci* **111**:2819–2830.
- Sinha B, Köster D, Ruez R, Gonnord P, Bastiani M, Abankwa D, Stan RV, Butler-Browne G, Vedie B, Johannes L, et al. (2011) Cells respond to mechanical stress by rapid disassembly of caveolae. *Cell* **144**:402–413.
- Smith TH, Coronel LJ, Li JG, Dores MR, Nieman MT, and Trejo J (2016) Protease-activated receptor-4 signaling and trafficking is regulated by the clathrin adaptor protein complex-2 independent of β -arrestins. *J Biol Chem* **291**:18453–18464.

- Smyth EM, Austin SC, Reilly MP, and FitzGerald GA (2000) Internalization and sequestration of the human prostacyclin receptor. *J Biol Chem* **275**: 32037–32045.
- Spiess K, Bagger SO, Torz LJ, Jensen KHR, Walsler AL, Kvam JM, Møgelmoose AK, Daugvilaite V, Junnila RK, Hjortø GM, et al. (2019) Arrestin-independent constitutive endocytosis of GPR125/ADGRA3. *Ann N Y Acad Sci* **1456**: 186–199.
- Staus DP, Hu H, Robertson MJ, Kleinhenz ALW, Wingler LM, Capel WD, Latorraca NR, Lefkowitz RJ, and Skiniotis G (2020) Structure of the M2 muscarinic receptor- β -arrestin complex in a lipid nanodisc. *Nature* **579**:297–302.
- Sundborger A, Soderblom C, Vorontsova O, Evergren E, Hinshaw JE, and Shupliakov O (2011) An endophilin-dynamin complex promotes budding of clathrin-coated vesicles during synaptic vesicle recycling. *J Cell Sci* **124**:133–143.
- Syme CA, Zhang L, and Bisello A (2006) Caveolin-1 regulates cellular trafficking and function of the glucagon-like peptide 1 receptor. *Mol Endocrinol* **20**:3400–3411.
- Szakadati G, Tóth AD, Oláh I, Erdélyi LS, Balla T, Várnai P, Hunyady L, and Balla A (2015) Investigation of the fate of type I angiotensin receptor after biased activation. *Mol Pharmacol* **87**:972–981.
- Tang Y, Hu LA, Miller WE, Ringstad N, Hall RA, Pitcher JA, DeCamilli P, and Lefkowitz RJ (1999) Identification of the endophilins (SH3p4/p8/p13) as novel binding partners for the β 1-adrenergic receptor. *Proc Natl Acad Sci USA* **96**: 12559–12564.
- Thorn H, Stenkula KG, Karlsson M, Ortegren U, Nystrom FH, Gustavsson J, and Stralfors P (2003) Cell surface orifices of caveolae and localization of caveolin to the necks of caveolae in adipocytes. *Mol Biol Cell* **14**:3967–3976.
- Tian X, Kang DS, and Benovic JL (2014) β -arrestins and G protein-coupled receptor trafficking. *Handb Exp Pharmacol* **219**:173–186.
- Traub LM (2009) Tickets to ride: selecting cargo for clathrin-regulated internalization. *Nat Rev Mol Cell Biol* **10**:583–596.
- Traub LM (2011) Regarding the amazing choreography of clathrin coats. *PLoS Biol* **9**: e1001037.
- Trejo J, Altschuler Y, Fu HW, Mostov KE, and Coughlin SR (2000) Protease-activated receptor-1 down-regulation: a mutant HeLa cell line suggests novel requirements for PAR1 phosphorylation and recruitment to clathrin-coated pits. *J Biol Chem* **275**:31255–31265.
- Trejo J, Hammes SR, and Coughlin SR (1998) Termination of signaling by protease-activated receptor-1 is linked to lysosomal sorting. *Proc Natl Acad Sci USA* **95**: 13698–13702.
- Vehlow A, Soong D, Vizcay-Barrena G, Bodo C, Law AL, Perera U, and Krause M (2013) Endophilin, Lamellipodin, and Mena cooperate to regulate F-actin-dependent EGF-receptor endocytosis. *EMBO J* **32**:2722–2734.
- Vines CM, Revankar CM, Maestas DC, LaRusch LL, Cimino DF, Kohout TA, Lefkowitz RJ, and Prossnitz ER (2003) N-formyl peptide receptors internalize but do not recycle in the absence of arrestins. *J Biol Chem* **278**:41581–41584.
- Vögler O, Bogatkewitsch GS, Wriske C, Krummenerl P, Jakobs KH, and van Koppen CJ (1998) Receptor subtype-specific regulation of muscarinic acetylcholine receptor sequestration by dynamin. Distinct sequestration of m2 receptors. *J Biol Chem* **273**:12155–12160.
- Walker JKL, Premont RT, Barak LS, Caron MG, and Shetzline MA (1999) Properties of secretin receptor internalization differ from those of the β 2-adrenergic receptor. *J Biol Chem* **274**:31515–31523.
- Wei H, Ahn S, Shenoy SK, Karnik SS, Hunyady L, Luttrell LM, and Lefkowitz RJ (2003) Independent β -arrestin 2 and G protein-mediated pathways for angiotensin II activation of extracellular signal-regulated kinases 1 and 2. *Proc Natl Acad Sci USA* **100**:10782–10787.
- Xiao K, Shenoy SK, Nobles K, and Lefkowitz RJ (2004) Activation-dependent conformational changes in β -arrestin 2. *J Biol Chem* **279**:55744–55753.
- Xu W, Reith MEA, Liu-Chen LY, and Kortagere S (2019) Biased signaling agonist of dopamine D3 receptor induces receptor internalization independent of β -arrestin recruitment. *Pharmacol Res* **143**:48–57.
- Yeow I, Howard G, Chadwick J, Mendoza-Topaz C, Hansen CG, Nichols BJ, and Shvets E (2017) EHD proteins cooperate to generate caveolar clusters and to maintain caveolae during repeated mechanical stress. *Curr Biol* **27**:2951–2962.e5.
- Yin W, Li Z, Jin M, Yin YL, de Waal PW, Pal K, Yin Y, Gao X, He Y, Gao J, et al. (2019) A complex structure of arrestin-2 bound to a G protein-coupled receptor. *Cell Res* **29**:971–983.
- Yin W, Liu H, Peng Z, Chen D, Li J, and Li JD (2014) Mechanisms that underlie the internalization and extracellular signal regulated kinase 1/2 activation by PKR2 receptor. *Cell Signal* **26**:1118–1124.
- Zhang J, Ferguson SSG, Barak LS, Ménard L, and Caron MG (1996) Dynamin and β -arrestin reveal distinct mechanisms for G protein-coupled receptor internalization. *J Biol Chem* **271**:18302–18305.
- Zhuang Z, Marshansky V, Breton S, and Brown D (2011) Is caveolin involved in normal proximal tubule function? Presence in model PT systems but absence in situ. *Am J Physiol Renal Physiol* **300**:F199–F206.
- Zimmerman B, Beautrait A, Aguila B, Charles R, Escher E, Claing A, Bouvier M, and Laporte SA (2012) Differential β -arrestin-dependent conformational signaling and cellular responses revealed by angiotensin analogs. *Sci Signal* **5**:ra33.

Address correspondence to: Thor C. Möller, Universitetsparken 2, 2100 Copenhagen, Denmark. E-mail: thor.moller@sund.ku.dk