

**Bi-directional Regulation of CaMKII Activity by Dopamine D4 Receptors in Prefrontal
Cortex**

Zhenglin Gu and Zhen Yan[†]

Dept. of Physiology and Biophysics, State University of New York at Buffalo, School of
Medicine and Biomedical Sciences, Buffalo, NY, 14214.

Running Title: D4 regulation of CaMKII Activity in PFC

† Address correspondence to: Dr. Zhen Yan, Department of Physiology and Biophysics, State University of New York at Buffalo, 124 Sherman Hall, Buffalo, NY, 14214. Tel: 716-829-3058. Fax: 716-829-2699. Email: zhenyan@buffalo.edu.

Text: 20 pages

Figures: 5

Tables: 0

References: 40

Abstract: 192 words

Introduction: 490 words

Discussion: 505 words

ABBREVIATIONS: CaMKII, Ca²⁺/calmodulin-dependent protein kinase II; PLC, phospholipase C; CNQX, 6-cyano-7-nitroquinoxaline-2,3-dione; APV, D(-)-2-amino-5-phosphonopetanoic acid; 2APB, 2-aminoethoxydiphenylborane; DHBP, 1,1'-diheptyl-4-4'-bipyridinium; NEM, N-ethylmaleimide; IP₃Rs, inositol-1,4,5-triphosphate receptors; RyRs, ryanodine receptors.

Abstract

The dopamine D₄ receptor in prefrontal cortex (PFC) plays a key role in normal mental functions and neuropsychiatric disorders. However, the cellular mechanisms and physiological actions of D₄ receptors remain elusive. In this study, we found that activation of D₄ receptors in PFC exerts a complex regulation of Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) – a multifunctional enzyme critically involved in synaptic plasticity that is fundamental for cognitive and emotional processes. In PFC slices with high neuronal activity, application of the D₄ receptor agonist PD168077 produced a potent reduction of the CaMKII activity, whereas in PFC slices with low neuronal activity, PD168077 caused a marked increase of the CaMKII activity. The D₄ up-regulation of CaMKII activity was through the stimulation of phospholipase C (PLC) pathway and elevation of intracellular Ca²⁺ via IP₃ receptors. These results reveal a bi-directional regulation of CaMKII activity by PFC D₄ receptors in response to changes in neuronal activity, and a non-classical signaling pathway underlying the D₄ up-regulation of CaMKII activity. This modulation provides a unique and flexible mechanism for D₄ receptors to regulate CaMKII activity, which could lead to dynamic regulation of many targets of CaMKII by D₄ receptors.

Introduction

Prefrontal cortex (PFC) is a brain region critically involved in the control of cognition, reasoning, perception and emotion (Goldman-Rakic, 1995). Dysfunction of PFC has been implicated in a variety of neuropsychiatric disorders including schizophrenia (Andreasen et al., 1997; Lewis and Lieberman, 2000). PFC functions are highly influenced by the dopaminergic input from the ventral tegmental area (Brozoski et al., 1979; Berger et al., 1988). Aberration of the dopaminergic system in PFC is considered as a major factor to the pathophysiology of schizophrenia (Grace, 1991; Carlsson et al., 2001).

Dopamine D₄ receptors are highly enriched in PFC neurons (Mrzljak et al., 1996; Wedzony et al., 2000). The elevated D₄ receptors found in PFC of schizophrenia patients (Seeman et al., 1993) and the high affinities of D₄ receptors for antipsychotic drugs (Van Tol et al., 1991; Kapur and Remington, 2001) suggest that D₄ receptors may be critically involved in PFC functioning and neuropsychiatric disorders (Oak et al., 2000). In agreement with this, D₄ receptor antagonists ameliorate cognitive deficits caused by the psychotomimetic drug phencyclidine (Jentsch et al., 1997; 1999). Moreover, mice lacking D₄ receptors exhibit supersensitivity to psychomotor stimulants (Rubinstein et al., 1997) and reduced exploration of novel stimuli (Dulawa et al., 1999). To understand how D₄ receptors regulate PFC functions under normal and pathological conditions, we need to determine what are the potential targets of D₄ receptors that are critically involved in the regulation of cognitive and emotional processes subserved by PFC.

One potential target of such for D₄ receptors is the Ca²⁺/calmodulin-dependent protein kinase II (CaMKII). CaMKII is highly expressed in the forebrain and concentrated at postsynaptic densities at glutamatergic synapses (Kennedy et al., 1983). This ideal position allows the multifunctional enzyme to play a central role in regulating several key postsynaptic targets required for synaptic plasticity that is integral for learning and memory (Malenka and Nicoll, 1999; Soderling et al., 2001). Mice with deficient CaMKII exhibit impairments in spatial learning

(Silva et al., 1992) and permanent memory retention (Frankland et al., 2001). In addition to the cognitive deficit, these CaMKII mutant mice exhibit a spectrum of behavioral abnormalities associated with emotional disorders including a decreased fear response and an increase in defensive aggression (Chen et al., 1994).

The function of CaMKII is shaped by its autoregulation and subcellular localization (Hudmon and Schulman, 2002). CaMKII is autophosphorylated at Thr²⁸⁶ when the enzyme is activated in the presence of Ca²⁺/calmodulin, leading to the appearance of a sustained, Ca²⁺-independent activity (Miller and Kennedy, 1986). This autoregulatory property enables CaMKII to act as a molecular memory device to detect synaptic activity, and to coordinate and execute Ca²⁺ signal transduction. CaMKII also dynamically alters its subcellular distribution following NMDA receptor stimulation through a mechanism involving Ca²⁺/calmodulin binding and autophosphorylation (Shen and Meyer, 1999). Given the convergent involvement in PFC functions for D₄ receptors and CaMKII, we sought to understand their interactions by examining the D₄ regulation of CaMKII activity in this study.

Materials and Methods

Western blot analysis. PFC slices were prepared as previously described (Gu et al., 2003). After treatment with different agents as indicated in the text, equal amounts of protein from slice homogenates were separated on 7.5% acrylamide gels and transferred to nitrocellulose membranes. The blots were blocked with 5% nonfat dry milk for 1 hour at room temperature. Then the blots were incubated with the anti-Thr²⁸⁶-phosphorylated α -CaMKII antibody (Santa Cruz, 1:2,000) for 1 hour at room temperature. After being rinsed, the blots were incubated with horseradish peroxidase-conjugated anti-rabbit antibodies (Amersham, 1:2,000) for 1 hour at room temperature. Following 3 washes, the blots were exposed to the enhanced chemiluminescence substrate. Then the blots were stripped for 1 hour at 50 °C followed by saturation in 5% nonfat

dry milk and incubated with an anti- α -CaMKII antibody (Upstate Biotechnology, 1:5,000) for the detection of the total α -CaMKII. Quantification was obtained from densitometric measurements of immunoreactive bands on films.

Dopamine receptor ligands PD168077 maleate (PD), L-745870 trihydrochloride (Tocris, Ballwin, MO), quinpirole, sulpiride and SCH23390 (Sigma, St. Louis, MO), as well as second messenger reagents U73122, genistein, BAPTA/AM, 2-aminoethoxydiphenylborane (2APB), 1,1'-diheptyl-4-4'-bipyridinium (DHBP), thapsigargin (Calbiochem, San Diego, CA), AMPA/KA receptor antagonist 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX), and NMDA receptor antagonist D(-)-2-amino-5-phosphonopentanoic acid (D -APV) (Sigma) were made up as concentrated stocks and stored at -20°C . The final DMSO concentration in all applied solutions was less than 0.1%. Stocks were thawed and diluted immediately prior to use.

Results

Bi-directional regulation of CaMKII activity by D_4 receptors in PFC neurons

CaMKII is activated by the binding of Ca^{2+} /calmodulin, and then it undergoes autophosphorylation at Thr²⁸⁶, which renders the enzyme to obtain Ca^{2+} -independent autonomous activity (Miller and Kennedy, 1986). Thus, the activated CaMKII (Thr²⁸⁶-phosphorylated) should be sensitive to stimuli that can change cellular Ca^{2+} , such as neuronal activity. So we first examined whether the activation levels of CaMKII might be dynamically regulated by D_4 receptors in response to different patterns of neuronal activity. PFC slices were incubated for 1 hour with either bicuculline (BIC, 10 μM) to increase activity through block of inhibitory transmission, or with CNQX (10 μM) and APV (20 μM) to decrease activity through block of excitatory transmission, followed by a 10-min treatment with the specific D_4 receptor agonist PD168077 (Glase et al., 1997; Wang et al., 2002). As shown in **Figure 1A**, the basal level of activated CaMKII in PFC slices was higher after BIC treatment, compared to CNQX/APV treatment.

PD168077 (20 μ M) caused a significant decrease of the activated CaMKII in PFC with high neuronal activity (BIC-treated), but caused a potent increase of the activated CaMKII in PFC with low neuronal activity (CNQX/APV-treated). In contrast to the bi-directional effect of PD168077, the D₂ receptor agonist quinpirole (20 μ M) only produced a reduction of the activated CaMKII irrespective of the neuronal activity. Total CaMKII levels exhibited no change with any of the treatment. Quantitative data from a series of experiments are summarized in **Figure 1B**.

PD168077 decreased Thr²⁸⁶-phosphorylated CaMKII by $65 \pm 11\%$ (n = 8) in BIC-treated PFC slices, while increased Thr²⁸⁶-phosphorylated CaMKII by $270 \pm 52\%$ (n = 8) in CNQX/APV-treated PFC slices. Quinpirole reduced Thr²⁸⁶-phosphorylated CaMKII by $72 \pm 12\%$ (n = 6) or $36 \pm 6\%$ (n = 6) in PFC slices treated with BIC or CNQX/APV, respectively.

Similar experiments were performed in PFC slices pretreated with TTX (0.5 μ M, 1 hr) to suppress spike activity. As shown in **Figure 1C** and **1D**, PD168077 caused a significant decrease ($59 \pm 10\%$, n = 8) of the Thr²⁸⁶-phosphorylated CaMKII in PFC with high neuronal activity (no TTX pretreatment), but caused a marked increase ($240 \pm 48\%$, n = 8) of the Thr²⁸⁶-phosphorylated CaMKII in PFC with low neuronal activity (TTX-pretreated). Quinpirole reduced the level of Thr²⁸⁶-phosphorylated CaMKII by $64 \pm 12\%$ (n = 6) or $31 \pm 5\%$ (n = 6) in PFC slices pretreated without or with TTX, respectively. These results indicate that D₄ receptors exert a dynamic bi-directional regulation of CaMKII activity depending on the neuronal activity.

To confirm that neuronal activity is manipulated by drugs that affect synaptic transmission, glutamatergic excitation or GABAergic inhibition, we compared the level of activated (Thr²⁸⁶-phosphorylated) CaMKII in PFC slices treated with saline or various drugs. As shown in **Figure 1E** and **1F**, comparing with saline-treated slices (ctl), slices treated with TTX, CNQX/APV, or the non-selective glutamate receptor antagonist kynurenic acid (1 mM) showed a substantial decrease of the activated CaMKII (TTX: $67 \pm 12\%$, n = 10; CNQX/APV: $72 \pm 13\%$, n = 10; kynurenic acid: $70 \pm 13\%$, n = 8). Moreover, the reduction of CaMKII activity by TTX or

CNQX/APV was not blocked by the PKA activator cpt-cAMP (100 μ M, cpt-cAMP + TTX: $58 \pm 13\%$, $n = 8$; cpt-cAMP + CNQX/APV: $63 \pm 10\%$, $n = 8$), suggesting that it is not mediated by PKA inhibition. On the other hand, comparing with saline-treated slices (ctl), slices treated with bicuculline caused little change on the activated CaMKII ($20 \pm 4\%$, $n = 8$), and slices treated with glutamate/glycine (100/10 μ M) or high KCl (30 mM) further increased the level of activated CaMKII (glutamate: $89 \pm 17\%$, $n = 8$; KCl: $92 \pm 17\%$, $n = 8$). These data suggest that PFC neurons are switched to the “low activity” state by the treatment with TTX, CNQX/APV or kynurenic acid, while they are at the “high activity” state in saline-treated (ctl), bicuculline-treated, glutamate/glycine-treated or high KCl-treated slices.

We further compared the effect of PD168077 on CaMKII activity in PFC slices at the “high activity” state. As shown in **Figure 1G** and **1H**, PD168077 caused a potent reduction of the activated CaMKII in saline-treated, glutamate/glycine-treated or high KCl-treated slices (saline: $58 \pm 12\%$, $n = 8$; glutamate: $69 \pm 15\%$, $n = 8$; KCl: $68 \pm 15\%$; $n = 8$), similar to the effect of PD168077 in bicuculline-treated slices (Figure 1A and 1B). These data further indicate that D₄ receptors decrease the level of CaMKII activation in PFC with high neuronal activity.

Mediation by D₄ receptors of the up-regulation of CaMKII activity in PFC slices

Since D₄ receptors couple to the “classical” inhibition of PKA pathway in transfected cell lines (Chio et al., 1994), it is surprising that D₄ receptors increased the activation level of CaMKII in PFC neurons with low neuronal activity. Thus, in subsequent experiments, we further examined the mechanisms underlying D₄ up-regulation of CaMKII activity in TTX-pretreated PFC slices.

The dose-dependence of PD168077-induced CaMKII activation is shown in **Figure 2A** and **2B**. A small effect could be detected following a 10-min exposure to 5 μ M of PD168077, and a saturating effect was seen at 20 μ M of PD168077. Quantification data exhibited a 3.4 ± 0.8 fold increase of CaMKII activity ($n = 8$, $p < 0.001$, ANOVA) by PD168077 (20 μ M, 10 min). The

kinetics of PD168077-induced activation of CaMKII was also tested. As demonstrated in **Figure 2C** and **2D**, the CaMKII activation induced by PD168077 (40 μ M) showed rapid and transient kinetics, reaching a peak at 10 minutes and declining to basal levels within 30-60 minutes.

To verify that D₄ receptors were mediating the PD168077 activation of CaMKII, we examined the ability of selective D₄ receptor antagonists to prevent the action of PD168077. As shown in **Figure 3A** and **3B**, PD168077 (40 μ M) produced a potent increase (3.7 ± 0.9 fold, $n = 12$) of activated CaMKII in PFC slices, and this effect was significantly ($p < 0.001$, ANOVA) blocked by L-745870 (20 μ M, 0.98 ± 0.2 fold, $n = 8$), a highly selective D₄ antagonist (Patel et al., 1997). In contrast to the strong effect of PD168077 on CaMKII activation in PFC slices, PD168077 failed to regulate CaMKII activity in striatal slices (1.1 ± 0.2 fold, $n = 6$, **Figure 3A** and **3B**), consistent with the highly enriched expression of D₄ receptors in PFC, but not in striatum.

To further confirm the involvement of D₄ receptors in the up-regulation of CaMKII activity, we tested the effect of dopamine (50 μ M) on CaMKII in the presence of D₁/D₅ antagonist SCH23390 (10 μ M) and D₂/D₃ antagonist sulpiride (10 μ M). As shown in **Figure 3C** and **3D**, when D₁/D₅ and D₂/D₃ receptors were blocked, dopamine produced an enhancement of CaMKII activity (3.4 ± 0.8 fold, $n = 8$), mimicking the PD168077 effect. Moreover, this effect of dopamine on CaMKII was blocked by the D₄ antagonist L-745870 (1.15 ± 0.2 fold, $n = 6$). These results suggest that dopamine released on PFC neurons could indeed elevate CaMKII activity via D₄ receptors.

Signaling mechanisms underlying the D₄ enhancement of CaMKII activity in PFC slices

Our previous study has shown that D₄ receptors decrease CaMKII activity in PFC slices (no TTX pretreatment) through a cascade involving the inhibition of PKA and ensuing disinhibition of protein phosphatase 1 (PP1) (Wang et al., 2003). We next examined the signal transduction

pathways mediating the increase of CaMKII activity by D₄ receptors in PFC slices when the neuronal activity was suppressed by TTX pretreatment. As shown in **Figure 4A**, application of the phospholipase C (PLC) inhibitor U73122 (1 μM), but not the broad-spectrum tyrosine kinase inhibitor genistein (100 μM), blocked the PD168077-induced increase of CaMKII activity. Moreover, application of the G_{i/o} protein alkylating agent N-ethylmaleimide (NEM, 30 μM) failed to prevent PD168077 from elevating CaMKII activity. As summarized in **Figure 4B**, the PD168077-induced activation of CaMKII (3.6 ± 0.8 fold, n = 14) was abolished in the presence of U73122 (0.9 ± 0.2 fold, n = 8), but was intact in the presence of genistein (3.5 ± 1.0 fold, n = 8) or NEM (3.9 ± 1.3 fold, n = 8). These results suggest that the D₄ enhancement of CaMKII activity is through a mechanism dependent on the stimulation of PLC pathway, but not the activation of tyrosine kinases or the coupling to G_{i/o} proteins.

Given the dependence of CaMKII activation on Ca²⁺, we then examined whether the D₄ enhancement of CaMKII activity required the Ca²⁺ entry from extracellular regions or the Ca²⁺ elevation from intracellular stores. To test this, PFC slices were incubated in Ca²⁺-free solutions or with the membrane-permeable Ca²⁺ chelator BAPTA/AM. As shown in **Figure 5A** and **5B**, PD168077 induced a potent increase (4.1 ± 1.2 fold, n = 8) in CaMKII activity under Ca²⁺-free conditions, similar to the PD168077 effect in normal Ca²⁺-containing solutions (3.8 ± 0.7 fold, n = 15). However, when intracellular Ca²⁺ increase was blocked by BAPTA/AM (50 μM), PD168077 failed to enhance CaMKII activity (1.12 ± 0.25 fold, n = 8). These results suggest that an increase of intracellular Ca²⁺ from internal stores is required for the D₄ enhancement of CaMKII activity.

In neurons, a major source of internal calcium is the stores present in the endoplasmic reticulum (ER) network. Both inositol-1,4,5-triphosphate receptors (IP₃Rs) and ryanodine receptors (RyRs) on ER are responsible for releasing calcium from this internal source (Kostyuk and Verkhatsky, 1994; Simpson et al., 1995). To determine which one was involved in the D₄

activation of CaMKII, PFC slices were pretreated with pharmacological agents to block these receptors. As shown in Figure 5A and 5B, application of 2APB (30 μ M), a membrane-permeable IP₃R antagonist (Hamada et al., 1999), abolished the D₄ effect on CaMKII activation (0.9 ± 0.2 fold, $n = 10$). In contrast, DHBP (30 μ g/ml), a potent RyR antagonist (Kang et al., 1994), failed to alter the D₄ enhancement of CaMKII activity (3.6 ± 0.9 fold, $n = 6$). Pretreatment of PFC slices with the intracellular calcium pump inhibitor thapsigargin (5 μ M, 30 min) to deplete internal stores of Ca²⁺ also eliminated the D₄ effect on CaMKII activation (1.0 ± 0.2 fold, $n = 8$). These results suggest that D₄ receptors elevate intracellular calcium via IP₃Rs to increase CaMKII activity.

As a control, we also examined the involvement of PLC/IP₃R signaling in D₄ reduction of CaMKII activity in high neuronal activity conditions. As shown in Figure 5C and 5D, application of the PLC inhibitor U73122 (1 μ M) or IP₃R antagonist 2APB (30 μ M) failed to block the PD168077-induced decrease of activated CaMKII ($62 \pm 13\%$, $n = 8$; U73122: $74 \pm 12\%$, $n = 8$; 2APB: $71 \pm 11\%$, $n = 8$), suggesting that the PLC/IP₃R signaling is not involved in D₄ reduction of CaMKII activity.

Discussion

CaMKII has been regarded as a cognitive kinase because of its involvement in regulating learning and memory and its autoregulatory properties that can be viewed as a type of molecular memory (Hudmon and Schulman, 2002). A variety of extracellular signals triggers the activation of CaMKII by elevating the intracellular Ca²⁺ level through Ca²⁺ influx or Ca²⁺ release from internal stores. We show here that stimulation of D₄ receptors can lead to either up- or down-regulation of CaMKII activity depending on basal neuronal activity. In PFC slices with suppressed neuronal activity, the level of activated CaMKII was increased by D₄ receptors, while in PFC slices with elevated neuronal activity, the level of activated CaMKII was decreased by D₄ receptors. This

dual regulation of CaMKII activity was unique for D₄ receptors, as it was not observed with D₂ receptor activation. By regulating CaMKII activity in such a dynamic activity-dependent fashion, D₄ receptors could fine-tune the functions of CaMKII flexibly and precisely.

How can D₄ receptors either decrease or increase CaMKII activity? Emerging evidence has suggested that G protein-mediated signal transduction is a complex signaling network with divergent and convergent pathways intimately intertwined (Gudermann et al., 1997). The “classical” signaling cascade for D₄ receptors is to couple to G_{i/o}-type G proteins to inhibit adenylyl cyclase and cAMP formation (Chio et al., 1994). The inhibition of PKA could cause the activation of PP1 via decreased phosphorylation of the inhibitory protein I-1 (Ingebritsen & Cohen, 1983), leading to the dephosphorylation of CaMKII and a decrease of CaMKII activity. Our previous study confirmed this mechanism for the D₄ down-regulation of CaMKII activity (Wang et al., 2003). In this study, we show that the D₄ up-regulation of CaMKII activity is through the stimulation of PLC pathway and elevation of intracellular Ca²⁺ via IP₃ receptors. How D₄ receptors activate the PLC pathway is not clear. One potential mechanism is that activation of D₄ receptors in PFC neurons leads to the release of G protein βγ subunits and thus potentiates the stimulation of PLC by βγ subunits (Camps et al., 1992).

This study mechanistically links together D₄ receptors and CaMKII, both of which have been implicated in cognitive and emotional processes associated with PFC. The D₄ regulation of CaMKII activity enables D₄ receptors to affect many aspects of cellular function via changing numerous CaMKII substrates, such as K⁺ channels, glutamate receptors, synapsin, CREB, tau and so on (Hudmon and Schulman 2002). A novel feature of this D₄ modulation CaMKII is that it is bi-directional depending on neuronal activity, and a “dual signaling”, i.e. inhibition of adenylyl cyclase and stimulation of PLC, underlies the D₄-induced suppression or potentiation of CaMKII activity. This supports the notion that many neuromodulators, like dopamine and serotonin, can have dual roles not only because they can act on a variety of different targets, but also because

they can act differently on the same target under different physiological conditions (Cai et al., 2002). This mechanism ensures that the modulation provides a feedback system to effectively maintain normal neuronal activity.

References

- Andreasen NC, O'Leary DS, Flaum M, Nopoulos P, Watkins GL, Boles Ponto LL, and Hichwa RD (1997) Hypofrontality in schizophrenia: distributed dysfunctional circuits in neuroleptic-naive patients. *Lancet* **349**:1730-4.
- Berger BS, Trottier C, Verney P, Gaspar P, and Alvarez C (1988) Regional and laminar distribution of the dopamine and serotonin innervation in the macaque cerebral cortex: A radioautographic study. *J. Comp. Neurol.* **273**:99-119.
- Brozoski TJ, Brown RM, Rosvold HE, and Goldman PS (1979) Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science* **205**: 929-32.
- Cai X, Flores-Hernandez J, Feng J, Yan Z (2002) Activity-dependent bi-directional regulation of GABA_A receptor channels by serotonin 5-HT₄ receptors in pyramidal neurons of the prefrontal cortex. *J. Physiology* **540**:743-59.
- Camps M, Carozzi A, Schnabel P, Scheer A, Parker PJ, Gierschik P (1992) Isozyme-selective stimulation of phospholipase C-beta 2 by G protein beta gamma-subunits. *Nature* **360**:684-6.
- Carlsson A, Waters N, Holm-Waters S, Tedroff J, Nilsson M, and Carlsson ML (2001) Interactions between monoamines, glutamate, and GABA in schizophrenia: new evidence. *Annu Rev Pharmacol Toxicol* **41**:237-60.
- Chen C, Rainnie DG, Greene RW, and Tonegawa S (1994) Abnormal fear response and aggressive behavior in mutant mice deficient for alpha-calcium-calmodulin kinase II. *Science* **266**: 291-4.

- Chio CL, Drong RF, Riley DT, Gill GS, Slightom JL, and Huff RM (1994) D₄ dopamine receptor-mediated signaling events determined in transfected Chinese hamster ovary cells. *J Biol Chem* **269**:11813-9.
- Dulawa SC, Grandy DK, Low MJ, Paulus MP, and Geyer MA (1999) Dopamine D₄ receptor-knock-out mice exhibit reduced exploration of novel stimuli. *J Neurosci* **19**:9550-6.
- Frankland PW, O'Brien C, Ohno M, Kirkwood A, Silva AJ (2001) Alpha-CaMKII-dependent plasticity in the cortex is required for permanent memory. *Nature* **411**:309-13.
- Glase SA, Akunne HC, Georgic LM, Heffner TG, MacKenzie RG, Manley PJ, Pugsley TA, Wise LD (1997) Substituted [(4-phenylpiperazinyl)-methyl]benzamides: selective dopamine D₄ agonists. *J Med Chem* **40**:1771-2.
- Goldman-Rakic PS (1995) Cellular basis of working memory. *Neuron* **14**:477-85.
- Grace AA (1991) Phasic versus tonic dopamine release and the modulation of dopamine system responsivity: a hypothesis for the etiology of schizophrenia. *Neuroscience* **41**:1-24.
- Gu Z, Zhong P, and Yan Z (2003) Activation of Muscarinic Receptors Inhibits β -Amyloid Peptide – Induced Signaling in Cortical Slices. *J. Biol. Chem.* **278**:17546-56.
- Gudermann T, Schoneberg T, Schultz G (1997) Functional and structural complexity of signal transduction via G-protein-coupled receptors. *Annu Rev Neurosci.* **20**:399-427.
- Hamada T, Liou SY, Fukushima T, Maruyama T, Watanabe S, Mikoshiba K, Ishida N (1999) The role of inositol trisphosphate-induced Ca²⁺ release from IP₃-receptor in the rat suprachiasmatic nucleus on circadian entrainment mechanism. *Neurosci Lett.* **263**:125-8.
- Hudmon A, Schulman H (2002) Neuronal CA²⁺/calmodulin-dependent protein kinase II: the role of structure and autoregulation in cellular function. *Annu Rev Biochem.* **71**:473-510.
- Ingebritsen TS, and Cohen P (1983) Protein phosphatases: properties and role in cellular regulation. *Science* **221**:331-8.

Jentsch JD, and Roth RH (1999) The neuropsychopharmacology of phencyclidine: from NMDA receptor hypofunction to the dopamine hypothesis of schizophrenia.

Neuropsychopharmacology **20**:201-25.

Jentsch JD, Redmond DE Jr., Elsworth JD, Taylor JR, Youngren KD, and Roth RH (1997)

Enduring cognitive deficits and cortical dopamine dysfunction in monkeys after long-term administration of phencyclidine. *Science* **277**:953-5.

Kang JJ, Hsu KS, Lin-Shiau SY (1994) Effects of bipyridylium compounds on calcium release from triadic vesicles isolated from rabbit skeletal muscle. *Br J Pharmacol.* **112**:1216-22.

Kapur S, and Remington G (2001) Atypical antipsychotics: new directions and new challenges in the treatment of schizophrenia. *Annu Rev Med* **52**:503-17.

Kennedy MB, Bennett MK, Erondy NE (1983) Biochemical and immunochemical evidence that the "major postsynaptic density protein" is a subunit of a calmodulin-dependent protein kinase. *Proc Natl Acad Sci U S A.* **80**:7357-61.

Kostyuk P, Verkhratsky A (1994) Calcium stores in neurons and glia. *Neuroscience* **63**:381-404.

Lewis DA, and Lieberman JA (2000) Catching up on schizophrenia: Natural history and neurobiology. *Neuron* **28**:325-334.

Malenka RC, Nicoll RA (1999) Long-term potentiation — a decade of progress? *Science* **285**:1870-4.

Miller SG, and Kennedy MB (1986) Regulation of brain type II Ca²⁺/calmodulin-dependent protein kinase by autophosphorylation: a Ca²⁺-triggered molecular switch. *Cell* **44**:861-70.

Mrzljak L, Bergson C, Pappy M, Huff R, Levenson R, and Goldman-Rakic PS (1996)

Localization of dopamine D₄ receptors in GABAergic neurons of the primate brain. *Nature* **381**:245-8.

Oak JN, Oldenhof J, and Van Tol HH (2000) The dopamine D(4) receptor: one decade of research. *Eur J Pharmacol* **405**:303-27.

- Patel, S., Freedman, S., Chapman, K.L., Emms, F., and Fletcher, A.E., et al. (1997) Biological profile of L-745,870, a selective antagonist with high affinity for the dopamine D₄ receptor. *J Pharmacol Exp Ther* **283**, 636-47.
- Rubinstein M, Phillips TJ, Bunzow JR, Falzone TL, Dziewczapolski G, Zhang G, Fang Y, Larson JL, McDougall JA, Chester JA, Saez C, Pugsley TA, Gershanik O, Low MJ, and Grandy DK (1997) Mice lacking dopamine D₄ receptors are supersensitive to ethanol, cocaine, and methamphetamine. *Cell* **90**:991-1001.
- Seeman P, Guan HC, and Van Tol HH (1993) Dopamine D₄ receptors elevated in schizophrenia. *Nature* **365**:441-5.
- Shen K, Meyer T (1999) Dynamic control of CaMKII translocation and localization in hippocampal neurons by NMDA receptor stimulation. *Science* **284**:162-6.
- Silva AJ, Stevens CF, Tonegawa S, and Wang Y (1992) Deficient hippocampal long-term potentiation in alpha-calcium-calmodulin kinase II mutant mice. *Science* **257**:201-6.
- Simpson PB, Challiss RA, Nahorski SR (1995) Neuronal Ca²⁺ stores: activation and function. *Trends Neurosci.* **18**:299-306.
- Soderling TR, Chang B, Brickey D (2001) Cellular signaling through multifunctional Ca²⁺/calmodulin-dependent protein kinase II. *J Biol Chem.* **276**:3719-22.
- Van Tol HH, Bunzow JR, Guan HC, Sunahara RK, Seeman P, Niznik HB, and Civelli O (1991) Cloning of the gene for a human dopamine D₄ receptor with high affinity for the antipsychotic clozapine. *Nature* **350**:610-4.
- Wang X, Zhong P, Yan Z (2002) Dopamine D₄ receptors modulate GABAergic signaling in pyramidal neurons of prefrontal cortex. *J. Neurosci.* **22**:9185-93.
- Wang X, Zhong P, Gu Z, Chen G, Yan Z (2003) Regulation of NMDA receptors by dopamine D₄ signaling in prefrontal cortex. *J. Neurosci.* **23**:9852-61.

Wedzony K, Chocyk A, Mackowiak M, Fijal K, and Czyrak A (2000) Cortical localization of dopamine D4 receptors in the rat brain--immunocytochemical study. *J Physiol Pharmacol* **51**: 205-21.

Footnotes

This work was supported by NIH grants MH63128 and AG21923 (Z.Y.), NSF grant IBN-0117026 (Z.Y.), and Howard Hughes Medical Institute Biomedical Research Support Program grant 53000261 (SUNY at Buffalo).

Legends for Figures

Fig. 1. The D₄ receptor agonist PD168077 induced an up- and down-regulation of CaMKII activity depending on the neuronal activity in PFC slices. **A, C.** Immunoblots of phospho-CaMKII and CaMKII. PFC slices were incubated in bicuculline (BIC, 10 μ M) or CNQX (10 μ M)/APV (20 μ M) for 1 hour (A), or pretreated without or with TTX (0.5 μ M, 1 hour, C), followed by a 10-min treatment with PD168077 (PD, 20 μ M) or quinpirole (Quin, 20 μ M). Extracts of slices were immunoblotted with an anti-phospho- α -CaMKII antibody. Following stripping out signals, membranes were reblotted with an antibody recognizing the total α -CaMKII. **B, D.** Percentage changes of p-CaMKII induced by PD168077 or quinpirole in PFC slices incubated with bicuculline- or CNQX/APV (B), or pretreated without or with TTX (D). **E.** Immunoblots of phospho-CaMKII. Top: PFC slices were treated with TTX, CNQX/APV (C+A), kynurenic acid (kyn, 1 mM), bicuculline for 30 min, or glutamate/glycine (Glu, 100/10 μ M), KCl (30 μ M) for 20 min. Bottom: cpt-cAMP (100 μ M) was added 10 min before TTX or CNQX/APV treatment. **G.** Slices were pretreated with saline (ctl), glutamate/glycine, or KCl for 10 min, followed by a 10-min treatment with PD168077. **F, H.** Percentage changes of p-CaMKII induced by treatments corresponding to E and G, respectively. **: $p < 0.01$; *: $p < 0.001$, ANOVA.

Fig. 2. PD168077 increased CaMKII activity in a dose- and time-dependent manner. **A.** Dose-dependence of the PD168077-induced activation of CaMKII. PFC slices (TTX-pretreated) were treated with PD168077 for 10 min at the indicated concentrations. **B.** Quantification of p-CaMKII induced by different concentrations of PD168077. **C.** Time course of the PD168077-induced activation of CaMKII. PFC slices (TTX-pretreated) were treated with PD168077 (40 μ M) for the indicated durations. **D.** Quantification of p-CaMKII induced by

PD168077 treatment for different lengths of time. **: $p < 0.01$; *: $p < 0.001$, ANOVA, compared to control (-).

Fig. 3. The PD168077-induced up-regulation of CaMKII activity was mediated by D₄ receptors.

A. Immunoblots of phospho-CaMKII. PFC or striatal slices (TTX-pretreated) were incubated in the absence or presence of the selective D₄ antagonist L-745870 (20 μ M, 15 min), followed by a 10-min treatment with PD168077. **B.** Quantification of p-CaMKII induced by PD168077 in PFC or striatal slices. **C.** Immunoblots of phospho-CaMKII. PFC slices (TTX-pretreated) were incubated with SCH23390 (10 μ M) and sulpiride (10 μ M) for 30 min to block D₁/D₅ and D₂/D₃ receptors. Then they were incubated in the absence or presence of L-745870 (20 μ M, 15 min), followed by a 10-min treatment with dopamine (50 μ M). **D.** Quantification of p-CaMKII induced by dopamine (co-applied with SCH23390 plus sulpiride) in the absence or presence of L-745870. *: $p < 0.001$, ANOVA, compared to the effect under control conditions (-).

Fig. 4. The D₄ potentiation of CaMKII activity was dependent on the stimulation of PLC









pathway. **A.** Immunoblots of phospho-CaMKII. PFC slices (TTX-pretreated) were incubated in the absence or presence of various agents for 30 min, followed by a 10-min treatment with PD168077. Agents included: the PLC inhibitor U73122 (1 μ M), the tyrosine kinase inhibitor genistein (100 μ M), and the G_{i/o} protein alkylating agent NEM (30 μ M). **B.** Quantification of p-CaMKII induced by PD168077 under different treatments. *: $p < 0.001$, ANOVA, compared to the effect under control conditions (-).




Fig. 5. D₄ receptors augmented CaMKII activity through elevation of intracellular Ca²⁺ via IP₃

receptors. **A.** Immunoblots of phospho-CaMKII. PFC slices (TTX-pretreated) were incubated

MOL (1404R)

in the absence or presence of various agents or in a Ca^{2+} -free solution for 30 min, followed by a 10-min treatment with PD168077. Agents included: the membrane-permeable Ca^{2+} chelator BAPTA/AM (50 μM), the IP_3R antagonist 2APB (30 μM), the RyR antagonist DHBP (30 $\mu\text{g/ml}$), and the intracellular calcium pump inhibitor thapsigargin (5 μM). **C.** PFC slices (bicuculline-pretreated) were pretreated with saline, U73122 (1 μM), or 2APB for 30 min, followed by a 10-min treatment with PD168077. **B, D.** Quantification of p-CaMKII induced by PD168077 under different treatments. *: $p < 0.001$, ANOVA, compared to the effect under control conditions (-).



A**B****C****D**