

MOL 4077

Mechanisms of agonist action at D₂ dopamine receptors

David J Roberts, Hong Lin and Philip G Strange*

School of Animal and Microbial Sciences, University of Reading, Whiteknights,
Reading, RG6 6AJ, UK

MOL 4077

running title: agonist mechanisms

corresponding author, Philip G Strange, School of Animal and Microbial Sciences,
University of Reading, Whiteknights, Reading, RG6 6AJ, UK, telephone: 44-118-378-
8015, fax: 44-118-378-6537, e-mail: p.g.strange@reading.ac.uk

Text pages: 16

Tables: 3

Figures: 5

References: 36

Words in abstract: 218

Words in introduction: 741

Words in Discussion: 1699

Abbreviations: GPCR: G protein-coupled receptor; 7-OH DPAT : 7-hydroxy-2-
dipropylaminotetralin; NPA: N-propylnorapomorphine

MOL 4077

Abstract

In this study we investigated biochemical mechanisms of agonist action at the G protein-coupled D₂ dopamine receptor expressed in CHO cells. Stimulation of [³⁵S]GTPγS binding by full and partial agonists was determined at different concentrations of [³⁵S]GTPγS (0.1 and 10 nM) and in the presence of different concentrations of GDP. At both concentrations of [³⁵S]GTPγS, increasing GDP decreased the [³⁵S]GTPγS binding observed with maximally stimulating concentrations of agonist, with partial agonists exhibiting greater sensitivity to the effects of GDP than full agonists. The relative efficacy of partial agonists was greater at the lower GDP concentrations. Concentration/response experiments were performed for a range of agonists at the two [³⁵S]GTPγS concentrations and with different concentrations of GDP. At 0.1 nM [³⁵S]GTPγS the potency of both full and partial agonists was dependent on the GDP concentration in the assays. At 10 nM [³⁵S]GTPγS the potency of full agonists exhibited a greater dependence on the GDP concentration whereas the potency of partial agonists was virtually independent of GDP. It is concluded that at the lower [³⁵S]GTPγS concentration the rate determining step in G protein activation is the binding of [³⁵S]GTPγS to the G protein. At the higher [³⁵S]GTPγS concentration, for full agonists [³⁵S]GTPγS binding remains the slowest step whereas for partial agonists another (GDP-independent) step, probably ternary complex breakdown, becomes rate determining.

MOL 4077

Introduction

There is much interest in understanding mechanisms of action of agonists at receptors (agonist efficacy) and the mechanistic distinction between full and partial agonists ((Black and Leff, 1983; Clarke and Bond, 1998; Colquhoun, 1998; Kenakin, 2002; Strange, 1999). For the G protein-coupled receptors (GPCRs), an influential biochemical model of GPCR action has been the ternary complex model and its recent extensions (De Lean et al., 1980; Samama et al., 1993; Weiss et al., 1996). The model describes a ground state of the receptor (R) which can isomerise to a partially activated form (R*) which is able to couple better to the G protein to form the active (R*G) state.

The ternary complex model accounts for differences in the relative efficacy of full and partial agonists in terms of different extents of stabilisation of active (AR*G) and inactive (AR) states of the receptor. Full agonists stabilise R*G better than partial agonists so that relative efficacy is explained in terms of the differential stabilisation of a single activated state. G protein activation and GDP/GTP exchange follow accordingly.

This model has been examined using ligand-binding studies to determine affinities of agonists for G protein coupled (higher affinity, K_h) and uncoupled (lower affinity, K_l) forms of the receptor. Some studies report a correlation between the K_l/K_h ratio for agonists and their relative efficacy (Alder et al., 2003; De Lean et al., 1980; Egan et al., 2000; Kearn et al., 1999; Payne et al., 2002; Watson et al., 2000), other studies do not (Gardner and Strange, 1998; Gardner et al., 1997; Payne et al., 2002). It seems that there may be additional factors influencing relative efficacy such as differential

MOL 4077

abilities of some agonists to induce G protein activation within AR*G. Different agonists may stabilise different activated states of receptors leading to differential activities (Seifert et al., 2001; Waelbroeck, 2001)

G protein activation occurs, however, as part of a cycle of reactions (Figure 1) (Mosser et al., 2002; Waelbroeck, 2001; Zhong et al., 2003) and the overall rate of G protein activation may be dependent on several of the component processes although the slowest of these will limit the overall rate. The reactions of the cycle are as follows:

- (a) *agonist (A) binds to receptor to stabilise AR**.
- (b) *AR* and G_{GDP} combine to form AR*G*. For some agonists AR*G stability is a guide to agonist relative efficacy; some agonists can produce a stable AR*G complex but are partial agonists (Gardner and Strange, 1998; Gardner et al., 1997; Payne et al., 2002) and so their activity must be limited by another event.
- (c) *GDP release*. This is typically considered to be the rate-determining step in GPCR activation in the absence of agonist (Ross, 1989). In the presence of agonist, GDP dissociation is accelerated, and GDP association decreased (Florio and Sternweis, 1989). GDP release could, however, be the slowest step in the cycle for some agonists, despite strong stabilisation of AR*G.
- (d) *GTP binding*. Cells contain high concentrations of GTP (~50 μ M) (Jinnah et al., 1993; Otero, 1990) so that this step will be fast, and another step rate-determining. This step may be examined using the GTP analogue

MOL 4077

($[^{35}\text{S}]\text{GTP}\gamma\text{S}$). Typically these assays are performed at low concentrations of $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ and this step may become rate determining (Waelbroeck, 2001).

(e) *AR*G dissociates releasing AR, $G_{\alpha}\text{GTP}$ and $G_{\beta\gamma}$* . This step may be agonist-dependent for some receptors (Hausdorff et al., 1990; Van Koppen et al., 1994) and could be rate determining if an agonist were unable to mediate rapid breakdown of AR*G.

(f) *The intrinsic GTPase of the G protein hydrolyses GTP to GDP and deactivates G_{α}* . This step itself is independent of agonist as it is an intrinsic activity of the G protein but is unlikely to be rate determining as, in the presence of proteins with GTPase accelerating activity, this step is fast (Ross and Wilkie, 2000).

There are, therefore, several steps in the cycle that are regulated by agonists and which could determine the relative efficacy of agonists. It is not known whether the rate determining step in the cycle is the same for all agonists. In this study, therefore, we have examined the ability of a range of full and partial agonists to mediate G protein activation via the D₂ dopamine receptor. We have perturbed the function of the G protein cycle by altering the concentrations of both GDP and GTP γ S in order to understand which step in the cycle is rate limiting for different agonists.

MOL 4077

Materials and Methods

Materials. [³⁵S]GTP γ S (~37 TBq.mmol⁻¹) and [³H]spiperone (~600 GBq.mmol⁻¹) were purchased from Amersham Biosciences (Buckinghamshire, UK). Optiphase HiSafe-3 scintillation fluid was purchased from Perkin-Elmer Life Sciences (Cambridge, UK). Dopamine, bromocriptine and (\pm)-7-OH-DPAT were purchased from TOCRIS (Bristol, UK). NPA, β -phenylethylamine, m-tyramine and p-tyramine were purchased from Sigma (Dorset, UK).

Cell culture. CHO cells stably expressing native D_{2short} dopamine receptors (Wilson et al., 2001) were grown in DMEM containing 5% foetal bovine serum and 400 μ g ml⁻¹ active geneticin (to maintain selection pressure). Cells were grown at 37°C in an humidified atmosphere of 5 % CO₂.

Membrane preparation. Membranes were prepared from CHO cells expressing D_{2short} dopamine receptors as described previously (Castro and Strange, 1993). Briefly, confluent 175 cm² flasks of cells were washed once with 5ml HEPES buffer (20mM HEPES, 1 mM EGTA, 1 mM EDTA, 10 mM MgCl₂; pH 7.4). Cells were then removed from the surface of the flasks using 5ml HEPES buffer and glass balls (2mm diameter) and were then homogenised using an Ultra-Turrax homogeniser (two 5 second treatments). The homogenate was centrifuged at 1700 g (10 mins, 4°C) after which the supernatant was centrifuged at 48000 g (60 mins; 4°C). The resulting pellet was resuspended in HEPES buffer at a concentration of 3 - 5 mg protein ml⁻¹

MOL 4077

(determined by the method of (Lowry et al., 1951) and stored in aliquots at -70°C until use.

Radioligand binding experiments.

Cell membranes (25 µg) were incubated with [³H]spiperone (0.35 nM) and competing drugs in HEPES buffer (20 mM HEPES, 1 mM EGTA, 1 mM EDTA, 10 mM MgCl₂, 100 mM NaCl; pH 7.4 (using KOH) containing 0.1 mM dithiothreitol) in a final volume of 1 ml for 3 h at 25°C. The assay was terminated by rapid filtration (through Whatman GF/C filters) using a Brandel cell harvester followed by four washes with 4 ml ice-cold phosphate-buffered saline (0.14 M NaCl, 3 mM KCl, 1.5 mM KH₂PO₄, 5 mM Na₂HPO₄; pH 7.4) to remove unbound radioactivity. Filters were soaked in 2 ml of scintillation fluid for at least 5 h and bound radioactivity was determined by liquid scintillation counting. Non-specific binding of [³H]-spiperone was determined in the presence of 3 µM (+)-butaclamol.

[³⁵S]GTPγS binding assays. Cell membranes (25 µg) were incubated with various concentrations of GDP and agonist for 20 min before addition of 0.1 nM [³⁵S]GTPγS for 30 min or 10 nM [³⁵S]GTPγS (0.5 nM [³⁵S]GTPγS with 9.5 nM GTPγS) for 3 min in HEPES buffer at 30°C containing 0.1 mM dithiothreitol. In the absence of GDP, incubation times with 0.1 nM [³⁵S]GTPγS were reduced to 15 min. The assay was terminated by rapid filtration (through Whatman GF/C filters) using a Brandel cell harvester followed by four washes with 4 ml ice-cold phosphate-buffered saline (0.14 M NaCl, 3 mM KCl, 1.5 mM KH₂PO₄, 5 mM Na₂HPO₄; pH 7.4) to remove unbound radioactivity. Filters were soaked in 2 ml Optiphase HiSafe-3 for at least 5 h and bound radioactivity was determined by liquid scintillation counting.

MOL 4077

Data analysis. Radioligand binding and [³⁵S]GTPγS binding data were analysed by non-linear regression using Prism (Graphpad San Diego, USA). Statistical significance over multiple data sets was determined using an unpaired 2-way ANOVA followed by a Bonferroni post-test whilst that for two groups was determined using a t-test. Statistical significance was determined as $P < 0.05$.

MOL 4077

Results

Effects of different concentrations of GDP and [³⁵S]GTPγS on agonist stimulation of [³⁵S]GTPγS binding

Maximal agonist-stimulated effect and relative agonist efficacy

The maximal agonist-stimulated effect and relative agonist efficacy were determined from the stimulation of [³⁵S]GTPγS binding by agonists in membranes of CHO cells expressing the D₂ receptor (Wilson et al., 2001), (expression level of D₂ receptor 1-1.5 pmol/mg protein). Agonist-stimulated [³⁵S]GTPγS binding is due to the D₂ receptor as there is no stimulation in untransfected cells (Gardner et al., 1996). The agonist-stimulated response is completely inhibited after pertussis toxin treatment (100 ng/ml, 18h, data not shown) indicating a role for G_{i/o} proteins. The principal G_{i/o} proteins in CHO cells are G_{i2} and G_{i3} (Gettys et al., 1994; Raymond et al., 1993). The stimulation of [³⁵S]GTPγS binding by two full agonists (dopamine and NPA) and two partial agonists (p-tyramine and (±)-7-OH-DPAT) was assessed in the presence of increasing concentrations of GDP (0.3 - 30 μM) and using two concentrations of [³⁵S]GTPγS (0.1 and 10 nM). The four agonists were used at maximally stimulating concentrations and total [³⁵S]GTPγS binding was corrected for the agonist-independent binding to give the agonist-stimulated binding (Figure 2). The association rate of [³⁵S]GTPγS binding stimulated by dopamine was much faster at the higher [³⁵S]GTPγS concentration (10 nM) (t_{1/2}: 1-2 min, data not shown) compared with the rate at the lower concentration of [³⁵S]GTPγS (0.1 nM) (t_{1/2}: 10-15 min, (Gardner et al., 1996)). Assays were, therefore, terminated after 3 min for experiments at 10 nM [³⁵S]GTPγS, compared to 30 min at 0.1 nM [³⁵S]GTPγS in

MOL 4077

order to ensure that the determinations were on the linear part of the time course. In this way the determinations of agonist-stimulated [35 S]GTP γ S binding correspond to rate measurements.

Maximal stimulated [35 S]GTP γ S binding (over basal) was decreased for both full and partial agonists with increasing GDP concentration, although full agonists required higher concentrations of GDP than partial agonists to reduce their stimulation below that observed at the lowest concentration of GDP (Figure 2 a,c). Maximal agonist-stimulated [35 S]GTP γ S bound was also increased by ~10 fold by increasing the [35 S]GTP γ S concentration from 0.1 to 10 nM, and, given the difference in assay time for the two concentrations of [35 S]GTP γ S, this corresponded to a ~ 100 fold increase in the rate of [35 S]GTP γ S binding.

Increasing the concentration of GDP reduced the relative efficacy of partial agonists compared to full agonists in an almost linear fashion (from ~75 to ~50% relative efficacy at 0.1 nM [35 S]GTP γ S and from ~60 to ~30% relative efficacy at 10 nM [35 S]GTP γ S) (Figure 2 b,d). When relative agonist efficacies were compared at the two concentrations of [35 S]GTP γ S, it was seen that the relative efficacy of the partial agonists was lower at the higher [35 S]GTP γ S concentration (Figure 2 b,d).

Given that the relative efficacy of partial agonists was higher at the lower GDP concentrations, we performed some assays in the absence of GDP. Under these conditions, both full and partial agonists were still able to promote [35 S]GTP γ S binding over basal levels (Table 1; Figure 3). Both full and partial agonists, however, now stimulated [35 S]GTP γ S binding to the same extent. There was no change in the rate or extent of basal [35 S]GTP γ S binding in the presence of (+)-butaclamol if the

MOL 4077

membranes were pre-treated with an agonist, suggesting that there is no pre-bound GDP present in the preparation (data not shown). Therefore, the stimulation of [³⁵S]GTPγS binding in the absence of GDP is not a reflection of release of bound GDP.

Agonist potency

In order to probe the relationship between the EC₅₀ for agonist stimulation of [³⁵S]GTPγS binding and GDP concentration, a series of agonist/concentration experiments was performed at different concentrations of GDP. The four agonists used above were assessed at two different concentrations of [³⁵S]GTPγS (0.1 nM and 10 nM) and additional agonists were also tested at the lower concentration of [³⁵S]GTPγS. Representative data are shown in Figure 4 and the derived EC₅₀ values are given in Table 3 and Figure 5. At 0.1 nM [³⁵S]GTPγS, increasing concentrations of GDP caused a similar rightward shift in agonist EC₅₀ for all of the agonists tested with the exception of bromocriptine. This shift falls between the agonist binding affinities calculated for G-protein coupled and uncoupled receptors determined under the conditions used for the [³⁵S]GTPγS binding assays (see below and Table 2). In contrast, bromocriptine, which displays no observable affinity preference for coupled or uncoupled receptors (Gardner et al., 1997), shows no change in EC₅₀ with changing GDP concentration. With the exclusion of bromocriptine, when the pEC₅₀ was plotted against log[GDP], there was no significant difference between the slopes of the lines ($P > 0.05$) and a mean slope of -0.31 was obtained.

When the [³⁵S]GTPγS concentration was increased to 10 nM, the EC₅₀ for the two full agonists tested was shifted rightward as before as the GDP concentration was

MOL 4077

increased, whereas for the two partial agonists the EC_{50} was much less affected by GDP. This effect is emphasised in the pEC_{50} versus $\log[GDP]$ plots (Figure 5). Linear relationships between pEC_{50} and $\log[GDP]$ were still observed but there were significantly greater slopes for the full agonists, NPA and dopamine, compared to the partial agonists (\pm)-7-OH-DPAT and p-tyramine ($P < 0.05$) (Figure 5, Table 3).

Binding of full and partial agonists to D₂ dopamine receptors

The binding of the agonists (dopamine, NPA, p-tyramine and (\pm)-7-OH-DPAT) was determined in competition versus the binding of [³H]spiperone using assay buffer containing 100mM NaCl as in the [³⁵S]GTP γ S binding experiments (see above). In each case the competition data were fitted best by a two-binding site model and the derived dissociation constants (K_h , K_l) are given in Table 2.

Discussion

In this study we have examined some basic mechanisms of agonist action using the D₂ dopamine receptor as a model GPCR, with the aim of understanding the mechanistic distinction between full and partial agonists. For the GPCRs, differences in the relative efficacy of agonists have been explained using the ternary complex model (De Lean et al., 1982; Lefkowitz et al., 1993; Weiss et al., 1996) whereby partial agonists stabilise the ternary complex (AR*G) less well than full agonists. The model does not always account for relative efficacy and this may relate to the fact that GPCR activation depends on a cycle of reactions (Figure 1) (Mosser et al., 2002; Waelbroeck, 2001; Zhong et al., 2003). During receptor activation the reactions of the cycle will not be at equilibrium and different agonists may influence steps in the cycle differentially. In this report we have examined how agonists with different relative efficacies influence the steps in the cycle using the D₂ dopamine

MOL 4077

receptor as a model system. From the data we have been able to show that full and partial agonists differ in their abilities to modulate different reactions in the G protein cycle. The study, therefore, provides a mechanistic basis for the distinction between full and partial agonists.

We used stimulation of [³⁵S]GTPγS binding by agonists as a measure of their relative efficacy and examined the effect of different concentrations of GDP on both the maximal [³⁵S]GTPγS binding and relative efficacy. Assays were performed at two concentrations of [³⁵S]GTPγS (0.1 nM, 10 nM) with several full and partial agonists. In the [³⁵S]GTPγS binding assays, two principal parameters were determined for each agonist, the concentration of agonist achieving half the maximal stimulated effect (potency, EC₅₀) and the maximal rate of [³⁵S]GTPγS binding stimulated by saturating agonist concentrations.

The effects of GDP on the potency (EC₅₀) for agonists to stimulate [³⁵S]GTPγS binding were assessed. At low concentrations of [³⁵S]GTPγS (0.1 nM) the potency of each of the agonists tested, with the exception of bromocriptine, was reduced as the GDP concentration was increased. The effect of GDP was similar for each agonist, independent of its relative efficacy, as shown by the similar slopes of the lines relating pEC₅₀ and log[GDP]. The effects of GDP here reflect binding of GDP to the AR*G state leading to its breakdown and sequestration of G protein as G_{GDP}. Higher concentrations of agonist are then required to stabilise AR*G in which [³⁵S]GTPγS binding occurs and the EC₅₀ for the agonist is increased. Simulations of these effects

MOL 4077

have been reported (McLoughlin and Strange, 2000). The slope of the line relating pEC_{50} and $\log[GDP]$ reflects the affinities of the agonist for the G protein-coupled and uncoupled states of the receptor and the sensitivity of the agonist/receptor/G protein complex to GDP. Bromocriptine has been shown have similar affinities for the coupled and uncoupled states (Gardner et al., 1997) so it is not surprising that it is insensitive to GDP.

When higher concentrations of [^{35}S]GTP γ S (10 nM) were used, the potencies of the two full agonists (NPA, dopamine) tested were sensitive to the effects of GDP. Indeed the pEC_{50} was more sensitive to $\log[GDP]$ than at the lower concentration of [^{35}S]GTP γ S ($P < 0.05$). In contrast the potencies of the two partial agonists ((\pm)-7-OH-DPAT, p-tyramine) were virtually independent of $\log[GDP]$, when assays were performed at the higher [^{35}S]GTP γ S concentration. A change in the mechanism of [^{35}S]GTP γ S binding, from a GDP-dependent rate-determining step to a GDP-independent rate-determining step, may have occurred for the partial agonists at the higher concentration of [^{35}S]GTP γ S. The 100-fold increase in [^{35}S]GTP γ S concentration also leads to a substantial increase (~100 fold) in the maximal rate of [^{35}S]GTP γ S binding stimulated by dopamine. This increase in the maximal rate of binding of [^{35}S]GTP γ S as the concentration is increased suggests that the rate-determining step in the cycle at the lower concentration of [^{35}S]GTP γ S is the [^{35}S]GTP γ S binding event. This reflects the rather low concentration of [^{35}S]GTP γ S that is being used and similar conclusions have been reached by others using other approaches (Waelbroeck, 2001). Agonists are able, therefore, to modulate directly the rate of binding of [^{35}S]GTP γ S as has been suggested by Florio and Sternweis

MOL 4077

(Florio and Sternweis, 1989). Indeed, in the present study, agonists were able to stimulate [³⁵S]GTPγS binding in the absence of GDP supporting these ideas. This underlines the idea that agonists are able to regulate the rate of [³⁵S]GTPγS binding and suggests that the G protein associated with receptor is a better substrate for [³⁵S]GTPγS binding than free G protein. In the present system, however, in the absence of GDP the agonists all mediated the same maximal [³⁵S]GTPγS binding i.e. all appear as full agonists. There may, therefore, be a limit on the stimulation of [³⁵S]GTPγS binding possible in this system and in the absence of GDP this is achieved by all the agonists tested here.

If, therefore, [³⁵S]GTPγS binding is the rate determining step at the lower concentration of [³⁵S]GTPγS for all agonists, then the effect of the increase in [³⁵S]GTPγS concentration, and concomitant increase in overall rate may have led to a change in the rate-determining step for some agonists. For the partial agonists, at the higher concentration of [³⁵S]GTPγS, another, GDP-independent step may have become rate-determining. In order for this process to be GDP-independent, it must be subsequent to formation of AR*G and a likely candidate could be ternary complex breakdown. For the full agonists, the rate-determining step at the higher concentration of [³⁵S]GTPγS is GDP dependent and most likely is the [³⁵S]GTPγS binding event, although we cannot rule out that GDP release has become rate-determining.

MOL 4077

The maximal agonist-stimulated [^{35}S]GTP γ S binding was reduced at both concentrations of [^{35}S]GTP γ S as the GDP concentration was increased. The maximal [^{35}S]GTP γ S binding reflects the rate of the slowest process in the cycle and the amount of the different species in the cycle. At the lower concentration of [^{35}S]GTP γ S, where the rate reflects directly the [^{35}S]GTP γ S binding process, the effect of GDP on the maximal rate of [^{35}S]GTP γ S binding may result from a reduction in the level of AR*G species following sequestration of G proteins as G_{GDP} when the G protein cycle is at steady state. Although the potency of bromocriptine is unaffected by changes in GDP (see above), the maximal [^{35}S]GTP γ S binding for this compound is reduced by increased GDP supporting the above ideas. At the higher [^{35}S]GTP γ S concentration, for the full agonists the binding event is still the slowest process, whereas for the partial agonists, ternary complex breakdown may be the slowest process in the cycle. Effects of GDP on maximal [^{35}S]GTP γ S binding rate presumably reflect sequestration of G proteins reducing levels of AR*G and hence the overall rate of the cycle.

As well as the effects of GDP on the maximal rates of [^{35}S]GTP γ S binding there were effects on the relative efficacies of the partial agonists. The maximal rates of [^{35}S]GTP γ S binding of the partial agonists were more sensitive to GDP than those of the full agonists, resulting in a reduction in relative efficacy for the two partial agonists as GDP was increased. This suggests that differences in relative efficacy may reflect differences in GDP sensitivity of different agonist/receptor/G protein species i.e. full agonists are able to overcome G protein sequestration more than partial agonists. For other GPCRs it has been shown that agonists may affect the

MOL 4077

affinity of the receptor for the G protein (Tota and Schimerlik, 1990). Thus different AR*G complexes may appear differentially sensitive to GDP. The relative efficacies of the partial agonists were also generally lower at higher [³⁵S]GTPγS and this may reflect the change in rate determining step to ternary complex breakdown for which the partial agonists are deficient relative to the full agonists.

In the present study bromocriptine stands out as having unusual properties in that its potency for stimulation of [³⁵S]GTPγS binding, when measured at 100 pM [³⁵S]GTPγS, is insensitive to GDP unlike the potencies of the other agonists tested (Figure 5). Bromocriptine exhibits similar behaviour in ligand binding assays in that its binding is insensitive to guanine nucleotides unlike other agonists (see for example (Gardner and Strange, 1998; Gardner et al., 1997)). We have suggested that this reflects stabilisation by bromocriptine, in the absence of G protein coupling, of a conformation of the receptor that is close to the conformation in the fully active G protein-coupled state (Strange, 1999). Hence there is little difference in affinity between the uncoupled and G protein-coupled forms of the receptor.

This study has, therefore, provided new information on the biochemical basis of the distinction between full and partial agonists. Several steps in the G protein cycle are agonist-dependent: GDP release from the G protein, [³⁵S]GTPγS binding to the G protein, breakdown of the ternary complex. In the experiments performed in this report at the low concentration of [³⁵S]GTPγS, the binding of [³⁵S]GTPγS is the slowest step and differential effects of agonists on this step reflect full and partial agonism. This conclusion is likely to apply to all [³⁵S]GTPγS binding assays

MOL 4077

performed on all GPCRs at these low concentrations of [³⁵S]GTPγS. At the higher concentrations of [³⁵S]GTPγS, the rate of [³⁵S]GTPγS binding increases ~100 fold and, for the full agonists, ternary complex breakdown is still fast enough for a GDP-dependent event, most likely the [³⁵S]GTPγS binding event, to be rate determining. For the partial agonists, however, another step, probably ternary complex breakdown is slower and becomes rate determining. These conclusions are of some significance in that the concentration of GTP in the cell is high (~50μM,(Jinnah et al., 1993; Otero, 1990)) so that in cells partial agonism may be apparent because of this limitation of the rate of ternary complex breakdown.

Given that the experiments conducted here at the higher concentration of [³⁵S]GTPγS reflect more closely cellular conditions it should be possible to relate effects of agonists on [³⁵S]GTPγS binding performed under these conditions to experiments performed on whole cells, for example examining effects of agonists to inhibit cAMP accumulation (see for example (Payne et al., 2002)). The present set of data for the higher [³⁵S]GTPγS concentration is not extensive enough to allow this correlation to be examined but this will be an important aim for future work in relating these in vitro assays to cellular assays.

MOL 4077

References

- Alder JT, Hacksell U and Strange PG (2003) Analysis of molecular determinants of affinity and relative efficacy of a series of R- and S-2-(dipropylamino)tetralins at the 5-HT_{1A} serotonin receptor. *Br J Pharmacol* **138**:1129-39.
- Black JW and Leff P (1983) Operational models of pharmacological agonism. *Proc R Soc Lond B Biol Sci* **220**:141-62.
- Castro SW and Strange PG (1993) Differences in the ligand binding properties of the short and long versions of the D₂ dopamine receptor. *J Neurochem* **60**:372-5.
- Clarke WP and Bond RA (1998) The elusive nature of intrinsic efficacy. *Trends Pharmacol Sci* **19**:270-6.
- Colquhoun D (1998) Binding, gating, affinity and efficacy: the interpretation of structure-activity relationships for agonists and of the effects of mutating receptors. *Br J Pharmacol* **125**:924-47.
- De Lean A, Kilpatrick BF and Caron MG (1982) Dopamine receptor of the porcine anterior pituitary gland. Evidence for two affinity states discriminated by both agonists and antagonists. *Mol Pharmacol* **22**:290-7.
- De Lean A, Stadel JM and Lefkowitz RJ (1980) A ternary complex model explains the agonist-specific binding properties of the adenylate cyclase-coupled beta-adrenergic receptor. *J Biol Chem* **255**:7108-17.
- Egan C, Grinde E, Dupre A, Roth BL, Hake M, Teitler M and Herrick-Davis K (2000) Agonist high and low affinity state ratios predict drug intrinsic activity and a revised ternary complex mechanism at serotonin 5-HT_{2A} and 5-HT_{2C} receptors. *Synapse* **35**:144-50.

MOL 4077

Florio VA and Sternweis PC (1989) Mechanisms of muscarinic receptor action on Go in reconstituted phospholipid vesicles. *J Biol Chem* **264**:3909-15.

Gardner B, Hall DA and Strange PG (1996) Pharmacological analysis of dopamine stimulation of [35S]-GTP gamma S binding via human D2short and D2long dopamine receptors expressed in recombinant cells. *Br J Pharmacol* **118**:1544-50.

Gardner B and Strange PG (1998) Agonist action at D2(long) dopamine receptors: ligand binding and functional assays. *Br J Pharmacol* **124**:978-84.

Gardner BR, Hall DA and Strange PG (1997) Agonist action at D2(short) dopamine receptors determined in ligand binding and functional assays. *J Neurochem* **69**:2589-98.

Gettys TW, Sheriff-Carter K, Moomaw J, Taylor IL and Raymond JR (1994) Characterization and use of crude alpha-subunit preparations for quantitative immunoblotting of G proteins. *Anal Biochem* **220**:82-91.

Hausdorff WP, Hnatowich M, O'Dowd BF, Caron MG and Lefkowitz RJ (1990) A mutation of the beta 2-adrenergic receptor impairs agonist activation of adenylyl cyclase without affecting high affinity agonist binding. Distinct molecular determinants of the receptor are involved in physical coupling to and functional activation of Gs. *J Biol Chem* **265**:1388-93.

Jinnah HA, Page T and Friedmann T (1993) Brain purines in a genetic mouse model of Lesch-Nyhan disease. *J Neurochem* **60**:2036-45.

Kearn CS, Greenberg MJ, DiCamelli R, Kurzawa K and Hillard CJ (1999) Relationships between ligand affinities for the cerebellar cannabinoid receptor CB1 and the induction of GDP/GTP exchange. *J Neurochem* **72**:2379-87.

MOL 4077

- Kenakin T (2002) Drug efficacy at G protein-coupled receptors. *Annu Rev Pharmacol Toxicol* **42**:349-79.
- Lefkowitz RJ, Cotecchia S, Samama P and Costa T (1993) Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins. *Trends Pharmacol Sci* **14**:303-7.
- Lowry O, Rosebrough N, Farr A and Randall R (1951) Protein measurement with the folin phenol reagent. *J Biol Chem* **193**:265-275.
- McLoughlin DJ and Strange PG (2000) Mechanisms of agonism and inverse agonism at serotonin 5-HT_{1A} receptors. *J Neurochem* **74**:347-57.
- Mosser VA, Amana IJ and Schimerlik MI (2002) Kinetic analysis of M2 muscarinic receptor activation of G_i in Sf9 insect cell membranes. *J Biol Chem* **277**:922-31.
- Otero AD (1990) Transphosphorylation and G protein activation. *Biochem Pharmacol* **39**:1399-404.
- Payne SL, Johansson AM and Strange PG (2002) Mechanisms of ligand binding and efficacy at the human D₂(short) dopamine receptor. *J Neurochem* **82**:1106-17.
- Raymond JR, Olsen CL and Gettys TW (1993) Cell-specific physical and functional coupling of human 5-HT_{1A} receptors to inhibitory G protein alpha-subunits and lack of coupling to G_s alpha. *Biochemistry* **32**:11064-73.
- Ross EM (1989) Signal sorting and amplification through G protein-coupled receptors. *Neuron* **3**:141-52.

MOL 4077

- Ross EM and Wilkie TM (2000) GTPase-activating proteins for heterotrimeric G proteins: regulators of G protein signaling (RGS) and RGS-like proteins. *Annu Rev Biochem* **69**:795-827.
- Samama P, Cotecchia S, Costa T and Lefkowitz RJ (1993) A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model. *J Biol Chem* **268**:4625-36.
- Seifert R, Wenzel-Seifert K, Gether U and Kobilka BK (2001) Functional differences between full and partial agonists: evidence for ligand-specific receptor conformations. *J Pharmacol Exp Ther* **297**:1218-26.
- Strange PG (1999) G-protein coupled receptors: conformations and states. *Biochem Pharmacol* **58**:1081-8.
- Tota MR and Schimerlik MI (1990) Partial agonist effects on the interaction between the atrial muscarinic receptor and the inhibitory guanine nucleotide-binding protein in a reconstituted system. *Mol Pharmacol* **37**:996-1004.
- Van Koppen CJ, Sell A, Lenz W and Jakobs KH (1994) Deletion analysis of the m4 muscarinic acetylcholine receptor. Molecular determinants for activation of but not coupling to the Gi guanine-nucleotide-binding regulatory protein regulate receptor internalization. *Eur J Biochem* **222**:525-31.
- Waelbroeck M (2001) Activation of guanosine 5'-[gamma-(35)S]thio-triphosphate binding through M(1) muscarinic receptors in transfected Chinese Hamster ovary cell membranes: 2. Testing the "two-states" model of receptor activation. *Mol Pharmacol* **59**:886-93.
- Watson J, Collin L, Ho M, Riley G, Scott C, Selkirk JV and Price GW (2000) 5-HT(1A) receptor agonist-antagonist binding affinity difference as a measure

MOL 4077

of intrinsic activity in recombinant and native tissue systems. *Br J Pharmacol* **130**:1108-14.

Weiss JM, Morgan PH, Lutz MW and Kenakin TP (1996) The cubic ternary complex receptor-occupancy model. III. resurrecting efficacy. *J Theor Biol* **181**:381-97.

Wilson J, Lin H, Fu D, Javitch JA and Strange PG (2001) Mechanisms of inverse agonism of antipsychotic drugs at the D(2) dopamine receptor: use of a mutant D(2) dopamine receptor that adopts the activated conformation. *J Neurochem* **77**:493-504.

Zhong H, Wade SM, Woolf PJ, Linderman JJ, Traynor JR and Neubig RR (2003) A spatial focusing model for G protein signals. Regulator of G protein signaling (RGS) protein-mediated kinetic scaffolding. *J Biol Chem* **278**:7278-84.

MOL 4077

Footnote

We thank the Wellcome Trust for financial support.

MOL 4077

Figure Legends

Figure 1 G protein cycle. See text for discussion of the constituent steps.

Figure 2. Stimulation by agonists of [³⁵S]GTPγS binding to membranes of CHO cells expressing the D₂ dopamine receptor: effects of GDP on maximal [³⁵S]GTPγS binding and relative agonist efficacy for full and partial agonists. Membranes were incubated with either 0.1nM (A, B) or 10 nM (C, D) [³⁵S]GTPγS in the presence of varying concentrations of GDP and maximal stimulatory concentrations of agonist as described in the Methods section. Agonist concentrations were 100 μM dopamine (■), 10 μM (±)-7-OH-DPAT (◆), 10 μM NPA (▲) and 1 mM para-tyramine (▼). Incubation times were 30 min (0.1 nM [³⁵S]GTPγS) or 3 min (0.1 nM [³⁵S]GTPγS). Data shown are mean ± sem, of 3-4 experiments with basal subtracted.

* $P < 0.05$; ** $P < 0.01$ for comparison of relative efficacy between 0.1 nM and 10 nM [³⁵S]GTPγS (B and D) or for comparison of pmol.mg⁻¹ [³⁵S]GTPγS binding from that observed at 300 nM GDP (A and C).

MOL 4077

Figure 3. Stimulation of [³⁵S]GTPγS binding by agonists in the absence of GDP. Membranes of CHO cells expressing the D₂ dopamine receptor were incubated with either dopamine (■), (±)7-OH-DPAT (◆), NPA (▲) or p-tyramine (▼) in the presence of 0.1 nM [³⁵S]GTPγS and the absence of added GDP as described in the Methods section. The data shown represent a single experiment replicated as in Table 1. Concentration/response curves are fitted best by sigmoidal curves with Hill coefficients of one.

Figure 4. Stimulation of [³⁵S]GTPγS binding to membranes of CHO cells expressing the D₂ dopamine receptor by agonists, effects of GDP on potency of dopamine (A,C) and p-tyramine (B,D). Membranes were incubated with either 0.1nM (A, B) or 10 nM (C, D) [³⁵S]GTPγS in the presence of varying concentrations of GDP and a range of concentrations of agonist as described in the Methods section. The data shown represent single experiments replicated as shown in Figure 5. Concentration/response curves are fitted best by sigmoidal curves with Hill coefficients of one.

MOL 4077

Figure 5. Stimulation of [³⁵S]GTPγS binding to membranes of CHO cells expressing the D₂ dopamine receptor by agonists, the relationship between GDP concentration and agonist pEC₅₀ at (A) 0.1 nM and (B) 10 nM [³⁵S]GTPγS. Concentration-response curves were constructed for dopamine (■), bromocriptine (□), NPA (▼), (±)-7-OH-DPAT (×), quinpirole (●), β-phenylethylamine (◆), m-tyramine (▲) and p-tyramine (◇) at the indicated GDP concentrations and the potency (EC₅₀) was determined as in Figure 4 and the Methods section.

Data shown are mean ± sem of 3-5 experiments performed in triplicate. For statistical analysis see Table 3.

MOL 4077

Tables

TABLE 1. Stimulation of [³⁵S]GTPγS binding by agonists in the absence of GDP.

	pEC ₅₀ (EC ₅₀ , nM)	Stimulation of [³⁵ S]GTPγS binding (pmol.mg protein ⁻¹)
dopamine	7.84 ± 0.04 (14)	0.056 ± 0.012
(±)-7-OH-DPAT	8.21 ± 0.09 (6)	0.067 ± 0.013
NPA	10.00 ± 0.12 (0.1)	0.067 ± 0.018
p-tyramine	5.48 ± 0.10 (3310)	0.057 ± 0.013

The stimulation of [³⁵S]GTPγS binding by agonists was determined as described in the Methods section in the absence of GDP. pEC₅₀ values were determined and maximal effects expressed as the stimulation of [³⁵S]GTPγS binding over basal in pmol.mg protein⁻¹. Maximal effects were not statistically different (*P*>0.05)

MOL 4077

TABLE 2. The binding of agonists to D₂ dopamine receptors

	pK _h (K _h , nM)	pK _l (K _l , nM)	%R _h	K _l /K _h
dopamine	7.61 ± 0.01 (24.5)	5.55 ± 0.02 (2820)	49 ± 7	115
(±)-7-OH-DPAT	7.83 ± 0.15 (14.8)	6.49 ± 0.08 (320)	42 ± 4	22
NPA	10.03 ± 0.16 (0.09)	8.21 ± 0.12 (6.2)	35 ± 5	69
p-tyramine	5.10 ± 0.25 (7940)	3.61 ± 0.07 (245000)	40 ± 10	31

Agonist binding was determined in competition versus [³H]spiperone (~0.3 nM) in a buffer containing Na⁺ (100mM) as described in the Methods section. Data were fitted best by a two-binding site model and analysed by non-linear regression to derive dissociation constants for the higher (K_h) and lower (K_l) affinity sites and the percentage of receptors in the high affinity state (%R_h). Data represent mean ± sem of 3 experiments performed in triplicate.

TABLE 3. The relationship between GDP concentration and agonist potency

	Slope	
	0.1 nM [³⁵ S]GTPγS	10 nM [³⁵ S]GTPγS
dopamine	-0.31 ± 0.06	-0.47 ± 0.04
bromocriptine	0.02 ± 0.04	ND
(±)-7-OH-DPAT	-0.31 ± 0.04	-0.20 ± 0.07
NPA	-0.37 ± 0.06	-0.55 ± 0.03
quinpirole	-0.29 ± 0.05	ND
β-phenylethylamine	-0.24 ± 0.05	ND
p-tyramine	-0.32 ± 0.03	-0.04 ± 0.09
m-tyramine	-0.30 ± 0.04	ND

The slopes of the lines shown in Fig 5 were determined by linear regression. ND = Not determined.

In the experiments at 0.1 nM [³⁵S]GTPγS all compounds had similar slopes (*P*>0.05) with the exception of bromocriptine.

In the experiments at 10nM [³⁵S]GTPγS, the slope for 7-OH-DPAT was different from 0 (*P*<0.05) and different from slopes for other compounds (*P*<0.05); the slope for p-tyramine was not different from 0 (*P*>0.05), the slopes for dopamine and NPA were different from slopes for 7-OH-DPAT and p-tyramine .

Slopes for NPA or dopamine were different at the two concentrations of [³⁵S]GTPγS.




Figure 1

Figure 2




Figure 3




Figure 4a, b




Figure 4c,d





Figure 5

