

MOLPHARM/2004/008409

Organotin compounds promote adipocyte differentiation as agonists of the peroxisome proliferator-activated receptor (PPAR) γ / retinoid X receptor (RXR) pathway.

Tomohiko Kanayama, Naoki Kobayashi, Satoru Mamiya, Tsuyoshi Nakanishi,
Jun-ichi Nishikawa

Department of Environmental Biochemistry (T.K., N.K., S.M., J.N.) and Department of Toxicology (T.N.), Graduate School of Pharmaceutical Sciences, Osaka University, Osaka, Japan

MOLPHARM/2004/008409

<Running title>

a) Running title: Organotins are activators of PPAR γ /RXR in adipogenesis

b) Corresponding Author: Jun-ichi Nishikawa

Department of Environmental Biochemistry, Graduate School of Pharmaceutical

Sciences, Osaka University, 1-6 Yamada-oka, Suita, Osaka 565-0871, Japan

Tel: 81-6-6879-8241, Fax: 81-6-6879-8244, E-mail: nisikawa@phs.osaka-u.ac.jp

c) Number of text pages: 30

Number of table: 1

Number of figures: 6

Number of references: 38

Number of words in *Abstract*: 113

Number of words in *Introduction*: 426

Numbers of words in *Discussion*: 1045

d) ABBREVIATIONS: GST, glutathione-S-transferase; IPTG, isopropyl- β -D-thiogalactopyranoside; PPAR, peroxisome proliferator-activated receptor; TIF2, transcriptional intermediary factor 2.

MOLPHARM/2004/008409

Abstract

Nuclear receptors play important roles in the maintenance of the endocrine system, regulation of organ differentiation, and fetal development. Endocrine disruptors exert their adverse effects by disrupting the endocrine system via various mechanisms. To assess the effects of endocrine disruptors on nuclear receptors, we developed a high-throughput method for identifying activators of nuclear receptors. Using this system, we found that triphenyltin and tributyltin were activators of peroxisome proliferators-activated receptor (PPAR) γ and retinoid X receptor. Because PPAR γ is a master regulator of adipocyte differentiation, we assessed the effect of organotin compounds on preadipocyte, 3T3-L1 cells. We found that organotin compounds stimulated differentiation of 3T3-L1 cells as well as expression of adipocyte marker genes.

MOLPHARM/2004/008409

Introduction

An endocrine disruptor is an exogenous substance or mixture that alters functions of the endocrine system and consequently causes adverse health effects in an intact organism, its progeny, or (sub)populations (WHO, 1996). Many naturally occurring and synthetic compounds, including DDT and its metabolites, PCBs, and some alkylphenols, have hormonal activities (Sohoni et al., 1998; Nishihara et al., 2000; Gray et al., 2001; Sanderson et al., 2002). Although the levels of natural hormones are precisely regulated metabolically, synthetic chemicals elude this regulation to stimulate organs by mechanisms different from those of natural hormones.

The importance of nuclear receptors in endocrine function has been well established by many studies. The human genome contains at least 48 members of the nuclear receptor family (Chawla et al., 2001), and various chemicals bind to nuclear receptors and influence the expression of target genes (Blair et al., 2000; Sultan et al., 2001). To evaluate the effects of numerous synthetic chemicals on many nuclear receptors, we developed the CoA-BAP system, a high-throughput method for identifying nuclear receptor ligands (Kanayama et al., 2003). In the present study, we applied the CoA-BAP system to the evaluation of 16 human nuclear receptors and 40 suspected endocrine disruptors. We found that organotin compounds such as triphenyltin (TPT) and tributyltin (TBT) strongly activated retinoid X receptor (RXR) and PPAR γ .

Organotin compounds have been used as agricultural fungicides, rodent

MOLPHARM/2004/008409

repellents, and molluscicides and in antifouling paints for ships and fishing nets (Piver, 1973; Fent, 1996). These widespread uses have resulted in the release of increasing amounts of organotins into the environment. Although the toxicity of organotins has been reviewed extensively (Boyer, 1989), the molecular target of organotins has not yet been identified.

Here we show that TPT and TBT are high-affinity ligands for RXR and PPAR γ . Organotin compounds act as agonists of both RXR α and PPAR γ in mammalian reporter gene assays and induce the expression of PPAR γ target genes. PPAR γ forms a heterodimer with RXR and binds to a defined DNA sequence in the promoter region of target genes (Mangelsdorf et al., 1995). PPAR γ is activated by a variety of fatty acids and a class of synthetic antidiabetic agents, the thiazolidinediones (Lehmann et al., 1995). PPAR γ serves as an essential regulator for adipocyte differentiation and lipid storage in mature adipocytes (Tontonoz et al., 1994). In light of these previous findings, we evaluated the effects of TPT and TBT on adipogenesis and found that organotins stimulate the differentiation of preadipocyte 3T3-L1 cells to adipocytes. Our data suggest that organotins exert their toxic effects through activation of the PPAR γ /RXR signaling pathway.

MOLPHARM/2004/008409

Materials and Methods

Plasmids.

The ligand-binding domains (LBDs) of the human nuclear receptors PPAR α (codons 168–468; GenBank accession no., L02932), PPAR γ 1(177–477; L40904), PPAR δ (139–441; L07592), LXR α (167–447; U22662), and LXR β (155–461; U07132) were amplified by RT-PCR from human liver mRNA as the template; the LBDs of human FXR (193–472; U68233) and human ERR γ (194–458; AF094518) were amplified similarly from human kidney mRNA, and that of human ERR β (195–434; AF094517) was amplified from human testis mRNA. The DNA sequences of the amplified fragments were confirmed by sequencing after subcloning into pGEX-4T (Amersham Biosciences, Piscataway, NJ). The expression vectors for the human nuclear receptors ER α/β , TR α , RAR α/γ , RXR α/γ , VDR, and hTIF2 were described previously (Kanayama et al., 2003). For expression in mammalian culture cells, the LBD of hRXR α was fused to the C-terminal end of the GAL4 DNA binding domain (amino acids 1–97) in the pBK-CMV expression vector (Stratagene, La Jolla, CA). The expression plasmid of (GAL4-DBD)-PPAR γ (pM-mPPAR γ 1) and the luciferase reporter plasmid p4xUAS-tk-luc (Kamei et al., 2003) were kind gifts from Dr. Y. Kamei.

Chemical reagents.

Diethyl phthalate, triphenyltin chloride, nitrofen, 4-nonylphenol, octachlorostyrene,

MOLPHARM/2004/008409

permethrin, triphenylmethane and triphenylethylene were purchased from Kanto Chemical (Tokyo, Japan). Amitrole, 2,4-dichlorophenoxy acetic acid, 1,2-dibromo-3-chloropropane, γ -HCH (lindane), pentachlorophenol, dihexyl phthalate, di-*n*-pentyl phthalate, dipropyl phthalate, 2,4-dichlorophenol, 4-nitrotoluene, and bisphenol A were purchased from Tokyo Kasei (Tokyo, Japan). Chenodeoxycholic acid, 1 α ,25-dihydroxy cholecalciferol, lithocholic acid, *all-trans* retinoic acid, 9-*cis* retinoic acid, and 3, 3', 5-triiodo-L-thyronine were purchased from Sigma (St. Louis, MO). 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂, rosiglitazone and TO-901317 were purchased from Cayman Chemical (Ann Arbor, MI). GW501516 was purchased from Calbiochem (San Diego, CA). All other chemicals were purchased from Wako Pure Chemical (Osaka, Japan). The 40 chemicals tested and the abbreviations used for them are listed in Table 1.

Preparation of proteins.

The histidine-tagged fusion protein hTIF2 NID-BAP, in which the nuclear receptor interaction domain of TIF2, was ligated to the bacterial alkaline phosphatase, was expressed in *Escherichia coli* BL21 (DE3) cells and purified on Ni-nitrilotriacetic acid agarose resin (QIAGEN, Valencia, CA). Except for LXR α/β and FXR, the GST fusion proteins were expressed in the *E. coli* BL21 (DE3) pLysS cells; LXR α/β and FXR were expressed in *E. coli* JM109 pRIL cells. The GST fusion proteins were purified by using Glutathione Sepharose 4B (Amersham Biosciences).

MOLPHARM/2004/008409

CoA-BAP system.

Detection of ligand-dependent interaction between nuclear receptors and TIF2 was carried out as described previously (Kanayama et al., 2003) but with slight modification. Briefly, 2 µg of nuclear receptor protein diluted in 100 µl of carbonate buffer (100 mM NaHCO₃, pH 8.4) was incubated in the well of a 98-well polystyrene microtiter plate (MaxiSorp, Nalge-Nunc International, Rochester, NY) at 4°C overnight. The plate was washed three times with 120 µl of buffer A (20 mM Tris-HCl, 100 mM KCl, 0.25 mM EDTA, 5% glycerol, 0.5 mM DTT, 0.05% Tween 20; pH 7.4), and then 100 µl of TIF2-BAP fusion protein (30 µg/ml) in buffer A was added to a well with the test chemical. After 1 h incubation at 4°C, the plate was washed three times with 120 µl of buffer B (50 mM Tris-HCl, 100 mM KCl, 5 mM MgCl₂, 0.10% Nonidet P-40; pH 7.2). The enzyme reaction was started by the addition of 100 µl of substrate solution (10 mM *p*-nitrophenyl phosphate in 100 mM Tris-HCl; pH 8.0). After incubation at 37°C for 30 to 90 min, the reaction was stopped by addition of 25 µl of 0.5 N NaOH. Finally, the absorbance at 405 nm was measured with a plate reader (MultiskanJX, Thermo LabSystems, Helsinki, Finland).

Cell culture.

Mouse 3T3-L1 (Dainippon Pharmaceutical, Osaka, Japan) and mouse NIH-3T3 (clone 5611, JCRB0615, Japanese Cancer Research Resources Bank) fibroblasts

MOLPHARM/2004/008409

were maintained at 37°C in 5% CO₂ in Dulbecco's modified Eagle's Medium (DMEM) (Nissui Pharm, Tokyo, Japan) supplemented with 10% calf serum (ICN Biomedicals, Aurora, OH). Mouse F9 embryonic carcinoma cells were maintained in 5% CO₂ at 37°C in DMEM supplemented with 10% fetal bovine serum (FBS) (ICN Biomedicals).

Transient transfection assays.

One day before transfection, 1 x 10⁵ cells were plated in a 35-mm dish containing phenol red-free Eagle's Minimum Essential Medium (Nissui Pharm) supplemented with 10% charcoal/dextran-treated FBS. The cells were transfected by lipofection using FuGENE 6 Transfection Reagent (Roche Diagnostics, Indianapolis, IN) with pBK-CMV-GAL4-hRXR α or pM-mPPAR γ 1 (300 ng/dish), p4xUAS-tk-luc (600 ng/dish), and RSV- β gal (100 ng/dish). Fresh medium with or without test chemical was added the day after transfection. After incubation for 24 h, cells were harvested and assayed for luciferase and β -galactosidase activity.

Adipocyte differentiation assays.

Mouse 3T3-L1 preadipocyte cells were used for the differentiation experiments. The day after the cells reached confluence, the medium was replaced with DMEM containing 10% FBS, 10 μ g/ml insulin, 0.5 mM 3-isobutyl-1-methylxanthine (Mix), and 1 μ M dexamethasone (Dex). At the same time, the cells were treated with a test chemical (rosiglitazone, 9-*cis* retinoic acid, or an organotin compound). After 60

MOLPHARM/2004/008409

hours, the medium was replaced with DMEM containing 10% FBS, 5 µg/ml insulin, and the test chemical. After 6 days, cells were fixed with 4% paraformaldehyde and stained with 0.5% Oil Red O. The amount of triglyceride was determined by Triglyceride E Test (Wako Pure Chemical).

RNA isolation, Northern blotting and RT-PCR analyses.

The 3T3-L1 cells were grown in DMEM containing 10% calf serum. The day after the cells became confluent, they were treated with vehicle (DMSO) only, rosiglitazone (Rosi), TPT, or TBT in DMEM containing 10% FBS and 5 µg/ml insulin. The cells were harvested at various times after treatment, and total RNA was isolated using TRIzol (Invitrogen, Carlsbad, CA). For Northern blot analyses, 25 µg of total RNA was electrophoresed through a 1% agarose gel containing 2% formaldehyde and then transferred to a Hibond-N+ nylon membrane (Amersham Biosciences). The filter was hybridized with each probe, which was labeled with [α -³²P] dCTP by using a random labeling kit (TaKaRa, Shiga, Japan). For RT-PCR, cDNA was synthesized using ReverTra Ace (Toyobo, Osaka, Japan), and PCR was performed using AmpliTaq Gold (Applied Biosystems). The primers used for amplification of the aP2 gene (a marker for adipocyte differentiation) were 5'-AAAATGTGTGATGCCTTTGTGGG-3' and 5'-TCATGCCCTTTCATAAACTCTTGTGG-3'.

MOLPHARM/2004/008409

Results

Application of CoA-BAP system to endocrine disruptors.

Reproductive abnormalities in wildlife can be associated with exposure to environmental pollutants capable of mimicking the action of natural hormones. Because the nuclear receptors of intrinsic hormone systems are likely to be targets of industrial chemicals, information on their ability to bind these chemicals is valuable for environmental risk assessment. To determine whether suspected endocrine disruptors can bind to members of the nuclear receptor family, we constructed assay systems for human nuclear receptors, including ER α/β , RAR α/γ , TR α , VDR, RXR α/γ , PPAR $\alpha/\gamma/\delta$, FXR, LXR α/β , and ERR α/γ , on the basis of the previously described CoA-BAP system (Kanayama et al., 2003). The cognate ligand for each nuclear receptor enhanced alkaline phosphatase activity in a dose-dependent manner (Fig. 1). In the ERR systems, 4-hydroxy tamoxifen-dependent dissociations between ERR and coactivator were observed, as previously reported (Coward et al., 2001; Tremblay et al., 2001).

Using these systems, we evaluated 40 suspected endocrine disruptors (Table 1) recognized by various organizations (e.g., WHO, Ministry of the Environment in Japan). The effects of the tested chemicals on the interaction between nuclear receptors and TIF2 (Fig. 2) suggest that several compounds possess agonistic activities for multiple receptors simultaneously. Butyl benzyl phthalate (no. 8), hexachlorocyclohexane (no. 26), maneb (no. 22), mancozeb (no.

MOLPHARM/2004/008409

23), and alkylphenols (nos. 10 and 11) were weakly agonistic for multiple receptors, including estrogen receptor (ER). One intriguing finding was that the effect of TBT on RXR α was as strong as that of its endogenous ligand, 9-*cis* retinoic acid (Fig. 3), and the agonist effect of TPT on PPAR γ was as strong as that of its well-known ligand, Rosi (Fig. 3). The EC₅₀ values of TBT on RXR α (7.4×10^{-8} M) and TPT on PPAR γ (9.5×10^{-8} M) were almost same as those of 9-*cis* retinoic acid (4.3×10^{-8} M) and Rosi (1.1×10^{-7} M), respectively. As triphenylmethane and triphenylethylene were not agonistic for RXR α and PPAR γ , the tin moiety was important for activity (Fig. 3).

Organotin compounds potentiated transactivation by RXR and PPAR γ .

The observations that organotin compounds enhanced the protein–protein interaction between the coactivator TIF2 and RXR α or PPAR γ suggested to us that these compounds activate transcription via these receptors. To confirm the results we obtained from the CoA-BAP system, we performed a reporter gene assay in mammalian culture cells using an expression vector for (GAL4-DBD)-RXR α or (GAL4-DBD)-PPAR γ and a reporter plasmid containing the luciferase gene along with GAL4 upstream activating sequence. Both TPT and TBT induced the transactivation function of RXR α or PPAR γ in a dose-dependent manner (Fig. 4). The effectiveness of these organotin compounds was comparable to that of known ligands. In addition, dibutyltin chloride, a TBT metabolite *in vivo*, also activated reporter activity in the PPAR γ system (data not shown).

MOLPHARM/2004/008409

Induction and promotion of adipocyte differentiation by organotin compounds in 3T3-L1 cells.

Recent studies indicate that PPAR γ plays a central role in adipocyte gene expression and differentiation (Tontonoz et al., 1994). PPAR γ is abundantly expressed in adipocytes, and its ligands induce the efficient conversion of fibroblastic cells to adipocytes, as measured by induction of adipocyte-specific genes and lipid accumulation (Lehmann et al., 1995). If organotin compounds can function as activators for PPAR γ /RXR *in vivo*, these compounds likely induce adipocyte differentiation. To investigate this possibility, we treated 3T3-L1 cells with TPT or TBT in two types of differentiation medium, a complete differentiation medium that contained the inducers Mix, Dex, insulin, and FBS, and an incomplete differentiation medium that lacked Mix and Dex. Although insulin is not always necessary for induction of differentiation, it efficiently enhances adipocyte development. Adipocyte differentiation was confirmed by staining with Oil Red O for lipid droplet accumulation. As expected, treatment of 3T3-L1 cells with either TPT or TBT in complete differentiation medium promoted adipocyte differentiation as well as did Rosi (Fig. 5, a–d). Even in incomplete differentiation medium, addition of organotin compounds induced adipocyte differentiation in contrast with the lack of induction after treatment with vehicle only (Fig. 5, e–h). Moreover, mRNA expression of the adipocyte differentiation marker aP2 was induced in a dose-dependent manner by addition of

MOLPHARM/2004/008409

organotin compounds (Fig. 6a). PPAR γ mRNA also was induced during the differentiation process (Fig. 6a), in agreement with the results of a previous study (Tontonoz et al., 1994). Induction of aP2 mRNA expression occurred late in adipogenesis (Fig. 6b), and organotin-treated cells demonstrated accumulation of triglyceride (Fig. 6c). Taken together, these data provide strong evidence that the two organotin compounds TPT and TBT can function as inducers of adipocyte differentiation through PPAR γ .

Discussion

Our study was designed to evaluate the effects of suspected endocrine disruptors on various nuclear receptors. The data show that several compounds have simultaneous effects on multiple nuclear receptors. In particular, organotin compounds (e.g., TBT and TPT) showed strong effects on RXR or PPAR γ , at levels comparable to those of 9-cis retinoic acid, an endogenous RXR ligand, and rosiglitazone, a known agonist of PPAR γ . In CoA-BAP systems, TBT showed strong effect on protein-protein interaction between RXR α and TIF2, but TPT did slight effect (Fig. 3a). TPT showed strong effect on protein-protein interaction between PPAR γ and TIF2, but TBT did not (Fig. 3b). On the contrary, when tested the transactivation assay, both TBT and TPT activated not only RXR α but also PPAR γ (Fig. 4). This discrepancy might reflect the diversity of coactivators. To date, many coactivators have been identified as nuclear receptor-interacting proteins. These

MOLPHARM/2004/008409

coactivators are supposed to have cell- or tissue-specific functions *in vivo* (Smith et al., 2004). In addition, PPAR γ reportedly change its interaction partners dependent on ligands (Kodera et al., 2000). We used only TIF2 in CoA-BAP system, while cells used for transactivation assays have many coactivators. The discrepancy of results from CoA-BAP systems and tranactivation assays might be explained by this difference of coactivators. Because *in vitro* screening methods tend to produce false-positive or -negative results like this, positive compounds should be further examined by other studies in a physiological context. Accordingly, we examined the effects of organotin compounds on the transcriptional regulation and adipogenesis which is a famous physiological event related to PPAR γ /RXR pathway.

Expoaure of rats *in utero* to TBT induces a dramatic increase in the incidence of low-birth-weight fetuses because of maternal hypothyroidisim (Adeeko et al., 2003). Further, the RXR agonist bexarotene causes clinically significant hypothyroidism in patients with cutaneous T-cell lymphoma (Duvic et al., 2001), and experimental exposure of rats to LG100268 (a selective RXR agonist) induces the acute phase of hypothyroidism (Liu et al., 2002). The similarities between the toxicities of TBT and selective RXR agonists suggested to us that at least some of the toxic effects of oraganotin compounds are mediated by RXR.

Most of the toxic effects of oraganotin compounds on sexual development and reproductive function have been documented in mollusks (Matthiessen et al., 1998). In gastropods, TBT and TPT cause imposex (Morcillo et al., 1999), an

MOLPHARM/2004/008409

irreversible syndrome in which male genital tracts (mainly a penis and a vas deferens) are imposed on female organisms (Smith, 1971). Although the physiological functions of organotin compounds have been studied extensively, the molecular target of organotin compounds had been unclear. To this end, we found that TPT and TBT were agonists for RXR and PPAR γ . It has been thought that the sexual toxicity of organotin compounds results from increased androgen levels because of inhibition of the aromatase enzyme complex that catalyzes conversion of androgen to estrogen. This enzyme complex consists of microsomal CYP19 and the reduced form of the flavoprotein nicotinamide adenine dinucleotide phosphate reductase. TBT-induced imposex in neogastropods reportedly is mediated by inhibition of aromatase (Bettin et al., 1996), and TBT inhibits the catalytic activity of aromatase derived from transfected cells (Heidrich et al., 2001; Cooke, 2002). However, the effective concentrations of enzyme inhibition were relatively high (above 10^{-6} M). In this study, we found that TBT and TPT induced the transactivation function of RXR α and PPAR γ at 10^{-8} M. It is reasonable that the effective concentration on gene expression was different from that on enzyme inhibition. In consistent with this, Nakanishi *et al.* demonstrated that 10^{-8} M of TBT or TPT induced hCG or aromatase activity along with mRNA expression in placental cells (Nakanishi et al., 2002). In ovarian granulosa cells, 20 ng/ml (about 6×10^{-8} M) of TBT or TPT suppress the P450_{aroma} gene expression (Saitoh et al., 2001). We have to consider about toxicities of organotin compounds in distinguishing the low-dose effect from

MOLPHARM/2004/008409

high-dose effect. Recently, we reported that RXR plays an important role in the development of gastropod imposex, by showing the cloning of RXR homolog from marine gastropod, binding of organotins to that receptor, and imposex induction by injection of RXR ligand, 9-cis retinoic acid (Nishikawa et al., *in press*). Gastropod imposex is known to be typically induced by very low concentrations of TBT and/or TPT (Bryan et al., 1986; Gibbs et al., 1986; Horiguchi et al., 1997). Although it has been theorized that organotins increases androgen levels through inhibition of aromatase activity and/or a suppression of androgen excretion, the inhibitory concentration of organotins is not enough low for explaining imposex induction. The low-dose effects are likely to be mediated by receptors. However, the study about organotin effects in mammals is still important, because the compositions of nuclear receptor family members are quite different between vertebrates and invertebrates (Escriva et al., 1997; Laudet, 1997). For example, there are no known homologues of steroid hormone receptors in *Drosophila* and *Caenorhabditis elegans* genome, and the group members of TR, RAR, VDR and PPAR appear to be late acquisitions during the evolution of the superfamily. Therefore, we examined the effects of suspected endocrine disruptors on human nuclear receptor family members. As a result, PPAR γ was identified as a new target molecule of organotin compounds in addition to RXR. This finding might introduce new insights in physiological functions of organotin compounds in mammals.

We were surprised to find that organotin compounds were high-affinity

MOLPHARM/2004/008409

ligands for RXR and PPAR γ . Until recently, it had been thought that among synthetic compounds, only hormone analogs could bind hormone receptors, because the relationships between hormones and their cognate receptors are very specific. However, some industrial chemicals do have unexpected effects on hormone receptors. Nuclear receptors are the likely targets, because their intrinsic ligands are fat-soluble, low-molecular-weight agents, as are the environmental pollutants. In fact, organotin compounds promote the adipocyte differentiation as agonists for PPAR γ /RXR. Recently, the ligands of PPAR γ and RXR are expected for antidiabetic agents, but they have some side effects at the same time (Mukherjee et al., 1977; Yki-Jarvinen, 2004). Although they may be good medicines in the case of using under doctor's control, wild lives are exposed to man-made chemicals in uncontrolled manner. It is possible that TBT and TPT cause adverse health effects on the organisms by disturbing the endocrine process mediated by PPAR γ /RXR.

ACKNOWLEDGMENTS

We are grateful to Dr. Y. Kamei (National Institute of Health and Nutrition, Japan) for providing the GAL4-reponsive reporter plasmid and pM-mPPAR γ 1.

MOLPHARM/2004/008409

References

- Adeeko A, Forsyth DS, Casey V, Cooke GM, Barthelemy, J Cyr DG, Trasler JM, Robair B and Hales BF (2003) Effects of in utero tributyltin chloride in the rat on pregnancy outcome. *Toxicol Sci* **74**: 407-415.
- Bettin C, Oehlmann J and Stroben E (1996) TBT-induced imposex in marine neo-gastropods is mediated by an increasing androgen level. *Hylgol Meeresunters* **50**: 299-317.
- Blair RM, Fang H, Branham WS, Hass BS, Dial SL, Moland CL, Tong, W Shi L, Perkins R and Sheehan DM (2000) The estrogen receptor relative binding affinities of 188 natural and xenochemicals: structural diversity of ligands. *Toxicol Sci* **54**: 138-153
- Boyer IJ (1989) Toxicity of dibutyltin, tributyltin and other organotin compounds to human and to experimental animals. *Toxicology* **55**: 253-298
- Bryan GW, Gibbs PE, Hummerstone LG and Burt GR (1986) The decline of the gastropod *Nucella lapillus* around south-west England: evidence for the effect of tributyltin from anti-fouling paints. *J Mar Biol Assoc UK* **66**: 611-640
- Chawla A, Repa JJ, Evans RM and Mangelsdorf DJ (2001) Nuclear receptors and lipid physiology: opening the X-files. *Science* **294**: 1866-1870.
- Cooke GM (2002) Effect of organotins on human aromatase activity in vitro. *Toxicol Lett* **126**: 121-130.
- Coward P, Lee D., Hull MV and Lehmann JM (2001) 4-Hydroxytamoxifen binds to

MOLPHARM/2004/008409

and deactivates the estrogen-related receptor gamma. *Proc Natl Acad Sci USA* **98**: 8880-8884.

Duvic M, Hymes K, Heald P, Breneman D, Martin AG, Myskowski P, Crowley C and Yocum RC (2001) Bexarotene is effective and safe for treatment of refractory advanced-stage cutaneous T-cell lymphoma: multinational phase II-III trial results. *J Clin Oncol* **19**: 2456-2471

Escriva H, Safi R, Hanni C, Langlois MC, Saumitou-Laprade P, Stehelin D, Capron A, Pierce R and Laudet V (1997) Lignad binding was acquired during evolution of nuclear receptors. *Proc Natl Acad Sci USA* **94**: 6803-6808.

Fent K (1996) Ecotoxicology of organotin compounds. *Crit Rev Toxicol* **26**: 1-117.

Gibbs PE and Bryan GW (1986) Reproductive failure in populations of the dog-whelk, *Nucella lapillus*, caused by imposex induced by tricetyl tin from antifouling paints. *J Mar Biol Assoc UK* **66**: 767-777

Gray LE, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, Veeramachaneni DN, Wilson V, Price M, Hotchkiss A, Oriando E and Guillette L (2001) Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update* **7**: 248-264.

Heidrich D D, Steckelbroek S and Klingmuller D (2001) Inhibition of human cytochrome P450 aromatase activity by butyltins. *Steroids* **66**: 763-769.

Horiguchi T, Shiraishi H, Shimizu M and Morita M (1997) Effects of triphenyltin chloride and five other organotin compounds on the development of imposex in

MOLPHARM/2004/008409

the rock shell, *Thais clavigera*. *Pollut* **95**: 85-91.

Kamei Y, Ohizumi H, Fujitani Y, Nemoto T, Tanaka T, Takahashi N, Kawada T,

Miyoshi M, Ezaki O and Kakizuka A (2003) PPAR γ coactivator 1 β /ERR ligand 1 is an ERR protein ligand, whose expression induces a high-energy expenditure and antagonizes obesity. *Proc Natl Acad Sci USA* **100**: 12378-12383.

Kanayama T, Mamiya S, Nishihara T and Nishikawa J (2003) Basis of a high-throughput method for nuclear receptor ligands. *J Biochem* **133**: 791-797.

Kodera Y, Takeyama K, Murayama A, Suzawa M, Masuhiro Y, Kato S (2000) Ligand type-specific interactions of peroxisome proliferator-activated receptor γ with transcriptional coactivators. *J Biol Chem* **275**: 33201-33204.

Laudet V (1997) Evolution of the nuclear receptor superfamily: early diversification from an ancestral orphan receptor. *J Mol Endocrinol* **19**: 207-226.

Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM and Kliewer SA (1995) An antidiabetic thiazolidinedione is a high-affinity ligand for peroxisome proliferator-activated receptor gamma. *J Biol Chem* **270**: 12953-12956.

Liu S, Ogilvie KM, Klausung K, Lawson MA, Jolley D, Li D, Bilakovics J, Pascual B, Hein N., Urcan M and Leibowitz MD (2002) Mechanism of selective retinoid X receptor agonist induced hypothyroidism in the rat. *Endocrinology* **143**: 2880-2885.

Mangelsdorf DJ and Evans RM (1995) The RXR heterodimers and orphan receptors. *Cell* **83**: 841-850.

MOLPHARM/2004/008409

Matthiessen P and Gibbs PE (1998) Critical appraisal of the evidence for tributyltin mediated endocrine disruption in mollusks. *Environ Toxicol Chem* **17**: 37-43.

Morcillo Y and Porte C (1999) Evidence of endocrine disruption in the imposex-affected gastropod *Bolinus brandaris*. *Environ Res* **81**: 349-354.

Mukherjee R, Davies PJA, Crombie DL, Bischoff ED, Cesario RM, Jow L, Hamann LG, Boehm MF, Mondon CE, Nadzan AM, Paterniti Jr JR and Heyman RA (1997) Sensitization of diabetic and obese mice to insulin by retinoid X receptor agonists. *Nature* **386**: 407-410.

Nakanishi T, Kohroki J, Suzuki S, Ishizaki J, Hiromori Y, Takasuga S, Itoh N., Watanabe Y, Utoguchi N and Tanaka K (2002) Trialkyltin compounds enhance human CG secretion and aromatase activity in human placental choriocarcinoma cells. *J Clin Endocrinol Metab* **87**: 2830-2837.

Nishihara T, Nishikawa J, Kanayama T, Dakeyama F, Saito K, Imagawa M, Takatori S, Kitagawa Y, Hori S and Utsumi H (2000) Estrogenic activities of 517 chemicals by yeast two-hybrid assay. *J Health Sci* **46**: 282-298.

Nishikawa J, Mamiya S, Kanayama T, Nishikawa T, Shiraishi F and Horiguchi T (2004) Involvement of the retinoid X receptor in the development of imposex caused by organotins in gastropod. *Environ Sci Technol* **38**: 6271-6276.

Piver WT (1973) Organotin compounds: industrial applications and biological investigation. *Environ Health Perspect* **4**: 61-79.

Saitoh M, Yanase T, Morinaga H, Tanabe M, Mu YM, Nishi Y, Nomura M, Okabe T,

MOLPHARM/2004/008409

- Goto K, Takayanagi R and Nawata H (2001) Tributyltin or triphenytin inhibits aromatase activity in human granulosa-like tumor cell line KGN. *Biochem Biophys Res Commun* **289**: 198-204.
- Sanderson JT, Boerma J, Lansbergen GW and van den Berg M (2002) Induction and inhibition of aromatase (CYP19) activity by various classes of pesticides in H295R human adrenocortical carcinoma cells. *Toxicol Appl Pharmacol* **182**: 44-54.
- Smith BS (1971) Sexuality in the American mud snail, *Nassarius obsoletus* Say. *Proc Malac Soc Lond* **39**: 377-378.
- Smith CL, O'Malley BW (2004) Coregulator function: a key to understanding tissue specificity of receptor modulators. *Endocr. Rev.* **25**: 45-71.
- Sohoni P and Sumpter JP (1998) Several environmental estrogens are also anti-androgens. *J Endocrinol* **158**: 327-339.
- Sultan C, Balaguer P, Terouanne B, Geoget V, Paris F, Jeandel C, Lumbroso S and Nicolas J (2001) Environmental xenoestrogens, antiandrogens and disorders of male sexual differentiation. *Mol Cell Endocrinol* **10**: 99-105.
- Tontonoz P, Hu E and Spiegelman BM (1994) Stimulation of adipogenesis in fibroblasts by PPAR γ 2, a lipid-activated transcription factor. *Cell* **79**: 1147-1156.
- Tremblay GB, Kunath T, Bergeron D, Lapointe L, Champigny C, Bader JA, Rossant J and Giguere V (2001) Diethylstilbestrol regulates trophoblast stem cell differentiation as a ligand of orphan nuclear ERR β . *Genes Dev* **15**: 833-838.

MOLPHARM/2004/008409

Yaki-Jarvinen H (2004) Thiazolidinediones. *N Engl J Med* **351**: 1106-1118.

MOLPHARM/2004/008409

Footnotes

- a) This work was supported by grants from the Ministry of the Environment, Japan.
- b) Address correspondence to: Dr. Jun-ichi Nishikawa, Department of Environmental Biochemistry, Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamada-oka, Suita, Osaka 565-0871, Japan. E-mail: nisikawa@phs.osaka-u.ac.jp

MOLPHARM/2004/008409

Legends for Figures

Fig. 1. Ligand-dependent interaction of nuclear receptor and TIF2 *in vitro*.

Ligand-dependent interactions between nuclear receptors and TIF2-BAP were determined as relative alkaline phosphatase (AP) activity (vertical axis). The receptor–ligands pairs tested were estrogen receptor (ER α/β) – 17 β -estradiol (E₂); retinoic acid receptor (RAR α/γ) – all-trans retinoic acid; thyroid hormone receptor (TR α) – 3,5,3'-triiodo-L-thyronine (T₃); vitamin D receptor (VDR) – 1 α , 25-dihydroxy cholecalciferol (1,25(OH)₂D₃); retinoid X receptor (RXR α/γ) – 9-cis retinoic acid; peroxisome proliferator-activated receptor (PPAR α/δ) – 15-deoxy-^{12,14} Δ -prostaglandin J₂ (PGJ₂); PPAR γ – rosiglitazone; liver X receptor (LXR α/β) – TO-901317; farnesoid X receptor (FXR) – chenodeoxy cholic acid (CDCA); and estrogen-related receptor (ERR α/γ) – 4-hydroxytamoxifen. Data shown are means \pm standard deviation of three independent experiments.

Fig. 2. Agonistic activities of suspected endocrine disruptors for various nuclear receptors.

The effects of chemicals on the interaction between nuclear receptors and the coactivator TIF2 were assessed using the CoA-BAP system. The numbers in the far-left column correspond to the chemicals listed in Table 1. The lowest effective concentrations (LOEC) of test chemicals were determined and compared with LOEC of cognate ligands shown in Fig. 1, red; 1~10 times as much

MOLPHARM/2004/008409

as cognate ligand, yellow; 10~100, green; 100~1000, gray; 1000~10000 times, white; not detected. Triphenyltin (no. 13) and tributyltin (no. 14) showed strong activity on PPAR γ and RXR α , respectively.

Fig. 3. Dose-response curves of the effects of organotin compounds on hRXR α and hPPAR γ in the CoA-BAP system. (A) Tributyltin (TBT; filled triangles) showed strong agonistic activity for hRXR α at as low a concentration as that of 9-*cis* retinoic acid (9cRA; filled diamonds). (B) Triphenyltin (TPT; open squares) showed strong agonistic activity to hPPAR γ at as low a concentration as that of rosiglitazone (Rosi; filled diamonds). Triphenylmethane (TPM; open circles) and triphenylethylene (TPE; open triangles) did not show any agonistic activity. Activity of the vehicle control (DMSO)-only is shown by filled circles.

Fig. 4. Organotin compounds induce transcriptional activity through RXR α and PPAR γ .

Ligand-dependent transactivation of RXR α and PPAR γ were detected as luciferase activity. (a) F9 cells were cotransfected with a GAL4-DBD-hRXR α expression plasmid and a GAL4-responsive reporter plasmid. (b) NIH-3T3 cells were cotransfected with a GAL4-DBD-mPPAR γ 1 expression plasmid and a GAL4-responsive reporter plasmid. The luciferase activities relative to the β -galactosidase activity is shown and represents the fold stimulation compared with

MOLPHARM/2004/008409

the activity of the vehicle-only control. Data shown are the means \pm standard deviation of three independent experiments. $**p < 0.01$, $***p < 0.001$ significantly different from vehicle controls.

Fig. 5. Enhancement of lipid accumulation by organotin compounds. 3T3-L1 cells were maintained in DMEM containing 10% calf serum. One day after reaching confluence, the cells were treated for 60 h with vehicle only (a, e), 100 nM rosiglitazone (b, f), 100 nM TPT (c, g) or 100 nM TBT (d, h) in complete differentiation medium (a–d) or incomplete differentiation medium (e–h). The cells received fresh medium every 48 h. On the 10th day after induction of differentiation, the cells were fixed with paraformaldehyde and stained with Oil Red O.

Fig. 6. Induction of adipocyte differentiation markers by organotin compounds. (a) Induction of adipocyte marker genes by organotin compounds in incomplete differentiation medium. 3T3-L1 cells were maintained in DMEM containing calf serum. One day after reaching confluence, the cells were treated with vehicle only, rosiglitazone (10 – 30 nM), TPT (10 – 30 nM), TBT (10 – 30 nM), or 9-cis retinoic acid (100 nM) in DMEM containing 10% FBS and 10 μ g/ml insulin. Total RNA was isolated at 10 days after treatment, and mRNA expression of the aP2 and PPAR γ genes was detected by Northern blot analysis. The ethidium-bromide staining for ribosomal RNAs is shown as a control. (b) Time course of aP2 gene

MOLPHARM/2004/008409

expression. 3T3-L1 cells were treated with vehicle only, rosiglitazone (100 nM), TPT (100 nM), or TBT (100 nM) in incomplete differentiation medium. The cells were harvested at the indicated time after treatment, and mRNA expression of the aP2 gene was analyzed by RT-PCR. (c) Lipid accumulation in differentiated 3T3-L1 cells. The cells were treated with 1, 10, 30, or 100 nM of chemicals. Ten days later, the amount of triglyceride was determined as described in Materials and Methods section.

Table 1. Suspected endocrine disruptors tested in this study.

No.	Compound	Abbreviation	CAS No.	No.	Compound	Abbreviation	CAS No.
1	Diethyl phthalate	DEP	84-66-2	21	Methomyl		16752-77-5
2	Dipropyl phthalate	DPrP	131-16-8	22	Maneb		12427-38-2
3	Di- <i>n</i> -butyl phthalate	DBP	84-74-2	23	Mancozeb		8018-01-7
4	Di- <i>n</i> -pentyl phthalate	DPP	131-18-0	24	Ziram		137-30-4
5	Dihexyl phthalate	DHP	84-75-3	25	Methoxychlor	MXC	72-43-5
6	Diethylhexyl phthalate	DEHP	117-81-7	26	Hexachlorocyclohexane	γ -HCH	58-89-9
7	Dicyclohexyl phthalate	DCHP	84-61-7	27	Permethrin		54645-53-1
8	Butyl Benzyl phthalate	BBP	85-68-7	28	2,4-D		94-75-7
9	Diethylhexyl adipate	DEHA	103-23-1	29	2,4,5-T		93-76-5
10	4-Nonylphenol	4-NP	25154-53-3	30	Simazine	CAT	122-34-9
11	<i>p</i> -Octylphenol	<i>p</i> -OP	1806-26-4	31	Alachlor		15972-60-8
12	Bisphenol A	BPA	80-05-7	32	PCP		87-86-5
13	Triphenyltin	TPT	639-58-7	33	Amitrole		61-82-5
14	Tributyltin	TBT	1461-22-9	34	Nitrofen	NIP	1836-75-5
15	4-Nitrotoluene	4-NT	99-99-0	35	Trifluralin		1582-09-8
16	Benzophenone	BZP	119-61-9	36	1,2-dibromo-3-chloropropane	DBCP	96-12-8
17	Benzo [a] pyrene	B[a]P	50-32-8	37	Malathone		121-75-5
18	Aldicarb		116-06-3	38	Kelthane		115-32-2
19	Vinclozolin		50471-44-8	39	2,4-Dichlorophenol	DCP	120-83-2
20	Carbaryl	NAC	63-25-2	40	Octachlorostyrene	OCS	29082-74-4

MOLPHARM/2004/008409











Fig. 1**(a) ERs****(b) RARs****(c) TR****(d) VDR****(e) RXRs****(f) PPARs****(g) FXR****(h) LXRs****(i) ERRs**

Fig. 3

(a) RXR α



(b) PPAR γ








Fig. 4**(a) RXR α** **(b) PPAR γ** 

Fig. 5


(a) DMSO (complete medium)




(b) Rosi (complete medium)




(c) TPT (complete medium)




(d) TBT (complete medium)




(e) DMSO (incomplete medium)



(f) Rosi (incomplete medium)



(g) TPT (incomplete medium)



(h) TBT (incomplete medium)






Fig. 6

(a) Northern blot



(b) RT-PCR (aP2 mRNA)



(c) Lipid accumulation

