

MOL 13847

Title page

**Functional complementation and the analysis of opioid receptor homo-
dimerization**

Geraldine Pascal and Graeme Milligan*

Molecular Pharmacology Group

Division of Biochemistry and Molecular Biology

Institute of Biomedical and Life Sciences

University of Glasgow

Glasgow G12 8QQ

Scotland, U.K.

MOL 13847

Running Title Page

Running title: opioid receptor dimerization

* correspondence to:

Graeme Milligan, Davidson Building, University of Glasgow, Glasgow G12 8QQ

Scotland, U.K.

Tel 44 141 330 5557

FAX 44 141 330 4620

e-mail g.milligan@bio.gla.ac.uk

Manuscript Information

Text pages 45

Tables 7

Figures 10

References 38

Abstract 247

Introduction 647

Discussion 1650

Abbreviations: DADLE, D-ala², leu⁵-enkephalin; DAMGO, [D-Ala²,NMe-Phe⁴,Gly-ol⁵]-enkephalin; DOP, delta opioid peptide; DPDPE, [D-Pen^{2,5}]-enkephalin; GPCR, G protein-coupled receptor; GTPγS, guanosine 5'-(γ-thio) triphosphate; KOP, kappa opioid peptide; MOP, mu opioid peptide; U69593, (+)-(5α,7α,8β)-N-methyl-N-[-7-(1-pyrrolodiny)-1-oxaspiro[4,5]dec-8-yl]benzeneacetamide.

MOL 13847

Abstract

Complementation of function following co-expression of pairs of non-functional G protein-coupled receptors that contain distinct inactivating mutations supports the hypothesis that such receptors exist as dimers. Chimeras between members of the metabotropic glutamate receptor-like family have been particularly useful because the N-terminal ligand binding and heptahelical transmembrane elements can be considered distinct domains. To examine the utility of a related approach for opioid receptors, fusion proteins were generated in which a Pertussis toxin-resistant (Cys³⁵¹Ile) variant of the G protein G_{i1}α was linked to the C-terminal tail of the DOP, KOP and MOP receptors. Each was functionally measured by agonist stimulation of [³⁵S]GTPγS binding in G_iα immunoprecipitates from membranes of Pertussis toxin-treated HEK293 cells. Agonist function was eliminated either by fusion of the receptors to G_{i1}αGly²⁰²Ala,Cys³⁵¹Ile or mutation of a pair of conserved Val residues in intracellular loop 2 of each receptor. Co-expression, but not simple mixing, of the two inactive fusion proteins reconstituted agonist-loading of [³⁵S]GTPγS for each receptor. At equimolar amounts, reconstitution of DOP receptor function was more extensive than KOR or MOR. Reconstitution of DOP function required two intact receptors and was not achieved by provision of extra G_{i1}αCys³⁵¹Ile membrane anchored by linkage to DOP transmembrane domain 1. Inactive forms of all G protein α subunits can be produced by mutations equivalent to G_{i1}αGly²⁰²Ala. As the amino acids modified in the opioid receptors are highly conserved in most rhodopsin-like receptors this approach should be widely applicable to study the existence and molecular basis of receptor dimerization.

MOL 13847

Introduction

An extensive literature now exists on the capacity of a wide range of G protein-coupled receptors (GPCRs) to form dimers and/or higher-order oligomers (Lee et al., 2003, Breitwieser, 2004, Milligan, 2004). Despite this, many of the reports have been predominantly descriptive and provide limited insights into the proportion of different GPCRs that may exist as dimers, the relative propensity of different GPCRs to oligomerize, the molecular basis of dimerization and whether there are differences in the details of how closely related GPCRs form dimers/oligomers.

The ability of the DOP, KOP and MOP opioid receptor subtypes to form homo-dimers and/or higher-order oligomers has previously been investigated using both co-immunoprecipitation and resonance energy transfer techniques (Cvejic and Devi, 1997, George et al., 2000, McVey et al, 2001, Li-Wei et al., 2002, Ramsay et al., 2002). Despite this, there is little information available on the issues noted above, although informatic analysis has suggested potential interfaces in transmembrane helices that may contribute to opioid receptor subtype homo-dimerization (Filizola and Weinstein, 2002).

If the co-expression of two non-equivalent and non-functional mutants of a GPCR is both able and required to reconstitute receptor ligand binding and/or function this can provide evidence in favour of direct protein-protein interactions and quaternary structure for the active receptor (Milligan and Bouvier, 2005). For example, co-expression of two forms of the angiotensin AT1 receptor that were unable to bind angiotensin II or related ligands due to point mutations in either transmembrane III or transmembrane region V restored ligand binding (Monnot et al., 1996). Such an approach has also been used to explore mechanisms of dimerization. Theoretical

MOL 13847

models of GPCR dimerization include both 'contact' and 'domain swap' dimers.

Using the histamine H1 receptor as a model, Bakker et al., (2004) showed that although single point mutations in both transmembrane region III and transmembrane region VI prevented binding of antagonist radioligands, including [³H]mepyramine, co-expression of the two mutants resulted in reconstitution of [³H]mepyramine binding sites with the anticipated pharmacological characteristics. Conceptually this should not be possible for a 'contact' dimer in which transmembrane domains are not exchanged but simply appose each other.

In addition to the restoration of ligand binding, studies that have used pairs of non-functional mutants to restore GPCR signalling have produced data consistent with GPCR-GPCR interactions. By generating mutants of the luteinizing hormone receptor that were either unable to bind ligand or unable to signal, although able to bind the agonist, Lee et al., (2002) were able to reconstitute agonist-mediated regulation of cAMP levels following co-expression of the two mutants. The luteinizing hormone receptor, as with other GPCRs with related ligands, has an extended N-terminal region involved in ligand binding. As such, Lee et al., (2002) were able to consider the N-terminal 'exo-domain' and the seven transmembrane element 'endo-domain' as distinct entities in a manner equivalent to the extracellular and transmembrane elements of class C GPCRs that have allowed elegant chimeric receptor approaches to understand the mechanism of signal transduction through obligate hetero-dimers (Pin et al., 2005).

Recently, as a variant of this, functional complementation was observed following the co-expression of pairs of α_{1b} -adrenoceptor- $G_{11}\alpha$ and histamine H1 receptor- $G_{11}\alpha$ GPCR-G protein fusion proteins that were both inactive when expressed individually

MOL 13847

because they contained specific mutations in either the GPCR or G protein elements (Carrillo et al., 2003). All G protein α subunits contain a conserved Gly that, when mutated, prevents effective GDP-GTP exchange and hence activation (Milligan et al., 2005). Furthermore, nearly all class A, rhodopsin-like GPCRs have either one, or more usually two, hydrophobic residues in the second intracellular loop homologous to those mutated to generate inactive forms of the α_{1b} -adrenoceptor and histamine H1 receptor (Milligan et al., 2005). We thus wished to test if equivalent pairs of inactive opioid receptor-G α fusion proteins could be produced and to assess if variations in pharmacology and/or reconstitutive capacity could provide insights into the basis of opioid receptor subtype dimerization.

Materials and Methods

Materials/Ligands

Diprenorphine [15,16-³H] (50 Ci/mmol) and guanosine 5'-(γ -thio) triphosphate [³⁵S] (1250 mCi/mmol) were from PerkinElmer life science, Inc. (Boston, USA).

DADLE (D-ala², leu⁵-enkephalin), DAMGO ([D-Ala²,NMe-Phe⁴,Gly-ol⁵]-enkephalin, DPDPE ([D-Pen^{2,5}]-enkephalin), naloxone and Pertussis toxin were from SIGMA-Aldrich Company Ltd., Poole, Dorset, UK. U69593 ((+)-(5 α ,7 α ,8 β)-N-methyl-N-[7-(1-pyrrolodiny)-1-oxaspiro[4,5]dec-8-yl)benzeneacetamide) was from Tocris. Recombinant, myristoylated rat G_{i1 α} subunit was from Calbiochem.

MOL 13847

Antibodies/antisera

The anti-G_{α1-2} antiserum (SG) has been described previously (Green et al., 1990). The mouse monoclonal anti-Flag antibody (M2) was from SIGMA-Aldrich Company Ltd., Poole, Dorset, UK. The rabbit polyclonal antiserum anti-c-myc antiserum was from Cell Signalling Technology, Nottingham, UK

Molecular constructs

hDOP-G_{i1α}C³⁵¹I in pcDNA3.1 was generated previously (Moon et al., 2001) and used as a template to introduce mutations in the 2nd intracellular loop of the receptor to produce **hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I** using the QuikChange kit (Stratagene) and the following primers:

Sense primer: 5'-GAC CGC TAC ATC GCT GAG TGC CAC CCT GAC AAG GCC CTG GAC TTC-3'

Antisense primer: 5'-GAA GTC CAG GGC CTT GTC AGG GTG GCA **CTC** AGC GAT GTA GCG GTC-3'

Bold letters indicate bases altered. The PCR product was then digested with *DpnI* and transformed into bacteria.

hDOP-G_{i1α}G²⁰²A,C³⁵¹I

In a similar manner hDOP-G_{i1α}C³⁵¹I was used to introduce the G²⁰²A mutation in G_{i1α} using the following primers:

Sense primer: 5'-G TTT GAC GTG GGA GCC CAG AGA TCA GAG C-3'

Antisense primer: 5'-G CTC TGA TCT CTG GGC TCC CAC GTC AAA C-3'

The PCR product was then digested by *DpnI* and was transformed into bacteria.

MOL 13847

Flag-hDOPV¹⁵⁰E,V¹⁵⁴D-G_{ii}αC³⁵¹I

was constructed using the following primers:

Sense primer: 5' ACT AGT GCT AGC ATG GAC TAC AAG GAC GAC GAT GAT AAG
GAA CCG GCC CCC TCC GCC GGC-3'

Antisense primer: 5'-GAA TTT GGA TCC GGC GGC AGC GCC ACC GCC GGG-3'

The sense primer contains a Flag sequence (in italics) and an *NheI* restriction site (underlined) and corresponds to the N-terminal region of hDOP. The antisense primer contains a *BamHI* site (underlined) and corresponds to the C-terminal region of hDOP. The PCR product and pcDNA3.1 vector containing hDOPV¹⁵⁰E,V¹⁵⁴D-G_{ii}αC³⁵¹I were digested by *NheI* and *BamHI*. The digested products were then ligated.

c-myc-hDOP-G_{ii}αG²⁰²A,C³⁵¹I

was constructed using the following primers:

Sense primer: 5'-CCC TTT GCT AGC ATG GAA CAA AAG CTT ATT TCT GAA GAA
GAT CTG GAA CCG GCC CCC TCC GCC-3'

Antisense primer: 5'-GAA TTT GGA TCC GGC GGC AGC GCC ACC GCC GGG-3'

hDOP-G_{ii}αG²⁰²A,C³⁵¹I was amplified by these primers. The sense primer contains a c-myc sequence (in italics) and *NheI* restriction site (underlined), the antisense primer contains a *BamHI* site (underlined). The PCR product and pcDNA3.1 containing hDOP-G_{ii}αG²⁰²A,C³⁵¹I were digested with *NheI* and *BamHI*. The digested products were then ligated.

hMOPV¹⁶⁹EV¹⁷³D-G_{ii}αC³⁵¹I

MOL 13847

hMOR-G_{11α}C³⁵¹I cDNA in pcDNA3 was generated previously (Massotte et al.,2002) and was used as a template to introduce mutations in the 2nd intracellular loop of the receptor using the following primers:

Sense primer: 5'-GAT CGA TAC ATT GCA **GAG** TGC CAC CCT GAC AAG GCC TTA
GAT TTC-3'

Antisense primer: 5'-GAA ATC TAA GGC CTT GTC AGG GTG GCA **CTC** TGC AAT
GTA TCG ATC-3'

The appropriate valines were mutated into glutatmate (GAG) and aspartate (GAC) respectively. Altered bases mutated are in bold. The PCR product was digested by *DpnI* and was transformed into bacteria.

hMOP-G_{11α}G²⁰²A,C³⁵¹I

was produced as for hDOP-G_{11α}G²⁰²A,C³⁵¹I but using hMOP-G_{11α}C³⁵¹I cDNA as the template .

rKOP-G_{11α}C³⁵¹I

rKOP-G_{11α}C³⁵¹I was constructed using the following primers:

Sense primer: 5'-CCC AAA AAG CTT ATG GAG TCC CCC ATC CAG ATT TTC C-3'

Antisense primer: 5'-GGC ATC GGT ACC TAC TGG CTT ATT CAT CCC ACC CAC
ATC CCT CAT GGA-3'

Rat KOP was amplified between these primers corresponding to the N and C-termini of rKOP and containing *HindIII* and *KpnI* restriction sites (underlined). The PCR product and pcDNA3 containing G_{11α}C³⁵¹I were digested by the above enzymes. Because rKOP

MOL 13847

contains an internal *Hind*III site, a two-way ligation was performed to ligate the vector and the two elements of the digested PCR product.

rKOP V¹⁶⁰E,V¹⁶⁴D-Gi_{1α}C³⁵¹I

rKOP-Gi_{1α}C³⁵¹I cDNA as above was used as a template to introduce mutations in the 2nd intracellular loop of the receptor, using the following primers:

Sense primer: 5'-GAC CGC TAC ATT GCC **GAG** TGC CAC CCT GAC AAA GCT TTG GAT TTC-3'

Antisense prime: 5'-GAA ATC CAA AGC TTT GTC AGG GTG GCA **CTC** GGC AAT GTA GCG GTC-3'

Bases mutated are in bold.

rKOP-Gi_{1α}G²⁰²A,C³⁵¹I

rKOP-Gi_{1α}C³⁵¹I cDNA was used as a template to introduce the mutation in Gi_{1α} as for hDOP and hKOP.

Flag-Nt-TM1-Gi_{1α} C³⁵¹I

Flag-Nt-TM1-Gi_{1α} C³⁵¹I was constructed using the following primers:

Sense primer: 5' ACT AGT GCT AGC ATG GAC TAC AAG GAC GAC GAT GAT AAG GAA CCG GCC CCC TCC GCC GGC-3'

Antisense primer: 5'-CCC ATT GGA TCC GGT GGC CGT CTT CAT CTT AGT GTA CCG-3'

Flag-hDOP-Gi_{1α} C³⁵¹I was used as template for PCR. The first 252bp were amplified by PCR and were then digested using *Bam*HI and *Nhe*I (restriction sites underlined). The same

MOL 13847

digestion was used on the template, *NheI* being situated at the end of the receptor sequence.

PCR products and vector were ligated.

Cell transfection and treatment

HEK293 cells were transfected using Lipofectamine reagent (Gibco Life Technologies) or Gene Juice (Novagen) and the appropriate cDNA(s) according to the manufacturers' instructions. Cells were treated with Pertussis toxin (25ng/ml) for 16 to 18 h prior to harvest.

[³H]Diprenorphine binding

The expression of GPCR-G protein fusions was assessed by measuring the specific binding of [³H]diprenorphine in cell membrane preparations. Non-specific binding was assessed by the addition of 100μM naloxone. Samples were incubated for 1h at 25°C and bound ligand separated from free by vacuum filtration through GF/B filters pre-treated with 0.3% polyethyleneimine in TEM (10mM Tris/HCL, 0.1mM EDTA, 10mM MgCl₂, pH adjusted to 7.5). Bound ligand was estimated by liquid scintillation spectroscopy. Competition studies were conducted with 1nM [³H]diprenorphine and a range of concentrations of other ligands. Data were analysed using GraphPad Prism software (San Diego, CA). Saturation data were fit to non-linear regression curves.

[³⁵S]GTPγS binding studies

Experiments were initiated by adding the assay buffer mix (20 mM HEPES (pH 7.4), 3 mM MgCl₂, 100mM NaCl, 10 μM GDP, 0.2 mM ascorbic acid) containing 50 nCi [³⁵S]GTPγS

MOL 13847

in the presence or absence of agonist to a defined amount of membranes. Non-specific binding was determined in the presence of 100 μ M GTP γ S. The reaction was incubated for 15 min at 30°C and terminated by adding 1ml of ice-cold stop buffer. The samples were centrifuged for 15 min at 16000 x g at 4°C and the resulting pellets were resuspended in solubilization buffer (100 mM Tris HCl, 200 mM NaCl, 1 mM EDTA, 1.25% NP40, pH adjusted to 7.4) plus 0.2% SDS. Samples were pre-cleared with Pansorbin for 1h at 4°C and centrifuged for 2 min at 16000 x g. Supernatant was added to a mix of protein G and the anti-G_{11 α} /G_{12 α} antiserum, SG (Green et al., 1990) and left rotating overnight at 4°C for immunoprecipitation. The immunocomplexes were washed twice with ice-cold solubilization buffer and bound [³⁵S]GTP γ S measured.

Co-immunoprecipitation

Cells were resuspended in 1ml of 1 x RIPA (radio-immunoprecipitation assay) buffer and rotated for 60 min at 4°C to allow lysis. The samples were centrifuged at 14,000 x g for 10min at 4°C and the supernatant retained. 50 μ l of a protein G-sepharose/phosphate buffered saline slurry was added to the supernatant and rotated for further 60 min at 4°C to pre-clear. Samples were centrifuged at 14000 x g for 10 min at 4°C. Supernatant was conserved and protein concentration was measured using the BCA assay method. Samples were equalized to 1 μ g/ μ l. Target proteins were then immunoprecipitated from 500 μ l samples by incubation with 20 μ l of protein G-sepharose and the appropriate antibody/antiserum overnight at 4°C on a rotating wheel. Immune complexes were isolated by centrifugation at 14000 x g for 1 min and washed twice with RIPA buffer. Proteins were eluted from the protein G-sepharose by the addition of 30-50 μ l Laemmli buffer and heated for 4 min at 85°C. The eluates were then loaded onto SDS-PAGE gels.

MOL 13847

Quantitation of Flag-Nt-TM1-G_{11α}C³⁵¹I expression levels

Varying amounts (12.5-50ng) of recombinantly expressed, myristoylated rat G_{11α} were run on SDS-PAGE alongside membranes of HEK293 cells transfected to co-express Flag-Nt-TM1-G_{11α}C³⁵¹I and hDOP-G_{11α}G²⁰²A,C³⁵¹I. Following immunoblotting with the anti-G_{11α}/G_{12α} antiserum SG, densitometry indicated that the signal corresponding to the recombinant G_{11α} increased in a linear fashion over this range. Interpolation of the immuno-signal corresponding to Flag-Nt-TM1-G_{11α}C³⁵¹I (molecular mass 60.57kDa) in different amount of transfected cell membranes allowed estimation of expression levels.

Results

A fusion protein was constructed between the human DOP (hDOP) receptor and a form of the α subunit of the G protein G₁₁ that was rendered resistant to the ADP-ribosyltransferase activity of Pertussis toxin by conversion of Cys³⁵¹ to Ile (G_{11α}C³⁵¹I). The hDOP-G_{11α}C³⁵¹I fusion protein was expressed transiently in HEK293 cells that were also treated with Pertussis toxin (25ng/ml, 16h) prior to harvest to cause ADP-ribosylation of the endogenously expressed forms of the G_i/G_o group of G proteins. Membranes prepared from these cells were used in saturation [³H]diprenorphine ligand binding assays to measure expression levels of the construct and its affinity for the ligand (Table 1). Expression levels were 1816 ± 209 fmol/mg membrane protein and the pK_d for [³H]diprenorphine 9.20 ± 0.03 (n = 4, means ± S.E.M.). The functionality of hDOP-G_{11α}C³⁵¹I was assessed by the capacity of the synthetic opioid peptide D-ala², leu⁵ enkephalin (DADLE) to stimulate binding of [³⁵S]GTPγS in membranes containing the construct that were subsequently

MOL 13847

immunoprecipitated with the anti-G_{i1α}/G_{i2α} antiserum, SG (Figure 1a). Virtually no [³⁵S]GTPγS was recovered in immunoprecipitates from membranes of mock-transfected cells treated with either DADLE or vehicle (Figure 1a). By contrast, although binding of [³⁵S]GTPγS in immunoprecipitates from hDOP-G_{i1α}C³⁵¹I-expressing cell membranes was greatly increased by DADLE, the construct was also able to load [³⁵S]GTPγS in the absence of agonist (Figure 1a). When membrane amounts corresponding to varying levels of hDOP-G_{i1α}C³⁵¹I were used, DADLE stimulation of [³⁵S]GTPγS binding was linear with fusion protein amount over the full range tested and up to at least 60fmol (Figure 1b).

We have previously demonstrated that mutation of Gly²⁰⁸ to Ala in the G protein G_{i1α} prevents receptor-mediated guanine nucleotide exchange and hence [³⁵S]GTPγS binding (Carrillo et al., 2002). The α subunit of all hetero-trimeric G proteins contains Gly at the equivalent position. To test the general effect of mutating this Gly on the capacity of receptors to enhance guanine nucleotide exchange we thus generated hDOP-G_{i1α}G²⁰²A,C³⁵¹I. When this was expressed in HEK293 cells and membranes prepared from Pertussis toxin-treated cells, neither the level of expression of this construct nor the binding affinity for [³H]diprenorphine was different from hDOP-G_{i1α}C³⁵¹I (Table 1). However, although 10μM DADLE caused a 5.2 +/- 0.3 fold (n = 4 mean ± S.E.M.) increase in levels of [³⁵S]GTPγS binding compared to vehicle-treated controls in samples immunoprecipitated from membranes expressing 15 fmol of hDOP-G_{i1α}C³⁵¹I (Figure 2), no significant DADLE stimulation of [³⁵S]GTPγS binding was observed in immunoprecipitated samples derived from membranes containing 15 fmol of hDOP-G_{i1α}G²⁰²A,C³⁵¹I (Figure 2). Furthermore,

MOL 13847

[³⁵S]GTPγS loading in the absence of DADLE was substantially reduced (Figure 2). Mutation of hydrophobic residues in the second intracellular loop of family A GPCRs can essentially eliminate G protein activation without major effects on antagonist ligand binding (Carrillo et al., 2003, Milligan et al., 2005). To test this for hDOP we thus also generated hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I. This also was expressed as well as hDOP-G_{i1α}C³⁵¹I (Table 1) but bound [³H]diprenorphine with 3-fold lower affinity than hDOP-G_{i1α}C³⁵¹I (Table 1). [³⁵S]GTPγS binding studies demonstrated this construct also to have much reduced basal guanine nucleotide exchange and not to produce a statistically significant increase in binding of [³⁵S]GTPγS in response to DADLE (Figure 2). When hDOP-G_{i1α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I were co-expressed and membranes containing 15 fmol of [³H]diprenorphine binding sites were used in [³⁵S]GTPγS binding studies, DADLE stimulation was partially reconstituted (Figure 2). With membranes from these cells containing 30 fmol of [³H]diprenorphine binding sites, DADLE-stimulated [³⁵S]GTPγS binding was 60% of that achieved in membranes expressing 15 fmol of the wild type hDOP-G_{i1α}C³⁵¹I fusion construct (Figure 2). Reconstitution of DADLE-stimulated [³⁵S]GTPγS binding required the co-expression of hDOP-G_{i1α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I and not simply the presence of both in the assay. When membranes containing 15 fmol of individually expressed hDOP-G_{i1α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I were simply mixed prior to the assay to provide 30 fmol of fusion proteins in the assay no reconstitution of DADLE-stimulated [³⁵S]GTPγS binding was observed (Figure 2). These data are consistent with a requirement for hDOP interactions to generate function.

MOL 13847

Interestingly, the affinity of [³H]diprenorphine binding in membranes co-expressing hDOP-G_{i1α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I was equivalent to the individually expressed hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I construct (Table 1). Although this observation might indicate the presence of a substantially greater proportion of hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I than hDOP-G_{i1α}G²⁰²A,C³⁵¹I in the membranes from co-expressed cells this is not consistent with the functional reconstitution data (Figure 2) or with the equivalent levels of expression of these two constructs when expressed individually (Table 1). However, to examine this further and to confirm direct interactions between hDOP-G_{i1α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I we performed co-immunoprecipitation studies using membranes of HEK293 cells transfected to express individually or co-express N-terminally modified Flag-hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I and/or c-myc-hDOP-G_{i1α}G²⁰²A,C³⁵¹I. Immunoprecipitation with anti-Flag antibody followed by SDS-PAGE and immunoblotting with anti-c-myc antibody resulted in detection of specific c-myc immunoreactivity only when the two fusion constructs were co-expressed (Figure 3), consistent with a physical interaction between the two variants. To further explore aspects of pharmacology of the fusion proteins, membranes from Pertussis toxin-treated HEK 293 cells transfected to express hDOP-G_{i1α}C³⁵¹I; hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I; hDOP-G_{i1α}G²⁰²A,C³⁵¹I or the combination of hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I + hDOP-G_{i1α}G²⁰²A,C³⁵¹I were used in [³H]-diprenorphine/DADLE competition binding experiments (Table 2). Two-site binding curves reflecting higher and lower affinity binding sites for the agonist DADLE were best fitted in each case. Introduction of the G²⁰²A mutation in the G-protein subunit did not alter DADLE binding properties substantially as similar pK_h and pK_i values

MOL 13847

were observed for hDOP-G_{11α}G²⁰²A,C³⁵¹I as for hDOP-G_{11α}C³⁵¹I (Table 2). In contrast, the double mutation in the 2nd intracellular loop of hDOP receptor did alter the binding affinity of DADLE with some 30 fold loss of affinity in both high and low affinity binding sites (hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I pK_h = 7.4 ± 0.2, pK_i 5.0 ± 0.4, hDOP-G_{11α}C³⁵¹I pK_h = 9.0 ± 0.2, pK_i 6.8 ± 0.42). In membranes co-expressing hDOP-G_{11α}G²⁰²A,C³⁵¹I and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I there was no significant difference in the percentage of high and low site numbers compared to the wild type hDOP-G_{11α}C³⁵¹I fusion protein (P>0.05, 1 way ANOVA) (Table 2). A similar reduction in affinity of the high affinity site for the DOP selective peptide agonist [D-Pen^{2,5}]-enkephalin (DPDPE) was also observed when comparing hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I to hDOP-G_{11α}C³⁵¹I or hDOP-G_{11α}G²⁰²A,C³⁵¹I (Table 3) and although a similar trend was observed for the low affinity site (Table 3), this did not achieve statistical significance due to relatively imprecise estimates of pK_i. Wild type DPDPE binding characteristics were again restored following co-expression of hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I and hDOP-G_{11α}G²⁰²A,C³⁵¹I (Table 3). Assuming the predominant form of the hDOP is as a dimer, rather than a higher-order oligomer, co-expression of hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I and hDOP-G_{11α}G²⁰²A,C³⁵¹I must be expected to generate hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I dimers and hDOP-G_{11α}G²⁰²A,C³⁵¹I dimers, which as shown in Figure 2 are inactive, as well as the functionally reconstituted hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I + hDOP-G_{11α}G²⁰²A,C³⁵¹I dimer. Ligand binding studies must reflect the full population of these different hDOP ‘homo-dimers’ in the cell membrane. By contrast, in functional assays, only hDOP-G_{11α}C³⁵¹I homo-dimers and hDOPV¹⁵⁰E,V¹⁵⁴D-G_{11α}C³⁵¹I + hDOP-

MOL 13847

$G_{i1\alpha}G^{202}A,C^{351}I$ 'homo-dimers' are reported (Figure 2). The potency of DADLE to stimulate [^{35}S]GTP γ S binding via the hDOP- $G_{i1\alpha}C^{351}I$ dimer and the reconstituted hDOPV $^{150}E,V^{154}D-G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ dimer was not different (Figure 4a). Equally, the prototypic opioid receptor antagonist naloxone was equipotent in its ability to prevent DADLE-stimulated [^{35}S]GTP γ S binding via the hDOP- $G_{i1\alpha}C^{351}I$ dimer and the reconstituted hDOPV $^{150}E,V^{154}D-G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ dimer (Figure 4b).

To assess if the reconstitution of function observed upon co-expression of hDOPV $^{150}E,V^{154}D-G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ could possibly be accounted for simply by the provision of the $G_{i1\alpha}C^{351}I$ attached to the inactive hDOPV $^{150}E,V^{154}D$ receptor rather than specifically requiring interactions between hDOPV $^{150}E,V^{154}D$ and hDOP we generated and expressed a construct (Flag-Nt-TM1- $G_{i1\alpha}C^{351}I$) in which $G_{i1\alpha}C^{351}I$ was linked to a sequence comprising the N-terminal domain, transmembrane region 1 and the first intracellular loop of hDOP. This construct did not bind [3H]diprenorphine (data not show) but its expression as an apparent 48 kDa polypeptide could clearly be detected by immunoblotting transfected HEK293 membranes with the anti- $G_{i1\alpha}/G_{i2\alpha}$ antiserum (Figure 5a). Parallel SDS-PAGE and immunodetection of varying amounts of recombinantly expressed $G_{i1\alpha}$ followed by densitometry of the signals allowed production of a standard curve for $G_{i1\alpha}$ expression that was linear over the range (0-50 ng) employed. Based on the anti- $G_{i1\alpha}$ immunological signal in membranes corresponding to Flag-Nt-TM1- $G_{i1\alpha}C^{351}I$ and its calculated molecular mass (60.57 kDa) we estimated levels of this construct to be 11.2 pmol/mg membrane protein. Therefore, this construct was present at some 6

MOL 13847

times the level of the hDOP-G_{i1α} fusion proteins. Co-transfection of Flag-Nt-TM1-G_{i1α}C³⁵¹I with hDOP-G_{i1α}G²⁰²A,C³⁵¹I resulted in very low, but statistically significant, increases in levels of [³⁵S]GTPγS binding in anti-G_{i1α}/G_{i2α} antiserum immunoprecipitates when DADLE was added to such membranes (Figure 5b). These very small signals did not reflect the possibility that although hDOP-G_{i1α}G²⁰²A,C³⁵¹I and Flag-Nt-TM1-G_{i1α}C³⁵¹I were co-expressed they were present in distinct membrane compartments. Co-expression of Flag-Nt-TM1-G_{i1α}C³⁵¹I with c-myc-hDOP-G_{i1α}G²⁰²A,C³⁵¹I allowed their co-immunoprecipitation (Figure 6a), indicating not only proximity but their capacity for physical interactions. Equally, co-expression of c-myc-Nt-TM1 with the isolated Flag-hDOP allowed their co-immunoprecipitation, indicating interactions were not a reflection of contacts between the two copies of the G protein (Figure 6b).

To extend these reconstitution studies to the other opioid receptors we generated equivalent fusion proteins incorporating the human MOP-1 (hMOP) receptor. hMOP-G_{i1α}C³⁵¹I, hMOP-G_{i1α}G²⁰²A,C³⁵¹I and hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I were expressed individually in HEK293 cells and following Pertussis toxin-treatment and membrane preparation, expression levels and affinity for [³H]diprenorphine were assessed via saturation binding studies. No significant differences between the three constructs were noted in either parameter (Table 4). Equally, following co-expression of hMOP-G_{i1α}G²⁰²A,C³⁵¹I and hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I the characteristics of [³H]diprenorphine binding were equivalent. In functional [³⁵S]GTPγS binding studies (Figure 7), the selective MOP receptor agonist [D-Ala²,NMe-Phe⁴,Gly-ol⁵]-enkephalin (DAMGO) (10μM) caused a 5.28 ± 0.24 fold (n = 4, mean ± S.E.M.) stimulation in end of assay anti-G_{i1α}/G_{i2α} antiserum

MOL 13847

immunoprecipitates. As with the related hDOP constructs, membranes expressing equal amounts of either hMOP-G_{11α}G²⁰²A,C³⁵¹I or hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I did not result in DAMGO stimulation of [³⁵S]GTPγS binding (Figure 7). Co-transfection of hMOP-G_{11α}G²⁰²A,C³⁵¹I and hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I did result in partial reconstitution of DAMGO-stimulated [³⁵S]GTPγS binding (Figure 7), an effect not achieved by simple mixing of membranes individually expressing hMOP-G_{11α}G²⁰²A,C³⁵¹I or hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I (Figure 7). In comparison to the 60% reconstitution of hDOP function, membranes expressing twice as many hMOP receptor [³H]diprenorphine binding sites following co-expression of the two inactive mutant fusion proteins allowed only 40% of the amount of agonist-stimulated [³⁵S]GTPγS binding as generated by the wild type hMOP-G_{11α}C³⁵¹I fusion (Figure 7). A potential explanation for this was uncovered on examining the potency of DAMGO to stimulate [³⁵S]GTPγS binding in membranes expressing hMOP-G_{11α}C³⁵¹I and co-expressing hMOP-G_{11α}G²⁰²A,C³⁵¹I and hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I. The potency of this ligand was reduced ($p < 0.05$) by some 2 fold at the functionally reconstituted dimer ($pEC_{50} = 6.1 \pm 0.07$) compared to the wild type dimer ($pEC_{50} = 6.5 \pm 0.04$). Interestingly, although both hMOP-G_{11α}C³⁵¹I and hMOP-G_{11α}G²⁰²A,C³⁵¹I displayed both high and low affinity binding sites for DAMGO when this ligand was allowed to compete with [³H]diprenorphine (Figure 8, Table 5), only a low affinity binding component could be detected for hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I (Figure 8, Table 5) akin to what might be anticipated if GPCR and G protein were uncoupled. When hMOPV¹⁶⁹E,V¹⁷³D-G_{11α}C³⁵¹I and hMOP-G_{11α}G²⁰²A,C³⁵¹I were co-expressed the characteristics of DAMGO binding were akin to a mixture of the two mutant constructs

MOL 13847

(Figure 8, Table 5) and analysis of the binding curves was consistent with the two constructs being present in a close to 1:1 ratio.

Studies were also performed on the rat (r)KOP receptor. rKOP-G_{i1α}C³⁵¹I, rKOP-G_{i1α}G²⁰²A,C³⁵¹I and rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I fusions were generated and expressed. These also all bound [³H]diprenorphine with high affinity and expressed to similar levels (Table 6) but, as with the hDOP constructs, a reduction in affinity was recorded for the rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I construct that incorporated mutations into the second intracellular loop of the receptor. As with the equivalent hDOP and hMOP constructs rKOP-G_{i1α}C³⁵¹I allowed a large increase in [³⁵S]GTPγS binding in response to agonist treatment (Figure 9). Individual expression of neither rKOP-G_{i1α}G²⁰²A,C³⁵¹I nor rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I resulted in stimulation of [³⁵S]GTPγS binding in the presence of the KOP receptor-selective agonist U69593 whereas co-expression of rKOP-G_{i1α}G²⁰²A,C³⁵¹I and rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I did (Figure 9). At a maximally effective concentration of U69593 (10μM) membranes expressing twice as many rKOP [³H]-diprenorphine binding sites following co-expression of the two inactive mutants, allowed some 50% of the amount of agonist-stimulated [³⁵S]GTPγS binding as generated by the wild type rKOP-G_{i1α}C³⁵¹I fusion (Figure 9). Akin to the hMOP constructs, in competition studies between [³H]diprenorphine and U69593, both rKOP-G_{i1α}C³⁵¹I and rKOP-G_{i1α}G²⁰²A,C³⁵¹I displayed both high and low affinity binding sites for the agonist. However, rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I displayed only a single, low affinity site for U69593 (Figure 10, Table 7). Also, as with the hMOP constructs, co-expression of rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I and rKOP-G_{i1α}G²⁰²A,C³⁵¹I resulted in a pattern of U69593 binding consistent with a mixture of the pharmacology of the two constructs (Figure 10,

MOL 13847

Table 7). The potency of U69593 to activate rKOP-G_{i1α}C³⁵¹I (pEC₅₀ = 7.3 ± 0.08) was higher (p < 0.05) than for the reconstituted rKOP dimer (pEC₅₀ = 6.8 ± 0.13).

Discussion

Fusion proteins between GPCRs and G protein α subunits have been used to examine a wide range of function of these polypeptides (Milligan, 2002, Milligan et al., 2004) with the defined 1:1 stoichiometry of the partner proteins being of particular use in measures of agonist-induced GTPase turnover number (Moon et al., 2001), the regulation, co-ordinated (Stevens et al., 2001) or otherwise (Barclay et al., 2005), of post-translational thio-acylation of GPCR and G protein and the effects of mutations in either partner that alter protein steady-state expression levels (Ward and Milligan, 2002). In the current studies we have generated and explored the function and pharmacology of fusions between each of the DOP, KOP and MOP opioid receptors with G_{i1α}. The functionality of each of these mutants was established in [³⁵S]GTPγS binding studies in which at assay termination, immunoprecipitation with an anti-G_{i1α}/G_{i2α} antiserum limited non-specific binding of the nucleotide. All commonly used cell lines express members of the G_{iα} G protein family that are substrates for pertussis toxin-catalysed ADP-ribosylation. To ensure agonist-driven [³⁵S]GTPγS binding reflected only binding to the fusion proteins under study, these were constructed using G_{i1α}C³⁵¹I (Bahia et al., 1998), which is insensitive to the actions of the toxin but able to be effectively activated by receptors, and by treating cells with Pertussis toxin prior to cell harvest to modify the endogenous G_{iα} pool. Mutation of Gly²⁰² to Ala in G_{i1α} resulted in a form of the G protein that was unable to exchange guanine nucleotide and bind [³⁵S]GTPγS in response to receptor agonists. All G protein α subunits have a Gly residue in the equivalent position and mutation should therefore be anticipated to produce

MOL 13847

equivalent lack of function mutants as previously shown for G_{11α} (Carrillo et al., 2002, 2003). Fusion of wild type G_{11α} to forms of the α_{1b}-adrenoceptor and the histamine H1 receptor containing hydrophobic to acidic residue mutations in intracellular loop 2 also results in lack of agonist-mediated [³⁵S]GTPγS binding without destruction of the ligand binding pocket (Carrillo et al., 2003). As most rhodopsin-like GPCRs have a pair of homologous hydrophobic residues (Milligan et al., 2005) and in the DOP, KOP and MOP receptors both are Val, we converted each of these to either Glu or Asp. This did not alter construct expression levels and had either no or only small effects on the binding affinity of [³H]diprenorphine. We were thus able to measure and equalise construct expression levels in preparation for functional studies. In each case, co-expression of the pair of non-functional opioid receptor-fusion proteins was able to partially reconstitute agonist-mediated [³⁵S]GTPγS binding. Reconstitution did require co-expression, simple mixing of membranes expressing the potentially complementary pairs did not generate agonist function. We have previously argued that such results require receptor dimerization (Carrillo et al., 2003) and provided evidence that the reconstitution reflects an ‘inter-’ rather than ‘intra-’ molecular interaction between GPCR and G protein (Carrillo et al., 2003). Although expression of a single fusion protein, wild type in both GPCR and G protein sequence, allows agonist mediated signal transduction, as with expression of a single GPCR cDNA, this does not allow direct exploration of GPCR quaternary structure. Indeed, the knowledge that a single cDNA was generally sufficient to generate the anticipated function and pharmacology of a GPCR played a significant part in the expectation that GPCRs would be single polypeptide, monomeric structures (Milligan, 2004). Previous studies by Molinari et al. (2003) have also noted a capacity of co-expressed pairs of inactive DOP-G protein fusions to reconstitute a signal. However, although they also concluded that this

MOL 13847

reflected inter-molecular interactions between the co-expressed forms, they did not specifically suggest this to require dimerization between the pair of DOP receptors. At least in part this may be because they also observed an ability of a DOP-G protein fusion to activate a G protein that was membrane anchored simply by linkage to transmembrane 1 of the vasopressin V2 receptor. In contrast with these observations, we observed only a very limited capacity of the hDOP-G_{i1α}G²⁰²A,C³⁵¹I construct to activate co-expressed G_{i1α}C³⁵¹I when it was tethered to the membrane by linkage to the N-terminal domain and transmembrane domain 1 of hDOP, even though the G protein was provided at some 5-6 times higher levels in this scenario than when provided by co-expression of the potentially complementary fusion protein. The basis for these differences is unclear but may relate to the high expression levels of the fusion proteins achieved and employed by Molinari et al. (2003) that were in the range in which so called ‘bystander’ interactions and effects have been observed (Mercier et al., 2002) likely due simply to physical proximity rather than direct protein-protein interactions. Although hDOP-G_{i1α}G²⁰²A,C³⁵¹I was unable to activate co-expressed Nt-TM1-G_{i1α}C³⁵¹I to any substantial extent these two constructs were able to interact because they could be co-immunoprecipitated following co-expression. This suggests that interaction between two complete receptors might be required for GPCR function and would support other evidence for conformational alterations in the partner GPCR in a dimer induced by ligand binding (Mesnier and Baneres, 2004, El-Asmar et al., 2005). As Nt-TM1 could also be co-immunoprecipitated with full length hDOP this suggests that TM1 and/or the N-terminal region of hDOP provides a protein-protein interaction interface. Although not explored in detail in these studies, for the α_{1b}-adrenoceptor symmetrical TM1-TM1 interactions provide key contributions to the quaternary organization of this GPCR (Carrillo et al., 2004) and a series of other reports

MOL 13847

have supported an important contribution of TM1 to the dimer interface(s) in other GPCRs (Overton and Blumer, 2002, Klco et al., 2003, Stanasila et al., 2003). Because ‘non-specific’ effects, potentially arising from high level expression in heterologous transfection studies, are an inherent concern, in the current experiments we maintained fusion construct expression in the range of 1-2 pmol/mg membrane protein and all ‘functional reconstitution’ experiments were performed under conditions in which agonist-stimulated [³⁵S]GTPγS binding increased linearly with construct amount. This was a key requirement for data analysis because if opioid receptors exist and function predominantly as dimers, the reconstitution strategy suggests that with 1:1 expression of the two mutant constructs then, in stochastic terms, 50% of the ligand binding sites should reflect ‘hetero’ interactions that can generate a functional response. A hypothesis was therefore that when using membranes co-expressing a pair of potentially suitable mutants, double the number of binding sites would be required to result in the same level of agonist-stimulated [³⁵S]GTPγS binding as with the wild type fusion. In all cases this was not achieved, with the level of reconstitution ranging from 40% for the MOP receptor to 60% for the DOP receptor. This may imply that not all cellular copies of a particular GPCR are present within dimers. This has been an extremely difficult issue to assess quantitatively. The proportion of a GPCR that migrates through SDS-PAGE as an SDS-resistant dimer is almost certainly a lower limit for the native state and although resonance energy transfer-based estimates of ‘dimer’ proportions have ranged from 25-85% (Mercier et al., 2002, Dinger et al, 2003) a considerable number of assumptions are required to allow such calculations (Milligan and Bouvier, 2005). Equally, there is growing evidence for a requirement of GPCR dimerization for productive signal transduction that is not restricted to the examples of the GABA_B and other family C receptors and for greater than dimeric, higher-order quaternary structure (Klco et al., 2003,

MOL 13847

Carrillo et al., 2004, Fotiadis et al., 2004). Equally, however, the basic strategy used herein might be restrictive as a pair of hydrophobic residues from the second intracellular loop were mutated to acidic residues and this might compromise the effectiveness of GPCR dimerization. It is worth noting, however, that the cytoplasmic face of the opioid receptor subtypes is very highly conserved between DOP, KOP and MOP and despite making the equivalent mutations in each, significant differences in reconstitutive effectiveness were observed. This may imply differences in the details of the homo-dimerization process. Although homo-dimerization of each of these three receptors has previously been recorded (Cvejcic and Devi, 1997, George et al., 2000, McVey et al, 2001, Li-Wei et al., 2002, Ramsay et al., 2002) there is no useful information on the similarities or differences in mechanisms of these interactions that have involved direct experimental study, although this topic has been considered via an informatic approach (Filizola and Weinstein, 2002). Although the mutation of hydrophobic residues in intracellular loop 2 may have limitations in producing an inactive GPCR, a marked advantage over certain other reconstitutive studies (Monnot et al., 1996, Bakker et al., 2004) is that the orthosteric GPCR ligand binding site was not destroyed. This allowed antagonist binding studies to confirm not only expression of each construct but that each inactive mutant was expressed at the same level as the wild type fusion. This was central to the 'stochastic' calculations of the potential makeup of the GPCR dimer population generated following co-expression of different proteins. Importantly, the complete conservation in G protein α subunits of the Gly residue modified herein to generate one of the pair of inactive fusions and the very high conservation of the pair of GPCR intracellular loop hydrophobic residues suggest that this strategy should be widely applicable (Milligan et al., 2005). It is likely, for example, to be of considerable use in mutational studies designed to identify key residues involved in the

MOL 13847

dimerization interface(s) (Hernanz-Falcon et al., 2004). Equally, there is no reason to limit such studies to GPCR homo-dimerization and the effectiveness of functional reconstitution may provide quantitative data on the propensity of GPCRs to hetero-dimerize. Indeed, this has been initiated by studies showing that the histamine H1 receptor and the α_{1b} -adrenoceptor are very poor interaction partners (Carrillo et al., 2003). Finally, as only the reconstituted 'hetero-dimer' is an active signalling unit, then in true GPCR hetero-dimerization studies the functional pharmacology of the hetero-dimer could be examined without interfering signals generated by the corresponding co-expressed homo-dimers, which, as shown herein, are essentially inactive.

MOL 13847

References

- Bahia DS, Wise A, Fanelli F, Lee M, Rees S, and Milligan G (1998) Hydrophobicity of residue³⁵¹ of the G protein G_{i1} alpha determines the extent of activation by the alpha_{2A}-adrenoceptor. *Biochemistry* **37**:11555-11562.
- Bakker RA, Dees G, Carrillo JJ, Booth RG, Lopez-Gimenez JF, Milligan G, Strange PG, and Leurs R. (2004) Domain swapping in the human histamine H1 receptor. *J Pharmacol Exp Ther* **311**:131-318.
- Barclay E, O'Reilly M, and Milligan G (2005) Activation of an α_{2A} -adrenoceptor-G α_{o1} fusion protein dynamically regulates the palmitoylation status of the G protein but not of the receptor. *Biochem J* **385**,197-206.
- Breitwieser GE (2004) G protein-coupled receptor oligomerization: implications for G protein activation and cell signaling. *Circ Res* **94**:17-27.
- Carrillo JJ, López-Gimenez JF, and Milligan G (2004) Multiple interactions between transmembrane helices generate the oligomeric α_{1b} -adrenoceptor. *Mol Pharmacol* **66**:1123-1137.
- Carrillo JJ, Pediani J, and Milligan G (2003) Dimers of class A G protein-coupled receptors function via agonist-mediated trans-activation of associated G proteins. *J Biol Chem* **278**:42578-42587.
- Carrillo JJ, Stevens PA, and Milligan G (2002) Measurement of agonist-dependent and-independent signal initiation of α_{1b} -adrenoceptor mutants by direct analysis of guanine nucleotide exchange on the G protein G α_{11} . *J Pharmacol Exp Ther* **302**:1080-1088.
- Cvejic S, Devi LA (1997) Dimerization of the delta opioid receptor: implication for a role in receptor internalization. *J Biol Chem* **272**:26959-26964.

MOL 13847

Dinger MC, Bader JE, Kobor AD, Kretschmar AK, and Beck-Sickinger AG (2003) Homodimerization of neuropeptide y receptors investigated by fluorescence resonance energy transfer in living cells. *J Biol Chem* **278**:10562-10571.

El-Asmar L, Springael JY, Ballet S, Andrieu EU, Vassart G, and Parmentier M (2005) Evidence for negative binding cooperativity within CCR5-CCR2b heterodimers. *Mol Pharmacol* **67**:460-469.

Filizola M, and Weinstein H (2002) Structural models for dimerization of G-protein coupled receptors: the opioid receptor homodimers. *Biopolymers* **66**: 317-325.

Fotiadis D, Liang Y, Filipek S, Saperstein DA, Engel A, and Palczewski K (2004) The G protein-coupled receptor rhodopsin in the native membrane. *FEBS Lett* **564**: 281-288.

George SR, Fan T, Xie Z, Tse R, Tam V, Varghese G, and O'Dowd BF (2000) Oligomerization of mu- and delta-opioid receptors. Generation of novel functional properties. *J Biol Chem* **275**:26128-26135.

Green A, Johnson JL, and Milligan G (1990). Down-regulation of G_i-subtypes by prolonged incubation of adipocytes with an A1 adenosine receptor agonist. *J Biol Chem* **265**:5206-5210.

Hernanz-Falcon P, Rodriguez-Frade JM, Serrano A, Juan D, del Sol A, Soriano SF, Roncal F, Gomez L, Valencia A, Martinez-A C, and Mellado M (2004) Identification of amino acid residues crucial for chemokine receptor dimerization. *Nature Immunol* **5**: 216-223.

Klco JM, Lassere TB, and Baranski TJ (2003) C5a receptor oligomerization. I. Disulfide trapping reveals oligomers and potential contact surfaces in a G protein-coupled receptor. *J Biol Chem* **278**:35345-35353.

MOL 13847

Lee C, Ji I, Ryu K, Song Y, Conn PM, and Ji TH (2002) Two defective heterozygous luteinizing hormone receptors can rescue hormone action. *J Biol Chem* **277**:15795-15800.

Lee SP, O'Dowd BF, and George SR (2003) Homo- and hetero-oligomerization of G protein-coupled receptors. *Life Sci* **74**, 173-180.

Li-Wei C, Can G, De-He Z, Qiang W, Xue-Jun X, Jie C, and Zhi-Qiang C (2002) Homodimerization of human mu-opioid receptor overexpressed in Sf9 insect cells. *Protein Pept Lett* **9**:145-152.

McClue SJ, Selzer E, Freissmuth M, and Milligan G (1992) Gi3 does not contribute to the inhibition of adenylate cyclase when stimulation of an alpha 2-adrenergic receptor causes activation of both Gi2 and Gi3. *Biochem J* **284**:565-568.

McVey M, Ramsay D, Kellett E, Rees S, Wilson S, Pope AJ and Milligan G (2001) Monitoring receptor oligomerization using time-resolved fluorescence resonance energy transfer and bioluminescence resonance energy transfer: The human δ opioid receptor displays constitutive oligomerization at the cell surface which is not regulated by receptor occupancy. *J Biol Chem* **276**:14092-14099.

Massotte D, Brillet K, Kieffer B, and Milligan G (2002) Agonists activate $G_{i1}\alpha$ or $G_{i2}\alpha$ at fused to the human mu opioid receptor differently. *J Neurochem* **81**:1372-1382.

Mercier JF, Salahpour A, Angers S, Breit A, and Bouvier M (2002) Quantitative assessment of beta 1- and beta 2-adrenergic receptor homo- and heterodimerization by bioluminescence resonance energy transfer. *J Biol Chem* **277**:44925-44931.

MOL 13847

Mesnier D, and Baneres JL (2004) Cooperative conformational changes in a G-protein-coupled receptor dimer, the leukotriene B(4) receptor BLT1. *J Biol Chem* **279**:49664-49670.

Milligan G (2002) The use of receptor-G protein fusion proteins for the study of ligand activity. *Receptors Channels* **8**: 309-317.

Milligan G (2004) G protein-coupled receptor dimerization: function and ligand pharmacology. *Mol Pharmacol* **66**: 1-7

Milligan G, and Bouvier M (2005) Methods to monitor the quaternary structure of G protein-coupled receptors. *FEBS J* **272**: 2914-2925.

Milligan G, Carrillo JJ, and Pascal G (2005) Functional complementation and the analysis of GPCR dimerization. In: *The G protein-coupled receptors handbook* (Ed. Devi, LA) pp 267-286. Humana press, Totowa NJ.

Milligan G, Feng G-J, Ward RJ, Sartania N, Ramsay D, McLean AJ, and Carrillo JJ (2004) G protein-coupled receptor fusion proteins in drug discovery. *Current Pharmaceutical Design* **10**:1989-2001.

Molinari P, Ambrosio C, Riitano D, Sbraccia M, Gro MC, and Costa T (2003) Promiscuous coupling at receptor-Galpha fusion proteins. The receptor of one covalent complex interacts with the alpha-subunit of another. *J Biol Chem* **278**: 15778-15788

Monnot C, Bihoreau C, Conchon S, Curnow KM, Corvol P, and Clauser E (1996) Polar residues in the transmembrane domains of the type 1 angiotensin II receptor are required for binding and coupling. Reconstitution of the binding site by co-expression of two deficient mutants. *J Biol Chem* **271**:1507-1513.

MOL 13847

- Moon HE, Cavalli A, Bahia DS, Hoffmann M, Massotte D, and Milligan G (2001) The human δ opioid receptor activates $G_{i1}\alpha$ more efficiently than $G_{o1}\alpha$. *J Neurochem* **76**:1805-1813.
- Overton MC, and Blumer KJ (2002) The extracellular N-terminal domain and transmembrane domains 1 and 2 mediate oligomerization of a yeast G protein-coupled receptor. *J Biol Chem* **277**:41463-41472.
- Pin J-P, Kniazeff J, Liu J, Binet V, Goudet C, Rondard P, and Prezeau L (2005) Allosteric functioning of dimeric class C G protein coupled receptors. *FEBS J* **272**: 2947-2955.
- Ramsay D, Kellett E, McVey M, Rees S, and Milligan G (2002) Homo - and hetero-oligomeric interactions between G protein-coupled receptors in living cells monitored by two variants of bioluminescence resonance energy transfer. Hetero-oligomers between receptor subtypes form more efficiency than between less closely related sequences. *Biochem J* **365**:429-440.
- Stanasila L, Perez JB, Vogel H, and Cotecchia S (2003) Oligomerization of the alpha 1a- and alpha 1b-adrenergic receptor subtypes. Potential implications in receptor internalization. *J Biol Chem* **278**:40239-40251.
- Stevens PA, Pediani J, Carrillo JJ, and Milligan G (2001) Co-ordinated agonist-regulation of receptor and G protein palmitoylation and functional rescue of palmitoylation-deficient mutants of the G protein $G_{i1}\alpha$ following fusion to the α_{1b} -adrenoceptor. Palmitoylation of $G_{i1}\alpha$ is not required for interaction with β/γ complex. *J Biol Chem* **276**: 35883-35890.

MOL 13847

Ward RJ, and Milligan G (2002) Reciprocal mutations of highly conserved residues in transmembrane helices 2 and 7 of the α_{2A} -adrenoceptor restore agonist activation of $G_{i1}\alpha$. *Cell Signalling* **14**, 139-144

Footnotes

These studies were supported, in part, by a Scottish Enterprise 'Proof of concept' award to GM.

Legends for Figures

Figure 1. A hDOP- $G_{i1}\alpha$ C³⁵¹I fusion protein is functional

1A. 10 (1,3) or 20 (2,4) μ g of Pertussis toxin-treated, HEK 293 cell membranes expressing (3,4) or not (1,2) hDOP- $G_{i1}\alpha$ C³⁵¹I were used to measure the binding of [³⁵S]-GTP γ S in the absence (open bars) or presence (filled bars) of 10 μ M DADLE. At assay termination samples were immunoprecipitated with an anti- $G_{i1}\alpha$ / $G_{i2}\alpha$ antiserum and counted.

1B. Membranes, as above, expressing different amounts of hDOP- $G_{i1}\alpha$ C³⁵¹I were used to measure DADLE (10 μ M) stimulation of [³⁵S]-GTP γ S binding.

Data are means \pm S.E.M. of triplicate assays. 2 further experiments produced similar data.

Figure 2. Reconstitution of hDOP- $G_{i1}\alpha$ C³⁵¹I function by co-expression of two non-functional mutants

MOL 13847

Membranes of Pertussis toxin-treated HEK 293 cells expressing 15fmol of hDOP- $G_{i1\alpha}C^{351}I$ (1), hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ (2), hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ (3) or hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ (4) were used to measure basal (open bars) and 10 μ M DADLE (filled bars) binding of [³⁵S]GTP γ S as in Figure 1A. Membranes co-expressing a total of 30 fmol hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ (5) were also analysed as were membranes expressing 15fmol of hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ or 15fmol of hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ that were mixed prior to assay (6). Data represent n = 5 experiments performed in triplicate. * significant (p < 0.05) stimulation by DADLE.

Figure 3. Interactions between co-expressed hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ and hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ monitored by co-immunoprecipitation

3A. Membranes from control HEK 293 cells (1) and cells transiently expressing Flag-hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ (2), c-myc-hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ (3), Flag-hDOPV¹⁵⁰E,V¹⁵⁴D- $G_{i1\alpha}C^{351}I$ + c-myc-hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$ (4) or a mix of membranes from lanes 2 and 3 (5) were immunoprecipitated with anti-Flag antibody and after resolution by SDS-PAGE were immunoblotted to detect c-myc immunoreactivity.

3B. Samples equivalent to 3A were directly resolved by SDS-PAGE and immunoblotted to detect Flag immunoreactivity.

Figure 4. Similar functional pharmacology of hDOP- $G_{i1\alpha}C^{351}I$ and the reconstituted dimer

MOL 13847

- 4A.** Membranes of Pertussis toxin-treated HEK 293 cells expressing 15fmol of hDOP-G_{i1α}C³⁵¹I (open symbols) or hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I + hDOP-G_{i1α}G²⁰²A,C³⁵¹I (closed symbols) were used to measure the ability of increasing concentrations of DADLE to enhance [³⁵S]GTPγS binding as in Figure 1A. Because the absolute amount of [³⁵S]GTPγS bound was less per [³H]diprenorphine binding site in membranes expressing the functionally reconstituted dimer (see Figure 2) data are shown as % maximal signal.
- 4B.** The ability of varying concentrations of naloxone to inhibit [³⁵S]GTPγS binding stimulated by 100nM DADLE is shown. Data are means +/- S.E.M. of n = 3 experiments.

Figure 5. Provision of Flag-Nt-TM1-G_{i1α}C³⁵¹I does not reconstitute substantial function to hDOP-G_{i1α}G²⁰²A,C³⁵¹I

- 5A.** Membranes from control, pertussis toxin-treated HEK 293 cells (1) and those transfected to express Flag-Nt-TM1-G_{i1α}C³⁵¹I (2) or Flag-Nt-TM1-G_{i1α}C³⁵¹I + hDOP-G_{i1α}G²⁰²A,C³⁵¹I (3, 4) were resolved by SDS-PAGE and immunoblotted using the anti-G_{i1α}/G_{i2α}antiserum. The polypeptide(s) migrating with apparent M_r close to 48 kDa is Flag-Nt-TM1-G_{i1α}C³⁵¹I whilst the polypeptide(s) with apparent M_r close to 40kDa is endogenously expressed G_{i1α}/G_{i2α}. Previous studies of HEK293 cells has shown this to be predominantly G_{i2α} which is expressed at some 50pmol/mg membrane protein (McClue et al., 1992).

MOL 13847

5B. Membranes expressing 15 fmol of hDOP-G_{i1α}C³⁵¹I (1), hDOP-G_{i1α}G²⁰²A,C³⁵¹I (2), 10 μg of membranes expressing Flag-Nt-TM1-G_{i1α}C³⁵¹I (estimated to contain 112 fmol of this construct (see Results) (3), membranes co-expressing 15 (4), or 30 (5) fmol of hDOP-G_{i1α}G²⁰²A,C³⁵¹I + (estimated 116.5 fmol (4) or 233 fmol (5)) Flag-Nt-TM1-G_{i1α}C³⁵¹I or a mixture of 10 μg of membranes expressing Flag-Nt-TM1-G_{i1α}C³⁵¹I (112 fmol) + 30 fmol of hDOR-G_{i1α}G²⁰²A,C³⁵¹I (6) were used to measure the binding of [³⁵S]GTPγS in the absence (open bars) or presence of 10 μM (filled bars) or 100nM (checkered bars) DADLE. Data represent means ± S.E.M. of n = 5 experiments performed in triplicate. * significant (p < 0.05) stimulation by DADLE.

Figure 6. Co-expressed hDOP-G_{i1α}G²⁰²A,C³⁵¹I and Nt-TM1-G_{i1α}C³⁵¹I interact and can be co-immunoprecipitated

6A. Membranes from Pertussis toxin-treated HEK 293 cells (1) and equivalent cells transiently expressing c-myc-hDOP-G_{i1α}G²⁰²A,C³⁵¹I (2), Flag-Nt-TM1-G_{i1α}C³⁵¹I (3), or Flag-Nt-TM1-G_{i1α}C³⁵¹I + c-myc-hDOP-G_{i1α}G²⁰²A,C³⁵¹I (4), were immunoprecipitated with anti-Flag antibody and anti-c-myc immunoreactivity detected after separation of the samples by SDS-PAGE (**Upper panel**). The expression of Flag-Nt-TM1-G_{i1α}C³⁵¹I in the appropriate samples was confirmed by immunoblotting membranes with anti-Flag antibody (**Lower panel**).

6B. Membranes from Pertussis toxin-treated HEK 293 cells (1) or those transiently expressing Flag-hDOP (2), c-myc-Nt-TM1 (3), or Flag-hDOP + c-myc-Nt-TM1 (4) were immunoprecipitated with anti-Flag antibody and detected with anti-c-myc antibody after being resolved by SDS-PAGE.

MOL 13847

Figure 7. Reconstitution of hMOP function by co-expression of two non-functional hMOP-G_{i1α} mutants

Membranes of Pertussis toxin-treated HEK 293 cells expressing 15fmol of hMOP-G_{i1α}C³⁵¹I (1); hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I (2), hMOP-G_{i1α}G²⁰²A,C³⁵¹I (3), and either 15 (4) or 30 (5) fmol of co-transfected hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I + hMOP-G_{i1α}G²⁰²A,C³⁵¹I were used to measure [³⁵S]-GTPγS binding in the absence (open bars) or presence (filled bars) of 10 μM DAMGO as in Figure 2. A control was provided by mixing membranes expressing 15 fmol hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I and 15 fmol hMOPG_{i1α}G²⁰²A,C³⁵¹I prior to assay (6). Data represent means +/- S.E.M. of n = 4 experiments performed in triplicate. * Significant (p < 0.05) stimulation by DAMGO.

Figure 8. The characteristics of binding of DAMGO to individually expressed and co-expressed hMOP-G_{i1α} fusion proteins

Membranes expressing hMOP-G_{i1α}C³⁵¹I (squares); hMOP-G_{i1α}G²⁰²A,C³⁵¹I (inverted triangles), hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I (triangles) or both hMOP-G_{i1α}G²⁰²A,C³⁵¹I and hMOPV¹⁶⁹E,V¹⁷³D-G_{i1α}C³⁵¹I (circles) were used to measure the ability of varying concentrations of DAMGO to compete for binding with 1nM [³H]diprenorphine. Data represent n = 4 experiments performed in triplicate.

Figure 9. Reconstitution of rKOP function by co-expression of two non-functional rKOP-G_{i1α} mutants

MOL 13847

Membranes of HEK 293 cells expressing 15fmol of rKOP-G_{i1α}C³⁵¹I (1); rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I (2) , rKOP-G_{i1α}G²⁰²A,C³⁵¹I (3) and 15 (4) or 30 (5) fmol of [³H]diprenorphine binding sites following co-expression of rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I+ rKOP-G_{i1α}G²⁰²A,C³⁵¹I were used to measure [³⁵S]GTPγS binding in the absence (open bars) or presence of 10 μM (filled bars) or 100 nM (checked bars) U69593. A control was performed by mixing membranes expressing 15 fmol of rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I and 15 fmol of rKOP-G_{i1α}G²⁰²A,C³⁵¹I (6). Data represent means +/- S.E.M. of n = 4 experiments performed in triplicate. * Significant (p < 0.05) stimulation by U69593.

Figure 10. The characteristics of binding of U69593 to individually expressed and co-expressed rKOP-G_{i1α} fusion proteins

Membranes expressing rKOP-G_{i1α}C³⁵¹I (squares); rKOP-G_{i1α}G²⁰²A,C³⁵¹I (inverted triangles), rKOPV¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I (triangles) or both rKOP-V¹⁶⁰E,V¹⁶⁴D-G_{i1α}C³⁵¹I and rKOP-G_{i1α}G²⁰²A,C³⁵¹I (circles) were used to measure the ability of varying concentrations of U69593 to compete for binding with 1nM [³H]diprenorphine. Data represent n = 4 experiments performed in triplicate.

MOL 13847

Tables

Table 1. Expression levels and [³H]diprenorphine binding affinity of hDOP-G_{i1α}C³⁵¹I fusion proteins

Construct	B_{max} (fmol/mg)	pK_d
hDOP-G_{i1α}C³⁵¹I	1816 ± 209	9.20 ± 0.03
hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I	2181 ± 228	8.78 ± 0.01***
hDOP-G_{i1α}G²⁰²A,C³⁵¹I	1777 ± 285	9.19 ± 0.05
hDOPV¹⁵⁰E,V¹⁵⁴D-G_{i1α}C³⁵¹I + hDOP-G_{i1α}G²⁰²A,C³⁵¹I	2310 ± 301	8.85 ± 0.02***

Data represent means ± SEM of n=4 experiments performed on different membrane preparations. *** Significantly different from hDOP-G_{i1α}C³⁵¹I, P<0.001

MOL 13847

Table 2

Binding affinity of DADLE for individually expressed and co-expressed hDOP- $G_{i1\alpha}C^{351}I$ fusion proteins.

Construct	pK_h	% K_h sites	pK_i	Hill number
hDOP- $G_{i1\alpha}C^{351}I$	9.03 ± 0.18	63 ± 6	6.79 ± 0.42	-0.39 ± 0.03
hDOPV ¹⁵⁰ E,V ¹⁵⁴ D- $G_{i1\alpha}C^{351}I$	$7.40 \pm 0.24^{**}$	57 ± 8	$4.99 \pm 0.37^*$	-0.50 ± 0.02
hDOP- $G_{i1\alpha}G^{202}A,C^{351}I$	8.70 ± 0.12	59 ± 3	5.82 ± 0.23	-0.34 ± 0.04
hDOPV ¹⁵⁰ E,V ¹⁵⁴ D- $G_{i1\alpha}C^{351}I$ + hDOP- $G_{i1\alpha}G^{202}AC^{351}I$	8.69 ± 0.15	45 ± 9	6.29 ± 0.28	-0.41 ± 0.04

Data represent means \pm SEM of n=4 experiments performed in triplicate on different membrane preparations

* Significantly different from hDOP- $G_{i1\alpha}C^{351}I$, $P < 0.05$

** Significantly different from hDOP- $G_{i1\alpha}C^{351}I$, $P < 0.01$

MOL 13847

Table 3

Binding affinity of DPDPE for individually expressed and co-expressed hDOP-

G_{iiα}C³⁵¹I fusion proteins

Construct	pK _h	% K _h site	pK _i	Hill number
hDOP-G _{iiα} C ³⁵¹ I	8.79 ± 0.09	70 ± 3	5.89 ± 0.25	-0.41 ± 0.005
hDOPV ¹⁵⁰ E,V ¹⁵⁴ D- G _{iiα} C ³⁵¹ I	7.42 ± 0.11**	51±20	5.48 ± 0.41	-0.61 ± 0.04
hDOP- G _{iiα} G ²⁰² A,C ³⁵¹ I	8.88 ± 0.15	65 ± 4	5.82 ± 0.38	-0.40 ± 0.03
hDOPV ¹⁵⁰ E,V ¹⁵⁴ D- G _{iiα} C ³⁵¹ I + hDOP- G _{iiα} G ²⁰² A,C ³⁵¹ I	8.31 ± 0.28	55 ± 12	6.10 ± 0.35	-0.46 ± 0.01

Data represent means ± SEM of n=4 experiments performed in triplicate on different membrane preparations

** Significantly different from hDOR-G_{iiα}C³⁵¹I, P<0.01, 1 way ANOVA

MOL 13847

Table 4

Expression levels and [³H]diprenorphine binding affinity of hMOP-G_{iiα}C³⁵¹I fusion proteins

Construct	B_{max} (fmol/mg)	pK_d
hMOP-G_{iiα}C³⁵¹I	1217 ± 72	9.47 ± 0.08
hMOPV¹⁶⁹E,V¹⁷³D-G_{iiα}C³⁵¹I	901 ± 110	9.39 ± 0.17
hMOP-G_{iiα}G²⁰²A,C³⁵¹I	1251 ± 20	9.52 ± 0.08
hMOPV¹⁶⁹E,V¹⁷³D-G_{iiα}C³⁵¹I + hMOP-G_{iiα}G²⁰²A,C³⁵¹I	1285 ± 120	9.56 ± 0.10

Data represent means ± SEM from n=3 experiments performed in triplicate on different membrane preparations. Statistics were performed using 1 way ANOVA on B_{max} and pK_d numbers.

MOL 13847

Table 5

Binding affinity of DAMGO for individually expressed and co-expressed hMOP- $G_{i1\alpha}C^{351}I$ fusion proteins

Construct	pK_h	% high affinity site	pK_i	Hill number
hMOP-$G_{i1\alpha}C^{351}I$	8.71 ± 0.18	48 ± 4	6.91 ± 0.27	-0.53 ± 0.02
hMOPV¹⁶⁹E,V¹⁷³D- $G_{i1\alpha}C^{351}I$			6.02 ± 0.02*	-0.82 ± 0.05*
hMOP-$G_{i1\alpha}G^{202}A,C^{351}I$	8.68 ± 0.23	60 ± 2	6.69 ± 0.23	-0.52 ± 0.02
hMOPV¹⁶⁹E,V¹⁷³D- $G_{i1\alpha}C^{351}I$ + hMOP- $G_{i1\alpha}G^{202}A,C^{351}I$	8.47 ± 0.06	40 ± 2	6.18 ± 0.07	-0.34 ± 0.01

Data represent means ± SEM from n=3 experiments performed in triplicate on different membrane preparations.

Statistics were performed using 1 way ANOVA on pK_h and pK_i numbers.

* Significantly different from hMOP- $G_{i1\alpha}C^{351}I$, P<0.05

MOL 13847

Table 6

Expression levels and [³H]diprenorphine binding affinity of rKOP-G_{iiα}C³⁵¹I fusion proteins

Construct	B _{max} (fmol/mg)	pK _d
rKOP-G _{iiα} C ³⁵¹ I	2355 ± 193	9.30 ± 0.06
rKOPV ¹⁶⁰ E,V ¹⁶⁴ D-G _{iiα} C ³⁵¹ I	2391 ± 177	8.88 ± 0.04**
rKOP-G _{iiα} G ²⁰² AC ³⁵¹ I	2191 ± 148	9.32 ± 0.06
rKOPV ¹⁶⁰ E,V ¹⁶⁴ D-G _{iiα} C ³⁵¹ I + rKOP-G _{iiα} G ²⁰² AC ³⁵¹ I	2417 ± 187	9.17 ± 0.06

Data represent means ± SEM from n=3 experiments performed in triplicate on different membrane preparations.

Statistics were performed using 1 way ANOVA on B_{max} and pK_d numbers.

** significantly different P<0.01

MOL 13847

Table 7

**Binding affinity of U69593 for individually expressed and co-expressed rKOP-
 $G_{i1\alpha}C^{351}I$ fusion proteins**

Construct	pK_h	% high affinity site	pK_l	Hill number
rKOP- $G_{i1\alpha}C^{351}I$	8.85 ± 0.19	61 ± 2	7.10 ± 0.18	-0.56 ± 0.06
rKOPV ^{160E,V¹⁶⁴D} - $G_{i1\alpha}C^{351}I$			6.00 ± 0.06*	-0.82 ± 0.12
rKOP- $G_{i1\alpha}G^{202}A,C^{351}I$	8.91 ± 0.21	57 ± 3	7.12 ± 0.26	-0.54 ± 0.06
rKOPV ^{160E,V¹⁶⁴D} - $G_{i1\alpha}C^{351}I$ + rKOP- $G_{i1\alpha}G^{202}A,C^{351}I$	8.92 ± 0.31	53 ± 2	6.62 ± 0.23	-0.40 ± 0.04


Data represent means ± SEM of n=4 experiments performed in triplicate on different membrane preparations.

Statistics were performed using 1 way ANOVA on pK_h and pK_l numbers and on high affinity site numbers.

* Significantly different P<0.05

Figure 1

1A.



1B.




Figure 2




Figure 3






Figure 4

4A.



4B.





Figure 5

5A.



5B.





Figure 6

6A.



6B.




Figure 7




Figure 8




Figure 9




Figure 10

