

**G<sub>q</sub>-mediated activation of JNK by the GRP receptor is  
inhibited upon co-stimulation of the G<sub>s</sub>-coupled dopamine  
D<sub>1</sub> receptor in Cos-7 cells**

**Anthony S L Chan and Yung H Wong**

Department of Biochemistry, the Biotechnology Research Institute, and the Molecular  
Neuroscience Center, Hong Kong University of Science and Technology, Clear Water Bay,  
Kowloon, Hong Kong.

**Running title:**  $G_s$ /cAMP inhibits  $G_q$ -mediated JNK activation in Cos-7 cells

**Corresponding author:** Prof. Yung H Wong

Department of Biochemistry, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong.

Tel: (852) 2358 7328

Fax: (852) 2358 1552

Email: [boyung@ust.hk](mailto:boyung@ust.hk)

No. of text pages: 32

No. of tables: 0

No. of figures: 9

No. of references: 40

No. of words in the *Abstract*: 236

No. of words in the *Introduction*: 748

No. of words in the *Discussion*: 1500

**Abbreviations:** GRPR, gastrin-releasing peptide-preferring bombesin receptor; JNK, c-Jun N-terminal kinase; MAPK, mitogen-activated protein kinase; ORL<sub>1</sub>R, opioid receptor-like receptor; PI3K, phosphoinositide-3-kinase.

## **ABSTRACT**

G protein-coupled receptors (GPCRs) of  $G_i$  or  $G_q$  coupling specificity are effectively linked to activation of the JNK cascade. However, little is known with regard to the regulation of JNK by  $G_s$ -coupled receptors. In this report, we utilized Cos-7 cells transfected with the dopamine  $D_1$  receptor ( $D_1R$ ) to illustrate the signaling mechanism for  $G_s$ -mediated JNK activation. Stimulation of  $D_1R$  triggered a weak but significant elevation of JNK activity in a time- and dose-dependent manner. This  $D_1R$ -mediated JNK activation required the participation of  $G\beta\gamma$ , Src-like kinases and small GTPases, while disruptions of cAMP-, PI3K-, and EGFR-mediated signaling had no effect. Co-stimulation of  $D_1R$  with GPCRs of other coupling specificities resulted in differential activation profiles of JNK. Activation of  $G_s$ -coupled  $D_1R$  weakly potentiated the JNK activation induced by the  $G_i$ -coupled opioid receptor-like receptor ( $ORL_1R$ ), but exhibited a significant inhibitory effect on the kinase activity triggered by the  $G_q$ -coupled gastrin-releasing peptide-preferring bombesin receptor (GRPR). Administration of Sp-cAMPS (a cAMP analogue which mimics the  $G_s$ /cAMP signal) also suppressed the JNK activation mediated by  $G_q$ -coupled GRPR, as well as the  $Ca^{2+}$ -induced kinase activation upon thapsigargin treatment. Moreover, the  $Ca^{2+}$  signal from GRPR synergistically potentiated the  $D_1R$ -triggered cAMP elevation, when the two receptors were simultaneously stimulated. Taken together, our results demonstrated that stimulation of  $G_s$ -coupled receptors in Cos-7 cells not only enhanced the JNK activity, but also exhibited a “tuning” effect on the kinase activation mediated by GPCRs of other coupling specificities.

## **INTRODUCTION**

Mitogen-activated protein kinases (MAPKs) are expressed in nearly all eukaryotic cells, and the basic assembly of MAPK pathways is a three-component module conserved from yeast to human (i.e. MAPK kinase kinase  $\rightarrow$  MAPK kinase  $\rightarrow$  MAPK). At least three subtypes of MAPK have been identified so far, they are extracellular signal-regulated kinase (ERK), c-Jun N-terminal kinase (JNK) and p38 (Widmann *et al.*, 1999). Among them, the biological functions of JNK are relatively diverse, ranging from cell proliferation, differentiation, survival, to apoptosis (Dunn *et al.*, 2002). It is believed that JNK may exhibit its multi-functional characteristics by phosphorylating the transcription factors (e.g. c-Jun and ATF-2) and hence modulating cellular gene expression (Dunn *et al.*, 2002).

G protein-coupled receptors (GPCRs) are a major group of transmembrane receptors for detecting extracellular signals (Pfleger and Eidne, 2005). GPCRs, particularly those selectively coupled to the  $G_i$  and  $G_q$  families of G proteins, are efficiently linked to the activation of JNK.  $G_i$ -coupled receptors mainly require a  $G\beta\gamma$ /Src-dependent mechanism to stimulate the JNK cascade (Chan and Wong, 2004a; Kam *et al.*, 2004), while  $G_q$ -coupled receptors utilize both  $G\beta\gamma$ /Src and  $Ca^{2+}$  signals to regulate the kinase activity (Chan and Wong, 2004b). In addition to the participation of Src-like kinases, functional activities of PI3K isoforms and the trans-activation of epidermal growth factor receptors (EGFR) have been proposed as alternative routes for the  $G\beta\gamma$ -mediated pathway (Lopez-Illasaca *et al.*, 1998; Murga *et al.*, 2000; Pierce *et al.*, 2001). Despite the possible differential involvements of signaling intermediates,  $G\beta\gamma$  seems to play an important role for both  $G_i$  and  $G_q$ -mediated JNK activation in response to stimulation of GPCRs.

Receptors coupled to the G<sub>s</sub> family of G proteins are characterized by their abilities to trigger adenylyl cyclase-mediated cAMP formation (Balmforth *et al.*, 1986). Among all GPCRs which show coupling preferences toward a particular G protein family, much less is known with regard to the stimulation of JNK by G<sub>s</sub>-coupled receptor. It has been demonstrated that activation of dopamine D<sub>1</sub> receptors in SK-N-MC human neuroblastoma cells is linked to increased JNK activity in a cAMP and PKA-dependent manner (Zhen *et al.*, 1998), and is therefore readily suppressed by pretreatment with Rp-cAMPS or H89, which specifically inhibit the cAMP/PKA signaling. Another report on the β<sub>2</sub>-adrenergic receptor in DDT1 MF-2 smooth muscle cells also supports this idea, and further suggests the possible involvement of Rho family GTPases in the G<sub>s</sub>-mediated JNK stimulation (Yamauchi *et al.*, 2001). In contrast, numerous studies have demonstrated that G<sub>s</sub>-coupled receptors generally lack the ability to stimulate JNK. For instance, activation of G<sub>s</sub>-coupled adenosine A<sub>2A</sub> receptor in HMC-1 human mast cells does not enhance the JNK activity (Feoktistov *et al.*, 1999). Likewise, studies of Chinese hamster ovary (CHO) cells over-expressing G<sub>s</sub>-coupled corticotrophin-releasing factor receptors (Rossant *et al.*, 1999) or β-adrenergic receptors (Gerhardt *et al.*, 1999) illustrate that these receptors are incapable of stimulating JNK activity upon specific agonist treatment.

Despite the inability of many G<sub>s</sub>-coupled receptors to stimulate JNK, cAMP has been suggested as an activator for JNK activity. Administration of cAMP analogues (e.g. 8-Br-cAMP) and adenylyl cyclase stimulants (e.g. forskolin) have been reported to activate the JNK cascade in DDT1 MF-2 smooth muscle cells (Yamauchi *et al.*, 2001) and MC3T3-E1 preosteoblast cells (Kanno *et al.*, 2004), respectively. However, other groups showed that these cAMP-elevating agents have no effect on JNK activation in hepatocytes (Reinehr *et al.*, 2004), and are even associated with an inhibitory effect on the JNK activity triggered by

epidermal growth factor (McCawley *et al.*, 2000). The inhibitory effect of cAMP on the growth factor-induced JNK activation implies that activation of G<sub>s</sub>-coupled receptors, or administration of cAMP-elevating agents, may suppress the kinase activation triggered by GPCRs of other coupling specificities.

We have previously demonstrated that transfected Cos-7 cells transiently expressing GPCRs are useful cellular models to study the activation of JNK mediated by G<sub>i</sub>- and G<sub>q</sub>-coupled receptors (Chan and Wong, 2000; Chan *et al.*, 2002; Chan and Wong, 2004a), with the experimental results highly consistent with those obtained from endogenous cellular systems (Kam *et al.*, 2003; Kam *et al.*, 2004; Chan and Wong, 2004b). In this report, we utilized Cos-7 cells transfected with G<sub>s</sub>-coupled dopamine D<sub>1</sub> receptor (D<sub>1</sub>R) to illustrate the signaling mechanism for JNK activation, and further investigated the effects of co-stimulation of G<sub>s</sub>-coupled receptor with G<sub>i</sub>- or G<sub>q</sub>-coupled receptors on the JNK activity. Our results suggest that stimulation of G<sub>s</sub>-coupled receptor in Cos-7 cells not only enhances the JNK activity, but also exhibits differential regulatory effects on the kinase activation mediated by G<sub>i</sub>- and G<sub>q</sub>-coupled receptors.

## **MATERIALS AND METHODS**

**Materials.** The cDNAs encoding the GRPR and ORL<sub>1</sub>R were kindly provided by Dr. Jim Battey (National Institutes of Health, Rockville) and Dr. Gang Pei (Shanghai Institutes for Biological Sciences, Shanghai), respectively. The plasmid encoding D<sub>1</sub>R was obtained from Guthrie Research Institute (Sayre, PA). The cDNAs of dominant negative mutants RasS17N and RacT17N were generous gifts from Dr. Eric Stanbridge (University of California, Irvine), RhoT19N and Cdc42T17N were provided by Dr. Marc Symons (Picower Institute for Medical Research, New York). The plasmid of HA-tagged JNK was donated by Dr. Tatyana Voyno-Yasenetskaya (University of Illinois, Chicago). [ $\gamma$ -<sup>32</sup>P]ATP was purchased from DuPont NEN (Boston, MA). PTX and 12CA5 (Anti-HA) antibody were purchased from List Biological Laboratories (Campbell, CA) and Roche Molecular Biochemicals (Indianapolis, IN), respectively. Phospho-CREB antibody and CREB antibody were obtained from Cell Signaling Technology (Beverly, MA). Cell culture reagents including Lipofectamine PLUS<sup>TM</sup> were obtained from Invitrogen (Carlsbad, CA). Bombesin, dopamine and nociceptin were purchased from Sigma (St. Louis, MO). Thapsigargin, BAPTA-AM, AG1478, radicicol, wortmannin, calphostin C, Rp-cAMPs and Sp-cAMPS were obtained from Calbiochem (San Diego, CA).

**Cell culture and transfection.** Green monkey kidney fibroblast Cos-7 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) fetal calf serum (FCS), 50 units/ml penicillin and 50  $\mu$ g/ml streptomycin, and were maintained at 37°C in an environment of 5% CO<sub>2</sub>. Cos-7 cells were transferred to 6-well plates at 4x10<sup>5</sup> cells/well (for JNK assay) or to 12-well plates at 1.5x10<sup>5</sup> cells/well (for AC and PLC assays). Transfection was performed by means of Lipofectamine PLUS<sup>TM</sup> reagents following the supplier's instructions.

***In vitro* JNK assay.** 36 hours after transfection, Cos-7 cells were serum-starved overnight and then treated with various inhibitors if necessary. The cells were then stimulated with specific agonists for the indicated durations and the assays terminated by washing the cells with phosphate-buffered saline (PBS), followed by the addition of 500  $\mu$ l of ice-cold detergent-containing lysis buffer (50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 5 mM EDTA, 40 mM  $\text{NaP}_2\text{O}_7$ , 1% Triton X-100, 1 mM DTT, 200  $\mu$ M  $\text{Na}_3\text{VO}_4$ , 100  $\mu$ M PMSF, 2  $\mu$ g/ml leupeptin, 4  $\mu$ g/ml aprotinin and 0.7  $\mu$ g/ml pepstatin). Lysates obtained were subjected to JNK assay as described previously (Chan and Wong, 2000). 50  $\mu$ l of each supernatant was used for the detection of JNK-HA expression, and the remaining (450  $\mu$ l) was incubated for 1 hour at 4°C with 12CA5 (Anti-HA) antibody (2  $\mu$ g/sample), followed by incubation with 30  $\mu$ l of protein A-agarose (50% slurry) at 4°C for 1 h. The resulting immunoprecipitates were washed twice with lysis buffer and twice with kinase assay buffer (40 mM HEPES, pH 8.0, 5 mM  $\text{Mg}(\text{C}_2\text{H}_3\text{O}_2)_2$ , 1 mM EGTA, 1 mM DTT, 200  $\mu$ M  $\text{Na}_3\text{VO}_4$ ). Washed immunoprecipitates were resuspended in 40  $\mu$ l of kinase assay buffer containing 5  $\mu$ g of GST-c-Jun per reaction, and the kinase reactions were initiated by the addition of 10  $\mu$ l of ATP buffer (50  $\mu$ M ATP containing 2  $\mu$ Ci of  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  per sample). After 30 min incubation at 30°C with occasional shaking, the reactions were terminated by 10  $\mu$ l of 6X sample buffer, and the samples were resolved by 12% SDS-PAGE. The radioactivity incorporated to GST-c-Jun was detected by autoradiogram, and the signal intensity was quantified by PhosphorImager (Molecular Dynamics 445 SI).

**Phospholipase C (PLC) assay.** One day after transfection, Cos-7 cells were labeled for 18 h with 0.75 ml of inositol-free DMEM containing  $[\text{}^3\text{H}]\text{-myo-inositol}$  (5  $\mu$ Ci/ml) and 10% FCS (v/v), followed by serum starvation for 18 h. The cells were then pre-treated in assay medium (20 mM HEPES-buffered DMEM with 20 mM LiCl) for 10 min, and subsequently



stimulated in the presence or absence of indicated drugs for 30 min at 37°C. The reactions were terminated by aspiration of drug-containing medium, followed by the addition of ice-cold 20 mM formic acid solution. After 1 h incubation at 4°C, cell extracts were subjected to ion exchange chromatography as described previously (Chan and Wong, 2004a).

**Adenylyl cyclase (AC) assay.** Transfected Cos-7 cells were labeled with 2  $\mu$ Ci/ml of [<sup>3</sup>H]adenine in DMEM (10% FCS, v/v) for 18 h. After serum starvation for 18 h, cells were treated with the assay medium (DMEM containing 20 mM of HEPES and 1 mM of 1-methyl-3-isobutylxanthine) in the presence or absence of indicated drugs for 30 min at 37°C. The reactions were terminated by aspiration of drug-containing medium, followed by the addition of ice-cold 5% (v/v) trichloroacetic acid (TCA) solution with 1 mM ATP (1 ml/well) and kept at 4°C for 1 h. Intracellular levels of [<sup>3</sup>H]-cAMP were determined by sequential chromatography as described previously (Chan and Wong, 2004a).

## **RESULTS**

***Time/Dose-dependencies of D<sub>1</sub>R-mediated JNK activation.*** We have previously demonstrated that Cos-7 cells transiently expressing D<sub>1</sub>R serve as a reliable model for studying G<sub>s</sub>-mediated signaling, which is linked to a weak but significant activation of JNK upon dopamine treatment (Chan and Wong, 2004a). Hence, we began our study by challenging D<sub>1</sub>R-transfectants with a fixed dose of dopamine for increasing durations, or with increasing doses for a fixed period. The D<sub>1</sub>R-induced JNK activation gradually increased within the first 15 min of dopamine treatment, reaching the maximal level at 30 min and decreased to a near basal level after 60 min of receptor stimulation (Fig. 1A). Increasing agonist concentrations also gradually enhanced the kinase activity with the maximal effect occurring at 10  $\mu$ M of dopamine (Fig. 1B). To examine whether this induced kinase activation was contributed by the G<sub>s</sub>/cAMP signaling, the cells were pretreated with Rp-cAMPS, a cell-permeable cAMP analogue with inhibitory effects on cAMP/PKA-regulated cellular events. Our results showed that Rp-cAMPS (100  $\mu$ M) had no significant inhibitory effect on the D<sub>1</sub>R-induced JNK activity (Fig. 1C). Further investigation on the G<sub>s</sub>-coupled lutropin hormone receptor (LHR) transiently expressed in Cos-7 cells also produced similar results (Fig. 1D). Our control experiment indicated that Rp-cAMPS (100  $\mu$ M) effectively suppressed the cAMP/PKA-mediated phosphorylation of CREB upon stimulation with dopamine (Fig. 1E).

***Signaling intermediates of D<sub>1</sub>R-mediated JNK activation.*** Activation of JNK in response to G<sub>i</sub>- and G<sub>q</sub>-coupled receptors appears to depend on G $\beta\gamma$  subunits (Chan and Wong, 2004a, 2004b). Such a response can be suppressed by transducin which acts as an effective scavenger to remove free G $\beta\gamma$  subunits released upon G protein activation. For Cos-7 cells expressing G<sub>s</sub>-coupled D<sub>1</sub>R, the dopamine-induced JNK activation was almost completely inhibited when transducin was co-expressed in the cells (Fig. 2A). This is consistent with a previous report

wherein  $G\beta\gamma$  subunits are better activators than the  $\alpha$ -subunit of  $G_s$  ( $G\alpha_s$ ) in terms of JNK activation (Coso *et al.*, 1996). In order to reveal the identities of  $G\beta\gamma$ -regulated intermediates for the  $D_1R$ -mediated JNK activation, target-specific inhibitors (radicol and PP1 for Src-like kinases, wortmannin for PI3K isoforms, and AG1478 for EGFR) were employed. Pretreatment of radicol (Fig. 2A) and PP1 (data not shown) significantly suppressed the  $D_1R$ -mediated JNK activation, while wortmannin and AG1478 had no effect on the induced kinase response (Fig. 2A). Intermediates downstream of Src-like kinases include guanine nucleotide exchange factor (GEF)-regulated GTPase activities of the Ras and Rho family members (Kiyono *et al.*, 2000), and the study of their involvements are usually performed by expression of the corresponding dominant negative mutants of these small GTPases (i.e. RasS17N, RacT17N, RhoT19N and Cdc42T17N). When  $D_1R$  was co-expressed with either one of these mutants followed by subsequent dopamine treatment, the induced JNK activation was significantly inhibited in the presence of RasS17N, RacT17N or Cdc42T17N, but not RhoT19N (Fig. 2B). Activation of JNK by  $G_s$ -coupled LHR was also characterized by dependencies on  $G\beta\gamma$ , Src-like kinases and small GTPases (data not shown).

***Differential regulatory effects of  $G_s$ -coupled  $D_1R$  on the JNK activation triggered by  $G_i$ -coupled  $ORL_1R$  and  $G_q$ -coupled  $GRPR$ .*** The preceding experiments demonstrated that  $G_s$ -coupled  $D_1R$  activated JNK in a  $G\beta\gamma$ -, Src-like kinase- and small GTPase-dependent manner (Fig. 2). Such dependencies have also been observed in our previous studies for the same kinase activation mediated by  $G_i$ -coupled  $ORL_1R$  (Chan and Wong, 2000) and  $G_q$ -coupled  $GRPR$  (Chan and Wong, 2004b). In order to investigate the integrated JNK activities upon GPCR co-activation, we co-transfected  $G_s$ -coupled  $D_1R$  with either  $G_i$ -coupled  $ORL_1R$  (Fig. 3A), or  $G_q$ -coupled  $GRPR$  (Fig. 3B) in Cos-7 cells, followed by individual or co-administration of corresponding agonists. For cells co-expressing  $D_1R$  and  $ORL_1R$ , individual

treatment with appropriate agonists (dopamine for D<sub>1</sub>R and nociceptin for ORL<sub>1</sub>R) triggered JNK activation to ~1.5 fold and ~2.0 fold, respectively (Fig. 3A), as compared with the corresponding basal (defined as 1.0 fold). These magnitudes of JNK activities were consistent with the results shown in Fig. 1 for D<sub>1</sub>R, and with our previous report for ORL<sub>1</sub>R (Chan and Wong, 2000), using Cos-7 cells transiently expressing either one of the receptors. Co-stimulation of both D<sub>1</sub>R and ORL<sub>1</sub>R resulted in JNK activation in a roughly additive manner (Fig. 3A). These results in conjunction with our previous findings (Chan and Wong, 2000) showed that, although D<sub>1</sub>R and ORL<sub>1</sub>R utilize a similar mechanistic pathway (Gβγ → Src-like kinase → small GTPase → JNK cascade), D<sub>1</sub>R remained capable of complementing the kinase response triggered by ORL<sub>1</sub>R. To reveal if G<sub>s</sub>-coupled D<sub>1</sub>R also exhibits a similar effect on the JNK activity stimulated by G<sub>q</sub>-coupled GRPR, Cos-7 cells were co-transfected with these two receptors. Activation of D<sub>1</sub>R by dopamine retained the ability to activate JNK weakly, and the GRPR agonist (i.e. bombesin) was still capable of triggering a 6-7 fold kinase activity (Fig. 3B), similar to our results in the previous report (Chan and Wong, 2004b). Unexpectedly, when both D<sub>1</sub>R and GRPR were co-stimulated in the same cells, the induced JNK activation was significantly decreased as compared with the GRPR-mediated response (Fig. 3B). These results suggested that even though the G<sub>s</sub>-coupled D<sub>1</sub>R is linked to activation of JNK activity, it possesses an inhibitory effect on the same kinase activity triggered by G<sub>q</sub>-coupled GRPR in Cos-7 cells.

***Modulation of D<sub>1</sub>R-mediated adenylyl cyclase activation by G<sub>i</sub>-coupled ORL<sub>1</sub>R and G<sub>q</sub>-coupled GRPR.*** The differential regulatory effects of D<sub>1</sub>R on the ORL<sub>1</sub>R- and GRPR-induced JNK activity (Fig. 3) implied that, although G<sub>s</sub>-, G<sub>i</sub>-, and G<sub>q</sub>-coupled receptors utilize a common Gβγ-dependent mechanism to stimulate the JNK activity, G protein subfamily-specific signaling may influence each other and enable an “alternative” regulatory route for

their integrated activation of JNK. Measurements of the  $G_s$ /adenylyl cyclase-mediated cAMP elevation may serve as a useful means for examining this possibility. In Cos-7 cells co-expressing  $G_s$ -coupled  $D_1R$  and  $G_i$ -coupled  $ORL_1R$ , administration of dopamine triggered an elevated cAMP level, while nociceptin treatment induced no observable changes in cAMP production (Fig. 4A). When both of these two agonists were co-administered, a net increase of cAMP formation was produced, which was associated with a significant inhibition as compared to the dopamine-induced activity (Fig. 4A). This result agreed with the classical G protein signaling model that the inhibitory  $G_i$  signal opposed the stimulatory  $G_s$  signal on the adenylyl cyclase-mediated cAMP formation.

Similarly, when Cos-7 cells co-expressing  $G_s$ -coupled  $D_1R$  and  $G_q$ -coupled GRPR were stimulated by their specific agonists, dopamine treatment was capable of triggering enhanced cAMP formation, while bombesin did not significantly stimulate this activity (Fig. 4B). However, when dopamine and bombesin were co-administered, the induced cAMP production was nearly doubled as compared to the dopamine effect (Fig. 4B). This indicated that the  $G_q$  signal from GRPR synergizes with the  $G_s$ -mediated adenylyl cyclase activation by  $D_1R$ . To investigate the mechanism which gave rise to this synergistic response, a series of experiments was performed by target-specific inhibition on the  $G_q$  signaling, or by testing the  $G\beta\gamma$ -dependency of adenylyl cyclase activation. For cells co-transfected with transducin, removal of  $G\beta\gamma$  subunits (released from  $G_s$  and  $G_q$ ) was incapable of eliminating the synergistic response upon co-stimulation of  $D_1R$  and GRPR (Fig. 4C). When the cells were pretreated with calphostin C to inhibit PKC functions, dopamine treatment remained capable of increasing the cAMP level, and the subsequent synergistic effect with bombesin was still present (Fig. 4D). Chelation of intracellular  $Ca^{2+}$  by BAPTA-AM did not affect the dopamine or bombesin responses, but it almost completely eliminated the synergistic adenylyl cyclase

activation, by returning the cAMP level to nearly the same level as the dopamine treatment alone (Fig. 4E). The above results suggested that the  $G_q$ -induced  $Ca^{2+}$  activity is important for the GRPR-mediated potentiation of the  $D_1R$ -triggered cAMP elevation. In order to provide direct evidence for the synergism of  $G_s$ -mediated cAMP formation in response to the elevated  $Ca^{2+}$  activity, Cos-7 cells expressing  $D_1R$  alone were stimulated with dopamine and thapsigargin (an extensively used agent for elevating intracellular  $Ca^{2+}$  level) separately or simultaneously. Again, dopamine treatment significantly enhanced the cAMP level but thapsigargin failed to do so, while their co-application generated synergistic adenylyl cyclase activation (Fig. 4F). These results showed that the  $G_q$ -coupled GRPR may utilize a  $Ca^{2+}$  signal to potentiate the adenylyl cyclase activity induced by  $G_s$ -coupled  $D_1R$ .

***Co-stimulation of  $G_s$ -coupled  $D_1R$  with either  $G_q$ -coupled GRPR or  $G_i$ -coupled  $ORL_1R$  had no effect on the PLC activity.***  $G_q$ -coupled receptors are known to utilize both  $G\beta\gamma$ - as well as PLC-dependent pathways to regulate the JNK activity (Chan and Wong, 2004b). Thus, we examined if signal integration might have occurred along the  $G_q$ /PLC pathway. In Cos-7 cells co-expressing  $G_q$ -coupled GRPR and  $G_s$ -coupled  $D_1R$ , treatment with bombesin stimulated  $IP_3$  formation, while activation of  $D_1R$  by dopamine neither stimulated PLC, nor potentiated the GRPR-triggered response (Fig. 5A). Since the activities of PLC  $\beta$ -isoforms may be potentiated by  $G\beta\gamma$  subunits released upon  $G_i$  activation (Chan *et al.*, 2000), a similar assay was hence performed with Cos-7 cells co-expressing  $G_s$ -coupled  $D_1R$  and  $G_i$ -coupled  $ORL_1R$  to determine whether co-operative signaling between  $G_s$  and  $G_i$  is capable of elevating the  $IP_3$  formation. However, irrespective of whether these cells were treated with dopamine and nociceptin individually or simultaneously, no significant enhancement of  $IP_3$  formation was observed (Fig. 5B).

*cAMP serves as a suppressor of G<sub>q</sub>-mediated JNK activation in Cos-7 cells.* The preceding experiments demonstrated that co-stimulation of D<sub>1</sub>R and GRPR was associated with a synergistic elevation of cAMP (Fig. 4B) but a diminished magnitude of JNK activation (Fig. 3B). To examine if the cAMP signal can suppress G<sub>q</sub>-mediated JNK activation, Cos-7 cells were transfected with D<sub>1</sub>R and GRPR and then treated with Sp-cAMPS (a cell-permeable analogue with stimulatory effect on cAMP-mediated signaling) in the absence or presence of bombesin. Sp-cAMPS itself did not stimulate JNK, instead, it significantly suppressed the bombesin-mediated JNK activation (Fig. 6A). The inability of Sp-cAMPS to suppress the JNK activation mediated by G<sub>i</sub>-coupled ORL<sub>1</sub>R suggested that cAMP signaling specifically inhibited G<sub>q</sub>- but not G<sub>i</sub>-mediated JNK activation in Cos-7 cells (Fig. 6B). Further investigations showed that this selective inhibitory effect on GRPR-mediated JNK activation showed a dose-dependent character towards Sp-cAMPS (Fig. 7). In addition to the Gβγ-dependent pathway, G<sub>q</sub>-coupled receptors also require a Ca<sup>2+</sup> component to stimulate the JNK cascade (Chan and Wong, 2004b). It is possible that cAMP exerts its inhibitory effect on the Ca<sup>2+</sup>-mediated pathway, resulting in a G<sub>s</sub>-mediated suppression of the G<sub>q</sub>-induced JNK activation. In fact, our previous report has already demonstrated that when transfected Cos-7 cells were treated with thapsigargin to induce a Ca<sup>2+</sup>-dependent JNK activation, co-treatment with increasing concentration of Sp-cAMPS resulted in a gradual decrease of kinase activities (Chan and Wong, 2004a). All these data support the idea that co-operation of G<sub>s</sub> and G<sub>q</sub> signaling may result in decreased G<sub>q</sub>-mediated JNK activation, which is primarily caused by a suppressive effect of cAMP on the Ca<sup>2+</sup> signaling component of the G<sub>q</sub>-mediated JNK activity.

*Disruption of Ca<sup>2+</sup> signaling by BAPTA-AM suppressed the D<sub>1</sub>R/GRPR-mediated JNK activity to the D<sub>1</sub>R/ORL<sub>1</sub>R-induced level.* As discussed earlier, G<sub>s</sub>-, G<sub>i</sub>- and G<sub>q</sub>-coupled receptors utilize the Gβγ-dependent pathway to regulate the activity of JNK cascade in Cos-7

cells, with the  $G_q$ -mediated response associated with an additional involvement of  $Ca^{2+}$  signaling. These characteristics imply that in the absence of the elevated  $Ca^{2+}$  signal upon  $G_q$  activation, co-stimulation of  $G_q$  and  $G_s$  should be similar to that of  $G_i$  and  $G_s$ , in terms of JNK activation mediated by  $G\beta\gamma$  subunits from two different G protein sub-families. In order to examine this hypothesis, Cos-7 cells co-expressing  $G_q$ -coupled GRPR and  $G_s$ -coupled  $D_1R$  were pre-treated with BAPTA-AM to deplete intracellular  $Ca^{2+}$ , followed by co-stimulation with bombesin and dopamine. Cos-7 cells co-expressing  $G_i$ -coupled  $ORL_1R$  and  $G_s$ -coupled  $D_1R$  and co-stimulated with nociceptin and dopamine were employed as a control for comparison. Indeed, our results showed that disruption of  $Ca^{2+}$  signaling by BAPTA-AM attenuated the GRPR/ $D_1R$ -induced JNK activity to a level similar to that of the  $ORL_1R$ / $D_1R$ -mediated response (Fig. 8). This finding further demonstrates the importance of  $Ca^{2+}$  mobilization for the GRPR/ $D_1R$ -induced JNK response, whereas a cAMP signal (initiated from  $G_s$  and further potentiated by  $G_q$ ) exhibits an inhibitory effect on the activity of the JNK cascade (Fig. 9B).



## **DISCUSSION**

Receptors coupled to G<sub>s</sub> have been suggested to utilize cAMP to activate JNK. In the present report, we provide evidence that G<sub>s</sub>-linked receptors are also capable of stimulating this kinase via an alternative pathway, in which Gβγ subunits serve as the primary player in the signal transduction. Since the various studies were performed in different cellular models, we cannot rule out the possibility that such differential dependency is a function of cell type specificity. Irrespective of whether the G<sub>s</sub>-mediated JNK activation is processed in a cAMP-dependent (Zhen *et al.*, 1998) or cAMP-independent manner (as shown in this report), it seems that small GTPases serve as common intermediates for both signaling models.

The activities of small GTPases are regulated by guanine nucleotide exchange factors (GEFs). cAMP-responsive GEFs which can trigger ERK activation have been proposed (Laroche-Joubert *et al.*, 2002), but the corresponding candidates for JNK regulation remain unclear. An exchange protein activated by cAMP (Epac) has been suggested as a cAMP-responsive GEF in rat renal collecting duct tubule cells for the activation of the small GTPase, Rap, which activates the ERK (Laroche-Joubert *et al.*, 2002) rather than the JNK cascade (Mochizuki *et al.*, 2000). However, selective activations of PKA and Epac in various cell lines (including CHO, PC12 and HEK293 cells) revealed that the former, but not the latter, contributes to the cAMP-dependent ERK activation (Enserink *et al.*, 2002). A recent study even proposed that Epac is capable of activating the JNK cascade in a small GTPase-independent manner (Hochbaum *et al.*, 2003). These studies support opposing ideas and it remains unclear as to the identities of GEFs which are responsible for cAMP-mediated JNK activation.

In contrast, GEFs which are responsive to  $G\beta\gamma$  and the subsequent downstream Src-like kinases have been defined more clearly. The activity of Ras-GRF1 (a GEF for Ras and Rac) can be promoted through tyrosine phosphorylation by Src, which in turn leads to activation of JNK in a Rac-dependent manner (Kiyono *et al.*, 2000). In Cos-7 cells, both  $G_i$ - and  $G_q$ -coupled receptors trigger JNK activation via a  $G\beta\gamma$ /Src-dependent mechanism, which can be suppressed by dominant negative mutants of both Sos (i.e. Son of Sevenless) and small GTPases (Chan and Wong, 2004b; Kam *et al.*, 2004). These findings clearly support the involvement of GEFs and small GTPases in the activation of JNK by GPCRs. In addition to Src-like kinases, PI3K acts as a downstream effector for  $G\beta\gamma$  (Lopez-Illasaca *et al.*, 1998), and its phospholipid products are capable of regulating the binding of certain GEFs to Rac (Das *et al.*, 2000). EGFR transactivation has been proposed as a possible route which links GPCR signaling to the activation of MAPK subgroups (Luttrell *et al.*, 1997). However, our results clearly demonstrated that  $G_s$ -coupled receptors (as least for the  $D_1R$  and LHR), similar to other receptors linked to  $G_i$  and  $G_q$  (Chan and Wong, 2004b; Kam *et al.*, 2004), utilized Src-like kinases, rather than PI3K or EGFR as the primary intermediate to stimulate JNK activity in Cos-7 cells. It should be noted that further investigations on other  $G_s$ -coupled receptors are required in order to establish a more definitive conclusion on the  $G_s$ -induced activation of JNK.

Investigations on receptors with specific G protein coupling preferences allow us to understand how different G proteins utilize various intermediates to trigger a biological response. However, in a physiological environment, cells are likely to have their receptors co-stimulated by different arrays of extracellular stimulus, and the resulting outcomes are determined by the integration of signals that occur inside the cells. There is increasing evidence to demonstrate the existence of signaling “cross-talk” between GPCRs of different

coupling specificities (Selbie and Hill, 1998; Hur and Kim, 2002). By expressing different GPCRs in Cos-7 cells followed by stimulation with specific agonists individually or simultaneously, complex patterns are observed for the regulation of cAMP and IP<sub>3</sub> levels, as well as for JNK activities.

Co-stimulation of G<sub>s</sub>-coupled D<sub>1</sub>R and G<sub>i</sub>-coupled ORL<sub>1</sub>R resulted in an elevated cAMP level which was lower than that of the D<sub>1</sub>R response, but no significant enhancement of IP<sub>3</sub> level could be observed (Fig. 5). As illustrated in the present report, cAMP is not as effective as Gβγ subunits for the triggering of JNK activation in Cos-7 cells. Hence, the Gβγ subunits released from G<sub>s</sub> and G<sub>i</sub> may co-operate with each other, utilizing a Gβγ/Src-dependent pathway to mediate the JNK activation in a roughly additive manner. The reason for higher JNK activating capability associated with G<sub>i</sub> as compared to G<sub>s</sub> is currently unknown, but it should be borne in mind that the Gα subunits of different G protein members may preferably associate with specific isoforms of Gβγ subunits (Albert and Robillard, 2002), which may in turn mediate the downstream signaling in a similar, but not identical manner. Alternatively, the total amount of Gβγ subunits releasable from G<sub>i</sub>/G<sub>o</sub> (with five subtypes) is expected to be greater than those from G<sub>s</sub>. Hence, G<sub>s</sub> may weakly augment the G<sub>i</sub>-mediated JNK activation by contributing additional Gβγ subunits to the pathway (Fig. 9A).

The G<sub>s</sub> signal neither potentiates nor inhibits the G<sub>q</sub>-mediated IP<sub>3</sub> formation in Cos-7 cells. In contrast, G<sub>s</sub>-induced cAMP elevation is synergistically potentiated by the Ca<sup>2+</sup> signal from G<sub>q</sub>, indicating that the Ca<sup>2+</sup>-responsive adenylyl cyclase isoforms (e.g. type I, V and VI) may be predominantly expressed in the cells (Wayman *et al.*, 1994). Interestingly, G<sub>s</sub>-induced cAMP signaling is associated with an inhibitory function on the Ca<sup>2+</sup>-dependent JNK pathway. Such an effect may be magnified in the presence of Ca<sup>2+</sup>-mediated synergistic cAMP formation,

which in turn diminishes the co-operative effect between  $\text{Ca}^{2+}$  and  $\text{G}\beta\gamma$  of  $\text{G}_q$  (Chan and Wong, 2004b), resulting in a decreased JNK activation as compared to the normal  $\text{G}_q$ -mediated response (Fig. 9B). This idea is supported by our previous finding that, activation of vasopressin  $\text{V}_2$  receptor (with dual coupling specificities towards  $\text{G}_s$  and  $\text{G}_q$ ) triggers JNK activity with a magnitude higher than the  $\text{G}_s$ -, but lower than the  $\text{G}_q$ -mediated activity (Chan and Wong, 2004a). In fact,  $\text{G}_q$ -mediated JNK pathway may involve the  $\text{Ca}^{2+}$ -responsive focal adhesion kinase family (e.g. Pyk2) as important intermediates, and the possible cAMP-mediated inhibition towards this kinase signaling (Li *et al.*, 1997) supports our mechanistic model for the  $\text{G}_q/\text{G}_s$ -integrated JNK activity.

One might argue that the decreased GRPR-mediated JNK activation could be due to receptor desensitization induced by  $\text{D}_1\text{R}$  signaling components (e.g. the cAMP-dependent protein kinase). It should be noted that the elevated  $\text{IP}_3$  formation triggered by GRPR was not affected by simultaneous activation of  $\text{D}_1\text{R}$  (Fig. 5A). Hence, it is unlikely that  $\text{D}_1\text{R}$  signaling suppresses the GRPR-mediated JNK activation by down-regulating the components in the GRPR/ $\text{G}_q$ /phospholipase C pathway. Although the present study did not provide any information about the possible occurrence of receptor dimerization, however, the suppression on GRPR-mediated JNK activation by  $\text{D}_1\text{R}$  signaling (Fig. 3B) or direct administration of Sp-cAMPS (Fig. 6A) implied that, irrespective of possible receptor dimerization (between GRPR and  $\text{D}_1\text{R}$ ), the  $\text{G}_s/\text{cAMP}$  signal definitely possesses an inhibitory effect on the  $\text{G}_q/\text{Ca}^{2+}$ -mediated JNK activation in Cos-7 cells (Chan and Wong, 2004a).

We have previously demonstrated that GPCRs of different coupling specificities have differential abilities to activate JNK (i.e.  $\text{G}_q > \text{G}_i > \text{G}_s$ ), and signals from receptor tyrosine kinases (e.g. EGF receptor) selectively augment the  $\text{G}_i$ -mediated stimulation of JNK activity

in Cos-7 cells (Chan and Wong, 2004a). In this report, we further illustrate that modulation of JNK activity also exists in the integration of GPCR signals. Since the JNK activation induced by  $G_q$ -coupled receptors (including GRPR, bradykinin  $BK_2$ , muscarinic acetylcholine  $M_1$  and histamine  $H_1$  receptors) in Cos-7 cells are  $Ca^{2+}$ -dependent (Chan and Wong, 2004b), and cAMP serves as an effective inhibitor for the  $Ca^{2+}$ -induced JNK activity in the same cells (Chan and Wong, 2004a), it is possible that the JNK activation triggered by other  $G_q$ -coupled receptors are subject to similar inhibitory regulation by the  $G_s$ /cAMP signal. The JNK activity induced by  $G_q$ -coupled receptors was diminished by co-activation of  $G_s$ -coupled receptors, resulting in a kinase activity lower than  $G_q$ , but higher than the  $G_i$ - and  $G_s$ -mediated responses. On the other hand, the  $G_i$ -induced JNK activity was weakly enhanced upon co-stimulation of  $G_s$ -coupled receptor. Taking all these findings together, a graded activation profile of JNK could be achieved by linking GPCRs of different coupling preferences to a signaling network, in which specific (e.g. cAMP from  $G_s$  and  $Ca^{2+}$  from  $G_q$ ) as well as common (e.g.  $G\beta\gamma$ /Src) signals from different G proteins can be integrated with each other to modulate the activities of the JNK cascade (Fig. 9).

The present report illustrates the “fine tuning” character of JNK activation in response to cross-communication between GPCR of different coupling specificities. In fact, the magnitudes of activity as well as the duration of MAPK activation are required to trigger a sequential cascade of transcription factor induction, and their subsequent gene transcriptional activities (Harada *et al.*, 2001). The resulting gene products may in turn, serve as critical factors which determine cell fate such as proliferation, differentiation or apoptosis (Kobayashi and Tsukamoto, 2001). Further investigation should be focused on suitable endogenous cellular systems in order to define the possible biological consequence of this integrated JNK activity.

### **ACKNOWLEDGEMENTS**

We thank the kind donation of various cDNAs from Drs. Eric Stanbridge, Gang Pei, Jim Battey, Marc Symons and Tatyana Voyno-Yasenetskaya for our present study. We also thank Dr. David New for helpful discussion, Lydia Poon and Wendy Yeung for technical support in cell culture.

### **FOOTNOTES**

This work was supported in part by grants from the Research Grants Council of Hong Kong (HKUST 3/03C), the University Grants Committee (AoE/B-15/01), and the Hong Kong Jockey Club. YHW was a recipient of the Croucher Senior Research Fellowship.

## **REFERENCES**

Albert PR and Robillard L (2002) G protein specificity: traffic direction required. *Cell Signal* 14(5): 407-418.

Balmforth AJ, Ball SG, Freshney RI, Graham DI, McNamee HB and Vaughan PF (1986) D<sub>1</sub> dopaminergic and  $\beta$ -adrenergic stimulation of adenylate cyclase in a clone derived from the human astrocytoma cell line G-CCM. *J Neurochem* 47:715-719.

Chan JSC, Lee JWM, Ho MKC, and Wong YH (2000) Preactivation permits subsequent stimulation of phospholipase C by G<sub>i</sub>-coupled receptors. *Mol. Pharmacol.* 57: 700-708.

Chan AS, Wong YH (2000) Regulation of c-Jun N-terminal kinase by the ORL<sub>1</sub> receptor through multiple G proteins. *J Pharmacol Exp Ther.* 295(3):1094-1100.

Chan AS, Lai FP, Lo RK, Voyno-Yasenetskaya TA, Stanbridge EJ, Wong YH (2002) Melatonin MT<sub>1</sub> and MT<sub>2</sub> receptors stimulate c-Jun N-terminal kinase via pertussis toxin-sensitive and -insensitive G proteins. *Cell Signal.* 14(3):249-257.

Chan AS, Wong YH (2004a) Epidermal growth factor differentially augments G<sub>i</sub>-mediated stimulation of c-Jun N-terminal kinase activity. *Br J Pharmacol.* 142(4):635-646.

Chan AS, Wong YH (2004b) G $\beta\gamma$  signaling and Ca<sup>2+</sup> mobilization co-operate synergistically in a Sos and Rac-dependent manner in the activation of JNK by G<sub>q</sub>-coupled receptors. *Cell Signal.* 16(7):823-836.

Coso OA, Teramoto H, Simonds WF and Gutkind JS (1996) Signaling from G protein-coupled receptors to c-Jun kinase involves beta gamma subunits of heterotrimeric G proteins acting on a Ras and Rac1-dependent pathway. *J Biol Chem* 271(8):3963-3966.

Das B, Shu X, Day GJ, Han J, Krishna UM, Falck JR and Broek D (2000) Control of intramolecular interactions between the pleckstrin homology and Dbl homology domains of Vav and Sos1 regulates Rac binding. *J Biol Chem* 275(20):15074-15081

Disa J, Parameswaran N, Nambi P, Aiyar N (2000) Involvement of cAMP-dependent protein kinase and pertussis toxin-sensitive G-proteins in CGRP mediated JNK activation in human neuroblastoma cell line. *Neuropeptides*. 34(3-4):229-233.

Dunn C, Wiltshire C, MacLaren A, Gillespie DA (2002) Molecular mechanism and biological functions of c-Jun N-terminal kinase signalling via the c-Jun transcription factor. *Cell Signal*. 14(7):585-593.

Ellsworth BS, White BR, Burns AT, Cherrington BD, Otis AM, Clay CM (2003) c-Jun N-terminal kinase activation of activator protein-1 underlies homologous regulation of the gonadotropin-releasing hormone receptor gene in  $\alpha$  T3-1 cells. *Endocrinology*. 144(3):839-849.

Enserink JM, Christensen AE, de Rooij J, van Triest M, Schwede F, Genieser HG, Doskeland SO, Blank JL, Bos JL (2002) A novel Epac-specific cAMP analogue demonstrates independent regulation of Rap1 and ERK. *Nat Cell Biol*. 4(11):901-906.

Feoktistov I, Goldstein AE, Biaggioni I (1999) Role of p38 mitogen-activated protein kinase and extracellular signal-regulated protein kinase kinase in adenosine A<sub>2B</sub> receptor-mediated interleukin-8 production in human mast cells. *Mol Pharmacol*. 55(4):726-734.

Gerhardt CC, Gros J, Strosberg AD, Issad T (1999) Stimulation of the extracellular signal-regulated kinase 1/2 pathway by human  $\beta_3$  adrenergic receptor: new pharmacological profile and mechanism of activation. *Mol Pharmacol*. 55(2):255-262.

Harada T, Morooka T, Ogawa S and Nishida E (2001) ERK induces p35, a neuron-specific activator of Cdk5, through induction of Egr1. *Nat Cell Biol* 3(5):453-459.

Hochbaum D, Tanos T, Ribeiro-Neto F, Altschuler D, Coso OA (2003) Activation of JNK by Epac is independent of its activity as a Rap guanine nucleotide exchanger. *J Biol Chem*. 278(36):33738-33746.

Hur EM and Kim KT (2002) G protein-coupled receptor signaling and cross-talk achieving rapidity and specificity. *Cell Signal* 14:397-405.



Kam AY, Chan AS, Wong YH (2003) Rac and Cdc42-dependent regulation of c-Jun N-terminal kinases by the  $\delta$ -opioid receptor. *J Neurochem.* 84(3):503-513.

Kam AY, Chan AS, Wong YH (2004)  $\kappa$ -opioid receptor signals through Src and focal adhesion kinase to stimulate c-Jun N-terminal kinases in transfected COS-7 cells and human monocytic THP-1 cells. *J Pharmacol Exp Ther.* 310(1):301-310.

Kanno Y, Tokuda H, Nakajima K, Ishisaki A, Shibata T, Numata O, Kozawa O (2004) Involvement of SAPK/JNK in prostaglandin E<sub>1</sub>-induced VEGF synthesis in osteoblast-like cells. *Mol Cell Endocrinol.* 220(1-2):89-95.

Kiyono M, Kaziro Y and Satoh T (2000) Induction of rac-guanine nucleotide exchange activity of Ras-GRF1/CDC25(Mm) following phosphorylation by the nonreceptor tyrosine kinase Src. *J Biol Chem* 275:5441-5446.

Kobayashi K and Tsukamoto I (2001) Prolonged Jun N-terminal kinase (JNK) activation and the upregulation of p53 and p21(WAF1/CIP1) preceded apoptosis in hepatocytes after partial hepatectomy and cisplatin. *Biochim Biophys Acta* 1537:79-88.

Laroche-Joubert N, Marsy S, Michelet S, Imbert-Teboul M, Doucet A (2002) Protein kinase A-independent activation of ERK and H,K-ATPase by cAMP in native kidney cells: role of Epac I. *J Biol Chem.* 277(21):18598-18604.

Li X, Yu H, Graves LM and Earp HS (1997) Protein kinase C and protein kinase A inhibit calcium-dependent but not stress-dependent c-Jun N-terminal kinase activation in rat liver epithelial cells. *J Biol Chem* 272:14996-15002.

Lopez-Illasaca M, Gutkind JS and Wetzker R (1998) Phosphoinositide 3-kinase  $\gamma$  is a mediator of G $\beta\gamma$ -dependent Jun kinase activation. *J Biol Chem* 273:2505-2508.

Luttrell LM, Della Rocca GJ, van Biesen T, Luttrell DK, Lefkowitz RJ (1997) G $\beta\gamma$  subunits mediate Src-dependent phosphorylation of the epidermal growth factor receptor. A scaffold for G protein-coupled receptor-mediated Ras activation. *J Biol Chem.* 272(7):4637-4644.

McCawley LJ, Li S, Benavidez M, Halbleib J, Wattenberg EV, Hudson LG (2000) Elevation of intracellular cAMP inhibits growth factor-mediated matrix metalloproteinase-9 induction and keratinocyte migration. *Mol Pharmacol.* 58(1):145-151.

Mochizuki N, Ohba Y, Kobayashi S, Otsuka N, Graybiel AM, Tanaka S, Matsuda M (2000) Crk activation of JNK via C3G and R-Ras. *J Biol Chem.* 275(17):12667-12671.

Murga C, Fukuhara S and Gutkind JS (2000) A novel role for phosphatidylinositol 3-kinase  $\beta$  in signaling from G protein-coupled receptors to Akt. *J Biol Chem* 275:12069-12073.

Pfleger KD, Eidne KA (2005) Monitoring the formation of dynamic G-protein coupled receptor-protein complexes in living cells. *Biochem J.* 385(Pt 3):625-637.

Pierce KL, Tohgo A, Ahn S, Field ME, Luttrell LM, Lefkowitz RJ (2001) Epidermal growth factor (EGF) receptor-dependent ERK activation by G protein-coupled receptors: a co-culture system for identifying intermediates upstream and downstream of heparin-binding EGF shedding. *J Biol Chem* 276:23155-23160.

Reinehr R, Becker S, Hongen A, Haussinger D (2004) The Src family kinase Yes triggers hyperosmotic activation of the epidermal growth factor receptor and CD95. *J Biol Chem.* 279(23):23977-23987.

Rossant CJ, Pinnock RD, Hughes J, Hall MD, McNulty S (1999) Corticotropin-releasing factor type 1 and type 2 $\alpha$  receptors regulate phosphorylation of calcium/cyclic adenosine 3',5'-monophosphate response element-binding protein and activation of p42/p44 mitogen-activated protein kinase. *Endocrinology.* 140(4):1525-1536.

Selbie LA and Hill SJ (1998) G protein-coupled receptor cross-talk: the fine-tuning of multiple receptor signaling pathways. *Trends Pharmacol Sci* 19: 87-93.

Watanabe G, Pena P, Albanese C, Wilsbacher LD, Young JB, Pestell RG (1997) Adrenocorticotropin induction of stress-activated protein kinase in the adrenal cortex in vivo. *J Biol Chem.* 272(32):20063-20069.

Wayman GA, Impey S, Wu Z, Kindsvogel W, Prichard L and Storm DR (1994) Synergistic activation of the type I adenylyl cyclase by  $Ca^{2+}$  and  $G_s$ -coupled receptors in vivo. *J Biol Chem* 269: 25400-25405.

Widmann C, Gibson S, Jarpe MB, Johnson GL (1999) Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol Rev.* 79(1):143-180.

Yamanchi J, Hirasawa A, Miyamoto Y, Itoh H and Tsujimoto G (2001)  $\beta_2$ -adrenergic receptor/cycle adenosine monophosphate (cAMP) leads to JNK activation through Rho family small GTPases. *Biochem Biophys Res Commun* 284:1199-1203.

Zhen X, Uryu K, Wang HY and Friedman E (1998)  $D_1$  dopamine receptor agonists mediate activation of p38 mitogen-activated protein kinase and c-Jun N-terminal kinase by a protein kinase A-dependent mechanism in SK-N-MC human neuroblastoma cells. *Mol Pharmacol* 54:453-458.

## **FIGURE LEGENDS**

**Figure 1. Activation of JNK triggered by G<sub>s</sub>-coupled receptors.** Cos-7 cells were co-transfected with the cDNAs of JNK-HA, the G<sub>s</sub>-coupled D<sub>1</sub>R (A, B, C and E), or the G<sub>s</sub>-coupled LHR (D). D<sub>1</sub>R-expressing cells were stimulated with dopamine (10 μM) for increasing durations (A), or incubated with increasing concentrations of the agonist for 30 min (B). Cells expressing D<sub>1</sub>R (C and E) or LHR (D) were also pretreated in the absence (control) or presence of Rp-cAMP (100 μM, 30 min) before stimulation with their agonists (10 μM dopamine for D<sub>1</sub>R and 1 μg/ml hCG for LHR). JNK assay was performed as described in *Materials and Methods*. The values shown represent the mean ± S.E. from at least four separate experiments. \*Activation of D<sub>1</sub>R (C) or LHR (D) significantly activate the JNK activity as compared to the basal (Bonferroni paired *t*-test, P<0.05). (E) D<sub>1</sub>R-induced CREB phosphorylation was inhibited by pre-treatment with Rp-cAMPS.

**Figure 2. Gβγ subunits, Src-like kinases and small GTPases serve as important intermediates for D<sub>1</sub>R-mediated JNK activation.** Cos-7 cells were transfected with the cDNAs of JNK-HA and the G<sub>s</sub>-coupled D<sub>1</sub>R (A and B). The cells were either co-expressing transducin (Gα<sub>t</sub>) for Gβγ scavenging, pretreated with radicicol (10 μM, 1 h for Src-like kinases), wortmannin (100 nM, 15 min for PI3K) or AG1478 (500 nM, 30 min for EGFR) for specific signal disruption (A), or transfected with different dominant negative mutants of small GTPases (B) before agonist treatment. JNK assay was performed as described in *Materials and Methods*. Values shown represent the mean ± S.E. from three to six separate experiments. \*Administration of dopamine induced significant activation of JNK as compared to the corresponding basal. #Removal of Gβγ by transducin, pretreatment with radicicol, and

the presence of RasS17N, RacT17N and Cdc42T17N significantly inhibited the JNK activation as compared with the control group (Bonferroni paired *t*-test,  $P < 0.05$ ).

**Figure 3. Differential regulatory effects of  $G_s$ -coupled  $D_1R$  on the JNK activation triggered by  $G_i$ -coupled  $ORL_1R$  and  $G_q$ -coupled GRPR.** Cos-7 cells co-expressing JNK-HA and  $D_1R$  with  $ORL_1R$  (A) or GRPR (B) were stimulated with their corresponding agonists (10  $\mu$ M dopamine, DOP for  $D_1R$ , 100 nM nociceptin, OFQ for  $ORL_1R$ , and 100 nM bombesin, BBS for GRPR) separately or simultaneously. The JNK activities were determined at 30 min after individual or combinatory treatment of these agonists as indicated. Values shown represent the mean  $\pm$  S.E. from four separate experiments. \*Individual or combinatory treatment significantly increased the JNK activities as compared to the basal. #Co-stimulation of  $D_1R$  significantly inhibited the GRPR-induced JNK activation (Bonferroni paired *t*-test,  $P < 0.05$ ).

**Figure 4. Modulation of  $D_1R$ -mediated adenylyl cyclase activation by  $G_i$ -coupled  $ORL_1R$  and  $G_q$ -coupled GRPR.** Cos-7 cells co-expressing  $D_1R$  with either  $ORL_1R$  (A) or GRPR in the absence (B, D, E) or presence (C) of transducin ( $G\alpha_t$ ) were labeled with [ $^3H$ ]adenine (A – E). These cells were then pretreated with (D and E) or without (A, B and C) calphostin C (Cal. C, 10  $\mu$ M, 30 min) or BAPTA-AM (10  $\mu$ M, 30 min) for the inhibition of PKC and  $Ca^{2+}$  signaling, respectively. Cos-7 cells expressing  $D_1R$  alone were also labeled with [ $^3H$ ]adenine for cAMP assay (F). The assays for cAMP elevation which reflect the activity of adenylyl cyclase was determined at 30 min after individual or co-stimulation with specific agonists and thapsigargin (Thap, 5  $\mu$ M) as indicated. Values shown represent the mean  $\pm$  S.E. from three separate experiments. \*Dopamine (DOP, 10  $\mu$ M) treatment in the absence or presence of nociceptin (OFQ, 100 nM) significantly enhanced the cAMP level as

compared to the basal (A). \*Treatment with dopamine in the absence or presence of bombesin (B – E) or thapsigargin (F) significantly enhanced the cAMP level as compared to the corresponding basal. \*\*Co-treatment with bombesin (B, C and D) or thapsigargin (F) significantly increased the D<sub>1</sub>R-induced cAMP elevation. #Co-stimulation of ORL<sub>1</sub>R significantly inhibited the D<sub>1</sub>R-induced cAMP elevation (Bonferroni paired *t*-test, P<0.05).

**Figure 5. Co-stimulation of G<sub>s</sub>-coupled D<sub>1</sub>R with either G<sub>q</sub>-coupled GRPR or G<sub>i</sub>-coupled ORL<sub>1</sub>R had no effect on PLC.** Cos-7 cells co-expressing D<sub>1</sub>R with either GRPR (A) or ORL<sub>1</sub>R (B) were labeled with [<sup>3</sup>H]myo-inositol for IP<sub>3</sub> assay. The levels of IP<sub>3</sub> formation which reflect the activities of PLC were determined at 30 min after individual or co-stimulation with specific agonists as indicated. Values shown represent the mean ± S.E. from three separate experiments. \*Treatment of bombesin in the absence or presence of dopamine significantly enhanced the IP<sub>3</sub> levels as compared to the corresponding basal.

**Figure 6. Sp-cAMPS selectively suppressed the JNK activation mediated by GRPR rather than ORL<sub>1</sub>R.** Cos-7 cells co-expressing JNK-HA and D<sub>1</sub>R with either GRPR (A) or ORL<sub>1</sub>R (B) were treated with Sp-cAMPS (Sp, 100 μM) and the indicated agonists separately or simultaneously. The JNK activities were determined at 30 min after separate or combinatory treatment as indicated. Values shown represent the mean ± S.E. from three independent experiments. \*Individual or combinatory treatment resulted in significant activation of JNK activity. #Co-treatment with Sp-cAMPS (100 μM) significantly inhibited the bombesin-induced JNK activation (Bonferroni paired *t*-test, P<0.05).

**Figure 7. Sp-cAMPS suppressed the bombesin-induced JNK activation in a dose-dependent manner.** Cos-7 cells co-expressing JNK-HA, D<sub>1</sub>R and GRPR were stimulated by










bombesin (100 nM) accompanied with increasing concentrations of Sp-cAMPS (0 – 1000  $\mu$ M). The JNK activities were determined at 30 min after individual or combinatory treatment as indicated. Values shown represent the mean  $\pm$  S.E. from three separate experiments, with the bombesin-induced JNK activation in the absence of Sp-cAMPS defined as 100%. <sup>#</sup>Co-treatment with Sp-cAMPS significantly inhibited the bombesin-induced JNK activation (Bonferroni paired *t*-test, *P*<0.05).



**Figure 8. Disruption of Ca<sup>2+</sup> signaling by BAPTA-AM suppressed the D<sub>1</sub>R/GRPR-mediated JNK activity to the D<sub>1</sub>R/ORL<sub>1</sub>R-induced level.** Cos-7 cells co-expressing JNK-HA and D<sub>1</sub>R with either GRPR or ORL<sub>1</sub>R were pretreated in the absence (control) or presence of BAPTA-AM (10  $\mu$ M, 30 min) followed by addition of their specific agonists. The JNK activities were determined at 30 min after the combinatory drug treatment. Values shown represent the mean  $\pm$  S.E. from three separate experiments. \*Co-administration of agonists resulted in significant activation of JNK activities as compared to their corresponding basal. <sup>#</sup>Pre-treatment with BAPTA-AM significantly suppressed the JNK activity induced by co-stimulation with bombesin and dopamine (Bonferroni paired *t*-test, *P*<0.05).



**Figure 9. Schematic diagrams for the regulatory effects of G<sub>s</sub> on the G<sub>i</sub>- and G<sub>q</sub>-mediated JNK activation.** (A) G $\beta\gamma$  subunits from G<sub>s</sub> and G<sub>i</sub> converge their signals at Src-like tyrosine kinases and subsequently trigger the activation of JNK cascade, while cAMP has no inhibitory effect on the G<sub>i</sub>-mediated JNK activation. (B) G $\beta\gamma$  subunits from G<sub>s</sub> and G<sub>q</sub> converge their signals at Src-like kinases, and the G $\alpha_s$ -mediated cAMP elevation is synergistically potentiated in the presence of Ca<sup>2+</sup>-responsive adenylyl cyclase isoforms. The G $\beta\gamma$ /Src and Ca<sup>2+</sup> signals co-operate with each other to regulate the JNK activation (Chan and


Wong, 2004b), while cAMP suppresses the induced kinase activation by interfering the Ca<sup>2+</sup>-mediated pathway.






**A****B****C****D****E**

**A****B**

**A****B**





**A****B**