

Sustained Ah Receptor Activity Attenuates Liver Regeneration

Kristen A. Mitchell, Courtney A. Lockhart, Gengming Huang and Cornelis J. Elferink

Department of Pharmacology and Toxicology, University of Texas Medical Branch,

Galveston, TX 77555-1031

RUNNING TITLE: AhR activation suppresses liver regeneration

ADDRESS CORRESPONDENCE TO:

Cornelis Elferink, Ph.D.

301 University Boulevard

Department of Pharmacology and Toxicology

University of Texas Medical Branch

Galveston, TX 77555-1031

(409) 772-9624 (ph)

(409) 772-9642 (fax)

coelferi@utmb.edu

TEXT PAGES: 32

TABLES: 0

FIGURES: 8

REFERENCES: 37

ABSTRACT: 219

INTRODUCTION: 740

DISCUSSION: 1,488

ABBREVIATIONS: AhR, aryl hydrocarbon receptor; BrdU, 5-bromo-2'-deoxyuridine; CDK2, cyclin dependent kinase; PH, partial hepatectomy; qRT-PCR, quantitative real-time PCR; TCDD, 2,3,7,8-tetrachlorodibenzo-*p*-dioxin.

ABSTRACT

In hepatocyte-derived cell lines, either loss of aryl hydrocarbon receptor (AhR) function or treatment with a persistent AhR agonist such as 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) can disrupt G1 phase cell cycle progression. The present study used liver regeneration to explore mechanistically how AhR activity modulates hepatocyte proliferation in vivo. Treatment of mice with 20 µg/kg TCDD one day prior to 70% partial hepatectomy (PH) resulted in a 50-75% suppression in liver regeneration. Impaired proliferation was not associated with changes in levels of IL-6 or TNFα, which prime quiescent hepatocytes to enter G1 phase. In fact, administration of TCDD 12 h after PH, a period well beyond the priming phase, still induced the G1 arrest. Decreased proliferation in TCDD-treated mice correlated with reduced cyclin-dependent kinase-2 (CDK2) activity, a pivotal regulator of G1/S phase transition. In contrast to observations made in cell culture, suppressed CDK2 activity was not strictly associated with increased binding of the CDK2 inhibitors p21^{Cip1} or p27^{Kip1}. However, TCDD decreased levels of cyclin E binding to CDK2, despite normal cyclin E expression. The evidence also suggests that TCDD-induced hepatic growth arrest depends upon sustained AhR activity because transient AhR activation in response to endogenous queues failed to suppress the regenerative response. These findings establish a functional role for the AhR in regulating normal cell cycle control during liver regeneration.

INTRODUCTION

The aryl hydrocarbon receptor (AhR) is a conditionally activated bHLH/PAS transcription factor that mediates the toxicity of halogenated aromatic hydrocarbons and structurally related chemicals. Upon ligand binding, the receptor translocates to the nucleus and dimerizes with the AhR nuclear translocator protein (ARNT), forming a complex that binds to dioxin-response elements in the promoter region of selected genes and alters transcription (Rowlands and Gustafsson, 1997). Exposure to the most potent AhR ligand 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) elicits an array of biological and toxic effects related to cell and/or tissue homeostasis (Mandal, 2005).

Increasing evidence indicates that the AhR influences cell cycle progression in the absence of exogenous ligand. For example, AhR-defective mouse hepatoma (Hepa 1c1c7) cells proliferate more slowly than wild-type cells. This growth inhibition occurs during G1 phase of the cell cycle and can be reversed upon transfection with AhR cDNA (Ma and Whitlock, 1996). Moreover, mouse embryo fibroblasts (MEF) from AhR nullizygous mice exhibit lower proliferation rates than MEF from wild-type mice (Elizondo et al., 2000), providing compelling evidence that normal movement through the cell cycle requires a functional AhR.

Interestingly, treatment of certain cell lines with TCDD elicits changes in proliferation similar to those observed in AhR-defective or AhR-deficient cells. In 5L rat hepatoma cells, exposure to TCDD inhibits G1 progression (Gottlicher and Wiebel, 1991; Weiss et al., 1996). This inhibition requires the induction of p27^{Kip1} to inhibit cyclin-dependent kinase 2 (CDK2) activity (Kolluri et al., 1999). In both 5L cells and mouse hepatoma cells, G1 arrest depends on the interaction of the AhR with the retinoblastoma tumor suppressor protein (pRb) (Elferink et al., 2001; Ge and Elferink, 1998; Puga et al., 2000) and appears to involve distinct mechanisms

of action (Huang and Elferink, 2005; Marlowe et al., 2004). Therefore, delays in hepatocyte cell cycle progression occur when the AhR is activated by a poorly metabolized, persistent agonist such as TCDD or when the receptor is removed altogether. These seemingly contradictory findings are reconciled by the observation that impaired cell cycle progression also occurs when AhR activity is sustained in the absence of exogenous agonist by inhibiting cytochrome P4501A1 activity (Levine-Fridman et al., 2004). Thus, it appears that some level of basal or transient AhR activity, presumably through receptor interactions with endogenous ligands, promotes cell cycle progression, but prolonged AhR activity during G1 phase progression triggers a growth arrest. This receptor-regulated activity during cell cycle progression is distinct from the observation that transgenic mice harboring a constitutively active AhR develop tumors (Moennikes et al., 2004). Tumor formation results from loss of check-point control, and the incidence of tumor development is thought to be the consequence of selecting for cells that have escaped the growth-inhibitory effect resulting from sustained AhR activity.

Relatively little is known regarding how the AhR regulates cell cycle progression in vivo. An intriguing report by Bauman et al. (1995) demonstrated that prolonged (2-week) pretreatment of rats with TCDD decreased hepatocyte proliferation following 70% partial hepatectomy (PH), implicating that AhR-mediated regulation of cell cycle progression may occur in vivo. However, the mechanism of action remains unresolved and may differ markedly from the cell culture findings, in part because liver regeneration is influenced by multiple, systemically-derived signals and involves non-transformed cells. To elucidate how the AhR modulates cell cycle progression in vivo, we used a mouse model of liver regeneration following PH. Following resection of two-thirds of the liver, remaining hepatocytes enter the cell cycle as a largely synchronized population that divides until the original organ mass is restored (Mangnall et al.,

2003). Growth factors and cytokines such as $\text{TNF}\alpha$ and IL-6 essentially drive the otherwise quiescent hepatocytes into G1 phase, whereas continued progression through the cell cycle is regulated primarily at the G1/S phase transition, which is governed by the coordinated activity of CDKs (Albrecht et al., 1999). Activation of cyclin D/CDK4-6 and cyclin E/CDK2 during G1 phase results in the phosphorylation and inactivation of pRb (Hatakeyama and Weinberg, 1995), whereas negative regulation of CDK2 activity is achieved through the binding of CDK inhibitor proteins p27^{Kip1} and p21^{Cip1} (Albrecht et al., 1998; Sherr and Roberts, 1995).

In the present study, TCDD was used to disrupt normal, physiological AhR function and determine the contribution of sustained receptor activity to liver regeneration. Our findings demonstrate that, while many of the observations made in hepatoma cell lines are recapitulated during liver regeneration, AhR activity may regulate *in vivo* hepatocyte proliferation by mechanisms distinct from those observed *in vitro*.

MATERIALS AND METHODS

Animals

Female, C57Bl/6 mice (8-10 weeks old; The Jackson Laboratory, Bar Harbor, ME) were used for all experiments. Animals were housed in microisolator cages, maintained on a 12-hour light cycle, and provided with food and water *ad libitum* at all times.

TCDD

TCDD (Cerilliant, Round Rock, TX) was dissolved in anisole, diluted in peanut oil and administered via gavage at 20 µg/kg body weight. Control mice received peanut oil spiked with an equivalent amount of anisole. Mice were treated with TCDD or vehicle 24 hours prior to surgery unless otherwise indicated. The 20 µg/kg dose of TCDD is not overtly toxic (Pohjanvirta and Tuomisto, 1994) and was selected based on a dose-response study (data not shown), in which this was the lowest dose at which consistent, reproducible suppression of liver regeneration was observed.

Surgical Procedures

Mice were anesthetized with inhaled isoflurane, and PH was performed by surgically resecting 70% of the liver based on the procedure described by (Higgins and Anderson, 1931). It is noteworthy that the medial and left lateral hepatic lobes were resected individually to avoid damage to the gall bladder. The contribution of these hepatic lobes to the total liver mass was determined by wet weight to verify that removal of 70% of the liver was achieved. Control mice received a sham surgery in which the abdominal cavity was opened, and the liver was gently

manipulated but not resected. All mice recovered expediently, and mortality associated with surgical procedures was less than 2%. Mice were killed postoperatively by cervical dislocation at the indicated times. All experiments were performed in compliance with the standards set by the UTMB Animal Care and Use Committee.

BrdU labeling

To measure proliferation in the liver, mice were injected i.p. with 50 mg/kg 5-bromo-2'-deoxyuridine (BrdU; Sigma-Aldrich, St. Louis, MO) two hours prior to sacrifice. For continuous labeling studies, mice were provided drinking water containing BrdU (0.8 mg/ml) immediately following surgery, and water bottles containing BrdU were protected from light and replenished daily until mice were killed 72 h postoperatively. After sacrifice, fresh liver tissue was fixed in 10% buffered formalin for 18 h, then processed and stained for histological analysis as described below.

Histopathology/Immunohistochemistry

Slides containing liver sections were either stained with hematoxylin and eosin (H&E), or processed and stained to reveal BrdU incorporation using a biotinylated anti-BrdU antibody (Invitrogen, Carlsbad, CA) followed by incubation with avidin-conjugated horseradish peroxidase and the substrate 3,3'-diaminobenzidine (DAB). Tissue processing, staining and immunohistochemistry were performed by the Research Histopathology Core Facility at UTMB. For analysis of proliferation, the numbers of brown-stained nuclei (BrdU⁺) were counted in three separate low-power fields per animal and expressed as a percentage of the total number of nuclei.

Measurement of TNF α and IL-6 production

Blood was collected in heparin-coated tubes and plasma was recovered following centrifugation. Levels of TNF α and IL-6 were measured in unpooled plasma samples using sandwich enzyme-linked immunosorbant assay (ELISA) kits (Pierce Biotech, Rockford, IL). The assays were performed according to the manufacturer's protocols and all samples were run in duplicate. The lower limit of detection for the assays was 50 pg/ml.

Immunoprecipitation/Western Blotting

Frozen liver tissue was mechanically homogenized in 50 mM HEPES (pH 7.5) containing 150 mM NaCl, 10% glycerol, 0.1% Tween 20, 7.5 mM MgCl₂, 7.5 mM EGTA and the following protease/phosphatase inhibitors: 25 mM β -glycerolphosphate, 5 mM DTT, 1 mM NaF, 1 mM Na₃VO₄, 1 mM PMSF, 1 μ g/ml aprotinin, and 0.1 μ g/ml leupeptin. Protein concentrations were determined using a DC protein assay kit (Bio-Rad Laboratories, Inc., Hercules, CA). To detect cytochrome P4501A1 protein expression, liver homogenates were probed with a CYP1A1 antibody (BD Discovery Labware, Bedford, MA) by western blotting as described below. For all other studies, liver homogenates were immunoprecipitated prior to western blotting. For immunoprecipitation, 500 μ l of total cellular protein (1 μ g/ μ l) was incubated with 2 μ g of rabbit or goat anti-CDK2 antibody (sc-163 or sc-163G, Santa Cruz Biotechnology, Santa Cruz, CA) for 3 h at 4°C then incubated for an additional hour with either protein G or protein A sepharose (Sigma-Aldrich). For western blotting, sepharose beads were washed 5 times in NETN buffer (20 mM Tris-HCl pH 8.0, 100 mM NaCl, 0.5% NP-40, 1 mM EDTA), fractionated by SDS-PAGE on a 15% gel, and transferred to polyvinylidene difluoride membranes (Bio-Rad Laboratories, Inc.). Membranes were blocked for 1 h at RT in 5% (w/v) dry milk in Tris-buffered saline with 0.1%

(v/v) Tween 20, then incubated for 4 h at RT or overnight at 4°C with the following primary antibodies (all purchased from Santa Cruz Biotechnology): rabbit CDK2 (sc-163), goat CDK2 (sc-163G), rabbit cyclin E (sc-481), rabbit cyclin A (sc-596), mouse p21 (sc-6246), mouse p27 (sc-1641). Membranes were incubated with horseradish peroxidase-conjugated secondary antibodies for 1 h at RT and visualized using an enhanced chemiluminescence detection method according to the manufacturer's protocol (Amersham Biosciences, Piscataway, NJ). Densitometry on the cytochrome P4501A1 western blot was performed using ImageJ, a public domain NIH Image program (developed at the U.S. National Institutes of Health and available on the Internet at <http://rsb.info.nih.gov/nih-image/>).

Quantitative Real-Time RT-PCR

Total RNA was isolated from frozen liver tissue using RNAqueous (Ambion, Inc., Austin, TX) and quantitative real-time PCR (qRT-PCR) was performed by the Sealy Center for Cancer Cell Biology and Real-Time PCR Core Facility at the University of Texas Medical Branch. Reagents included Applied Biosystems assays-by-design 20x assay mix of primers and TaqMan® probes (FAM™ dye-labeled) for the target gene (mouse CYP1A1), and pre-developed 18S rRNA (VIC™-dye labeled probe) TaqMan® assay reagent (P/N 4319413E) for an internal control. The primers were designed to span exon-exon junctions so as not to detect genomic DNA. Primers and probe sequences were searched against the Celera Genomic database (www.celeradiscoverysystem.com) to confirm specificity. Validation experiments were performed to test the efficiency of the target and reference amplifications. The absolute value of the slope of log input amount versus CT were <0.1. Separate tubes (singleplex) for one-step RT-PCR were performed with 20 ng RNA for both target gene and endogenous control using the

TaqMan® one-step RT-PCR master mix reagent kit (Applied Biosystems, Foster City, CA). The cycling parameters for one-step RT-PCR were: reverse transcription 48°C for 30 min, AmpliTaq activation 95°C for 10 min, denaturation 95°C for 15 sec and annealing/extension 60°C for 1 min (repeat 40 times) on ABI7000. Triplicate CT values were analyzed in Microsoft Excel using the comparative CT method as described by the manufacturer (Applied Biosystems). The amount of target (2-CT) was obtained after normalization to an endogenous reference (18S rRNA).

Kinase Assay

To measure kinase activity, 500 µg of total cellular protein was immunoprecipitated with 2 µg CDK2 antibody as described above. Sepharose beads were washed 4 times in NETN buffer, once in kinase assay buffer (50 mM Tris-Cl, pH 7.5, 80 mM β-glycerolphosphate, 50 mM NaF, 20 mM EGTA, 15 mM Mg(OAc)₂, 1 mM DTT, 1 mM PMSF, 0.5 mM Na₃VO₄, 50 µM ATP, 5 µg/ml aprotinin, 1 µg/ml leupeptin), and resuspended in 15 µl kinase assay buffer containing 10 µg histone H1 and 5 µCi [γ-³²P]ATP (3000 Ci/mmol) (Amersham Biosciences). After 40 minutes at RT, the reaction was quenched by the addition of 15 µl of 2x SDS loading buffer. Reactions were fractionated by SDS-PAGE on a 15% gel, which was subsequently dried and autoradiographed.

Statistical Analyses

Statistical analyses were performed using Prism (version 4.0, GraphPad Software, San Diego, CA). Data were evaluated either by Student's *t*-Test or by two-way ANOVA followed by Bonferonni post-hoc test. Data were considered significantly different at $p \leq 0.05$.

RESULTS

Analysis of TCDD toxicity in regenerating liver.

To evaluate any overt toxic effects of TCDD treatment on the regenerating liver, a histological analysis was performed and liver wet weight was measured. PH-induced liver regeneration in vehicle-treated mice was accompanied by panlobular hydropic vacuolation of hepatic tissue (Fig. 1A). Treatment with TCDD generally increased hepatic vacuolation and also altered the integrity of hepatocyte nuclei. It is noteworthy that these effects varied considerably among animals in each treatment group. In contrast, vacuolation was not observed in liver tissue from sham-operated mice pretreated with either vehicle or TCDD, and no overt histological changes were detected in liver tissues from these mice. Consistent with the reported effects of TCDD on hepatomegaly, preoperative liver wet weight was about 15% greater in TCDD-treated mice as compared to vehicle-treated mice, and this increase was maintained postoperatively throughout the regenerative process (Fig. 1B). Increased liver: body weight ratios in TCDD-treated mice are attributed to increased liver mass, as body weight was consistent among vehicle- and TCDD-treated mice throughout the 5-day period following surgery.

AhR activation suppresses proliferation in the regenerating liver.

The influence of AhR activation on cell cycle progression in the regenerating liver was addressed by evaluating *in vivo* BrdU incorporation. Thirty-six hours after PH, which is when peak DNA synthesis typically occurs in the regenerating mouse liver (Mangnall et al., 2003), BrdU incorporation confirms that about 60% of all hepatic nuclei are in S phase in vehicle-treated mice (Fig. 2A). BrdU staining was uniform throughout the liver, consistent with a global

proliferative response aimed at restoring liver cell mass. Pretreatment with TCDD resulted in about a four-fold reduction in proliferation following PH (Fig. 2B). It is noteworthy that BrdU staining in these livers was also panlobular and was detected in all lobes. In contrast, BrdU incorporation was minimal (<4%) in sham-operated mice that were pretreated with either vehicle or TCDD.

To determine if reduced proliferation in TCDD-treated mice reflected a delay in cell cycle progression instead of suppression, BrdU staining was assessed at various times throughout the regenerative process. TCDD treatment suppressed proliferation at both 36h and 48h after PH, when peak proliferation was detected in vehicle-treated mice, but had no effect on proliferation at later times (Fig. 3A). Regeneration in TCDD-treated mice appeared to follow the same kinetics as regeneration in vehicle-treated mice. Moreover, in studies using continuous BrdU labeling, proliferation was detected in almost 100% of hepatic nuclei in vehicle-treated mice 72 hours after PH (Fig. 3B). In contrast, only 50-60% of the nuclei in livers from TCDD-treated mice had incorporated BrdU by this time, indicating that almost half of the hepatocytes in these mice actually fail to proliferate, rather than undergo a delay in cell cycle progression.

AhR activity during the regenerative process.

In keeping with our recent findings that sustained AhR activity is required to elicit a growth arrest (Levine-Fridman et al., 2004), we confirmed that TCDD-induced AhR activity persisted throughout the regenerative process. As shown in Figure 4, cytochrome P4501A1 expression was induced in TCDD-pretreated mice and persisted for at least 5 days following PH, consistent with sustained AhR activity throughout the regenerative process in these mice. Also noteworthy is that removal of a substantial portion of the liver does not impair AhR activity

elicited by TCDD. Interestingly, in vehicle-pretreated mice, cytochrome P4501A1 protein levels began to increase 36 h after PH and peaked at 72 h before tapering off. Analysis of CYP1A1 mRNA by qRT-PCR reveals that the increase in P4501A1 protein in vehicle-treated mice is preceded by a pronounced (42-fold) but transient increase in mRNA 24 h post-PH (Fig. 4B). This indicates that AhR activity is transiently induced during G1 phase of the regenerative process despite the absence of an exogenous ligand. The qRT-PCR data also confirm that the persistent expression of P4501A1 protein in livers from TCDD-treated mice is matched by sustained elevation of CYP1A1 mRNA levels. These observations complement our recent findings in cell culture (Levine-Fridman et al., 2004).

TCDD treatment does not alter the production of IL-6 or TNF α during liver regeneration.

Given that the initiation of liver generation depends on cytokines such as IL-6 and TNF α , which facilitate the movement of quiescent hepatocytes into G1 phase of the cell cycle, we tested the possibility that TCDD suppressed the production of these cytokines either before or shortly after PH. In vehicle-treated mice, peak levels of IL-6 were detected 2-4 h after PH (Fig. 5A), whereas TNF α levels peaked 8-12 h after PH (Fig. 5B). Although there was a trend towards increased production of these cytokines following PH in TCDD-treated mice, this effect was not statistically significant and was not consistently observed among experiments. Prior to surgery, neither cytokine was detected in the plasma of mice pretreated with either TCDD or vehicle control. Likewise, cytokine production in sham-operated mice was minimally induced and remained unaltered by TCDD treatment.

Decreased proliferation in TCDD-treated mice is not due solely to alterations in early regenerative processes.

Based on the observation that TCDD did not impair cytokine production, we addressed the overall involvement of such early regenerative processes to the decreased proliferation observed in TCDD-treated mice. We found that when TCDD was administered 12 h post-PH, well after the early "priming" phase of liver regeneration, TCDD still caused a marked decrease in proliferation, similar to that observed in mice pretreated for 24 h (Fig. 6). Likewise, a five-fold reduction was observed in hepatic tissue from mice when TCDD was administered concomitantly with PH. The data suggest that AhR-mediated effects occur subsequent to signaling events that prompt hepatocytes out of quiescence and into G1 phase of the cell cycle.

CDK2 activity is diminished during liver regeneration in TCDD-treated mice.

Given that early events important for moving hepatocytes into G1 phase were not altered by AhR activation, we tested the possibility that hepatocytes in TCDD-treated mice were stalling in G1 phase. Hence, attention was directed towards the G1 phase CDKs. An analysis of CDK4 failed to reveal TCDD-dependent alterations (data not shown), but examination of CDK2 activity exposed differences in response to TCDD treatment. In vehicle-treated mice, CDK2 phosphorylation of substrate histone H1 was readily detected 36 and 48 h after PH, whereas CDK2 activity was barely detectable 36 h after PH in TCDD-exposed mice (Fig. 7). Kinase activity in TCDD-treated mice exhibited some interindividual variability (data not shown), but generally increased at 48 h, consistent with the increase in BrdU incorporation observed between 36 h and 48 h in these mice (Fig. 3A). The evidence reveals a tight correlation between diminished BrdU incorporation and reduced CDK2 activity, demonstrating that the TCDD-

induced growth arrest is linked to inhibition of G1-phase CDK2 activity normally associated with proliferation.

AhR activation alters the association of CDK2 with cell cycle regulatory proteins.

To mechanistically determine why CDK2 kinase activity was impaired in TCDD-exposed mice, we conducted coimmunoprecipitation experiments to examine the association of CDK2 with cyclins E and A, regulatory proteins that bind to CDK2. CDK2/cyclin E complex formation was observed 36 and 48 h after PH (Fig. 8), consistent with cyclin E promoting CDK2 activity during cell cycle progression (Geisen and Moroy, 2002). In contrast, considerably less cyclin E was detected bound to CDK2 in the regenerating liver from TCDD-treated mice. The decrease in CDK2/cyclin E association is not reflected in total cyclin E expression, which was not noticeably affected by TCDD (data not shown). Consistent with the sequential role of cyclin E and A regulation of CDK2, cyclin A is first detected in association with the kinase at 48 h. However, reduced CDK2/cyclin A binding was observed in TCDD-treated mice, most notably at 72 h. CDK2 activity is not only a function of cyclin binding activity, but is also influenced by the action of the Cip/Kip inhibitors, p21^{Cip1} and p27^{Kip1}. Association of CDK2 with p27^{Kip1} is detectable in the quiescent liver from both vehicle- and TCDD-treated mice, but decreases transiently to a nearly undetectable level as cells enter G1 phase after PH, only to recover as cells complete the first cycle of cell division. p21^{Cip1} follows a reciprocal profile being transiently associated with CDK2. Although p21^{Cip1} and p27^{Kip1} binding to CDK2 appear slightly elevated in TCDD treated livers at 24 h and 48 h respectively, enhanced TCDD-dependent inhibitor binding to the kinase complex was neither pronounced nor consistent between animals, rendering their role in the inhibition of kinase activity uncertain. Much clearer is the marked TCDD-dependent decrease in

cyclin binding as an explanation for diminished CDK2 activity.

DISCUSSION

Studies using hepatocyte-derived transformed cell lines demonstrate a functional role for the AhR in regulating cell cycle progression at the G1/S phase transition. Because the establishment of immortalized cell lines relies on corrupting or bypassing cell cycle regulatory checkpoints to permit continued proliferation, it has been difficult to appreciate the relevance of such studies to receptor function in non-transformed cycling cells. Our findings using liver regeneration as a model recapitulate many of the observations made in previous cell culture studies and confirm that prolonged AhR activity can elicit a G1 arrest in non-transformed, cycling somatic cells *in vivo*. Yet these findings suggest that the mechanisms by which the AhR controls hepatocyte proliferation *in vivo* may be distinct from those identified *in vitro* that implicate an obligatory role for p21^{Cip1} and p27^{Kip1} (Barnes-Ellerbe et al., 2004; Kolluri et al., 1999).

The present findings demonstrate that TCDD treatment suppresses BrdU incorporation in the regenerating liver, suggesting that prolonged AhR activation can impair cell cycle progression into S phase and commitment to cell division. This study extends upon the previous finding by Bauman et al. (1995), in which treatment of rats with TCDD before PH reduced the percentage of cells in S phase. Importantly, failure of hepatocytes to transit to S phase in our studies was not due to delayed proliferation, as cumulative BrdU incorporation was decreased 40-50% in TCDD-treated hepatectomized mice. We also detected only minimal apoptosis when regenerating livers from vehicle- and TCDD-treated mice were stained by TUNEL (data not shown), indicating that the TCDD-induced inhibition in liver regeneration is not due to increased hepatic apoptosis. The conclusion that S phase progression is stalled in TCDD-treated mice is further supported by the detection of reduced association of CDK2 with cyclin A, a major

activating subunit of CDK2 during S phase. Collectively, our findings demonstrate that the G1 arrest observed in transformed hepatocyte-derived cell lines treated with TCDD (Gottlicher and Wiebel, 1991; Marlowe et al., 2004; Weiss et al., 1996) is recapitulated in vivo during physiologically relevant proliferation processes.

Given the suppressive effects of TCDD on hepatocyte proliferation, it is intriguing that the mass of the remaining liver was greater in TCDD-treated mice than vehicle controls. Increased liver mass was also observed preoperatively in TCDD-treated mice, consistent with numerous reports of hepatomegaly induced by TCDD (Morris et al., 1992; Silkworth et al., 1989). Recent evidence indicates that TCDD-induced hepatomegaly occurs by an AhR-mediated mechanism that requires nuclear localization of the receptor, but the precise mechanism is unclear (Bunger et al., 2003). Although it is conceivable that the hepatomegaly observed in TCDD-treated mice results from an AhR-mediated inhibition of apoptosis in the liver, we find no evidence that apoptosis levels are altered by TCDD following PH or sham surgery. Given the increase in hydropic vacuolation that we and others have observed in livers from TCDD-treated mice (Walisser et al., 2005), it is possible that such structural changes in hepatocytes could contribute to the increase in liver mass, although additional studies would be required to address this. It is also noteworthy that vacuolation varied considerably among mice in each treatment group and did not correlate with BrdU incorporation in the liver (data not shown).

Our observation that BrdU incorporation was minimal in sham-operated mice pretreated with either vehicle or TCDD indicates that TCDD does not possess intrinsic mitogenic activity. This is consistent with other studies that demonstrate that, whereas TCDD is a potent tumor promoter, no TCDD-mediated changes in proliferation occur in the absence of a tumor-initiating agent (Pitot et al., 1980). Key mitogenic stimuli that occur early (0.5-6 h) after PH include the

production of TNF α and IL-6, which essentially "drive" quiescent hepatocytes into G1 phase. Early speculation focused on the possibility that TCDD treatment decreased the production of these cytokines immediately following PH, consistent with reports in which production of IL-6 and TNF α is variously increased or decreased, depending on the model system (Fan et al., 1997; Ito et al., 2002; Jensen et al., 2003; Kerkvliet et al., 1996; Prell et al., 1995). However, we found that neither TNF α nor IL-6 levels were suppressed in TCDD-treated mice following either PH or sham-operation, indicating that these soluble mediators are not likely targets of TCDD-induced growth arrest in the regenerating liver. We also examined downstream targets of IL-6 signaling pathways and found no TCDD-mediated changes in phosphorylation of either Jak or Stat3 (data not shown). Since epidermal growth factor (EGF) is also a potent liver mitogen that signals through the Stat3 protein (Cressman et al., 1995), we infer that EGF signaling likewise is not altered by TCDD treatment. The observation that TCDD administration 12 h postoperatively still suppressed proliferation reinforces that disruption of early cytokine production or signaling is unlikely to play a significant role in the TCDD-induced growth arrest. Assuming that TCDD-mediated suppression of liver regeneration occurs independently of the priming phase, we infer that AhR activity disrupts events within G1 phase in cycling hepatocytes. In mice, the G1/S transition occurs 32-36 h after PH (Weglarz and Sandgren, 2000). Our observation that treatment with TCDD reduces CDK2 kinase activity 36 h after PH provides compelling evidence that prolonged AhR activity does indeed impair cell cycle progression through the G1/S boundary. Although individual variation existed among mice at later time points, levels of CDK2 kinase activity closely correlated with BrdU incorporation in individual mice, strengthening the assumption that altered kinase activity in TCDD-treated mice contributes directly to suppressed hepatocyte proliferation.

Given that CDK2 activity during G1 phase is absolutely dependent on cyclin E, it is plausible that decreased CDK2 activity in TCDD-treated mice could be explained by the decreased formation of CDK2/cyclin E complexes observed 36 and 48 h after PH. However, this contrasts with the findings made in 5L hepatoma cells, in which decreased cyclin E-dependent kinase activity did not result from decreased association of CDK2 with cyclin E (Kolluri et al., 1999). Instead, in this report, decreased kinase activity was attributed to the direct TCDD-mediated induction of p27^{Kip1} and subsequent association of the inhibitor with the CDK2/cyclin E complex. Our findings did not reveal a strong correlation between TCDD treatment and p27^{Kip1} expression or CDK2/p27^{Kip1} (or p21^{Cip1}) complex formation. It is possible that CDK2 activity during liver regeneration is negatively regulated, at least in part, by the interaction between CDK2 and cyclin E. It is noteworthy that we found no evidence that TCDD altered either CDK2 or cyclin E protein levels in whole cell lysates (data not shown), which contrasts a recent report in which treatment of Hepa1c1c7 cells with TCDD decreased mRNA levels of both CDK2 and cyclin E (Marlowe et al., 2004). It is conceivable that the TCDD-induced decrease in cyclin E binding to CDK2 observed during liver regeneration may reside with enhanced turnover of complex-associated cyclin E. (Welcker et al., 2003) recently demonstrated that CDK2- and GSK3-mediated cyclin E phosphorylation promotes SCF^{Fbw7} ubiquitin ligase-dependent degradation of the CDK-bound cyclin. Future studies examining the capacity of cyclin E phospho-site specific mutants (refractory to ubiquitin-mediated proteolysis) designed to override the TCDD induced growth arrest should prove insightful.

At this time it is less clear why TCDD also reduced cyclin A binding to CDK2 at the later time point. The finding in C33A cells that AhR activity, in conjunction with pRb, can suppress cyclin A expression (Strobeck et al., 2000) is not supported by our observation that cyclin A

levels were unaffected by TCDD. A more plausible explanation rests with the sequential nature of cyclin E and A binding to the kinase as cells progress through G1 and S phase. The CDK2-cyclin A interaction is necessary to maintain kinase activity during S phase, hence, this complex would not be expected to form in G1 phase arrested cells.

The conclusion that sustained AhR activity impairs G1/S progression is particularly intriguing in light of the data demonstrating cytochrome P4501A1 expression in the regenerating liver. In TCDD-treated mice, induction of CYP1A1 persists throughout the regenerative process. In vehicle-treated mice, transient CYP1A1 induction in the absence of an exogenous agonist occurs 36 h after PH, which coincides with progression through G1/S phase. Clearly, such transient activation of the AhR is insufficient for eliciting a G1 arrest in cycling hepatocytes *in vivo*. Instead, it is conceivable that such transient activation is in fact required for normal cell cycle progression to occur. We have previously shown that in the absence of an exogenous agonist, AhR activity is transiently induced in serum-released 5L cells (Levine-Fridman et. al., 2004). This study highlighted the negative feedback between induced P4501A1 activity and subsequent termination of AhR activity through metabolic agonist depletion. Thus, it is plausible that CYP1A1 induction during normal cell cycle progression is a mechanism directed at downregulating receptor activity through the metabolic depletion of unidentified endogenous AhR ligands that promote cell cycle progression. Failure to engage the negative feedback results in prolonged AhR activity and induction of antiproliferative signaling.

ACKNOWLEDGEMENTS

We would like to thank Melanie Howell for excellent technical assistance and Dr. Steve Weinman for histopathological analysis of liver tissue. We are also grateful to Dr. Hui-Qun Wang and other members of the UTMB Research Histopathology Core Facility for preparing and staining the liver tissue for these studies.

REFERENCES

- Albrecht JH, Poon RY, Ahonen CL, Rieland BM, Deng C and Crary GS (1998) Involvement of p21 and p27 in the regulation of CDK activity and cell cycle progression in the regenerating liver. *Oncogene* **16**(16):2141-2150.
- Albrecht JH, Rieland BM, Nelsen CJ and Ahonen CL (1999) Regulation of G(1) cyclin-dependent kinases in the liver: role of nuclear localization and p27 sequestration. *Am J Physiol* **277**(6 Pt 1):G1207-1216.
- Bauman JW, Goldsworthy TL, Dunn CS and Fox TR (1995) Inhibitory effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on rat hepatocyte proliferation induced by 2/3 partial hepatectomy. *Cell Prolif* **28**(8):437-451.
- Bunger MK, Moran SM, Glover E, Thomae TL, Lahvis GP, Lin BC and Bradfield CA (2003) Resistance to 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity and abnormal liver development in mice carrying a mutation in the nuclear localization sequence of the aryl hydrocarbon receptor. *J Biol Chem* **278**(20):17767-17774.
- Cressman DE, Diamond RH and Taub R (1995) Rapid activation of the Stat3 transcription complex in liver regeneration. *Hepatology* **21**(5):1443-1449.
- Elferink CJ, Ge NL and Levine A (2001) Maximal aryl hydrocarbon receptor activity depends on an interaction with the retinoblastoma protein. *Mol Pharmacol* **59**(4):664-673.
- Elizondo G, Fernandez-Salguero P, Sheikh MS, Kim GY, Fornace AJ, Lee KS and Gonzalez FJ (2000) Altered cell cycle control at the G(2)/M phases in aryl hydrocarbon receptor-null embryo fibroblast. *Mol Pharmacol* **57**(5):1056-1063.

- Fan F, Yan B, Wood G, Viluksela M and Rozman KK (1997) Cytokines (IL-1beta and TNFalpha) in relation to biochemical and immunological effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in rats. *Toxicology* **116**(1-3):9-16.
- Ge NL and Elferink CJ (1998) A direct interaction between the aryl hydrocarbon receptor and retinoblastoma protein. Linking dioxin signaling to the cell cycle. *J Biol Chem* **273**(35):22708-22713.
- Geisen C and Moroy T (2002) The oncogenic activity of cyclin E is not confined to Cdk2 activation alone but relies on several other, distinct functions of the protein. *J Biol Chem* **277**(42):39909-39918.
- Gottlicher M and Wiebel FJ (1991) 2,3,7,8-Tetrachlorodibenzo-p-dioxin causes unbalanced growth in 5L rat hepatoma cells. *Toxicol Appl Pharmacol* **111**(3):496-503.
- Hatakeyama M and Weinberg RA (1995) The role of RB in cell cycle control. *Prog Cell Cycle Res* **1**:9-19.
- Higgins GM and Anderson RM (1931) Experimental pathology of the liver: Restoration of the liver of the white rat following partial surgical removal. *Arch Pathol* **12**:186-202.
- Huang G and Elferink CJ (2005) Multiple mechanisms are involved in Ah receptor-mediated cell cycle arrest. *Mol Pharmacol* **67**(1):88-96.
- Ito T, Inouye K, Fujimaki H, Tohyama C and Nohara K (2002) Mechanism of TCDD-induced suppression of antibody production: effect on T cell-derived cytokine production in the primary immune reaction of mice. *Toxicol Sci* **70**(1):46-54.
- Jensen BA, Leeman RJ, Schlezinger JJ and Sherr DH (2003) Aryl hydrocarbon receptor (AhR) agonists suppress interleukin-6 expression by bone marrow stromal cells: an immunotoxicology study. *Environ Health* **2**(1):16.

- Kerkvliet NI, Baecher-Steppan L, Shepherd DM, Oughton JA, Vorderstrasse BA and DeKrey GK (1996) Inhibition of TC-1 cytokine production, effector cytotoxic T lymphocyte development and alloantibody production by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Immunol* **157**(6):2310-2319.
- Kolluri SK, Weiss C, Koff A and Gottlicher M (1999) p27(Kip1) induction and inhibition of proliferation by the intracellular Ah receptor in developing thymus and hepatoma cells. *Genes Dev* **13**(13):1742-1753.
- Levine-Fridman A, Chen L and Elferink CJ (2004) Cytochrome P4501A1 promotes G1 phase cell cycle progression by controlling aryl hydrocarbon receptor activity. *Mol Pharmacol* **65**(2):461-469.
- Ma Q and Whitlock JP, Jr. (1996) The aromatic hydrocarbon receptor modulates the Hepa 1c1c7 cell cycle and differentiated state independently of dioxin. *Mol Cell Biol* **16**(5):2144-2150.
- Mandal PK (2005) Dioxin: a review of its environmental effects and its aryl hydrocarbon receptor biology. *J Comp Physiol [B]* **175**(4):221-230.
- Mangnall D, Bird NC and Majeed AW (2003) The molecular physiology of liver regeneration following partial hepatectomy. *Liver Int* **23**(2):124-138.
- Marlowe JL, Knudsen ES, Schwemberger S and Puga A (2004) The aryl hydrocarbon receptor displaces p300 from E2F-dependent promoters and represses S phase-specific gene expression. *J Biol Chem* **279**(28):29013-29022.
- Moennikes O, Loeppen S, Buchmann A, Andersson P, Ittrich C, Poellinger L and Schwarz M (2004) A constitutively active dioxin/aryl hydrocarbon receptor promotes hepatocarcinogenesis in mice. *Cancer Res* **64**(14):4707-4710.

- Morris DL, Snyder NK, Gokani V, Blair RE and Holsapple MP (1992) Enhanced suppression of humoral immunity in DBA/2 mice following subchronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Toxicol Appl Pharmacol* **112**(1):128-132.
- Pitot HC, Goldsworthy T, Campbell HA and Poland A (1980) Quantitative evaluation of the promotion by 2,3,7,8-tetrachlorodibenzo-p-dioxin of hepatocarcinogenesis from diethylnitrosamine. *Cancer Res* **40**(10):3616-3620.
- Pohjanvirta R and Tuomisto J (1994) Short-term toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in laboratory animals: effects, mechanisms, and animal models. *Pharmacol Rev* **46**(4):483-549.
- Prell RA, Oughton JA and Kerkvliet NI (1995) Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on anti-CD3-induced changes in T-cell subsets and cytokine production. *Int J Immunopharmacol* **17**(11):951-961.
- Puga A, Barnes SJ, Dalton TP, Chang C, Knudsen ES and Maier MA (2000) Aromatic hydrocarbon receptor interaction with the retinoblastoma protein potentiates repression of E2F-dependent transcription and cell cycle arrest. *J Biol Chem* **275**(4):2943-2950.
- Rowlands JC and Gustafsson JA (1997) Aryl hydrocarbon receptor-mediated signal transduction. *Crit Rev Toxicol* **27**(2):109-134.
- Sherr CJ and Roberts JM (1995) Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev* **9**(10):1149-1163.
- Silkworth JB, Cutler DS and Sack G (1989) Immunotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in a complex environmental mixture from the Love Canal. *Fundam Appl Toxicol* **12**(2):303-312.

- Strobeck MW, Fribourg AF, Puga A and Knudsen ES (2000) Restoration of retinoblastoma mediated signaling to Cdk2 results in cell cycle arrest. *Oncogene* **19**(15):1857-1867.
- Walisser JA, Glover E, Pande K, Liss AL and Bradfield CA (2005) Aryl hydrocarbon receptor-dependent liver development and hepatotoxicity are mediated by different cell types. *Proc Natl Acad Sci U S A* **102**(49):17858-17863.
- Weglarz TC and Sandgren EP (2000) Timing of hepatocyte entry into DNA synthesis after partial hepatectomy is cell autonomous. *Proc Natl Acad Sci U S A* **97**(23):12595-12600.
- Weiss C, Kolluri SK, Kiefer F and Gottlicher M (1996) Complementation of Ah receptor deficiency in hepatoma cells: negative feedback regulation and cell cycle control by the Ah receptor. *Exp Cell Res* **226**(1):154-163.
- Welcker M, Singer J, Loeb KR, Grim J, Bloecher A, Gurien-West M, Clurman BE and Roberts JM (2003) Multisite phosphorylation by Cdk2 and GSK3 controls cyclin E degradation. *Mol Cell* **12**(2):381-392.

FOOTNOTES

Supported by the National Institutes of Environmental Health Sciences (NIEHS) Grant R01ES07800 (C.J.E.) and F32ES013588 (K.A.M.) and by NIEHS grants T32ES07254 and ES06676.

FIGURE LEGENDS

Figure 1. Analysis of TCDD toxicity during PH-induced liver regeneration. (A)

Representative photomicrographs (200x) of H&E-stained liver tissue harvested 36 h after PH/sham surgery in mice that were pretreated for 24 h with vehicle or TCDD. (B) Ratio of liver weight (wet) to body weight in vehicle- and TCDD-treated mice at the indicated times after PH. For each time indicated, 3-5 mice were used per treatment group. Results are representative of three separate experiments. Asterisks depict a significant difference when compared to vehicle group at same time point ($p \leq 0.05$).

Figure 2. Exposure to TCDD suppresses hepatocyte proliferation following PH. Mice were

treated with vehicle or TCDD 24 h prior to PH or sham surgery. Mice were killed 36 h later, immediately following a 2-hour pulse with BrdU, and liver tissue was processed and stained for immunohistochemistry. (A) BrdU-positive cells are identified by the dark brown (DAB-stained) nuclei in the photomicrographs (200X). (B) Nuclei were counted in three random fields (>300 cells/field) in liver tissue from each mouse 36 h after surgery. Data represent the average percentage of BrdU+ nuclei (\pm SEM) in liver tissue from 5 mice per treatment group. Results are representative of six separate experiments. Asterisk depicts a significant difference when compared to vehicle treatment group at same time ($p \leq 0.05$).

Figure 3. Suppressed liver regeneration is not due to delayed hepatocyte proliferation in

TCDD-treated mice. Mice were treated with vehicle or TCDD 24 h before PH. (A) Mice were pulsed for 2 h with BrdU before being sacrificed postoperatively at the indicated time. BrdU

incorporation was determined as described in Figure 2. Data represent the average percentage of BrdU+ nuclei (\pm SEM) in liver tissue from 5 mice per treatment group. Results are representative of three separate experiments. (B) Mice were continuously provided BrdU, which was added to their drinking water (0.8 mg/ml) for the 72 h immediately following PH. Data represent the average percentage of BrdU+ nuclei (\pm SEM) in liver tissue from 6 mice per treatment group. Asterisks depict a significant difference when compared to vehicle-treated mice that received PH ($p \leq 0.05$).

Figure 4. AhR activity persists throughout the regenerative process in TCDD-treated mice and is transiently induced in vehicle-treated mice. Mice were treated with vehicle or TCDD 24 h before PH and sacrificed postoperatively at the indicated time. (A) Liver homogenates (20 μ g/lane) were probed for cytochrome P4501A1 protein by western blot, and expression was analyzed by densitometry. Data are representative of three similar experiments. (B) CYP1A1 mRNA levels were measured in liver by qRT-PCR. Data are normalized to 18S and represent one mouse per treatment group. Open bars represent data from vehicle-treated mice, and solid bars depict TCDD-treated mice.


Figure 5. TCDD treatment does not alter the production of IL-6 or TNF α during liver regeneration. Mice were treated with vehicle or TCDD 24 h before PH or sham surgery and sacrificed postoperatively at the indicated time. Data reflect the average plasma concentrations of IL-6 (A) or TNF α (B). Each treatment group consisted of 5-6 PH mice and 3 sham-operated mice. Results are representative of 3-4 separate experiments.

Figure 6. Decreased proliferation TCDD-treated mice after PH is not due solely to alterations in early regenerative processes. Mice (5-8 per treatment group) were treated with vehicle or TCDD 24 h before PH, 5 min before PH, or 12 h after PH. Mice were pulsed with BrdU and killed 36 h postoperatively. Data depict the average percentage of BrdU+ nuclei (\pm SEM) in the regenerating liver. Asterisk indicates a significant difference when compared to vehicle-treated mice treated at the same time point ($p \leq 0.05$).


Figure 7. CDK2 activity is diminished during liver regeneration in TCDD-treated mice. Mice were treated with vehicle or TCDD 24 h prior to PH and killed at the indicated time. Liver homogenates were immunoprecipitated with rabbit IgG (first lane) or CDK2 antibody (remaining lanes). Kinase activity was measured by phosphorylation of histone H1. Data depict CDK2 kinase activity in the regenerating liver from individual mice at various times after PH, and is correlated with the percentage of BrdU+ hepatocytes from the same mouse. An * indicates that BrdU incorporation was $<1\%$.

Figure 8. AhR activation alters the association of CDK2 with cyclins E and A. Mice were treated with vehicle or TCDD 24 h before PH and sacrificed postoperatively at the indicated time. Liver homogenates were immunoprecipitated with a CDK2 antibody, and the indicated protein probed for by western blotting. Data are representative of three separate experiments in which unpooled liver homogenates were assayed from 2-3 individual mice for each time point.

A



B







A**B**

Figure 3

A



B







A**B**

Figure 5

A



B




Figure 6




Figure 7




Figure 8

