

MOL # 26195

Focal adhesion kinase (FAK) and protein kinase B cooperate to suppress doxorubicin-induced apoptosis of breast tumor cells.

Maroesja J. van Nimwegen[#], Merei Huigsloot[#], Annamarie Camier, Ine B. Tijdens and Bob van de Water[†].

Division of Toxicology, Leiden Amsterdam Center for Drug Research, Leiden University, Leiden, The Netherlands.

MOL # 26195

Running title: FAK and PKB in tumor cell apoptosis

Corresponding author:

Bob van de Water

Division of Toxicology, LACDR Leiden University

Einsteinweg 55, P.O. Box 9502

2300 RA Leiden

The Netherlands

Tel.: 31-71-5276223; Fax: 31-71-5274277

E-mail: b.water@LACDR.LeidenUniv.nl

Number of text pages: 24

Number of tables: 1

Number of figures: 7

Number of references: 40

Number of words abstract: 191

Number of words introduction: 734

Number of words discussion: 1304

Abbreviations: AV, AnnexinV; CLSM, Confocal Laser Scanning Microscopy; Dox, doxorubicin; ECM, extracellular matrix; FAK, focal adhesion kinase; FRNK, focal adhesion kinase-related non-kinase; GFP, green fluorescent protein; Neo, neomycin-resistant cells; PI, propidium iodide; PI3K, phosphatidyl inositol 3-kinase; PS, phosphatidylserine; PY, phosphotyrosine; zVAD-fmk, benzyloxycarbonyl-Val-Ala-DL-Asp-fluoromethylketone

MOL # 26195

Abstract

Focal adhesion kinase (FAK) is upregulated in a variety of cancers including breast cancer, in association with poor disease prognosis. In the present study we examined the role of focal adhesion kinase (FAK) in the control of anticancer drug-induced apoptosis of mammary adenocarcinoma MTLn3 cells. Doxorubicin caused formation of well-defined focal adhesions and stress fibers early after treatment, which was later followed by their loss in association with the onset of apoptosis. Phosphorylation of FAK on tyrosine 397 decreased only during the onset of doxorubicin-induced apoptosis, in a Bcl-2 and caspase-independent manner. Doxorubicin also caused an early activation of protein kinase B (PKB). Expression of the dominant negative acting focal adhesion kinase-related non-kinase (FRNK) sensitized MTLn3 cells to apoptosis caused by doxorubicin. FRNK inhibited the doxorubicin-induced activation of PKB. Also, inhibition of phosphatidylinositide-3 (PI-3) kinase with wortmannin inhibited the activation of PKB by doxorubicin. Both wortmannin as well as transient overexpression of the dual lipid/protein phosphatase PTEN enhanced doxorubicin-induced cell death. Altogether these data fit with a model wherein FAK is involved in the doxorubicin-induced activation of the PI-3 kinase/PKB signaling route, thereby suppressing the onset of apoptosis caused by doxorubicin.

MOL # 26195

Introduction

Adhesion to the ECM is essential for normal functioning of epithelial cells and provides survival signaling cues (reviewed by Frisch and Screaton, 2001; Giancotti and Ruoslahti, 1999). In normal cells, detachment generally results in apoptosis, also termed anoikis (Frisch and Francis, 1994; Frisch and Ruoslahti, 1997). However, tumor cells are often resistant to anoikis. This resistance is linked to constitutive activation of survival signaling pathways, by oncogenic transformation, that otherwise depend on cell adhesion (McFall et al., 2001; Streuli and Gilmore, 1999). These survival signals may confer resistance not only to anoikis but also to apoptosis induced by anticancer drugs due to an overlap in signaling pathways that control cell death. Indeed, adhesion to ECM proteins via the $\beta 1$ integrin subunit protected small cell lung cancer cells against doxorubicin and etoposide-induced apoptosis, and breast cancer cells against paclitaxel-induced apoptosis (Aoudjit and Vuori, 2001; Sethi et al., 1999). This dysregulation of cell-ECM interaction mediated signaling in tumor cells, followed by suppression of, or resistance to, apoptosis may well contribute to metastasis formation. Likewise, metastatic tumor cells may already be (partly) resistant to anticancer agent-induced apoptosis. Therefore, it is important to know how focal adhesion dependent signaling is linked to protection against anticancer agent induced apoptosis.

Adhesion to the ECM occurs through the extracellular domain of integrins, resulting in recruitment and activation of several focal adhesion-associated proteins to the intracellular domains. One of these proteins is focal adhesion kinase (FAK), which, upon binding to the integrins, is autophosphorylated on tyrosine residue 397 (Schaller et al., 1994). This phosphorylation recruits and activates Src and, together, these two tyrosine

MOL # 26195

kinases induce a cascade of tyrosine phosphorylations that regulate the activity and interactions of several adapter and signaling proteins at the focal adhesions (Schlaepfer et al., 1994), for example paxillin, p130Cas and Grb2 (Tachibana et al., 1997; Thomas et al., 1999). In addition, the FAK autophosphorylation domain interacts with phosphatidylinositol-3 kinase (PI-3K), which plays a central role in the activation of several downstream signaling pathways including survival signaling (see below). Several reports indicate that FAK is involved in the survival signaling generated by adhesion to the ECM.

Amplification of the FAK gene has been observed in several tumor types and the increase in FAK expression is associated with an increase in metastatic potential (Agochiya et al., 1999; Owens et al., 1995). This may be associated with protection against anoikis since experimental expression of constitutively active FAK decreased anoikis and enhanced tumor formation (Frisch et al., 1996). Conversely, inhibition of FAK function by FAK antisense oligonucleotides, microinjection of FAK antibodies or by expression of the C-terminal domain of FAK, which competes with FAK for focal adhesion localization, results in detachment and apoptosis (Hungerford et al., 1996; Xu et al., 2000). Moreover, FAK is cleaved by caspases during apoptosis induced by a variety of stimuli (van de Water et al., 2001; Wen et al., 1997); the resulting C-terminal domain of FAK resembles FAK-related non-kinase (FRNK) and acts as a dominant inhibitor of FAK, thereby further blocking FAK-mediated anti-apoptotic signaling (Gervais et al., 1998; Wen et al., 1997). Although much is known on the role of FAK in the control of cell survival in general, little information is available on the potential role of the

MOL # 26195

mechanism by which FAK interferes with the apoptosis caused by the anticancer drug doxorubicin in tumor cells.

A major survival signaling route is mediated by the activation of PI-3 kinase. Activation of PI-3 kinase results in generation of the second messengers PI(3,4,5)P₃ and PI(3,4)P₂. These phospholipids recruit protein kinase B (PKB or Akt) to the plasma membrane, and subsequently PKB is phosphorylated by PDK. Activated PKB provides survival signaling by inactivation of a series of pro-apoptotic proteins, for example p21WAF, FKHR, Bad and GSK3 (Datta et al., 1997). This PI3-kinase/PKB survival route can be inhibited by the tumor suppressor gene PTEN, which dephosphorylates both PIP₃ lipids as well as the protein kinases Shc and FAK (Di Cristofano and Pandolfi, 2000). So far, little is known on the relationship between FAK and the PI-3 kinase/PKB route in the control of cytostatic-induced apoptosis of tumor cells. Since our preliminary experiments indicated that doxorubicin caused activation of the PKB signaling pathway, we further determined the potential relationship between FAK function and PKB-mediated survival signaling.

To study the relationship between FAK and the PI-3K/PKB pathway in the control of doxorubicin-induced apoptosis we employed the rat mammary adenocarcinoma cell line MTLn3. These cells are often used as a model to study molecular mechanisms of metastasis formation (Kiley et al., 1999) and responses to drug therapy both *in vitro* and *in vivo* (Welch et al., 1983; Huigsloot et al., 2001). We have previously characterized, in detail, the induction of apoptosis by doxorubicin in these cells (Huigsloot et al., 2001; Huigsloot et al., 2002). Our current combined observations demonstrate that doxorubicin causes the early formation of stress fibers and focal

MOL # 26195

adhesions in association with activation of PKB. Pharmacological and molecular biological approaches indicate that PKB and FAK cooperate to suppress doxorubicin-induced apoptosis.

MOL # 26195

Materials and Methods

Chemicals. Alpha modified minimal essential medium with ribonucleosides and deoxyribonucleosides (α -MEM), Fetal Bovine Serum (FBS), penicillin/streptomycin, Lipofectamin Plus and geneticin (G418 sulphate) were from Life Technologies (Rockville, MD). Collagen (type I, rat tail) was from Upstate Biotechnology (Lake Placid, NY). Doxorubicin, etoposide, propidium iodide (PI) and RNase A were from Sigma (St. Louis, MO). Benzyloxycarbonyl-Val-Ala-DL-Asp-fluoromethylketone (zVAD-fmk) was from Bachem (Bubendorf, Switzerland). Annexin V was from Boehringer Mannheim (Basel, Switzerland). Hoechst 33258 was from Molecular Probes (Leiden, The Netherlands). Allophycocyanin (APC) was from Prozyme (San Leandro, CA). All other chemicals were of analytical grade.

Cell culture. MTLn3 rat mammary adenocarcinoma cells were originally developed by Dr. D.R. Welch (Jake Gittlen Cancer Research Institute, The Pennsylvania State University College of Medicine, Hershey, PA, USA) and used between passages 48 and 58. MTLn3 cells overexpressing Bcl-2 or empty vector (Bcl-2 and Neo cells, respectively) were obtained and characterized as described previously (Farrelly et al., 1999). All cell lines were cultured in α -MEM supplemented with 5% (v/v) FBS (complete medium). For experiments, cells were plated at a density of 4×10^3 cells/cm² in Corning plates (Acton, MA) and grown for three days in complete medium supplemented. For transient transfection, subconfluent cells were transfected with pEGFP, pEGFP-FRNK, pEGFP-FAT or GFP-PTEN wt, GFP-PTEN mutant using Lipofectamin Plus reagents. tetFRNK-MTLn3 cells were created as described previously (van Nimwegen et al., 2005). Exposure to doxorubicin occurred at 24 hours after

MOL # 26195

transfection. Cells were exposed to doxorubicin for one hour in Hanks' Balanced Salt Solution (137 mM NaCl, 5 mM KCl, 0.8 mM MgSO₄·7H₂O, 0.4 mM Na₂HPO₄·2H₂O, 0.4 mM KH₂PO₄, 1.3 mM CaCl₂, 4 mM NaHCO₃, 25 mM HEPES, 5 mM D-glucose; pH 7.4). After removal of doxorubicin, cells were allowed to recover in α -MEM containing 2.5% (v/v) FBS and penicillin/streptomycin for the indicated periods. In some experiments cells were recovered in α -MEM containing 2.5% (v/v) FBS, penicillin/streptomycin and 100 μ M zVAD-fmk.

Determination of cell death. For Annexin V/propidium iodide (AV/PI) staining, cells were washed twice in PBS (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄·2H₂O, 1.4 mM KH₂PO₄; pH 7.4) containing 1 mM EDTA (PBS-EDTA) and subsequently trypsinized with 0.13 g/L Trypsin in PBS-EDTA. Medium, washes and cells were combined, centrifuged (5 min, 200 g, 4°C) and the pellet was washed once with PBS-EDTA. Cells were allowed to recover from trypsinization in complete medium (30 minutes, 37°C). Externalized phosphatidylserine (PS) was labeled (15 minutes, 0°C) with APC-conjugated Annexin V in AV-buffer (10 mM HEPES, 145 mM NaCl, 5 mM KCl, 1.0 mM MgCl₂·6H₂O, 1.8 mM CaCl₂·2H₂O; pH 7.4). Propidium iodide (2 μ M) in AV-buffer was added one minute prior to analysis by flow cytometry on a FACScalibur (Becton Dickinson).

For cell cycle analysis, trypsinized and floating cells were pooled, washed with PBS-EDTA and fixated in 70% (v/v) ethanol (24 hours, -20°C). After two washes with PBS-EDTA, cells were incubated with PBS-EDTA containing 50 μ g/ml RNase A and 7.5 μ M PI (45 minutes, RT) and subsequently analyzed by flow cytometry.

MOL # 26195

Immunoblotting. Attached cells were scraped in ice-cold TSE+ (10 mM Tris-HCl, 250 mM sucrose, 1 mM EGTA, pH 7.4, containing 1 mM dithiothreitol, 10 µg/ml leupeptin, 10 µg/ml aprotinin, 1 mM sodium vanadate, 50 mM sodium fluoride, 1 mM phenylmethylsulfonyl fluoride). Floating cells in the medium and in one wash of PBS were pelleted (5 min, 200 g, 4°C) and pooled with scraped cells in TSE+. The protein concentration in the supernatant was determined using the BioRad protein assay using IgG as a standard. Fifteen µg of total cellular protein was separated by SDS-PAGE and transferred to PVDF membrane (Millipore, Etten Leur, The Netherlands). Blots were blocked with 5% (w/v) non-fat dry milk in TBS-T (0.5 M NaCl, 20 mM Tris-HCl, 0.05% v/v Tween-20; pH 7.4) and probed for active caspase-3 (CM-1, kindly provided by Dr. A. Srinivasan), PY (4G10, Upstate, Lake Placid, NY), PY³⁹⁷ FAK (polyclonal, Biosource Europe S.A., Nivelles, Belgium), FAK N-terminal antibody (77, BD BioSciences, Heidelberg, Germany), FAK C-terminal antibody (polyclonal, Upstate, Lake Placid, NY) or PKCδ (δ14K, kindly provided by Dr. S. Jaken), PTEN (BD BioSciences, Heidelberg, Germany), PKB (polyclonal, Cell signaling Technology, Beverly, MA) and P-Ser⁴⁷³-PKB (polyclonal, Cell Signaling Technology, Beverly, MA) followed by incubation with secondary antibody containing HRP and visualization with ECL reagent (Amersham Pharmacia Biotech, Uppsala, Sweden). For all Western blotting experiments equal protein loading was verified by Ponceau S (blots) and coomassie staining (gels) unless otherwise stated in figure legends.

Immunocytochemistry. Cells were cultured on 12 mm collagen-coated glass coverslips and fixated in 4% (w/v) formaldehyde in PBS. Coverslips were blocked in TBP (0.5% (w/v) BSA and 0.1% (v/v) Triton-X100 in PBS; pH 7.4) (1 hr, RT) and

MOL # 26195

subsequently incubated with primary antibody in TBP (1 hr, RT or O/N, 4°C). Coverslips were washed three times in TBP and incubated with Alexa488, Cy3 or Cy5-conjugated secondary antibodies or 0.2 units/ml rhodamine 123-conjugated phalloidin (Molecular Probes, Eugene, OR) in TBP (1 hr, RT). After washing with TBP, coverslips were incubated with 2 µg/ml Hoechst 33258 in PBS (15 min, RT), washed in PBS and mounted in Aqua PolyMount (Polysciences, Warrington, PA). Primary antibodies used were P-Tyr (4G10, Upstate, Lake Placid, NY) and vinculin (Sigma, St. Louis, MO). Imaging occurred using a Bio-Rad Radiance 2100 MP confocal laser scanning system equipped with a Nikon Eclipse TE2000-U inverted fluorescence microscope and a 60X Nikon objective.

Statistical analysis. Student's t test was used to determine if there was a significant difference between two means ($p < 0.05$). When multiple means were compared, significance was determined by one-way analysis of variance (ANOVA; $p < 0.05$).

MOL # 26195

Results

Doxorubicin causes focal adhesion formation prior to apoptosis. Treatment of the rat mammary adenocarcinoma MTLn3 cell line with the cytostatic doxorubicin resulted in a time and concentration dependent increase in apoptosis as well as activation of caspase-3 (Fig. 1A and B). Doxorubicin treated cells appeared more rounded up indicative for cytoskeleton-mediated cell contraction before cell death. Doxorubicin increased the number of F-actin stress fibers (Fig. 1C), which was associated with more concentrated focal adhesions as indicated by staining for the focal adhesion-associated protein vinculin (Fig. 1C). The F-actin organization and focal adhesion formation occurred early, between eight and twelve hours after exposure of the cells; after 16 hours these stainings were less pronounced, although these cells did not show signs of apoptosis (i.e. either positive for active caspase-3 and/or a fragmented nucleus). Since the onset of doxorubicin-induced apoptosis was associated with the loss of focal adhesions and F-actin stress fibers, abrogation of survival signaling generated by cell-ECM interaction may increase the effectiveness of doxorubicin-induced apoptosis. Indeed, when cells were kept in suspension, the extent of apoptosis caused by doxorubicin was strongly increased compared to cells allowed to attach to collagen (Fig. 1D). Next we studied whether this effect of doxorubicin was more general. For this purpose, MTLn3 cells were treated with either etoposide or cisplatin, two cytostatics which cause apoptosis after 16 hours in MTLn3 cells, a similar timeframe as doxorubicin (Huigsloot et al., 2001). Both etoposide as well as cisplatin caused the increased formation of focal adhesions as well as F-actin stress fiber formation (Fig. 1E).

MOL # 26195

Doxorubicin decreases protein tyrosine phosphorylation at focal adhesions independent of caspase-3 activity. Our results indicated that cell adhesion to the ECM confers a survival signal to MTLn3 cells exposed to doxorubicin. Therefore, we next investigated whether doxorubicin itself induced a change in cell-ECM dependent signaling preceding or during apoptosis. Adhesion dependent signaling at focal adhesions is mediated by a number of tyrosine kinases (reviewed by (Giancotti and Ruoslahti, 1999)) and many tyrosine phosphorylated proteins are localized at focal adhesions. We first investigated the time-course of tyrosine phosphorylation (PY) after treatment with doxorubicin. Up to 8 hours after exposure to doxorubicin, no major changes in overall tyrosine phosphorylation were observed (Fig. 2A). However, at 16 hours tyrosine phosphorylation was decreased and at 24 hours little tyrosine phosphorylation remained; the latter coincided with the onset of apoptosis (Fig. 2A right panel and Huigsloot et al., 2001).

To investigate whether the decrease in PY was associated with a loss of tyrosine phosphorylation at the focal adhesions, PY localization was determined by immunofluorescence in combination with F-actin staining (Fig. 2B). In control cells, a large portion of PY staining was located at focal adhesion-like structures at the cell periphery. At eight hours after exposure to doxorubicin, more pronounced focal adhesions were observed compared to control cells; the PY staining was also more intense. The newly formed focal adhesions were predominantly located at the ends of F-actin stress fibers. Since there was no overall increase in tyrosine-phosphorylation (Fig. 2A), the increase in PY staining at the focal adhesions is most likely due to complexation and/or concentration of tyrosine phosphorylated proteins at these signaling sites. At 16

MOL # 26195

hours, cells were rounded up and focal adhesions had become much smaller and less numerous, while F-actin fibers had disappeared. Hoechst staining indicated that the nuclei of these (attached) cells still appeared normal (data not shown). Inhibition of caspase-3 activity by addition of the pan-caspase inhibitor zVAD-fmk inhibited doxorubicin induced apoptosis (data not shown, (Huigsloot et al., 2002)) but could not prevent rounding of the cells or the disappearance of the large focal adhesions after 16 hrs (Fig. 2B).

Doxorubicin-induced dephosphorylation of FAK is independent of cleavage by caspases. Focal adhesion kinase (FAK) is an important tyrosine phosphorylated protein that localizes at focal adhesions and is involved in survival signaling. The most essential site of tyrosine phosphorylation on FAK is the autophosphorylation site Tyr³⁹⁷, which is required for both kinase activity and the interaction with other signaling proteins (Schaller et al., 1994). Using a phosphorylation state specific antibody, we found that doxorubicin caused dephosphorylation of Tyr³⁹⁷-FAK after 16 and 24 hours (Fig 3A). The expression level of FAK decreased after 16 and 24 hours, which was associated with the formation of a 75 kD band. The formation of this cleavage product is most likely mediated by caspases, which are activated in a similar time-course (Fig 2A bottom and (Huigsloot et al., 2001)). The dephosphorylation and caspase-3-mediated proteolysis of FAK may be difficult to separate and cell stress dependent (Kabir et al., 2002). Using zVAD-fmk we excluded the possibility that the decrease in Tyr³⁹⁷-phosphorylation was caused by caspase-mediated cleavage of FAK. zVAD-fmk effectively inhibited apoptosis (Huigsloot et al., 2001) as well as caspase activation upon exposure of MTLn3 cells to doxorubicin: no active caspase-3 and PKC δ cleavage fragments were formed (Fig. 3B).

MOL # 26195

zVAD-fmk also inhibited the cleavage of FAK, indicating that formation of the cleavage fragment of FAK was indeed due to a caspase-dependent process. Despite the inhibition of apoptosis and FAK cleavage, a decrease in tyrosine phosphorylation of FAK still occurred.

Since dephosphorylation of FAK occurs upstream of caspase activation, the question was raised whether FAK dephosphorylation occurs upstream of the mitochondrial commitment to apoptosis. Therefore, we used MTLn3 cells overexpressing Bcl-2 (Fig. 4A), which reduced apoptosis from 33 % in Neo cells to 11 % in Bcl-2 cells at 24 hours after exposure to 17 μ M doxorubicin (Fig. 4B). Bcl-2 overexpression also prevented the activation of caspases by doxorubicin as well as the cleavage of PKC δ (Fig. 4C). Importantly, FAK cleavage was also inhibited by Bcl-2 (Fig. 4C). In agreement with the persistence of dephosphorylation in case of caspase-inhibition by zVAD-fmk, inhibition of caspase activation by Bcl-2 overexpression did not prevent the dephosphorylation of FAK occurring 16 hours after exposure; FAK levels in Bcl-2 cells did not decrease at these time points. Together these data suggest that dephosphorylation of FAK occurs upstream of the Bcl-2 checkpoint at the level of the mitochondria and independent of caspase activation and may contribute to the commitment to apoptosis in MTLn3 cells.

Loss of FAK function by expression of eGFP-FRNK sensitizes MTLn3 cells to doxorubicin induced apoptosis. Next we further investigated the role of FAK in the protection against doxorubicin-induced apoptosis. For this purpose, we interfered FAK function by expression of the FAK-related non-kinase (FRNK) and the FAT domain of FAK. eGFP-FAK and eGFP were included as controls. At 24 hours after transfection,

MOL # 26195

eGFP was located diffusely throughout the cell, whereas the other constructs localized mainly at focal adhesions (data not shown, (van de Water et al., 2001)). To test if FAK is required for cell survival of MTLn3 cells we determined whether expression of eGFP-FAK, eGFP-FRNK or eGFP-FAT induced apoptosis. At 40 hours after transfection, eGFP positive cells were selected and apoptosis was determined in these cells by additional staining with APC-coupled annexin V and propidium iodide. Expression of eGFP-FRNK alone was sufficient to increase the percentage of annexin V⁺/PI⁺ (i.e. apoptotic) cells to 18 %, whereas eGFP-FAT increased apoptosis to only 14 % (Table I). No significant effect on apoptosis was observed by eGFP-FAK. Next, we examined the effect of abrogation of FAK signaling by expression of eGFP-FRNK on doxorubicin-induced apoptosis. Here we used 10 μ M of doxorubicin which allows identification of a synergistic effect. At 24 hours after transfection of the different eGFP-constructs, cells were exposed to doxorubicin and after 16 hours the percentage of apoptotic cells was determined (Table I). When eGFP-FRNK transfected cells were exposed to doxorubicin the percentage of apoptosis was even further increased, suggesting that functional FAK can suppress doxorubicin-induced apoptosis. Expression of eGFP-FAT was less effective and full length eGFP-FAK did not protect against doxorubicin, suggesting that FAK-mediated survival signaling was already maximal in MTLn3 cells. The apoptotic effect of eGFP-FRNK was concentration-dependent; only in a sub-population of MTLn3 cells with high eGFP-FRNK expression increased apoptosis was observed (data not shown). Inhibition of FAK using eGFP-FRNK was not able to potentiate etoposide-induced apoptosis (data not shown). These data suggest expression of eGFP-FRNK abrogates survival signaling and thereby increases doxorubicin-induced apoptosis in MTLn3 cells.

MOL # 26195

Doxorubicin causes PKB activation in a FAK-dependent manner. To investigate the FAK-mediated downstream signaling events involved in the control of cell survival of breast tumor cells, we used an MTLn3 cell line that conditionally expresses HA-tagged FRNK in a doxycycline-dependent manner (tetFRNK-MTLn3). Expression of HA-FRNK inhibited endogenous FAK function since it reduced MTLn3 cell attachment and spreading as well as cell migration (van Nimwegen et al., 2005). In these cells expression of HA-FRNK itself did hardly affect cell survival, due to the relative low expression of this inducible HA-FRNK compared to the transient transfected eGFP-FRNK. Yet HA-FRNK potentiated the onset of doxorubicin-induced apoptosis (Fig. 5A). We further evaluated the cell adhesion-mediated signaling pathways that are controlled by FAK in MTLn3 cells. Therefore tetFRNK-MTLn3 cells were allowed to adhere to collagen. We primarily focused on protein kinase B (PKB) because of its central role in cell survival signaling. Adhesion of cells to collagen resulted in a rapid activation of protein kinase B (Fig. 5B). Expression of HA-FRNK delayed the attachment-mediated activation of PKB (Fig. 5B). No effect of HA-FRNK expression on the activation of ERK and JNK in response to attachment was observed (data not shown).

Next we studied the relationship between PKB activity and doxorubicin-induced apoptosis. To our surprise doxorubicin caused an activation of PKB at 2 and 4 hours (Fig. 5C, control) which occurred slightly before the formation of stress fibers and focal adhesions. To test whether HA-FRNK also affected the doxorubicin-induced activation of PKB, cells were pretreated with doxycycline followed by exposure to doxorubicin. HA-FRNK expression did not affect the activity of PKB under control conditions (data not shown). However, importantly, in the HA-FRNK expressing cells PKB was not

MOL # 26195

phosphorylated in response to doxorubicin (Fig. 5C, HA-FRNK). This suggests a role for FAK in the activation of the PI-3 kinase survival pathway in response to doxorubicin exposure of MTLn3 cells.

Inhibition of PI-3 kinase and overexpression of PTEN enhance doxorubicin-induced apoptosis. The above indicated that normal FAK function is required for doxorubicin-mediated activation of PKB. Therefore, we determined the involvement of PKB in the control of doxorubicin-induced apoptosis. For this purpose we used both pharmacological and molecular biological approaches. Treatment of MTLn3 cells with an inhibitor of PI-3 kinase, wortmannin, completely prevented the phosphorylation of PKB (Fig. 6A). This inhibition of PKB phosphorylation was associated with enhanced apoptosis of MTLn3 cells after doxorubicin treatment as measured by cell cycle analysis (Fig. 6B). In addition, we used the dual-protein/lipid-phosphatase PTEN as an alternative to prevent PKB activation. PTEN is a tumor suppressor gene that is mutated in various types of cancer, including breast cancer (Wu et al., 2003). PTEN dephosphorylates PIP3, which is formed after activation of PI-3 kinase. To study the effect of PTEN, cells were transiently transfected with GFP-PTEN wild type or GFP-PTEN-C124A, a phosphatase inactive mutant. Immunoblotting using antibodies directed against PTEN and GFP confirmed the presence of the transfected proteins (Fig. 6C). Transfected cells were treated with doxorubicin and the percentage apoptotic cells was determined using annexin-V-APC/PI staining followed by flow cytometric analysis. Overexpression of wild type PTEN enhanced doxorubicin-induced apoptosis almost two-fold; this effect was dependent on the phosphatase activity, since expression of the phosphatase inactive mutant PTEN-C124A did not significantly affect doxorubicin-induced apoptosis. Both

MOL # 26195

wild type and mutant PTEN itself did not affect the viability of MTLn3 cells (Fig. 6D). PTEN can also dephosphorylate FAK on tyrosine residue 397 (Tamura et al., 1998). To exclude this possibility we also determined the FAK-Y³⁹⁷ phosphorylation status in GFP positive cells after transfection. Neither wild type nor mutant PTEN affected FAK phosphorylation or the reorganization of focal adhesions (data not shown). These data provide evidence that the doxorubicin-induced signaling through the PI-3 kinase/PKB pathway suppresses the doxorubicin-induced apoptosis.

MOL # 26195

Discussion

In the present study, we examined the relationship between FAK and the activation of the PI-3 kinase/PKB survival signaling route in control of doxorubicin-induced apoptosis of mammary adenocarcinoma cells. Several important conclusions can be drawn. Firstly, doxorubicin caused the reorganization of the F-actin network in association with increased focal adhesion formation which is directly associated with the increased activation of PKB, but not focal adhesion kinase. These cell biological changes are independent of activation of the apoptotic machinery, since increased expression of Bcl-2 or inhibition of caspase activity do not protect against it. Secondly, FAK function is involved in the maintenance of survival signaling in the MTLn3 cells, and inhibition of FAK function by expression of FRNK accelerates the onset of doxorubicin-induced apoptosis in association with a FRNK-mediated inhibition of PKB activation by doxorubicin. Thirdly, the doxorubicin-induced apoptosis is suppressed by PI-3 kinase/PKB signaling since overexpression of PTEN enhances doxorubicin-induced apoptosis. The combined data support a paradigm, in which doxorubicin causes an increased formation of focal adhesions; FAK is then involved in the doxorubicin-induced activation of the PKB signaling pathway, thereby suppressing the onset of Bcl-2 dependent apoptosis.

Our data indicate a close relationship between FAK function and doxorubicin-induced PKB activation and suppression of apoptosis caused by doxorubicin. First of all, doxorubicin itself induced the activation of PKB which correlated with the increased formation of stress fibers and focal adhesions. Expression of FRNK in MTLn3 cells inhibited the cell adhesion-mediated activation of PKB. Secondly, inhibition of PI-3

MOL # 26195

kinase using wortmannin blocked the doxorubicin-mediated PKB activation and sensitized cells to apoptosis. Thirdly, overexpression of PTEN, which dephosphorylates PIP3, also increased the sensitivity of MTLn3 cells towards doxorubicin. Although in prostate tumor cells PY³⁹⁷ of FAK is also a target for PTEN (Tamura et al., 1998), no effect of PTEN overexpression on PY³⁹⁷-FAK was seen. This excludes the possibility that the effect of PTEN is related to combined effect of PIP3 and FAK dephosphorylation, and further supports the role of PKB signaling in the control of doxorubicin-induced apoptosis. Also in LLC-PK1 cells the nephrotoxicant dichlorovinyl-cysteine caused activation of PKB which is inhibited by the dominant negative acting deletion mutant FAT. Moreover, in these cells PTEN overexpression also facilitates the onset of apoptosis (van de Water et al., manuscript in preparation). These combined datasets suggest the existence of a general cellular stress-mediated signaling pathway through FAK that results in the activation of PKB thereby providing survival signals.

Focal adhesions are important anchoring sites for the F-actin cytoskeleton and many focal adhesion-associated proteins directly or indirectly regulate actin dynamics. We found that, initially, doxorubicin increased the bundling of F-actin into stress fibers, which was associated with the formation of larger focal adhesions with a concentration of tyrosine phosphorylated proteins. A possible pathway leading to stress fiber formation is through reduction of doxorubicin to a semiquinone free radical intermediate by complex I of the mitochondrial electron transport chain and subsequent generation of reactive oxygen species (ROS) (Wallace, 2003). A similar cytoskeletal response was found in endothelial and glioblastoma cells, which, upon exposure to the ROS-generating agent H₂O₂, formed stress fibers prior to an increase in tyrosine phosphorylation of FAK and

MOL # 26195

paxillin (Sonoda et al., 1999). One of the proteins that is known to induce these specific cytoskeletal changes is p21Rho, a member of the Rho GTPase family (Ridley and Hall, 1992), suggesting that p21Rho may be an intermediate in doxorubicin- and ROS-induced signaling. Indeed, pharmacological inhibition of a downstream effector of RhoA, Rho-kinase, using Y27632, inhibited F-actin stress fiber accumulation and focal adhesion organization in MTLn3 cells caused by doxorubicin; also no effect of Y27632 on FAK dephosphorylation caused by doxorubicin was observed (van Nimwegen et al., unpublished observation). Tyrosine phosphorylation of FAK correlates with protection against apoptosis, suggesting that doxorubicin initially promotes survival signaling, which is subsequently overridden by pro-apoptotic signaling. After the initial increase, tyrosine phosphorylation strongly decreased at 16 hours after exposure, suggesting that doxorubicin-induced cytotoxicity involves several signaling pathways with differential kinetics. The decrease in tyrosine phosphorylation at focal adhesions occurred concomitant with the disruption of the F-actin network, suggesting a direct relationship between tyrosine phosphorylation and cytoskeletal organization. Such a decrease may be directly related to potential inactivation of c-Src activity in the process leading to the onset of apoptosis as has been described for TGF-beta1-induced apoptosis (Park et al., 2004).

Our data indicate that interference of the function of FAK by expression of eGFP-FRNK abrogated the FAK-mediated suppression of apoptosis in adherent cells. These data are consistent with those obtained by other studies: expression of either FAT in fibroblasts or the C-terminal domain of FAK, which is similar to FRNK, in breast cancer cells caused apoptosis (Ilic et al., 1998). In addition to these studies, we now show that

MOL # 26195

FAK is also involved in the suppression of apoptosis induced by the anticancer agent doxorubicin in breast cancer cells. This suggests that these two stress response pathways may converge. For treatment of adherent tumor cells the protective function of FAK was only demonstrated for cisplatin in FAT expressing glioblastoma cells (Jones et al., 2001) and a link to the PI-3 kinase/PKB pathway was not demonstrated. Induction of apoptosis of BT474 by the FAK C-terminal domain was inhibited by EGFR, which was associated with activation of the PI-3 kinase/PKB route (Golubovskaya et al., 2002). We found that overexpression of full length FAK did not provide protection against doxorubicin-induced apoptosis. This suggests that endogenous FAK is present in excess in MTLn3 cells and that the expression of either focal adhesion-associated binding partners and/or downstream adapter or signaling proteins limit the extent of FAK induced survival signaling. In that case, exogenously added FAK would be unable to amplify FAK-mediated survival signaling. This is in contrast to the findings in HL-60 leukemia cells, which express little endogenous FAK and normally do not adhere, and in which overexpression of FAK protected against hydrogen peroxide or etoposide induced apoptosis (Sonoda et al., 2000).

FRNK-mediated apoptosis was clearly dependent on its expression levels. In our inducible tetFRNK-MTLn3 cell line expression of FRNK itself did not affect apoptosis, while inhibition of FAK potentiated doxorubicin-induced apoptosis. This indicates that at these levels of FRNK expression growth factor receptors are able to mediate survival signals that compensate the decrease in FAK mediated survival signals.

The degradation of FAK was a late event which occurred in parallel with dephosphorylation of FAK. Only when apoptosis was prevented by inhibition of caspase

MOL # 26195

activity with zVADfmk, we could define doxorubicin-mediated FAK dephosphorylation as a process that does not require caspase activation. Overexpression of Bcl-2, like previously shown, prevented doxorubicin-induced apoptosis but did not prevent dephosphorylation of FAK. Therefore, doxorubicin-induced dephosphorylation of FAK is a process that occurs upstream of the mitochondrial check-point of apoptosis. In a recent study in opossum kidney cells ATP depletion rapidly (within 15 min) caused caspase-dependent fragmentation of FAK, which was inhibited by caspase-3 (Mao et al., 2003). Moreover, in primary porcine aortic endothelial cells staurosporine, but not cycloheximide, caused a rapid dephosphorylation of FAK that preceded the onset of apoptosis (Kabir et al., 2002). These data together indicate that the regulation of FAK dephosphorylation and degradation is dependent on the cell type and cellular stress conditions. Our data indicate that FAK dephosphorylation by doxorubicin occurs independent from caspase activity.

In conclusion, doxorubicin caused an early activation of PKB which is dependent on the function/localization of FAK at focal adhesions. Both inhibition of FAK with FRNK or inhibition of the PI-3K/PKB pathway with wortmannin or PTEN increased the sensitivity of MTLn3 cells towards doxorubicin-induced apoptosis. Doxorubicin also caused an early formation of F-actin stress fibers and focal adhesions, followed by dephosphorylation of PY³⁹⁷FAK and loss of focal adhesions. Abrogation of the apoptotic pathway was possible at the level of the mitochondria by overexpression of Bcl-2 and at the level of the executioner caspases by addition of zVAD-fmk. These data suggest that pharmacological inhibition of FAK and/or the PI-3K/PKB pathway may represent a

MOL # 26195

possible means to enhance the treatment of cancer with classical anticancer drugs such as doxorubicin.

MOL # 26195

Acknowledgements

We thank Dusko Ilic and Kenneth Yamada for providing cDNA constructs and the members of our laboratory for helpful suggestions and discussions.

MOL # 26195

References

- Agochiya M, Brunton VG, Owens DW, Parkinson EK, Paraskeva C, Keith WN and Frame MC (1999) Increased dosage and amplification of the focal adhesion kinase gene in human cancer cells. *Oncogene* **18**:5646-5653.
- Aoudjit F and Vuori K (2001) Integrin signaling inhibits paclitaxel-induced apoptosis in breast cancer cells. *Oncogene* **20**:4995-5004.
- Datta SR, Dudek H, Tao X, Masters S, Fu H, Gotoh Y and Greenberg ME (1997) Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell* **91**:231-241.
- Di Cristofano A and Pandolfi PP (2000) The multiple roles of PTEN in tumor suppression. *Cell* **100**:387-390.
- Farrelly N, Lee YJ, Oliver J, Dive C and Streuli CH (1999) Extracellular matrix regulates apoptosis in mammary epithelium through a control on insulin signaling. *J Cell Biol* **144**:1337-1348.
- Frisch SM and Francis H (1994) Disruption of epithelial cell-matrix interactions induces apoptosis. *J Cell Biol* **124**:619-626.
- Frisch SM and Ruoslahti E (1997) Integrins and anoikis. *Curr Opin Cell Biol* **9**:701-706.
- Frisch SM and Screaton RA (2001) Anoikis mechanisms. *Curr Opin Cell Biol* **13**:555-562.
- Frisch SM, Vuori K, Ruoslahti E and Chan-Hui PY (1996) Control of adhesion-dependent cell survival by focal adhesion kinase. *J Cell Biol* **134**:793-799.
- Gervais FG, Thornberry NA, Ruffolo SC, Nicholson DW and Roy S (1998) Caspases cleave focal adhesion kinase during apoptosis to generate a FRNK-like polypeptide. *J Biol Chem* **273**:17102-17108.
- Giancotti FG and Ruoslahti E (1999) Integrin signaling. *Science* **285**:1028-1032.
- Golubovskaya V, Beviglia L, Xu LH, Earp HS, III, Craven R and Cance W (2002) Dual inhibition of focal adhesion kinase and epidermal growth factor receptor pathways cooperatively induces death receptor-mediated apoptosis in human breast cancer cells. *J Biol Chem* **277**:38978-38987.
- Huigslout M, Tijdens IB, Mulder GJ and van de Water B (2001) Differential regulation of phosphatidylserine externalization and DNA fragmentation by caspases in anticancer drug-induced apoptosis of rat mammary adenocarcinoma MTLn3 cells. *Biochem Pharmacol* **62**:1087-1097.
- Huigslout M, Tijdens IB, Mulder GJ and van de Water B (2002) Differential regulation of doxorubicin-induced mitochondrial dysfunction and apoptosis by Bcl-2 in mammary adenocarcinoma (MTLn3) cells. *J Biol Chem* **277**:35869-35879.
- Hungerford JE, Compton MT, Matter ML, Hoffstrom BG and Otey CA (1996) Inhibition of pp125FAK in cultured fibroblasts results in apoptosis. *J Cell Biol* **135**:1383-1390.
- Ilic D, Almeida EA, Schlaepfer DD, Dazin P, Aizawa S and Damsky CH (1998) Extracellular matrix survival signals transduced by focal adhesion kinase suppress p53-mediated apoptosis. *J Cell Biol* **143**:547-560.
- Jones G, Machado J, Jr., Tolnay M and Merlo A (2001) PTEN-independent induction of caspase-mediated cell death and reduced invasion by the focal adhesion targeting domain (FAT) in human astrocytic brain tumors which highly express focal adhesion kinase (FAK). *Cancer Res* **61**:5688-5691.

MOL # 26195

Kabir J, Lobo M and Zachary I (2002) Staurosporine induces endothelial cell apoptosis via focal adhesion kinase dephosphorylation and focal adhesion disassembly independent of focal adhesion kinase proteolysis. *Biochem J* **367**:145-155.

Kiley SC, Clark KJ, Goodnough M, Welch DR and Jaken S (1999) Protein kinase C delta involvement in mammary tumor cell metastasis. *Cancer Res* **59**:3230-3238.

Mao H, Li F, Ruchalski K, Mosser DD, Schwartz JH, Wang Y and Borkan SC (2003) hsp72 inhibits focal adhesion kinase degradation in ATP-depleted renal epithelial cells. *J Biol Chem* **278**:18214-18220.

McFall A, Ulku A, Lambert QT, Kusa A, Rogers-Graham K and Der CJ (2001) Oncogenic Ras blocks anoikis by activation of a novel effector pathway independent of phosphatidylinositol 3-kinase. *Mol Cell Biol* **21**:5488-5499.

Owens LV, Xu L, Craven RJ, Dent GA, Weiner TM, Kornberg L, Liu ET and Cance WG (1995) Overexpression of the focal adhesion kinase (p125FAK) in invasive human tumors. *Cancer Res* **55**:2752-2755.

Park SS, Eom YW, Kim EH, Lee JH, Min dS, Kim S, Kim SJ and Choi KS (2004) Involvement of c-Src kinase in the regulation of TGF-beta1-induced apoptosis. *Oncogene* **19**:6272-6281.

Ridley AJ and Hall A (1992) The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell* **70**:389-399.

Schaller MD, Hildebrand JD, Shannon JD, Fox JW, Vines RR and Parsons JT (1994) Autophosphorylation of the focal adhesion kinase, pp125FAK, directs SH2-dependent binding of pp60src. *Mol Cell Biol* **14**:1680-1688.

Schlaepfer DD, Hanks SK, Hunter T and van der GP (1994) Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase. *Nature* **372**:786-791.

Sethi T, Rintoul RC, Moore SM, MacKinnon AC, Salter D, Choo C, Chilvers ER, Dransfield I, Donnelly SC, Strieter R and Haslett C (1999) Extracellular matrix proteins protect small cell lung cancer cells against apoptosis: a mechanism for small cell lung cancer growth and drug resistance in vivo. *Nat Med* **5**:662-668.

Sonoda Y, Matsumoto Y, Funakoshi M, Yamamoto D, Hanks SK and Kasahara T (2000) Anti-apoptotic role of focal adhesion kinase (FAK). Induction of inhibitor-of-apoptosis proteins and apoptosis suppression by the overexpression of FAK in a human leukemic cell line, HL-60. *J Biol Chem* **275**:16309-16315.

Sonoda Y, Watanabe S, Matsumoto Y, Aizu-Yokota E and Kasahara T (1999) FAK is the upstream signal protein of the phosphatidylinositol 3-kinase-Akt survival pathway in hydrogen peroxide-induced apoptosis of a human glioblastoma cell line. *J Biol Chem* **274**:10566-10570.

Streuli CH and Gilmore AP (1999) Adhesion-mediated signaling in the regulation of mammary epithelial cell survival. *J Mammary Gland Biol Neoplasia* **4**:183-191.

Tachibana K, Urano T, Fujita H, Ohashi Y, Kamiguchi K, Iwata S, Hirai H and Morimoto C (1997) Tyrosine phosphorylation of Crk-associated substrates by focal adhesion kinase. A putative mechanism for the integrin-mediated tyrosine phosphorylation of Crk-associated substrates. *J Biol Chem* **272**:29083-29090.

Tamura M, Gu J, Matsumoto K, Aota S, Parsons R and Yamada KM (1998) Inhibition of cell migration, spreading, and focal adhesions by tumor suppressor PTEN. *Science* **280**:1614-1617.

MOL # 26195

Thomas JW, Cooley MA, Broome JM, Salgia R, Griffin JD, Lombardo CR and Schaller MD (1999) The role of focal adhesion kinase binding in the regulation of tyrosine phosphorylation of paxillin. *J Biol Chem* **274**:36684-36692.

van de Water B, Houtepen F, Huigsloot M and Tijdens IB (2001) Suppression of chemically induced apoptosis but not necrosis of renal proximal tubular epithelial (LLC-PK1) cells by focal adhesion kinase (FAK). Role of FAK in maintaining focal adhesion organization after acute renal cell injury. *J Biol Chem* **276**:36183-36193.

van Nimwegen MJ, Verkoeijen S, van Buren L, Burg D and van de Water B (2005) Requirement for focal adhesion kinase in the early phase of mammary adenocarcinoma lung metastasis formation. *Cancer Res* **65**:4698-4706.

Wallace KB (2003) Doxorubicin-induced cardiac mitochondrionopathy. *Pharmacol Toxicol* **93**:105-115.

Welch DR, Neri A and Nicolson GL (1983) Comparison of 'spontaneous' and 'experimental' metastasis using rat 13762 mammary adenocarcinoma metastatic cell clones. *Invasion Metastasis* **3**:65-80.

Wen LP, Fahrni JA, Troie S, Guan JL, Orth K and Rosen GD (1997) Cleavage of focal adhesion kinase by caspases during apoptosis. *J Biol Chem* **272**:26056-26061.

Wu H, Goel V and Haluska FG (2003) PTEN signaling pathways in melanoma. *Oncogene* **19**:3113-3122.

Xu LH, Yang X, Bradham CA, Brenner DA, Baldwin AS, Craven RJ and Cance WG (2000) The focal adhesion kinase suppresses transformation-associated, anchorage-independent apoptosis in human breast cancer cells. Involvement of death receptor-related signaling pathways. *J Biol Chem* **275**:30597-30604.

MOL # 26195

Footnotes

#These authors contributed equally to this work.

This work was supported by a fellowship from the Royal Netherlands Academy for Arts and Sciences (B.v.d.W.) and by grant RUL 2001-2477 from the Dutch Cancer Society (B.v.d.W.).

MOL # 26195

Legends to figures

Figure 1. Doxorubicin-induced focal adhesion formation occurs prior to caspase-3 activation and apoptosis. MTLn3 cells were exposed to vehicle (control) or different concentrations doxorubicin for one hour and allowed to recover. At the indicated time-points, cells were fixated and the percentage apoptosis was determined by flow cytometric analysis of cellular DNA content as described in Materials and Methods (A). Caspase-3 activity was determined as described in Materials and Methods (B). Attached cells were immunostained for vinculin and F-actin and pictures were taken using CLSM (C). MTLn3 cells were exposed to vehicle or 10 μ M doxorubicin while kept in suspension or allowed to attach to collagen. After 8 hours the percentage of apoptosis was determined by flow cytometric analysis of cellular DNA content (D). MTLn3 cells were exposed to vehicle, 100 μ M etoposide or 100 μ M cisplatin and immunostained for vinculin and F-actin and pictures were taken using CLSM (E). Data shown (A, B and D) are the mean of three independent experiments \pm SE.

Figure 2. Doxorubicin causes a decrease in tyrosine phosphorylation at focal adhesions concomitant with the activation of caspase-3. MTLn3 cells were exposed to vehicle (Con) or 17 μ M doxorubicin for one hour and allowed to recover for the indicated period prior to fixation. Cell lysates were immunoblotted against phospho-tyrosine (A, left panel) and active caspase-3 (A, right panel). MTLn3 cells were fixated and stained for phospho-tyrosine and F-actin (B). Where indicated, zVAD-fmk (100 μ M) was added during the recovery of the exposed cells. Immunofluorescence was evaluated by CLSM

MOL # 26195

and phase contrast images are included. Data shown are representative for three independent experiments.

Figure 3. Doxorubicin causes FAK dephosphorylation independent of caspase activation. MTLn3 cells were treated as described in Fig. 2 for the indicated period (A) or for 24 hours (B) and where indicated, zVAD-fmk (100 μ M) was added during the recovery period. Immunoblots were stained with antibodies directed against PY³⁹⁷-FAK, FAK, PKC δ and the active cleavage fragments of caspase-3 (act. casp-3). Data shown are representative for three independent experiments.

Figure 4. Doxorubicin-induced FAK dephosphorylation occurs independent of the Bcl-2 control point of apoptosis. Cell lysates of neomycin resistant (Neo) and Bcl-2 overexpressing MTLn3 clones (Bcl-2) were immunoblotted against Bcl-2 (A). Neo and Bcl-2 cells were exposed to 17 μ M doxorubicin for one hour and allowed to recover for the indicated period prior to fixation. Apoptosis was evaluated using cell cycle analysis (B). Total cell lysates were evaluated for active caspase-3, PKC delta cleavage and FAK phosphorylation by immunoblotting (C). Data shown are representative for three independent experiments or the mean of three independent experiments \pm SE.

Figure 5. Conditional expression of FRNK enhances doxorubicin-induced apoptosis and reduces activation of PKB. Doxycycline inducible HA-FRNK expressing MTLn3 cells were treated for 24 hours with doxycycline to express HA-FRNK and subsequently exposed to 10 μ M doxorubicin. The percentage of apoptosis was determined by cell cycle

MOL # 26195

analysis (A). HA-FRNK expressing and control cells were trypsinized, kept in suspension in the absence of serum for 1 hour and plated on collagen coated plastic culture dishes. At the indicated time-points attached cells were immunoblotted for P-Ser⁴⁷³-PKB (P-PKB), PKB and FAK. Note the conditional expression of HA-FRNK after doxycycline pretreatment (B). HA-FRNK expressing and control cells were exposed to 10 μ M doxorubicin and after different periods of recovery, cell lysates were immunoblotted for P-PKB, PKB, HA and tubulin as a loading control (C). Data shown are representative for three independent experiments or the mean of three independent experiments \pm SE.

Figure 6. Inhibition of PI3K and overexpression of PTEN enhance doxorubicin-induced apoptosis. MTLn3 cells were exposed to 10 μ M doxorubicin for one hour and allowed to recover in the absence or presence of 10 nM wortmannin. After 8 hours cells were immunoblotted for PKB and P-PKB (A, upper band indicated by arrow) and after 16 hours the percentage apoptosis was determined by cell cycle analysis (B). MTLn3 cells were transiently transfected with eGFP, GFP-PTEN or GFP-PTEN-C124A and 24 hours after transfection cell lysates were immunoblotted for PTEN (C, left panel) and eGFP (C, right panel). Transiently transfected cells were exposed to 10 μ M doxorubicin and allowed to recover for 16 hours. The percentage apoptosis in the eGFP positive cells was determined by staining with annexin V/propidium iodide followed by flow cytometric analysis (D). Data shown are representative for three independent experiments (A, C) or the mean of three independent experiments \pm SE (B, D).

MOL # 26195

Figure 7. Working model for doxorubicin-induced apoptotic signaling pathways in MTLn3 adenocarcinoma cells. An early response in doxorubicin-induced cell stress is increased focal adhesion and F-actin stress fiber formation in association with the activation of the PI-3 kinase pathway. Inhibition of FAK (by FRNK) or inhibition of PI-3 kinase (by PTEN or wortmannin) inhibits the activation of PKB and increases the apoptotic cell death. At a later stage doxorubicin causes loss of FAs in association with dephosphorylation of PY³⁹⁷FAK. These events occur in cooperation with classical doxorubicin-induced DNA damage responses. Apoptosis can be abrogated at the level of the mitochondria (Bcl-2) and at the level of caspase activation (zVAD-fmk) without affecting upstream reorganization of FAs, FAK phosphorylation events and actin reorganization.

MOL # 26195


Table 1: Expression of eGFP-FRNK enhances doxorubicin induced apoptosis.

MTLn3 cells were transiently transfected with eGFP, eGFP-FAK, eGFP-FAT and eGFP-FRNK and subsequently exposed to 10 μ M doxorubicin. 16 hours after exposure of the cells, the percentage of apoptotic cells was determined by annexin V/ propidium iodide staining followed by flow cytometric analysis.


	Control (% apoptosis)	Doxorubicin (% apoptosis)
eGFP	9.6 (\pm 2.7)	16.2 (\pm 1.6)
eGFP-FAK	10.1 (\pm 2.9)	18.6 (\pm 3.2)
eGFP-FAT	14 (\pm 5.4)	21 (\pm 3.3)
eGFP-FRNK	18.2 (\pm 6.6)	26.7 (\pm 4.1)

Figure 1ABC

A



B



C





Figure 1DE

D



E




Figure 2





Figure 3











Figure 5**A****B****C**

Figure 6**A****B****C****D**