

MOL 26658

**Role of ectodomain lysines in the subunits of the heteromeric P2X<sub>2/3</sub>  
receptor**

William J. Wilkinson, Lin-Hua Jiang<sup>1</sup>, Annmarie Surprenant, and R. Alan  
North

Faculty of Life Sciences, University of Manchester, Michael Smith Building,  
Oxford Road, Manchester M13 9PT, U.K. (W.J.W., R.A.N.) and Department of  
Biomedical Science, University of Sheffield, Sheffield S10 2TN, U.K. (L.-H.J.,  
A.S.)

MOL 26658

Running Title: Ectodomain lysines in heteromeric P2X<sub>2/3</sub> receptors

Correspondence: R. Alan North, Faculty of Life Sciences, University of  
Manchester, Michael Smith Building, Oxford Road, Manchester M13 9PT, U.K  
Tel: 44-161-275-1499, Fax: 44-161-275-1498, e-mail:  
r.a.north@manchester.ac.uk

Text pages: 20

Tables: 0

Figures: 4

References: 31

Words in Abstract 221

Words in Introduction 501

Words in Discussion 1092

ABBREVIATIONS:  $\alpha\beta$ meATP,  $\alpha\beta$ -methylene-ATP

MOL 26658

## Abstract

Lysine residues near each end of the receptor ectodomain (in rat P2X<sub>2</sub> Lys<sup>69</sup> and Lys<sup>308</sup>) have been implicated in ATP binding to P2X receptors. We recorded membrane currents from human embryonic kidney cells expressing P2X subunits, and found that lysine to alanine substitutions at equivalent positions in the P2X<sub>3</sub> receptor (Lys<sup>63</sup> and Lys<sup>299</sup>) also prevented channel function. Heteromeric P2X<sub>2/3</sub> receptors are formed when P2X<sub>2</sub> and P2X<sub>3</sub> subunits are expressed together; they can be distinguished by their relatively sustained response to  $\alpha\beta$ -methylene-ATP. By co-expression of wild type P2X<sub>3</sub> and mutated P2X<sub>2</sub> subunit, we found that the heteromeric P2X<sub>2/3</sub> channel functioned normally when either lysine in the P2X<sub>2</sub> subunit was mutated to alanine (*i.e.* [K69A] or [K308A]), but not when both lysines were mutated to alanine (*i.e.* [K69A, K308A]). However, co-expression of wild type P2X<sub>2</sub> with a mutated P2X<sub>3</sub> subunit ([K68A] or [K299A]) produced no functional heteromers. The rescue of the single lysine mutant P2X<sub>2</sub> subunit by wild type P2X<sub>3</sub> (but not the converse) suggests that the heteromeric channel contains one P2X<sub>2</sub> and two P2X<sub>3</sub> subunits, and that the receptor functions essentially normally so long as two subunits are not mutated. The failure to rescue function in the P2X<sub>2</sub> subunit with both lysines mutated by wild type P2X<sub>3</sub> suggests that these residues from two different subunits interact in agonist binding or channel opening.

MOL 26658

The heteromeric P2X<sub>2/3</sub> receptor is of interest because of its predominant expression on a subset of primary afferent nerves involved in the sensation of chronic noxious damage (Jarvis, 2003), visceral distension (Vlaskovska et al., 2001), hypoxia (Gourine, 2005) and taste (Finger et al., 2005). A key role for these receptor subunits is supported by experiments using P2X<sub>3</sub> gene knock-outs (Cockayne et al., 2000; 2005), P2X<sub>3</sub> RNA suppression (Barclay et al., 2002; Honoré et al., 2002), and pharmacological antagonists selective for P2X<sub>3</sub> subunit-containing receptors (Jarvis et al., 2002).

There is good evidence that P2X subunits form channels as trimers (Nicke et al., 1998; North, 2002; Jiang et al., 2003; Barrera et al., 2005). Like other ligand-gated channels, P2X receptors can form by homo- or hetero-oligomeric assembly of subunits (for review see North, 2002). Although both P2X<sub>2</sub> and P2X<sub>3</sub> subunits are able to form homomeric channels, the heteromeric P2X<sub>2/3</sub> receptor that is also formed can be distinguished by its unique functional properties (Lewis et al., 1996). P2X<sub>2</sub> receptors are activated by ATP but not by the analog  $\alpha\beta$ -methylene-ATP ( $\alpha\beta$ meATP), and the current is largely sustained through a 2 s agonist application. P2X<sub>3</sub> receptors are activated by both ATP and  $\alpha\beta$ meATP and the current desensitizes very rapidly (<100 ms). The current through heteromeric P2X<sub>2/3</sub> channels can be identified as the sustained component when  $\alpha\beta$ meATP is the agonist (Lewis et al., 1996).

The molecular operation of P2X subunits is not well understood. Each receptor subunit has intracellular amino- and carboxy- termini, and two transmembrane domains (TM1: approximately residues 30 to 50; TM2:

MOL 26658

approximately residues 330-354) that are joined by an ectodomain of about 280 residues. The ectodomain is glycosylated and disulfide-bonded. There are no obvious 'canonical' ATP-binding amino acid sequences in the ectodomain. In the case of the P2X<sub>2</sub> homomeric receptor, two conserved ectodomain lysine residues have been suggested to play a role in binding the negatively charged phosphate moiety of ATP (Lys<sup>69</sup> and Lys<sup>308</sup> in rat P2X receptor), with the evidence being most complete for Lys<sup>69</sup> (Jiang et al., 2000). A similar role for the corresponding lysines has also been identified in the case of the P2X<sub>1</sub> receptor (Lys<sup>68</sup> and Lys<sup>309</sup>; Ennion et al., 2000). These lysines are situated about 20 amino acids from the outer end of each TM. Experiments on the P2X<sub>1</sub> receptor using partial agonists also suggest a role for aromatic residues Phe<sup>185</sup> and Phe<sup>291</sup> to coordinate the binding of the adenine ring of ATP (Roberts and Evans, 2004; Vial et al., 2004). An alternative model of ATP binding has been proposed by Yan et al., (2005) in the case of the P2X<sub>4</sub> receptor. This is based on secondary structure homology between part of the receptor (Lys<sup>180</sup> to Lys<sup>326</sup>) and class II aminoacyl-tRNA synthetases (Freist et al., 1998).

The present experiments focus on the two juxtamembrane lysines proposed to take part in ATP binding. We have found that for the heteromeric P2X<sub>2/3</sub> channel their importance in channel function depends greatly on whether they are in the P2X<sub>2</sub> or the P2X<sub>3</sub> subunit.

MOL 26658

## Materials and Methods

**Molecular and cell biology** Rat P2X receptor cDNAs were used as previously described (Jiang et al., 2000). Mutations were introduced using Quickchange site-directed mutagenesis (Stratagene, La Jolla CA) and confirmed by sequencing. Wild type and mutated receptors were transiently expressed in human embryonic kidney (HEK) 293 cells using lipofectamine 2000 (Invitrogen, San Diego CA) according to the manufacturer's instructions. Transfected cells were harvested and re-plated on glass coverslips 24 - 48 h after transfection and kept at 37° C for 18 - 48 h prior to electrophysiological recording. P2X<sub>2</sub> plasmid (0.2 µg) was co-transfected with P2X<sub>3</sub> plasmid (1 µg), these ratios having been established to produce equal amounts of protein when detected by Western blot with a common C-terminal epitope tag (Jiang et al., 2003). Enhanced green fluorescent protein cDNA (0.1 µg) was also included. Immunocytochemistry of transfected cells was as described, using a C terminus EYMPME epitope (Jiang et al., 2000, 2003).

**Electrophysiology** Whole-cell patch clamp recordings were made at 19 - 22 °C using a HEKA EPC9 amplifier (Heka, Lambrecht, Germany). The holding potential was -60 mV. Patch pipettes (3 - 6 MΩ) contained (mM): 145 NaCl, 10 HEPES and 10 EGTA (pH 7.3 with NaOH). The external solution contained (mM): 147 NaCl, 2 KCl, 2 CaCl<sub>2</sub>, 2 MgCl<sub>2</sub>, 10 HEPES and 13 glucose (pH 7.3 with NaOH). Both internal and external solutions were 290 - 310 mOsm/l. Agonists were applied using the RSC 200 rapid solution changer (Biologic Science Instruments, Grenoble, France), with flow-pipe tips approximately 150 µm from cell. All chemicals were purchased from Sigma (Poole, UK), and were made up daily from aliquots stored at -20° C.

MOL 26658

MOL 26658

## Results

**Homomeric channels** Previous work on homomeric human P2X<sub>1</sub> (Ennion et al., 2000) and rat P2X<sub>2</sub> (Jiang et al., 2000) receptors has indicated that Lys<sup>69</sup> and Lys<sup>308</sup> (rat P2X<sub>2</sub> numbering) are essential for channel function. We found that alanine substitution at the equivalent positions in the P2X<sub>3</sub> subunit also led to a non-functional homomeric channel. We observed no current with applications of  $\alpha\beta$ meATP (up to 1 mM;  $n = 5$ ) (Fig. 1) or ATP (up to 1 mM;  $n = 7$ ). Immunocytochemistry of transfected cells showed no difference in the pattern or distribution between wild type and mutated subunits. Fig. 1 also shows that equivalent mutations in other rat P2X receptors also prevented any response to agonist (P2X<sub>1</sub>[K68A]: ATP (300  $\mu$ M)  $23 \pm 9$  pA,  $n = 8$ ;  $\alpha\beta$ meATP (300 mM) no current,  $n = 3$ ; P2X<sub>4</sub>[K67A]: ATP (1 mM) no current,  $n = 8$ ; P2X<sub>7</sub>[K64A]: ATP (3 mM) no current,  $n = 3$ ).

Neither P2X<sub>2</sub>[K69A] nor P2X<sub>2</sub>[K308A] gave any detectable current in response to 1 or even 3 mM ATP (Fig. 1). However, co-expression of these two mutated receptors clearly did (Fig. 2). The currents were small compared to wild type P2X<sub>2</sub> currents (>600 pA/pF), but not obviously different in other properties (Fig. 2). It was striking that co-expression of two 'dead' subunits resulted in formation of a channel that is responsive to 100 or 300  $\mu$ M ATP. If we assume that neither mutation affects expression or assembly, then one would expect three-eighths of all trimeric channels to be [K69A]<sub>2</sub>[K308A] and three-eighths to be [K69A].[K308A]<sub>2</sub>. This result therefore indicates that the effects of these two point mutations are not independent.

**Heteromeric channels** For the P2X<sub>2/3</sub> receptors there was a striking difference in the effect of a single point mutation of either of the lysine



MOL 26658

residues, depending whether it was present in the P2X<sub>2</sub> or P2X<sub>3</sub> subunit. The currents in cells expressing mutant P2X<sub>2</sub> receptors with wild type P2X<sub>3</sub> subunits were essentially the same as those observed in wild type heteromeric receptors (Fig. 3). The  $\alpha\beta$ meATP concentrations giving half-maximal currents (EC<sub>50</sub>) were: for P2X<sub>2/3</sub> wild type, 27 ± 3.3  $\mu$ M ( $n = 5$ ); for P2X<sub>2</sub>[K69A] + P2X<sub>3</sub>, 22 ± 2.6  $\mu$ M ( $n = 5$ ), and for P2X<sub>2</sub>[K308A] + P2X<sub>3</sub>, 44 ± 2.6  $\mu$ M ( $n = 8$ )(the last two values are significantly different,  $p < 0.0001$ ). In contrast, cells expressing wild type P2X<sub>2</sub> with P2X<sub>3</sub>[K63A] subunits showed no currents in response to  $\alpha\beta$ meATP up to 100  $\mu$ M. At such high concentrations P2X<sub>2</sub> homomeric receptors are sensitive to  $\alpha\beta$ meATP ( $n = 5$ ) (Fig. 3)(also Jiang et al., 2003) and the small current observed was presumably passing through such receptors. Similarly, co-expression of wild type P2X<sub>2</sub> subunits with P2X<sub>3</sub> [K299A] gave no currents in response to  $\alpha\beta$ meATP (up to 100  $\mu$ M,  $n = 6$ ) (Fig. 3). These results indicate that a wild type P2X<sub>3</sub> subunit can rescue function when it enters a heteromer with a mutated P2X<sub>2</sub> subunit, but not *vice versa*.

This rescue did not occur when *both* ectodomain lysines were mutated in the same P2X<sub>2</sub> subunit. Co-expression of P2X<sub>2</sub> [K69A,K308A] with P2X<sub>3</sub> produced no sustained currents to  $\alpha\beta$ meATP ( $n = 9$ )(Fig. 3). However, a fast desensitizing current typical of homomeric P2X<sub>3</sub> receptors was usually observed (Fig. 3). A schematic summary of the expression of the further P2X<sub>2</sub> and P2X<sub>3</sub> subunits (either wild type or mutant) that were co-expressed, and the results of the functional studies, is shown in Fig. 4.

MOL 26658

## Discussion

The two lysines that are the focus of the present study are very highly conserved among P2X subunits. Lys<sup>69</sup> is present in 41 of 43 vertebrate sequences (including seven human, seven rat, seven mouse, one guinea pig, three chick, two xenopus, one bullfrog, nine zebrafish, six pufferfish); the zebrafish P2X<sub>2</sub> receptor has Asp at this position, and it can not be activated by ATP when heterologously expressed (Kucenas et al., 2003). This lysine is also present in the one invertebrate (schistosome) sequence that is known to operate as an ATP-gated channel (Agboh et al., 2004). Lys<sup>308</sup> is even more highly conserved, being found in all the above 44 sequences, and also in two of the five related *Dictyostelium* sequences. Of the eight positively charged residues in the P2X<sub>2</sub> receptor ectodomain, Lys<sup>69</sup> and Lys<sup>308</sup> are the two where replacement by alanine causes the greatest loss of channel function: even replacement by arginine results in more than 100-fold reduction in sensitivity to ATP (Jiang et al., 2000). The present experiments confirm that the lysine in the first of these positions is critical also for function of other P2X receptors, including P2X<sub>3</sub>.

The heterotrimeric P2X<sub>2/3</sub> receptor must contain either one or two copies of the P2X<sub>3</sub> receptor subunit. The simplest interpretation of the one way rescue of P2X<sub>2</sub> mutants by P2X<sub>3</sub> (but not *vice-versa*, Fig. 3), is that the P2X<sub>2/3</sub> receptors activated by  $\alpha\beta$ meATP have two copies of P2X<sub>3</sub> and one copy of P2X<sub>2</sub>. In other words, a functional channel can be formed and activated by  $\alpha\beta$ meATP even though it contains one 'dead' P2X<sub>2</sub> subunit (e.g. P2X<sub>2</sub>[K69A]) but no opening can be elicited from a receptor comprising two dead P2X<sub>3</sub> subunits with a wild type P2X<sub>2</sub> subunit. The rescue of wild type

MOL 26658

P2X<sub>3</sub> was complete in the case of P2X<sub>2</sub>[K69A], and slightly less so for P2X<sub>2</sub>[K308A] (Fig. 3), indicating that the two P2X<sub>2</sub> lysines do not have precisely equivalent roles.

This result contrasts with earlier findings in which a highly conserved intracellular C-terminal lysine was mutated (Chaumont et al. 2004). This lysine is required in homomeric P2X<sub>2</sub> and P2X<sub>3</sub> receptors for retention in the plasma membrane. In that case, wild type P2X<sub>3</sub> subunits could restore the appearance and retention in the membrane of mutant P2X<sub>2</sub> subunits (P2X<sub>2</sub>[K366A]), and this rescue was fully reciprocal because wild type P2X<sub>2</sub> also restored the appearance of P2X<sub>3</sub>[K357A] (Chaumont et al., 2004). Taken together, these results imply that only a single wild type subunit is needed for successful retention in the membrane, but two wild type (P2X<sub>3</sub>) subunits are needed for channel function. The conclusion that the functional heteromer contains two P2X<sub>3</sub> subunits and one P2X<sub>2</sub> subunit agrees well with the results of a completely independent approach in which subunits were joined by disulfide bonds between engineered cysteine substitutions at the outer ends of the TMs (Jiang et al., 2003). The finding is also consistent with the observation that the limiting slope of the Hill plot for activation of the heteromeric P2X<sub>2/3</sub> receptor is close to two, whereas for the homomeric P2X<sub>3</sub> and homomeric P2X<sub>2</sub> receptors it is close to three (Jiang et al. 2003). However, the present analysis can not distinguish between different numbers of ligand binding sites and differing degrees of intersubunit cooperativity in the heteromeric versus the homomeric channels.

For the homomeric P2X<sub>1</sub> and P2X<sub>2</sub> receptors, there is evidence that Lys<sup>69</sup> and Lys<sup>308</sup> contribute to an ATP binding site (Ennion et al., 2000; Jiang

MOL 26658

et al., 2000; Roberts and Evans, 2004). Therefore, our present finding that the heteromeric channel operates normally even though it incorporates a mutated P2X<sub>2</sub> subunit may be interpreted in this context. This would imply that a heterotrimeric P2X receptor can open from a less than fully liganded state. For example, the tetrameric AMPA receptor (Rosenmund et al., 1998) and the pentameric glycine receptor (Beato et al., 2002) can open from less than fully liganded states. But other interpretations are possible. For example, the P2X<sub>3</sub> (though not P2X<sub>2</sub>) subunit lysines may be critical for multimeric channel assembly. Or the mutation might impede the conformational change of gating: a lysine to alanine substitution in a single P2X<sub>2</sub> subunit of the trimer might provide insufficient energetic impediment, whereas two such independent substitutions in the P2X<sub>3</sub> subunits might prevent it completely.

We found surprising the observation that cells co-transfected with P2X<sub>2</sub>[K69A,K308A] subunit and wild type P2X<sub>3</sub> subunit produced no detectable heteromeric current (Fig. 3). In other words, the wild type P2X<sub>3</sub> subunits can rescue function in a channel containing a P2X subunit with a single mutation ([K69A] or [K308A])(a 'dead' P2X<sub>2</sub>) but not in a channel containing both those mutations (a 'double-dead' P2X<sub>2</sub>). This could have a simple energetic explanation of the type introduced above; namely, both P2X<sub>3</sub> subunits become liganded by  $\alpha\beta$ meATP but the mutations provide too great an energy barrier for opening to occur. Or it could indicate that the effects of the lysines are not independent, and that an interaction occurs between Lys<sup>69</sup> and Lys<sup>308</sup>. This would be consistent with the finding with homomeric P2X<sub>2</sub>

MOL 26658

receptors, that ATP can elicit currents when P2X<sub>2</sub>[K69A] and P2X<sub>2</sub>[K308A] are co-expressed, but not when either is expressed alone.

If one assumes that at least two agonist binding sites are required for channel activation (see above, and Jiang et al., 2003; Ding and Sachs, 1999), then these observations can be explained if both Lys<sup>69</sup> of one subunit and Lys<sup>308</sup> of another (or equivalent for the P2X<sub>3</sub> subunits) are needed at one ATP binding site. The presence of a 'dead' P2X<sub>2</sub> subunit would still provide for two  $\alpha\beta$ meATP binding sites (Fig. 4B, *left* and *middle*), but a 'double-dead' P2X<sub>2</sub> subunit would prevent  $\alpha\beta$ meATP binding at two sites and therefore prevent channel function (Fig. 4B, *right*). The interpretation that the agonist site is contributed from two different subunits in a heteromeric channel has parallels with nicotinic acetylcholine receptors (Sine, 2002). Moreover, P2X<sub>2</sub> subunits carrying mutations of an allosteric zinc binding site have been covalently joined in tandem from concatenated cDNAs: these experiments indicated that histidines from different subunits contribute to the zinc binding site (Nagaya et al., 2005).

The conclusion that the  $\alpha\beta$ meATP binding sites forms at the P2X<sub>3</sub>/P2X<sub>3</sub> and P2X<sub>2</sub>/P2X<sub>3</sub> interfaces is consistent with the observations that both nucleotide (Virginio et al. 1998; Burggaard et al., 2000) and non-nucleotide antagonists (Jarvis et al., 2002, 2004) mostly fail to discriminate between P2X<sub>3</sub> homomeric receptors and P2X<sub>2/3</sub> heteromeric receptors. A more complete understanding of the agonist binding site will inform the further development of small molecules that act as competitive antagonists at the P2X<sub>2/3</sub> heteromeric receptors, which are likely to have value as pain-relieving drugs.

MOL 26658

### **Acknowledgements**

The authors would like to thank Helen Broomhead, Louise Almond and Sandy Dossi for their technical assistance.

MOL 26658

## References

- Agboh KC, Webb TE, Evans RJ, Ennion SJ (2004) Functional characterization of a P2X receptor from *Schistosoma mansoni*. *J Biol Chem* **279**:41650-41657.
- Barclay J, Patel S, Dorn G, Wotherspoon G, Moffatt S, Eunson L, Abdel'al S, Natt F, Hall J, Winter J, Bevan S, Wishart W, Fox A, Ganju P (2002) Functional downregulation of P2X<sub>3</sub> receptor subunit in rat sensory neurons reveals a significant role in chronic neuropathic and inflammatory pain. *J Neurosci* **22**:8139-8147.
- Barrera NP, Ormond SJ, Henderson RM, Murrell-Lagnado RD, Edwardson JM (2005) Atomic force microscopy imaging demonstrates that P2X<sub>2</sub> receptors are trimers but that P2X<sub>6</sub> receptor subunits do not oligomerize. *J Biol Chem* **280**:10759-10765.
- Beato M, Groot-Kormelink PJ, Colquhoun D, Sivilotti LG (2002) Openings of the rat recombinant  $\alpha 1$  homomeric glycine receptor as a function of the number of agonist molecules bound. *J Gen Physiol* **119**:443-466.
- Burgard EC, Niforatos W, van Biesen T, Lynch KJ, Kage KL, Touma E, Kowaluk EA, Jarvis MF (2000) Competitive antagonism of recombinant P2X<sub>2/3</sub> receptors by 2', 3'-O-(2,4,6-trinitrophenyl) adenosine 5'-triphosphate (TNP-ATP). *Mol Pharmacol* **58**:1502-1510.
- Chaumont S, Jiang LH, Penna A, North RA, Rassendren F (2004) Identification of a trafficking motif involved in the stabilization and polarization of P2X receptors. *J Biol Chem* **279**:29628-29638.
- Cockayne DA, Hamilton SG, Zhu QM, Dunn PM, Zhong Y, Novakovic S, Malmberg AB, Cain G, Berson A, Kassotakis L, Hedley L, Lachnit WG,

MOL 26658

Burnstock G, McMahon SB, Ford AP (2000) Urinary bladder hyporeflexia and reduced pain-related behaviour in P2X<sub>3</sub>-deficient mice. *Nature* **407**:1011-1015.

Cockayne DA, Dunn PM, Zhong Y, Rong W, Hamilton SG, Knight GE, Ruan HZ, Ma B, Yip P, Nunn P, McMahon SB, Burnstock G, Ford AP (2005) P2X<sub>2</sub> knockout mice and P2X<sub>2</sub>/P2X<sub>3</sub> double knockout mice reveal a role for the P2X<sub>2</sub> receptor subunit in mediating multiple sensory effects of ATP. *J Physiol* **567**:621-639.

Ding S, Sachs F (1999) Single channel properties of P2X<sub>2</sub> purinoceptors. *J Gen Physiol* **113**:695-720.

Ennion S, Hagan S, Evans RJ (2000) The role of positively charged amino acids in ATP recognition by human P2X<sub>1</sub> receptors. *J Biol Chem* **275**:29361-29367.

Finger TE, Danilova V, Barrows J, Bartel DL, Vigers AJ, Stone L, Hellekant G, Kinnamon SC (2005) ATP signalling is crucial for communication from taste buds to gustatory nerves. *Science* **310**:1495-1499.

Freist W, Verhey JF, Stuhmer W, Gauss DH (1998) ATP binding site of P2X channel proteins: structural similarities with class II aminoacyl-tRNA synthetases. *FEBS Lett* **434**:61-65.

Gourine AV (2005) On the peripheral and central chemoreception and control of breathing: an emerging role of ATP. *J Physiol* **568**:715-724.

Honore P, Kage K, Mikusa J, Watt AT, Johnston JF, Wyatt JR, Faltynek CR, Jarvis MF, Lynch K (2002) Analgesic profile of intrathecal P2X<sub>3</sub> antisense oligonucleotide treatment in chronic inflammatory and neuropathic pain states in rats. *Pain* **99**:11-19.



MOL 26658

Jarvis MF et al. (2002) A-317491, a novel potent and selective non-nucleotide antagonist of P2X<sub>3</sub> and P2X<sub>2/3</sub> receptors, reduces chronic inflammatory and neuropathic pain in the rat. *Proc Natl Acad Sci U S A* **99**:17179-17184.

Jarvis MF (2003) Contributions of P2X<sub>3</sub> homomeric and heteromeric channels to acute and chronic pain. *Expert Opin Ther Targets* **7**:513-522.

Jarvis MF, Bianchi B, Uchic JT, Cartmell J, Lee CH, Williams M, Faltynek C (2004) [<sup>3</sup>H]A-317491, a novel high-affinity non-nucleotide antagonist that specifically labels human P2X<sub>2/3</sub> and P2X<sub>3</sub> receptors. *J Pharmacol Exp Ther* **310**:407-416.

Jiang L-H, Rassendren F, Surprenant A, North RA (2000) Identification of amino acid residues contributing to the ATP-binding site of a purinergic P2X receptor. *J Biol Chem* **275**:34190-34196.

Jiang L-H, Kim M, Spelta V, Bo X, Surprenant A, North RA (2003) Subunit arrangement in P2X receptors. *J Neurosci* **23**:8903-8910.

Kucenas S, Li Z, Cox JA, Egan TM, Voigt MM (2003) Molecular characterization of the zebrafish P2X receptor subunit gene family. *Neuroscience* **121**:935-945.

Lewis C, Neidhart S, Holy C, North RA, Buell G, Surprenant A (1995) Coexpression of P2X<sub>2</sub> and P2X<sub>3</sub> receptor subunits can account for ATP-gated currents in sensory neurons. *Nature* **377**:432-435.

Nagaya N, Tittle RK, Saar N, Dellal SS, Hume RI (2005) An intersubunit zinc binding site in rat P2X<sub>2</sub> receptors. *J Biol Chem* **280**:25982-25993.

Nicke A, Baumert HG, Rettinger J, Eichele A, Lambrecht G, Mutschler E, Schmalzing G (1998) P2X<sub>1</sub> and P2X<sub>3</sub> receptors form stable trimers: a

MOL 26658

novel structural motif of ligand-gated ion channels. *EMBO J* **17**:3016-3028.

North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* **82**:1013-1067.

Roberts JA, Evans RJ (2004) ATP binding at human P2X<sub>1</sub> receptors. Contribution of aromatic and basic amino acids revealed using mutagenesis and partial agonists. *J Biol Chem* **279**:9043-9055.

Rosenmund C, Stern-Bach Y, Stevens CF (1998) The tetrameric structure of a glutamate receptor channel. *Science* **280**:1596-1599.

Sine SM (2002) The nicotinic receptor ligand binding domain. *J Neurobiol* **53**:431-446.

Vial C, Roberts JA, Evans RJ (2004) Molecular properties of ATP-gated P2X receptor ion channels. *Trends Pharmacol Sci* **25**:487-493.

Virginio C, Robertson G, Surprenant A, North RA (1998) Trinitrophenyl-substituted nucleotides are potent antagonists selective for P2X<sub>1</sub>, P2X<sub>3</sub>, and heteromeric P2X<sub>2/3</sub> receptors. *Mol Pharmacol* **53**:969-973.

Vlaskovska M, Kasakov L, Rong W, Bodin P, Bardini M, Cockayne DA, Ford AP, Burnstock G (2001) P2X<sub>3</sub> knock-out mice reveal a major sensory role for urothelially released ATP. *J Neurosci* **21**:5670-5677.

Yan Z, Liang Z, Tomic M, Obsil T, Stojilkovic SS (2005) Molecular determinants of the agonist binding domain of a P2X receptor channel. *Mol Pharmacol* **67**:1078-1088.

MOL 26658

### **Footnotes**

The work was supported by the Wellcome Trust

<sup>1</sup>Present address: Faculty of Biological Sciences, University of Leeds, Leeds

LS2 9JT, U.K.

MOL 26658

**Fig. 1. A,** Mutation of conserved ectodomain lysines causes loss of function at rat P2X receptors. Pairs of traces show currents at wild type and lysine mutant receptors. Agonists were: for P2X<sub>1</sub>  $\alpha\beta$ meATP (300  $\mu$ M); for P2X<sub>4</sub> ATP (1 mM); and for P2X<sub>7</sub> BzATP (300  $\mu$ M); each applied for 2 s. **B,** Mutation of Lys<sup>69</sup> or Lys<sup>308</sup> at P2X<sub>2</sub> causes a loss of function (ATP, 1 mM), and equivalent mutations to Lys<sup>63</sup> and Lys<sup>299</sup> in P2X<sub>3</sub> have a similar effect ( $\alpha\beta$ meATP, 300  $\mu$ M).

**Fig. 2.** ATP elicits significant currents when lysine-mutant P2X<sub>2</sub> subunits are co-expressed ( $n = 9$ ). However, receptors P2X<sub>2</sub>[K69A] and P2X<sub>2</sub>[K308A] expressed separately do not respond to ATP at concentrations up to 3 mM ( $n = 12$ ). HEK, non-transfected cells ( $n = 3$ ). Insert shows a representative trace elicited by 1 mM ATP when lysine mutants are co-expressed, scale bars represent 50 pA and 1 s.

**Fig. 3** Co-expression of P2X<sub>2</sub> and P2X<sub>3</sub> subunits. **A,** currents elicited by 2 s application of 30  $\mu$ M  $\alpha\beta$ meATP. P2X<sub>2</sub>[K69A] and P2X<sub>2</sub>[K308A] can be rescued to form functional channels by expression with wild type P2X<sub>3</sub>. The corresponding mutations in the P2X<sub>3</sub> subunits are not rescued by wild type P2X<sub>2</sub> subunits. The double mutant P2X<sub>2</sub>[K69A,K308A] subunit does not function when co-expressed with wild type P2X<sub>3</sub>. However, the rapidly-desensitizing homomeric P2X<sub>3</sub> responses were typically observed on the first application of  $\alpha\beta$ meATP. **B,** concentration-response curves for some of the

MOL 26658

combinations shown in A. Responses are expressed as a % of the maximal response (mean  $\pm$  s.e. mean, 5 - 8 cells at each point).

**Fig. 4. A**, Schematic illustration of the P2X<sub>2</sub> and P2X<sub>3</sub> subunits used, and summary results. Small solid circles indicate positions of lysine to alanine substitution near outer end of TM1 or TM2. **B**, Schematic interpretation. Channels with two P2X<sub>3</sub> subunits and one P2X<sub>2</sub> subunit might provide two subunit contacts with two lysine residues (arrows, putative binding site), so long as only one of the P2X<sub>2</sub> lysines is substituted by alanine (*left, middle*) but would have only one such contact when P2X<sub>2</sub> subunit has both lysines changed to alanine.

Notes: <sup>1</sup>tested with ATP; <sup>2</sup>tested with  $\alpha\beta$ meATP; <sup>3</sup>normal wild type currents were  $522 \pm 74$  pA/pF ( $n = 9$ ; measured 200 ms after beginning superfusion); <sup>4</sup>dead indicates no detectable current ( $<2$  pA/pF); <sup>5</sup>ATP also evoked a sustained current, suggesting that the involvement of these residues is not specific to  $\alpha\beta$ meATP.

Figure 1




Figure 2





Figure 3

**A**



**B**








Figure 4

**A**


*P2X<sub>2</sub>* homomers<sup>1</sup>



*P2X<sub>2/3</sub>* heteromers<sup>2</sup>



*P2X<sub>3</sub>* homomers<sup>2</sup>



**B**

*P2X<sub>2</sub>*[K69A]  
+ *P2X<sub>3</sub>*

*P2X<sub>2</sub>*[K308A]  
+ *P2X<sub>3</sub>*

*P2X<sub>2</sub>*[K69A,K308A]  
+ *P2X<sub>3</sub>*

