

TITLE PAGE

Altered G protein coupling in an mGluR6 point mutant associated with congenital stationary night blindness

Donald Beqollari, Matthew J. Betzenhauser, and Paul J. Kammermeier *

Department of Pharmacology and Physiology, University of Rochester School of Medicine and Dentistry, Rochester, New York 14642 (D. B., M. J. B., P. J. K.); Current address: Department of Physiology and Cellular Biophysics, Columbia University, New York, New York 10032 (M. J. B.).

RUNNING TITLE PAGE

G protein coupling in a disease causing mGluR6 point mutant.

*Corresponding author: Paul J. Kammermeier, 601 Elmwood Ave. Box # 711 Rochester,
New York 14642

Tel. number: 585-275-5606

Fax number: 585-273-2652

email: Paul_Kammermeier@urmc.rochester.edu

Text page number: 24

Table number: 0

Figure number: 5

Reference number: 39

Abstract word count: 137

Introduction word count: 536

Discussion word count: 830

Abbreviations:

BBS = Bungarotoxin Binding Site

BTX = Bungarotoxin

CSNB = Congenital Stationary Night Blindness

mGluR6 = Metabotropic Glutamate Receptor Type 6

SCG = Superior Cervical Ganglion

ABSTRACT

The highly specialized metabotropic glutamate receptor type 6 (mGluR6) is postsynaptically localized and expressed only in the dendrites of ON bipolar cells. Upon activation of mGluR6 by glutamate released from photoreceptors, a non-selective cation channel is inhibited causing these cells to hyperpolarize. Mutations in this gene have been implicated in the development of congenital stationary night blindness type 1 (CSNB1). We investigated 5 known mGluR6 point mutants that lead to CSNB1 to determine the molecular mechanism of each phenotype. In agreement with other studies, 4 mutants demonstrated trafficking impairment. However, mGluR6 E775K (E781K in humans) suggested no trafficking or signaling deficiencies measured by our initial assays. Most importantly, our results indicate a switch in G-Protein coupling, in which E775K loses G_o coupling, but retains coupling to G_i , which may explain the phenotype.

INTRODUCTION

The metabotropic glutamate receptors (mGluRs) are classified into three subfamilies based on their molecular identity, pharmacology, and G-Protein coupling profile (Schoepp, 2001). Group III mGluRs (4, 6, 7, 8) couple exclusively to $G_{i/o}$ proteins and negatively regulate Adenylyl Cyclase (AC) (Prezeau et al., 1994). mGluR6 is a highly specialized G-protein coupled receptor (GPCR) that is exclusively expressed in the post-synaptic dendritic region of retinal ON bipolar cells (Nakajima et al., 1993; Quraishi et al., 2007; Vardi et al., 2000). Upon activation, mGluR6 initiates a signaling cascade that ultimately results in the inhibition of a non-selective cation channel, leading to hyperpolarization of the ON bipolar cells (Masland, 2001; Nawy, 1999). Recent studies suggest that the molecular identity of the regulated cation channel is likely TRPM1 (Bellone et al., 2008; Nakajima et al., 2009; Shen et al., 2009), and a few reports suggest that regulation is mediated by $G\alpha_o$, rather than the $G\alpha_i$ subunits or by $G\beta\gamma$ (Dhingra et al., 2004; Dhingra et al., 2000; Nawy, 1999). Our lab and others have demonstrated strong coupling of mGluR6 to $G\alpha_o$ in reconstitution experiments (Tian and Kammermeier, 2006), and antibodies directed toward $G\alpha_o$ were able to block the effect of glutamate on the channel in retinal slice recordings (Nawy, 1999).

Genetic defects in the GRM6 gene (which encodes mGluR6) can lead to congenital stationary night blindness (CSNB), characterized by myopia and mild to severe impairment of night vision. In humans, mGluR6 signaling abnormalities, caused by autosomal recessive mutations, lead to complete loss of night vision, CSNB1 (Dryja et al., 2005; Zeitz et al., 2007; Zeitz et al., 2005), along with the inability to adapt to light-dark

or cycles and reduced responsiveness to light stimuli in animal models (Pinto et al., 2007; Takao et al., 2000). CSNB2, which is characterized by less severe phenotypes, resolves from mutations occurring elsewhere in the ON visual pathway (Chang et al., 2006; Zeitz et al., 2006). Genetic studies in humans with CSNB1 and deletion of the GMR6 gene in mouse models result in the absence of the electroretinogram (ERG) b-wave (Dhingra et al., 2002; Dhingra et al., 2000; Dryja et al., 2005; Pinto et al., 2007; Zeitz et al., 2005), which reflects the depolarization of ON bipolar cells after exposure of photoreceptors to light. Interestingly, the absence of the b-wave was also observed in mice lacking $G\alpha_o$ in bipolar cells (Dhingra et al., 2002; Dhingra et al., 2000), which correlates well with the hypothesis that mGluR6 strongly couples to $G\alpha_o$ to facilitate the modulation of the cation channel.

To date, there are at least five mGluR6 point mutations that lead to CSNB in humans (P46L, G58R, G150S, C522Y, and E781K). Recent work suggests that each of these mutants exhibit impaired plasma membrane expression in HEK293 cells (Zeitz et al., 2005). Based on the location of these mutations, we anticipated that these mutants would have impaired glutamate responses due to decreased affinity for glutamate or compromised G-protein coupling. Our results confirm that four of these molecules fail to express on the plasma membrane in neurons. The mGluR6 E775K (E781K in humans) expressed and functioned normally in our preliminary assays. However, calcium mobilization data suggest that E775K couples predominantly to $G\alpha_i$, rather than $G\alpha_o$, which may explain the CSNB phenotype in humans.

MATERIALS and METHODS

SCG neuron isolation, cDNA injection and molecular methods. Detailed descriptions of the isolation and injection procedures have been previously described (Ikeda, 1997). Briefly, the SCG were dissected from adult rats and incubated in Earle's balanced salt solution (Life Technologies, Rochelle, MD) containing 0.6 mg/ml trypsin (Worthington, Freehold, NJ), 0.8 mg/ml collagenase D (Boehringer Mannheim, Indianapolis, IN) for 1 hour at 35° C. Cells were centrifuged twice, transferred to minimum essential medium (Fisher Scientific, Pittsburgh, PA), plated and placed in an incubator at 37° C until cDNA injection. Injection of cDNA was performed with an Eppendorf 5247 microinjector and Injectman NI2 micromanipulator (Madison, WI) 4-6 hours following cell isolation. Plasmids were stored at -20 °C as a 1 µg/µl stock solution in TE buffer (10 mM TRIS, 1 mM EDTA, pH 8). mGluR6 insert was subcloned into pCDNA3.1- (Invitrogen). All receptor cDNAs were injected at 0.1 µg/µl. Neurons were co-injected with "enhanced" green fluorescent protein cDNA (0.005 µg/µl; pEGFPN1; Clontech Laboratories, Palo Alto, CA, USA) or other fluorescent marker if necessary for identification of injected cells. Point mutants (P40L, G52R, G144S, C516Y, and E775K) were generated using the QuickChange Site-Directed Mutagenesis Kit (Stratagene, cat# 200518) and sequence verified. BBS-mGluR6 was constructed by sub-cloning the bungarotoxin binding site sequence (TGGAGATACTACGAGAGCTCCCTGGAGCCCTACCCTGAC) between 22nd-23rd residues of the original mGluR6 clone using the overlapping extension PCR method. G_q chimeras were also constructed using the QuickChange strategy. Following injection, cells were incubated overnight at 37° C and experiments are performed the following day.

Electrophysiology and data analysis. Patch clamp recordings were made using 8250 glass (Garner Glass, Claremont, CA). Pipette resistances were 1-3 M Ω yielding uncompensated series resistances of 1-5 M Ω . Series resistance compensation of 70-80% was used in all recordings. Data was recorded using an Axopatch 200B patch-clamp amplifier from Molecular Devices (Foster City, CA). Voltage protocol generation and data acquisition were performed using custom data acquisition software (generously donated by Stephen R. Ikeda, NIAAA) on a Macintosh G3 computer with an Instrutech ITC18 data acquisition board (HEKA Elektronik, Germany). Currents were sampled at 0.5-5 kHz low-pass filtered at 3-5 kHz using the 4-pole Bessel filter in the patch clamp amplifier, digitized, and stored on the computer for later analysis. All patch-clamp experiments were performed at 21-24 °C (room temperature). Data analysis was performed using Igor software (WaveMetrics, Lake Oswego, OR). For calcium current recordings in SCG, the external (bath) solution contained (in mM): 145 tetraethylammonium (TEA) methanesulfonate (MS), 10 4-(2-Hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), 15 glucose, 10 CaCl₂, and 300 nM tetrodotoxin, pH 7.4, osmolality 320 mOsm/kg. The internal (pipette) solution contains: 120 N-methyl-Dglucamine (NMG) MS, 20 TEA, 11 EGTA, 10 HEPES, 10 sucrose, 1 CaCl₂, 4 MgATP, 0.3 Na₂GTP, and 14 tris-creatine phosphate, pH 7.2, osmolality 300 mOsm/kg. Pertussis toxin (PTX) (List Biological Lab. Inc. Campbell, CA) was applied at 500 ng/ml media overnight at 37° C.

PTX-insensitive reconstitution recordings. A good stoichiometric balance between the α and $\beta\gamma$ subunits was necessary to perform the reconstitution experiments. As previously described, the triple pulse protocol that produces the “pre” and the “post” currents is an effective way to detect an imbalance (Kammermeier et al., 2003). Excess

of G α subunits will result in a “pre” current that is slightly higher than the “post” current due to α -GDP serving as a “sponge” for G $\beta\gamma$. By contrast, surplus of G $\beta\gamma$ will result in a “pre” current that is already inhibited (Herlitz et al., 1996; Ikeda, 1996; Ikeda and Dunlap, 1999). To overcome this obstacle, we only analyzed cells that had a post/pre ratio between 1-1.3. Ratios between these values are indicative a good stoichiometric G-protein subunit balance (Kammermeier et al., 2003). The α subunit plasmids were injected at 5-6 ng/ μ l and G β_1 and G γ_2 plasmids were injected at 8-10 ng/ μ l final concentration.

Confocal Microscopy. Injected SCG neurons (in glass coverslips) with the desired plasmid along with “enhanced” green fluorescent protein cDNA to confirm expression were incubated overnight at 37° C. 16-18 hours post injection cells were transferred into a perfusion chamber and washed with PBS, followed by treatment with 1 μ M α -bungarotoxin AlexaFluor 647 conjugate (Molecular Probes, Eugene, OR) for 1 minute. Cells were washed again with PBS, and imaging was performed using an inverted Nikon microscope through a 40X oil immersion objective lens. Confocal images were obtained by utilizing the EZ-C1 3.60 software program (Nikon, Melville, NY). Cells were excited at 488 and 633 nm and signal was detected at 530 \pm 30 and 688 \pm 20 nm for GFP and AlexaFluor 647 respectively.

Digital imaging of intracellular Ca²⁺ in muscarinic type 3 (m3) stable HEK293 cells. M3 HEK293 cells, obtained from Dr. Trevor Shuttleworth, (University of Rochester, Rochester, NY) were loaded with 2 mM of the Ca²⁺-sensitive dye fura-2 AM at 37° C for 15-20 minutes. Transfected cells were loaded with Fura-2 AM then transferred into a perfusion chamber. Cells were perfused in HEPES-buffered physiological saline con-

taining (in mM) 137 NaCl, 0.56 MgCl₂, 4.7 KCl, 1 Na₂HPO₄, 10 HEPES, 5.5 glucose, and 1.26 CaCl₂, pH 7.4. Imaging was performed using an inverted Nikon microscope through a 40X oil immersion objective lens (numerical aperture, 1.3). Fura-2 AM loaded cells were excited alternately with light at 340 and 380 nm by using a monochromator-based illumination system (TILL Photonics, Pleasanton, CA), and the emission at 510 nm was captured by using a digital frame transfer CCD camera. Images were captured every 2 s with an exposure of 40 ms and 4 by 4 binning. Analysis was performed by TILL Vision software. Any signal below 0.1 ratio was designated as background noise. The receptors and the Gq chimeras were transfected at 1 µg/35 mm dish, β₁ and γ₂ plasmids were transfected at 0.5 µg/35 mm dish, and finally dSRed-nuc was transfected at 0.2-0.3 µg/35 mm dish (for the supplementary experiment), and cherry for the Ca²⁺ mobilization experiments.

RESULTS

Modulation N-type (Ca_{v2.2}) calcium channels in superior cervical ganglion neurons (SCGs) by mGluR6 and five CSNB point mutants. Our initial hypothesis regarding the CSNB point mutants evolved from the idea that these receptors have a reduced or diminished ability to respond to glutamate. Thus, we examined modulation of Ca_{v2.2} channels in SCG neurons by each point mutant to examine receptor activity. The triple pulse voltage protocol, shown in Fig. 1A (inset), was used to assess channel activity and consists of two test pulses to +10 mV from a -80 mV holding potential, separated by a strong depolarizing step to +80 mV. The +80 mV step partially reverses the well

characterized voltage-dependent, G β γ mediated inhibition (Elmslie et al., 1990; Ikeda, 1991; Ikeda, 1996).

To determine the effects of five CSNB point mutations on mGluR6 signaling, the corresponding mutants were made in the rat mGluR6 sequence and each was expressed in SCG neurons. Fig.1A, shows a typical time-course of an SCG neuron expressing mGluR6, illustrating inhibition of calcium currents during glutamate application. No effect was observed in uninjected or GFP only injected cells (data not shown).

In Fig. 1B, the glutamate concentration-response curve for the wild-type and each point mutant is shown. Four of the mutants, P40L, G52R, G144S, and C516Y showed no response to glutamate. By contrast, cells expressing mGluR6 E775K showed responses similar to those of the wild-type mGluR6. We noticed a slight right shift in the EC₅₀ values of mGluR6 E775K concentration-response curve (EC₅₀ for mGluR6 was 8 μ M and 20 μ M GLU for the mutant). However, at 100 μ M glutamate applications the responses were identical (45 \pm 5% and 44 \pm 2% current inhibition for mGluR6 and E775K respectively). These data suggest that four of the mGluR6 CSNB point mutants are non-functional. By contrast, mGluR6 E775K appears to function similarly to the wild-type.

Four CSNB point mutants lack plasma membrane expression. To determine why the non-functional mutant receptors lacked responses, the plasma membrane expression of each was examined. Zeitz et al. (2007), reported that all of the CSNB point mutants are retained in the ER when overexpressed in HEK293 cells. To evaluate plasma membrane expression in SCG neurons, wild-type mGluR6 and each CSNB mu-

tant were tagged with a N-terminal, extracellular bungarotoxin (BTX) binding site (BBS) (Sekine-Aizawa and Huganir, 2004). The plasma membrane expression of each receptor was then examined by application of a fluorescent tagged α -bungarotoxin to SCG neurons expressing each receptor. Fig. 2 illustrates representative SCG injected neurons with the indicated plasmid along with pEGFP. Cells expressing GFP alone showed no membrane fluorescence after treatment with α -bungarotoxin-AlexaFluor 647 (shown in red). Further, no detectable membrane labeling was evident in cells expressing P40L, G52R, G144S, or C516Y. However, cells expressing the wild-type or mGluR6 E775K mutant demonstrated comparable surface membrane labeling (Fig. 2). To verify that the BTX binding site tag did not interfere with receptor function, electrophysiological experiments were performed by monitoring calcium channel modulation by 100 μ M glutamate in cells expressing mGluR6 or BBS-mGluR6. Similar responses were observed in SCG neurons expressing each receptor (Fig. 2). Calcium currents were inhibited $43 \pm 4\%$ ($n=14$) and $48 \pm 5\%$ ($n=8$) in cells expressing mGluR6 and BBS-mGluR6, respectively.

E775K does not activate $G\alpha_o$. Although most of our labeling data correlated with the previous report (Zeitze et al., 2007), one CSNB mutant, mGluR6 E775K, was able to traffic to the plasma membrane and function similarly to the wild-type mGluR6 in SCG neurons. Because wild-type mGluR6 appears to act in the retina through $G\alpha_{oA}$ stimulation (Dhingra et al., 2002) it is possible that the E775K mutant may produce CSNB due to an inability to activate $G\alpha_{oA}$, rather than lack of expression or due to trafficking defects. Because the E775K residue resides in the third intracellular loop (i3), a change in

the G-protein coupling seemed possible (Francesconi and Duvoisin, 1998). Therefore, the G-protein coupling of mGluR6 E775K was examined more closely.

As with wild-type mGluR6, E775K responses were abolished by overnight pertussis toxin (PTX) treatment, indicating that the mutant coupled exclusively to $G_{i/o}$ proteins (see Fig. 3). Calcium currents were inhibited $45 \pm 4\%$ ($n=6$) by 100 μ M GLU and $6 \pm 1\%$ ($n=5$) with overnight PTX treatment when mGluR6 was expressed in SCG neurons. Similar responses were obtained ($42 \pm 3\%$, $n=11$ and $3 \pm 1\%$, $n=7$ when mGluR6 E775K was expressed).

Next, the possibility that the E775K mutation changes from a predominantly G_o -coupled to a predominantly G_i -coupled receptor was tested. Therefore, coupling of mGluR6 E775K to specific G-proteins was examined by reconstitution using PTX-insensitive $G_{i/o}$ protein mutations of the cysteine residue, near the end of C-terminus (351/2 position, the site for ADP ribosylation) to glycine, rendering the G-proteins insensitive to PTX, as described previously (Bahia et al., 1998).

SCG neurons expressing either mGluR6 wild-type or E775K were treated with PTX to inactivate endogenously expressed $G_{\alpha_{i/o}}$ proteins and calcium current modulation was examined when $G\beta_1$ and $G\gamma_2$ were co-expressed along with each $G_{\alpha_{i/o}}$ CG PTX-insensitive protein (see Fig. 3).

Consistent with previous results (Tian and Kammermeier, 2006), reconstitution with $G_{\alpha_{oA}}$ CG resulted in a strong coupling to mGluR6 wild-type ($32 \pm 1\%$ inhibition $n=7$). However, no coupling was detected with E775K mutant ($2 \pm 1\%$ inhibition $n=6$). Surprisingly, no coupling was observed with G_{i1} CG, G_{i2} CG, or G_{i3} CG either (3 ± 2 , 3 ± 2 , and $4 \pm 2\%$ inhibition respectively) (Fig. 4), despite a strong PTX-sensitive signal when

mGluR6 E775K was expressed alone. Thus, although the E-K mutation appeared to abolish signaling through G_{α_A} , it is possible that the reconstitution assay is not sufficiently sensitive, or that the C-G mutation may selectively disrupt coupling to the mutant receptor. Therefore, an alternative approach to investigate mGluR6 E775K signaling was necessary.

E775K couples predominantly to G_i proteins. As an alternative strategy to investigate mGluR6 E775K signaling, Ca^{2+} mobilization was examined in m3 (muscarinic type 3) HEK293 stable cell lines expressing $G_qG_{i/o}$ chimeric proteins in which the extreme C-terminus amino acids has been exchanged to allow activation by mGluR6 (Blahos et al., 1998). G_q chimeras have been widely used to study $G_{i/o}$ coupled GPCR signaling (Kowal et al., 2003; Kowal et al., 2002; Walker et al., 2005; Zhang et al., 2003).

To this end, $G_qG_{\alpha_A(9)}$ and $G_qG_{i1/2(9)}$ chimeras were generated (the last 9 residues for G_{i1} and G_{i2} are identical), and used to perform Ca^{2+} fura-2 AM imaging experiments in m3 HEK293 stable cell lines. M3 receptors couple well to native G_q , which would serve as an internal control when 300 nM carbachol was applied. These cells, were heterologously transfected with each receptor (mGluR6 or E775K), the G_q chimera, and $G\beta_1\gamma_2$ constructs, along with cherry (a red fluorescent protein) to identify expressing cells.

Fig.5A shows representative Ca^{2+} traces illustrating Ca^{2+} signals when each agonist was applied. In cells transfected with the receptor alone no signal was detected upon glutamate application (mGluR6 151 cells analyzed; E775K 149 cells). These results suggested that neither mGluR6 nor mGluR6 E775K could induce an intracellular Ca^{2+} signal via natively expressing G-proteins. When the wild-type receptor was coex-

pressed with $G_qG_{\alpha A(9)}$ and $G\beta_1\gamma_2$, a strong Ca^{2+} response was detected in 31 of 208 cells analyzed. By contrast only 1 of 82 cells expressing mGluR6 E775K produced a detectable response, and this response was weak (Fig. 5 A,B). These results correlated well with the electrophysiological data (Fig. 3). When $G_qG_{i1/2(9)}$ was expressed some signaling was detected with mGluR6 (3 out of 88 cells). However, this signal was weaker and less frequent than when $G_qG_{\alpha A(9)}$ was expressed. Interestingly, mGluR6 E775K was able to signal in response to glutamate in 15 out of 460 $G_qG_{i1/2(9)}$ expressing cells (Fig. 5A, B). These data confirm that mGluR6 couples strongly to $G\alpha_o$ and weakly to $G\alpha_i$, while demonstrating that E775K mutant receptor predominantly activates $G\alpha_i$. Further, since this mutation causes CSNB in humans, effectively resulting in loss of function, these data support the conclusion that mGluR6 in retinal ON bipolar cells produces its primary signal via $G\alpha_o$ activation, rather than $G\alpha_i$ or $G\beta\gamma$.

DISCUSSION

We report here that a disease causing mGluR6 mutant (E775K) functions and expresses similar to the wild-type receptor in SCG neurons. Further investigation of the signaling cascade revealed that mGluR6 E775K is incapable of activating $G\alpha_o$ and primarily couples to $G\alpha_i$ proteins. Overall, the data presented here suggest that mGluR6 primarily functions through $G\alpha_o$ proteins in retinal ON bipolar cells, and an inability of the receptor to activate $G\alpha_o$ results in loss of function, leading to CSNB.

Three of the CSNB mGluR6 point mutants examined here, P40L, G52R, and G144S are located in the Venus Fly-Trap Domain (VFD) near the agonist binding site (Bessis et al., 2000; Rosemond et al., 2002). The C516 residue is located in the Cys-

teine Rich Region (CRR) of the receptor, which separates the VFD from the 7TM region and is believed to be important in the intramolecular signal transmission (Rondard et al., 2006). Therefore, the mechanism of loss of function for each seemed apparent. Surprisingly, each of these mutants lacked function not due to a predicted deficit in ligand binding or intramolecular signal transduction, but due to loss of plasma membrane expression. These results are consistent with those of Zeitz et. al. in HEK293 cells, and support the notion that mGluR6 trafficking is tightly regulated such that even small structural perturbations can severely alter receptor trafficking.

The E775 residue is located in the third intracellular loop of the receptor. The second (i2) and the third (i3) of mGluRs participate in G-protein contact and activation. Interestingly the i3 loop of all mGluRs contains a conserved region, and single point mutations in this region have been known to impair receptor/G-protein coupling (Francesconi and Duvoisin, 1998).

According to the functional data presented here, E775K produced responses similar to the wild-type receptor at 100 μ M GLU with only a slight shift in potency (see Fig. 1). Membrane surface labeling indicated that the mutant receptor is present on the plasma membrane in SCGs following nuclear cDNA injection (Fig. 2), and in HEK293 cells when transfected (see supplementary Fig. 1). However, the observation that an mGluR6 point mutation that resulted in a nearly normal functioning receptor was puzzling.

Wild-type mGluR6 couples to G_o proteins in ON bipolar cells (Nawy, 1999) and in reconstitution studies in SCG neurons (Tian and Kammermeier, 2006). Further, the primary signal transduction induced by mGluR6 in retinal ON bipolar cells, the inhibition of

a non-selective cation conductance, presumably TRPM1 (Shen et al., 2009), appears to require activation of $G\alpha_o$, as disruption of neither $G\alpha_i$ nor $G\beta\gamma$ can occlude this effect (Dhingra et al., 2002; Nawy, 1999). Therefore, a change in G-protein coupling preference of mGluR6 from $G\alpha_o$ to $G\alpha_i$ could result in CSNB even if the mutant receptor is otherwise functional. This hypothesis was tested for mGluR6 E775K with two strategies. First, by reconstitution of signaling using PTX-insensitive mutant $G\alpha$ proteins in SCG neurons, and second by examining intracellular calcium signaling with fura-2AM in HEK293 cells expressing $G\alpha_qG\alpha_{i/o}$ chimeric proteins.

Unfortunately, reconstitution of signaling with the mutant receptor in SCG neurons with PTX-insensitive G-proteins could not produce a complete answer. The data suggested a loss of E775K coupling to $G_{oA}CG$. However, when the G_iCG mutants were utilized, no coupling was observed. These negative results may have resulted from a weak expression of the G-protein mutant in SCGs, low sensitivity of the assay (channel modulation), or the possibility that mGluR6 E775K coupled weakly to more than one G_i protein, but not robustly to any one in isolation.

To this end, a more sensitive assay, Ca^{2+} imaging experiments with fura-2AM, was employed. We constructed $G_qG\alpha_{i/o}$ chimeras in which the last 9 residues were swapped with that of the G_{oA} , G_{i1} , and G_{i2} (the last 9 residues of G_{i1} and G_{i2} are identical, so this construct was called $G_qG_{i1/2}$). Similar constructs have been used to study mGluR2 and mGluR4 signaling in CHO cells (Kowal et al., 2003). In accordance with our electrophysiological data, mGluR6 was able to strongly couple to G_qG_{oA} , with less robust coupling to $G_qG_{i1/2}$. However, mGluR6 E775K failed to couple to G_qG_{oA} , but elicited substantial Ca^{2+} signals when expressed with $G_qG_{i1/2}$. It is important to note that during

these experiments several plasmid constructs were required (receptor, chimera, $G\beta_1$, $G\gamma_2$, and cherry), so the probability of getting all the components in any one cell is low. Therefore, the low number of responding cells in each experiment was expected.

In conclusion, we demonstrate here an alternative explanation for the CSNB phenotype (for E781K mutant in humans) as has been previously proposed. These results indicate mGluR6 E775K functions and expresses like the wild-type receptor, but its ability to activate $G\alpha_{oA}$ has been diminished. The mutant receptor retains the capability to activate G_i proteins in heterologous and reconstitution assays, but that may not be sufficient to produce effective signaling in retinal ON bipolar cells.

ACKNOWLEDGMENTS

We thank D. I. Yule (University of Rochester) for use of equipment and reagents and Trevor Shuttleworth (University of Rochester) for m3 HEK293 cells.

REFERENCES

- Bahia DS, Wise A, Fanelli F, Lee M, Rees S and Milligan G (1998) Hydrophobicity of residue351 of the G protein Gi1 alpha determines the extent of activation by the alpha 2A-adrenoceptor. *Biochemistry* **37**(33):11555-11562.
- Bellone RR, Brooks SA, Sandmeyer L, Murphy BA, Forsyth G, Archer S, Bailey E and Grahn B (2008) Differential gene expression of TRPM1, the potential cause of congenital stationary night blindness and coat spotting patterns (LP) in the Appaloosa horse (*Equus caballus*). *Genetics* **179**(4):1861-1870.
- Bessis AS, Bertrand HO, Galvez T, De Colle C, Pin JP and Acher F (2000) Three-dimensional model of the extracellular domain of the type 4a metabotropic glutamate receptor: new insights into the activation process. *Protein Sci* **9**(11):2200-2209.
- Blahos J, 2nd, Mary S, Perroy J, de Colle C, Brabet I, Bockaert J and Pin JP (1998) Extreme C terminus of G protein alpha-subunits contains a site that discriminates between Gi-coupled metabotropic glutamate receptors. *J Biol Chem* **273**(40):25765-25769.
- Chang B, Heckenlively JR, Bayley PR, Brecha NC, Davisson MT, Hawes NL, Hirano AA, Hurd RE, Ikeda A, Johnson BA, McCall MA, Morgans CW, Nusinowitz S, Peachey NS, Rice DS, Vessey KA and Gregg RG (2006) The nob2 mouse, a null mutation in *Cacna1f*: anatomical and functional abnormalities in the outer retina and their consequences on ganglion cell visual responses. *Vis Neurosci* **23**(1):11-24.

- Dhingra A, Faurobert E, Dascal N, Sterling P and Vardi N (2004) A retinal-specific regulator of G-protein signaling interacts with Galpha(o) and accelerates an expressed metabotropic glutamate receptor 6 cascade. *J Neurosci* **24**(25):5684-5693.
- Dhingra A, Jiang M, Wang TL, Lyubarsky A, Savchenko A, Bar-Yehuda T, Sterling P, Birnbaumer L and Vardi N (2002) Light response of retinal ON bipolar cells requires a specific splice variant of Galpha(o). *J Neurosci* **22**(12):4878-4884.
- Dhingra A, Lyubarsky A, Jiang M, Pugh EN, Jr., Birnbaumer L, Sterling P and Vardi N (2000) The light response of ON bipolar neurons requires G[alpha]o. *J Neurosci* **20**(24):9053-9058.
- Dryja TP, McGee TL, Berson EL, Fishman GA, Sandberg MA, Alexander KR, Derlacki DJ and Rajagopalan AS (2005) Night blindness and abnormal cone electroretinogram ON responses in patients with mutations in the GRM6 gene encoding mGluR6. *Proc Natl Acad Sci U S A* **102**(13):4884-4889.
- Elmslie KS, Zhou W and Jones SW (1990) LHRH and GTP-g-S modify calcium current activation in bullfrog sympathetic neurons. *Neuron* **5**(1):75-80.
- Francesconi A and Duvoisin RM (1998) Role of the second and third intracellular loops of metabotropic glutamate receptors in mediating dual signal transduction activation. *J Biol Chem* **273**(10):5615-5624.
- Herlitze S, Garcia DE, Mackie K, Hille B, Scheuer T and Catterall W (1996) Modulation of Ca²⁺ channels by G-protein bg subunits. *Nature* **380**:258-262.
- Ikeda SR (1991) Double-pulse calcium channel current facilitation in adult rat sympathetic neurones. *J Physiol* **439**:181-214.

- Ikeda SR (1996) Voltage-dependent modulation of N-type calcium channels by G-protein $\beta\gamma$ subunits. *Nature* **380**:255-258.
- Ikeda SR (1997) Heterologous expression of receptors and signaling proteins in adult mammalian sympathetic neurons by microinjection., in *Methods in Molecular Biology* (Challis RAJ ed) pp 191-202, Humana Press, Inc., Totowa, NJ.
- Ikeda SR and Dunlap K (1999) Voltage-dependent modulation of N-type calcium channels: role of G protein subunits. *Adv in Sec Mess Phos Res* **33**:131-151.
- Kammermeier PJ, Davis MI and Ikeda SR (2003) Specificity of metabotropic glutamate receptor 2 coupling to G proteins. *Mol Pharmacol* **63**(1):183-191.
- Kowal D, Nawoschik S, Ochalski R and Dunlop J (2003) Functional calcium coupling with the human metabotropic glutamate receptor subtypes 2 and 4 by stable co-expression with a calcium pathway facilitating G-protein chimera in Chinese hamster ovary cells. *Biochem Pharmacol* **66**(5):785-790.
- Kowal D, Zhang J, Nawoschik S, Ochalski R, Vlattas A, Shan Q, Schechter L and Dunlop J (2002) The C-terminus of Gi family G-proteins as a determinant of 5-HT(1A) receptor coupling. *Biochem Biophys Res Commun* **294**(3):655-659.
- Masland RH (2001) The fundamental plan of the retina. *Nat Neurosci* **4**(9):877-886.
- Nakajima Y, Iwakabe H, Akazawa C, Nawa H, Shigemoto R, Mizuno N and Nakanishi S (1993) Molecular characterization of a novel retinal metabotropic glutamate receptor mGluR6 with a high agonist selectivity for L-2-amino-4-phosphonobutyrate. *J Biol Chem* **268**(16):11868-11873.

- Nakajima Y, Moriyama M, Hattori M, Minato N and Nakanishi S (2009) Isolation of ON bipolar cell genes via hrGFP-coupled cell enrichment using the mGluR6 promoter. *J Biochem* **145**(6):811-818.
- Nawy S (1999) The metabotropic receptor mGluR6 may signal through G(o), but not phosphodiesterase, in retinal bipolar cells. *J Neurosci* **19**(8):2938-2944.
- Pinto LH, Vitaterna MH, Shimomura K, Siepka SM, Balannik V, McDearmon EL, Omura C, Lumayag S, Invergo BM, Glawe B, Cantrell DR, Inayat S, Olvera MA, Vessey KA, McCall MA, Maddox D, Morgans CW, Young B, Pletcher MT, Mullins RF, Troy JB and Takahashi JS (2007) Generation, identification and functional characterization of the nob4 mutation of Grm6 in the mouse. *Vis Neurosci* **24**(1):111-123.
- Prezeau L, Carrette J, Helpap B, Curry K, Pin JP and Bockaert J (1994) Pharmacological characterization of metabotropic glutamate receptors in several types of brain cells in primary cultures. *Mol Pharmacol* **45**(4):570-577.
- Quraishi S, Gayet J, Morgans CW and Duvoisin RM (2007) Distribution of group-III metabotropic glutamate receptors in the retina. *J Comp Neurol* **501**(6):931-943.
- Rondard P, Liu J, Huang S, Malhaire F, Vol C, Pinault A, Labesse G and Pin JP (2006) Coupling of agonist binding to effector domain activation in metabotropic glutamate-like receptors. *J Biol Chem* **281**(34):24653-24661.
- Rosemond E, Peltekova V, Naples M, Thogersen H and Hampson DR (2002) Molecular determinants of high affinity binding to group III metabotropic glutamate receptors. *J Biol Chem* **277**(9):7333-7340.
- Schoepp DD (2001) Unveiling the functions of presynaptic metabotropic glutamate receptors in the central nervous system. *J Pharmacol Exp Ther* **299**(1):12-20.

- Sekine-Aizawa Y and Huganir RL (2004) Imaging of receptor trafficking by using alpha-bungarotoxin-binding-site-tagged receptors. *Proc Natl Acad Sci U S A* **101**(49):17114-17119.
- Shen Y, Heimel JA, Kamermans M, Peachey NS, Gregg RG and Nawy S (2009) A transient receptor potential-like channel mediates synaptic transmission in rod bipolar cells. *J Neurosci* **29**(19):6088-6093.
- Takao M, Morigiwa K, Sasaki H, Miyoshi T, Shima T, Nakanishi S, Nagai K and Fukuda Y (2000) Impaired behavioral suppression by light in metabotropic glutamate receptor subtype 6-deficient mice. *Neuroscience* **97**(4):779-787.
- Tian L and Kammermeier PJ (2006a) G protein coupling profile of mGluR6 and expression of G alpha proteins in retinal ON bipolar cells. *Vis Neurosci* **23**(6):909-916.
- Vardi N, Duvoisin R, Wu G and Sterling P (2000) Localization of mGluR6 to dendrites of ON bipolar cells in primate retina. *J Comp Neurol* **423**(3):402-412.
- Walker MW, Jones KA, Tamm J, Zhong H, Smith KE, Gerald C, Vaysse P and Branchek TA (2005) Use of *Caenorhabditis elegans* G α q chimeras to detect G-protein-coupled receptor signals. *J Biomol Screen* **10**(2):127-136.
- Zeit C, Forster U, Neidhardt J, Feil S, Kalin S, Leifert D, Flor PJ and Berger W (2007) Night blindness-associated mutations in the ligand-binding, cysteine-rich, and intracellular domains of the metabotropic glutamate receptor 6 abolish protein trafficking. *Hum Mutat* **28**(8):771-780.
- Zeit C, Kloeckener-Gruissem B, Forster U, Kohl S, Magyar I, Wissinger B, Matyas G, Borruat FX, Schorderet DF, Zrenner E, Munier FL and Berger W (2006) Muta-

tions in CABP4, the gene encoding the Ca²⁺-binding protein 4, cause autosomal recessive night blindness. *Am J Hum Genet* **79**(4):657-667.

Zeit C, van Genderen M, Neidhardt J, Luhmann UF, Hoeben F, Forster U, Wycisk K, Matyas G, Hoyng CB, Riemsdag F, Meire F, Cremers FP and Berger W (2005) Mutations in GRM6 cause autosomal recessive congenital stationary night blindness with a distinctive scotopic 15-Hz flicker electroretinogram. *Invest Ophthalmol Vis Sci* **46**(11):4328-4335.

Zhang JY, Nawoschik S, Kowal D, Smith D, Spangler T, Ochalski R, Schechter L and Dunlop J (2003) Characterization of the 5-HT₆ receptor coupled to Ca²⁺ signaling using an enabling chimeric G-protein. *Eur J Pharmacol* **472** (1-2): 33-38.

FOOTNOTES

None

LEGENDS FOR FIGURES

Fig. 1. Inhibition calcium channels by mGluR6 and CSNB point mutants. A, a representative time course and current trace (inset) of mGluR6-mediated inhibition of the native calcium currents in SCG neurons at indicated agonist (GLU) concentrations. B, concentration-response curve of mGluR6 and CSNB point mutants. The wild-type and the mutants are represented by the indicated symbol. The data is represented as average calcium current percent inhibition \pm SEM for mGluR6 (n=11), P40L (n=3), G52R (n=6), G144S (n=4), C516Y (n=7), and E775K 100 μ M (n=23), 30 μ M (n=12), 10 μ M (n=9), 1 and 0.1 μ M GLU (n=4).

Fig. 2. Surface membrane labeling of mGluR6 and CSNB point mutants in SCG neurons. BBS tagged wild type and point mutants were co-injected with GFP (green) and treated with α -bungarotoxin Alexa Fluor 647 conjugate (red). Average calcium current inhibition by glutamate for the wild-type (white bar) and BBS tagged mGluR6 (black bar) in SCGs is shown in the bar graph. Number of cells is indicated in parentheses.

Fig. 3. Reconstitution of coupling to mGluR6 (white bars) and mGluR6 E775K mutant (black bars) using PTX-insensitive $G_{oA}CG$. Average calcium current inhibition (\pm SEM) in SCG neurons expressing mGluR6 or E775K alone, following overnight PTX treatment, and in combination with PTX-insensitive $G_{oA}CG$ (and $G\beta_1$ and $G\gamma_2$) are shown. Number of cells is indicated in parentheses.

Fig. 4. Reconstitution of coupling to mGluR6 E775K using PTX-insensitive $G_{i1}CG$, $G_{i2}CG$, or $G_{i3}CG$. Average calcium current inhibition (\pm SEM) in SCG neurons expressing E775K alone or in combination with the indicated PTX-insensitive $G\alpha$ protein (plus β_1 and γ_2) are shown. Number of cells is indicated in parentheses.

Fig. 5. Reconstitution of mGluR6 and mGluR6 E775K coupling in m3 (muscarinic type 3) stable HEK293 cells. A, representative traces from cells expressing mGluR6 (left) and mGluR6 E775K (right) alone or in combination with the indicated chimera with $G\beta_1$ and $G\gamma_2$. The scale bars represent 2 ratio units and 60 sec. B, scatter plots showing the fluorescence intensity ratios (F340/F380) of mGluR6 or E775K alone or in combination with the indicated chimera (including $G\beta_1$ and $G\gamma_2$). Number of analyzed cells is indicated in the text.

Figure 1





Figure 2




Figure 3




Figure 4




Figure 5

