

MOL #53728

**Augmentation of fear extinction by infusion of glycine transporter blockers into the
amygdala**

Sheng-Chun Mao, Hui-Ching Lin and Po-Wu Gean

Institute of Basic Medical Sciences and Department of Pharmacology, Center for Gene
Regulation and Signal Transduction Research, National Cheng-Kung University, Tainan 701,
Taiwan

MOL #53728

Running Title: Glycine transporter inhibition augments extinction

Send correspondence to: Dr. Po-Wu Gean

Institute of Basic Medical Sciences and Department of Pharmacology, National Cheng-Kung University, Tainan 701, Taiwan

TEL: (886)-6-2353535 ext. 5507

FAX: (886)-6-2749296; E-mail: powu@mail.ncku.edu.tw

24 pages of text with 8 figures, 1 supplemental figure and 41 references

216 words in Abstract, 387 words in Introduction and 1083 words in Discussion

The abbreviations used are:

ACSF, artificial cerebrospinal fluid; AMPA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; ANOVA, analysis of variance; BLA, basolateral nucleus of the amygdala; CNQX, 6-cyano-2,3-dihydroxy-7-nitroquinoxaline; CS, conditioned stimulus; D-APV, D-2-amino-5-phosphonovalerate; DCS, D-cycloserine; DMSO, dimethyl sulfoxide; fEPSP, field excitatory postsynaptic potential; GABA, gamma-aminobutyric acid; GlyT1, glycine transporter type 1; LA, lateral nucleus of the amygdala; LTD, long-term depression ; LTP, long-term potentiation; NFPS, N[3-(4'-fluorophenyl)-3-(4'-phenylphenoxy)propyl]sarcosine ; NMDA, N-Methyl-D-aspartic Acid ; PSD95, postsynaptic density protein 95; SAP97, synapse-associated protein 97; TS, tetanic stimulation; US, unconditioned stimulus.

MOL #53728

Abstract

It is known that fear extinction is blocked by the NMDA receptor antagonist. In this study we investigate whether extinction could be facilitated by the enhancement of NMDA response, achieved by the blocking of glycine transporters. In amygdala slices, NMDA at a concentration that normally does not have a long-term effect was found to reduce the cellular levels of PSD-95 and synapse-associated protein 97 (SAP97), in addition to the surface expression of GluR1/2, in the presence of a glycine transporter blocker (N[3-(4-fluorophenyl)-3-(4'-phenylphenoxy)] propylsarcosine (NFPS). In *in vivo* experiments, extinction training applied 24 h after conditioning reduced startle potentiation without influencing the conditioning-induced increase in the surface expression of GluR1/2. However, NFPS augmented extinction and reversed the conditioning-induced increase in GluR1/2 when infused bilaterally into the amygdala before extinction training. The effects of NFPS were therefore blocked by the NMDA antagonist. In parallel, NFPS treatment in conjunction with extinction reversed the conditioning-induced AMPA/NMDA ratio. In behavioral tests, Tat-GluR2_{3Y}, a synthetic peptide that has been shown to block AMPA receptor endocytosis, inhibited only the additional reduction caused by NFPS treatment, rather than returning the fear potentiation levels to those of fear-conditioned animals that did not undergo extinction. These results suggest that NFPS in combination with extinction training reverses GluR1/2 surface expression and thus augments extinction of conditioned fear.

MOL #53728

Introduction

In the human brain, the functional MRI signal in the amygdala decreases over time during repeated presentations of emotional facial expressions (Wright et al., 2001). Analogous to animal studies, this process may represent a form of extinction in which the response to a feared cue decreases as the cue is repeatedly presented without any adverse consequences (Taylor et al., 2003; Hermans et al., 2006). This exposure-based psychotherapy is a standard treatment for a number of anxiety disorders (Yehuda, 2002). However, in some patients, the successful reduction of fear through exposure therapy is followed by a return of fear, because the aversive or fear memory was stronger and extinction training did not erase the original memory (Quirk et al., 2000; Myers and Davis, 2002; Maren and Quirk, 2004; Barad, 2006).

Demonstration of the blocking of fear extinction by the NMDA receptor antagonist (Falls et al., 1992; Lee and Kim, 1998; Lin et al., 2003) led to the hypothesis that pharmacological manipulation of the NMDA receptor may influence extinction. Later, it was shown that D-cycloserine (DCS), a partial agonist at the glycine-binding site of the NMDA receptor, facilitated extinction of conditioned fear in rats (Walker et al., 2002; Ledgerwood et al., 2003). In human studies, patients who received DCS had significantly better outcomes than the placebo group in enhancing fear reduction in cases of social phobia and fear of heights (Ressler et al., 2004; Davis et al., 2006). These results suggest that agents that augment NMDA responses may be useful in the treatment of anxiety disorders.

By binding to a strychnine-insensitive site on the NR1 NMDA receptor subunit, glycine acts as a necessary coagonist at the NMDA receptor (Johnson and Ascher, 1987; Thompson et

MOL #53728

al., 1989). The glycine concentration in the cerebrospinal fluid is estimated to be in the low micromolar range (Westergren et al., 1994), a concentration that is sufficient to saturate the site under physiological conditions. However, glycine may normally exist at nonsaturating concentrations within the synaptic regions because of the strategic placement of the high-affinity glycine transporter type 1 (GlyT1) around the synapses. Thus, NMDA responses could be enhanced after blocking GlyT1 (Bergeron et al., 1998). The purpose of this study was therefore to determine whether the pharmacological blocking of GlyT1 facilitates extinction. We also aimed to delineate the mechanism behind the action of GlyT1 blockers.

MOL #53728

Methods and Materials

Animals

All procedures were approved by the Institutional Animal Care and Use Committee of the College of Medicine, National Cheng-Kung University. Animals were housed four to a cage in a temperature (24°C)-controlled animal colony under a 12:12 light–dark cycle, with lights on at 0700 h, and pelleted rat chow and water were available *ad libitum*. All behavioral procedures took place during the light cycle.

Surgery

Rats (6–8-weeks-old) were anesthetized with sodium pentobarbital (50 mg/kg, i.p.) and subsequently mounted on stereotaxic apparatus. Two cannulae made of 22-gauge stainless steel tubing (C313G, Plastic Products) were implanted bilaterally into the lateral (LA) or basolateral (BLA) amygdala [anteroposterior (AP), –2.8 mm; mediolateral (ML), ± 4.5 mm; dorsoventral (DV), –7.0 mm], and a 28-gauge dummy cannula was inserted into each cannula in order to prevent clogging. The rats were monitored and handled daily and were given 7 days to recover from the procedure. NFPS (5 μ g/side dissolved in 50% DMSO for intra-amygdalar injection; 1 or 10 mg/kg dissolved in 25% 2-hydroxypropyl- β -cyclodextrin 75% saline for intraperitoneal injection) was purchased from Tocris Cookson Ltd (Northpoint, UK). Sarcosine (dissolved in saline) was obtained from Sigma (St. Louis, MO). Drugs were administered bilaterally to the amygdala in a volume of 0.8 μ l at a rate of 0.1 μ l/min. The infusion cannulae were left in place for 2 min before being withdrawn.

Behavioral apparatus and procedures

Rats were trained and tested in a stabilimeter device. The behavioral experiments of fear

MOL #53728

conditioning and extinction training were performed in a standard operant chamber (San Diego Instruments, San Diego, CA). The acoustic startle stimulus was 50 ms of white noise at an intensity of 95 dB; the visual CS was a 3.7-s light produced by an 8-W fluorescent bulb attached to the back of the stabilimeter; and the US was a 0.6-mA footshock of a duration of 0.5 s. Rats were placed in the training/testing chamber for 10 min and returned to their home cages on three consecutive days in order to habituate them to the test chamber and to minimize the effect of contextual conditioning. On the following two days, the rats were handled in the same chamber prior to fear conditioning for pre-exposure, during which the baseline startle was measured on each of the 2 days by administering 10 startle stimuli with a 2-min interstimulus interval (ISI). Rats with equivalent baseline mean startle amplitudes were then divided into separated matched groups. On the day of fear conditioning, the animal was taken to the room, allowed to habituate, and placed in the chamber as before. The CS-US pairing began after a 3-min acclimatization period in the chamber.

Experiment 1: Effect of NFPS on the NMDA receptor-mediated synaptic response

Male Sprague-Dawley rats (150–200 g) were decapitated and their brains rapidly removed and placed in cold oxygenated artificial cerebrospinal fluid (ACSF) solution. Subsequently, the brain was hemisected and cut transversely posterior to the first branch and anterior to the last branch of the superior cerebral vein. The resulting section was glued to the chuck of a Vibroslice tissue slicer. Transverse slices of a 450- μ m thickness were cut and the appropriate slices placed in a beaker of oxygenated ACSF at room temperature for at least one hour before recording. The ACSF solution was of the following composition (in mM): NaCl 117, KCl 4.7, CaCl₂ 2.5, MgCl₂ 1.2, NaHCO₃ 25, NaH₂PO₄ 1.2 and glucose 11. The ACSF

MOL #53728

was bubbled continuously with 95% O₂–5% CO₂ and had a pH of 7.4.

A single slice was transferred to the recording chamber, in which it was held submerged between two nylon nets and maintained at 32±1°C. The chamber consisted of a circular well of a low volume (1–2 ml) and was perfused constantly at a rate of 2–3 ml/min. Extracellular field potentials were created by electrical stimulation of the external capsule (EC), which contained fibers from the auditory cortex to the lateral amygdala, with a concentric bipolar stimulating electrode. Electrical stimuli (150 µsec in duration) were delivered at a frequency of 0.05 Hz, and baseline field potentials were adjusted to ~30–40% of the maximal responses. Bicuculline (10 µM) and CGP52432 (10 µM) were present in the perfusion solution to block GABA_A and GABA_B receptors, respectively. NMDA receptor-mediated fEPSP (fEPSP_{NMDA}) was isolated pharmacologically by using Mg⁺⁺-free solution in the presence of AMPA receptor antagonist CNQX (10 µM).

Experiment 2: Effects of GlyT1 blockers on extinction and reinstatement.

On day 1, rats were placed in the startle boxes and received 10 light–footshock pairings with an ITI of 2 min. On day 2, the rats were returned to the startle box and received 3 sessions of 10 presentations of the 3.7-sec light in the absence of either shock or the startle-elicited noise burst (extinction training). Each session was separated by 10 min, with an ITI of 1 min. The GlyT1 blocker NFPS (5 µg per side) was infused into the amygdala 30 min before extinction training. Twenty-four hours later (day 3), rats were returned to the startle chamber for testing. Thirty initial startle stimuli were presented in darkness at a 10-sec ISI (leader stimuli), followed by 30 startle-eliciting noise bursts presented alone in darkness (noise-alone trial) and 30 noise bursts presented 3.2 sec after onset of the 3.7-sec light

MOL #53728

(light–noise trials). The two trial types were presented in a balanced mixed order (ITI, 30 sec).

Startle amplitude was averaged over all of the light–noise or noise-alone stimuli within the test session, and the percentage of fear-potentiated startle was computed as follows: [(startle amplitude on CS–noise minus noise-alone trials)/(noise-alone trials)] x 100. Paired and unpaired rats were administered 10 light–shock pairings in a paired or pseudorandom manner and retention of memory was tested 48 h later with no extinction training administered. A separate group of rats that received NFPS infusion but no extinction training served as No-CS NFPS controls. To examine whether NFPS-treated rats exhibited reinstatement of fear, the rats were given US-alone trials (10 unpaired footshocks) 1 h after the test, and retention of memory was tested 24 h later.

Experiment 3: Effect of NFPS on the surface expression of GluR1 and GluR2

***In vitro* experiment:** Amygdala slices were treated with 20 μ M of NMDA, 10 μ M of NMDA, 10 μ M of NMDA plus 1 μ M NFPS, or 10 μ M of NMDA plus 1 mM sarcosine for 3 min. One hour later, LA and BLA tissues were dissected out, placed on ice and washed twice with ice-cold ACSF. Control slices were incubated in ACSF for 1 h. The slices were then incubated with ACSF containing 0.5mg/ml Sulfo-NHS-LC- Biotin (Pierce Chemical Co., Rockford, IL) for 1 h on ice. Next, the slices were rinsed in ACSF and then sonicated briefly in homogenizing buffer (1% Triton X-100, 0.1% SDS, 50 mM Tris-HCl, 0.3 M sucrose, 5 mM EDTA, 2 mM sodium pyrophosphate, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonyl fluoride (PMSF), 20 μ g/ml leupeptin, and 4 μ g/ml aprotinin; pH 7.5). After sonication, the samples were centrifuged at 18200 g for 30 min at 4°C and the supernatant was obtained. The protein concentration in the soluble fraction was then

MOL #53728

measured using a Bradford assay, with bovine serum albumin as the standard. Biotinylated protein (400 μ g) from the supernatant was precipitated with 50 μ l of 50% Neutravidin agarose (Pierce Chemical Co.) for 16 h at 4°C and washed 4 times with homogenizing buffer. Bound protein was re-suspended in 4 μ l of SDS sample buffer and boiled, then biotinylated protein was resolved in 8.5% SDS–polyacrylamide gel, blotted electrophoretically on PVDF membrane, and blocked overnight in TBS buffer containing 5% non-fatty milk. Surface GluR1 and GluR2 receptors and pan-cadherin (surface protein control) were detected by a biotinylation assay, followed by Western blot analysis using GluR1 (1:4000, Santa Cruz Biotech., Santa Cruz, CA), GluR2 (1:5000, Chemicon, Temecula, CA), PSD-95 (1:5000, Upstate, Lake Placid, NY), SAP97 (1:5000, Stressgen, Victoria, BC, Canada), or pan-cadherin (1:2500; Sigma) antibodies, followed by HRP-conjugated secondary antibody for 1 h. An enhanced chemiluminescence kit was used for detection, and Western blots were developed in the linear range used for densitometry. GluR1 and GluR2 levels in the NMDA- and NFPS-treated slices were expressed as a percentage of those in the control slices from rats that did not undergo drug treatment. Similarly, the GluR1 and GluR2 levels in the conditioned animals were expressed as a percentage of those in the naïve controls that did not receive light–shock pairings.

***In vivo* experiment:** On day 1, the rats were conditioned; 24 h later, they were treated with NFPS (5 μ g per side) before extinction training. LA and BLA tissues were dissected out 24 h after extinction training and the surface receptors were labeled with biotin.

Experiment 4: Effect of NFPS on the AMPA/NMDA ratio

Whole-cell patch-clamp recordings were made from the LA neurons. EPSCs were evoked at

MOL #53728

0.03 Hz by extracellular stimulation of fibers emerging from the internal capsule, which originate in the medial geniculate nucleus of the thalamus and project monosynaptically to the LA using a bipolar electrode. Patch electrodes were pulled from a thick-wall glass capillary (0.75 mm I.D., 1.5 mm O.D.) to a tip resistance of 3–5 M Ω . The composition of the internal solution was (in mM): Cs-gluconate 115, NaCl 5, EGTA 1, CaCl₂ 0.3, MgCl₂ 2, Na-ATP 5, Na-GTP 0.4, HEPES 10. The final pH of the internal solution was adjusted to 7.3 by adding 1 M KOH; the final osmolarity was adjusted to 280 mOsm by adding sucrose. Recordings were low-pass-filtered at 2.5–20 kHz and digitized at 5–50 kHz. The signal was monitored and recorded using an Axopatch 200B amplifier. On-line analysis and control of experimental acquisition was accomplished via a 586 (Intel)-based PC clone and a Digidata 1200 computer interface. AMPAR-mediated EPSC was evoked when the neurons were voltage-clamped at –70 mV, whereas NMDAR-mediated EPSC was determined as the current amplitude at 50 ms after the peak EPSC amplitude at a holding potential of +40 mV (Du et al., 2008).

Data analysis

All values in the text are mean \pm SEM. Differences between the groups were evaluated by one-way ANOVA followed by Newman–Keuls *post hoc* tests. The paired *t* test was used to analyze the differences in fEPSP before and after drug treatment, and the unpaired *t* test to analyze the differences in startle reflex between the drug-treated and vehicle control groups. The level of significance was $P < 0.05$.

MOL #53728

Results

Experiment 1: Effect of NFPS on the NMDA receptor-mediated synaptic response

NFPS is a potent and selective antagonist of GlyT1 (Bergeron et al., 1998; Aubrey and Vandenberg, 2001). The effect of NFPS (1 μ M) on fEPSP as a function of time is illustrated in Figure 1A. After the evoked responses had been stable for 20–30 min, NFPS was applied to the amygdala slices by bathing, which caused a reversible increase in the slope of fEPSP. On average, the magnitude of increase measured at 10 min after application was $85.2 \pm 11.2\%$ ($P < 0.01$). In order to determine the involvement of NMDA receptors in the action of NFPS, slices were pre-treated with NMDA antagonist D-APV (50 μ M) before the application of NFPS. Figure 1B shows that the effect of NFPS was abolished in the presence of D-APV ($100.8 \pm 2.3\%$ of control). We further examined the effect of NFPS on NMDA receptor-mediated fEPSP (fEPSP_{NMDA}), isolated pharmacologically using Mg⁺⁺-free solution in the presence of AMPA receptor antagonist CNQX (10 μ M). Ten minutes after the application of NFPS, the fEPSP_{NMDA} amplitude was increased by $83.7 \pm 9.0\%$. At end of the experiment, D-APV was applied to ensure that fEPSP_{NMDA} was indeed mediated by the NMDA receptors (Fig. 1C). Taken together, these results indicate that NFPS selectively enhanced the NMDA receptor-mediated synaptic response.

Experiment 2: Effect of NFPS on fear extinction

Rats were conditioned with 10 light–shock pairings and were randomly assigned to 4 groups, Paired, Extinction, Extinction+NFPS, and Extinction+vehicle. In the Paired group, the

MOL #53728

rats were tested for fear-potentiated startle 48 h later (day 3) without receiving extinction training or any drug infusion. The Extinction group was given light-alone trials (extinction training) on day 2 without drug infusion, while the Extinction+NFPS and Extinction+vehicle rats were infused with NFPS (5 μ g per side) or an equal amount of 50% DMSO (0.8 μ l) 30 min before extinction training. Memory retention was assessed 24 h after extinction training. As expected, extinction training decreased the fear-potentiated startle from $188.5 \pm 9.9\%$ to $74.5 \pm 6.6\%$ ($P < 0.001$, unpaired t -test). Furthermore, as can be seen from Figure 2, fear-potentiated startle was significantly lower in the NFPS-treated rats as compared with the vehicle-treated control and Ext groups ($F_{(2,15)} = 8.31$, $P < 0.01$). These results indicate that NFPS facilitated extinction. In a separate group of conditioned rats, HA-966 (6 mg/kg), an antagonist for the glycine-binding site of the NMDA receptor, was injected intraperitoneally followed by NFPS, 30 min before light-alone trials, and memory retention was assessed 24 h after extinction training. As shown in Figure 2, HA-966 blocked the effect of NFPS. The startle potentiation in rats treated with HA-966 plus NFPS was significantly higher than in those treated with NFPS alone and was comparable to those receiving extinction training only or extinction plus vehicle ($P > 0.5$).

The effect of NFPS might reflect a reduction in fear response, or, non-specifically, a disruption in the startle reflex independent of fear reduction. To distinguish between these possibilities, rats were treated with NFPS without extinction training, whereupon we found that the startle potentiation in these rats (NFPS w/o Ext) was comparable to those that had not been subjected to extinction training (Paired w/o Ext) ($t_{(10)} = 0.75$, $P = 0.47$), suggesting that NFPS augmented extinction and did not exert a general effect on the startle reflex.

MOL #53728

Furthermore, when NMDA receptor antagonist D-APV (12.5 nmol/site) was co-applied with NFPS, the effects of NFPS plus extinction training were completely abolished, indicating the requirement of NMDA receptor activation. A diagram showing the infusion cannulae tip locations is included as Supplemental Figure 1.

We next determined whether the NFPS-treated rats exhibited reinstatement. On day 1, rats were trained with 10 light–shock pairings and 24 h later received intra-amygdala infusion of NFPS (5 μ g) or vehicle 30 min before extinction training. Memory retention was tested 24 h later (day 3, Test 1). One hour later, the rats were given 10 US-alone trials and memory retention was tested 24 h later (day 4, Test 2). Figure 3 shows that in the NFPS-treated rats, the startle potentiation in Test 2 was significantly higher than in Test 1 ($P<0.01$) but was significantly less than in Test 2 of the vehicle-treated rats ($P<0.01$). These results indicate that the NFPS-treated rats exhibited reinstatement after the US presentations.

Experiment 3: Effect of NFPS on the surface expression of GluR1 and GluR2

Previous studies of the hippocampus have shown that NMDA-induced LTD is accompanied by a persistent reduction in the surface expression of GluR1 (Lee et al., 1998; Man et al., 2007). We tested whether this was the case in the amygdala slices by using cell surface biotinylation techniques, as described in the Methods section. Cell extracts were prepared and biotinylated surface proteins were isolated by Neutravidin precipitation 1 h after the application of NMDA (20 μ M, 3 min). Control slices were incubated in ACSF solution for 1 h. Figure 4A shows that NMDA (20 μ M) application caused a reduction in the GluR1 level ($65.5\pm 3.4\%$ of control, $P<0.001$ vs. control), whereas a lower concentration of NMDA (10

MOL #53728

μM) had no effect ($95.0\pm 3.4\%$). Only in the presence of NFPS ($1\ \mu\text{M}$) did $10\ \mu\text{M}$ NMDA induce a significant loss of surface GluR1 ($69.2\pm 4.2\%$, $P<0.001$ vs. NMDA $10\ \mu\text{M}$).

N-methylglycine (sarcosine) is an endogenous antagonist of glycine transporter-1 (McBain et al., 1989). We tested whether sarcosine produced a similar effect to NFPS. As illustrated in Figure 4B, in the presence of sarcosine ($1\ \text{mM}$), $10\ \mu\text{M}$ NMDA induced a significant loss of surface GluR1 ($69.0\pm 4.2\%$, $P<0.001$ vs. NMDA $10\ \mu\text{M}$). We also assessed the level of surface GluR2. As shown in Figure 4C, NMDA ($20\ \mu\text{M}$) application caused a reduction in the GluR2 level ($65.4\pm 4.7\%$, $P<0.001$ vs. control), whereas a lower concentration of NMDA ($10\ \mu\text{M}$) had no effect ($101.8\pm 5.8\%$). In the presence of NFPS ($1\ \mu\text{M}$), $10\ \mu\text{M}$ NMDA induced a significant loss of surface GluR2 ($61.1\pm 4.8\%$, $P<0.05$ vs. NMDA $10\ \mu\text{M}$). NFPS alone did not affect the surface GluR2 level ($96.3\pm 6.2\%$).

We examined whether NFPS affected GluR1 expression after extinction training. Rats were conditioned with 10 light–shock pairings and were randomly assigned to 4 groups. LA and BLA tissues were dissected out 24 h after extinction training and the level of GluR1 was determined. Extinction training did not significantly influence the conditioning-induced increase in GluR1 ($p=0.10$); however, the level was significantly lower in the NFPS-treated rats as compared with the vehicle-treated control and Ext groups ($F_{(2,15)}=6.8$, $P<0.01$) (Fig. 5A). Treatment with HA-966 ($6\ \text{mg/kg}$, i.p.) 30 min before the administration of NFPS prevented the effect of NFPS, such that the level of GluR1 in the HA966 plus NFPS-treated rats was not significantly different from that of rats that received extinction training only ($P>0.1$). Furthermore, NFPS-treated rats that did not undergo extinction training had a startle potentiation equivalent to that of the Paired or Ext rats ($P>0.5$), suggesting that NFPS by

MOL #53728

itself did not influence extinction. We also assessed the level of surface GluR2. As shown in Figure 5B, conditioning increased the surface expression of GluR2 to $141.2 \pm 8.5\%$ that of the naïve controls. Light-alone trials 24 h after training did not significantly influence the increase in GluR2 expression ($141.5 \pm 7.9\%$, $P > 0.5$); however, infusion of NFPS (5 μg per side) into the amygdala abolished the conditioning-induced increase in GluR2 ($102.3 \pm 2.3\%$, $P < 0.05$ vs. Extinction). Administration of NFPS without extinction training had no effect on the increase in GluR2 expression ($136.3 \pm 8.6\%$). Treatment with D-APV prevented the effect of NFPS, such that the level of GluR2 in the D-APV plus NFPS-treated was not significantly different to that of rats that received extinction training only ($139.2 \pm 7.6\%$, $P > 0.5$).

PSD-95 is a scaffolding protein of the postsynaptic density; it interacts with GluR indirectly through stargazin (Chen et al., 2000) and regulates GluR trafficking (El-Husseini Ael et al., 2002; Schnell et al., 2002). We therefore examined whether PSD-95 expression was altered after treatment with NFPS. Figure 6A shows that NMDA (20 μM) application caused a reduction in the PSD-95 level ($69.4 \pm 5.7\%$, $P < 0.001$ vs. control), whereas a lower concentration of NMDA (10 μM) had no effect ($98.3 \pm 3.7\%$). In the presence of NFPS (1 μM), 10 μM NMDA induced a significant loss of PSD-95 ($67.0 \pm 7.4\%$, $P < 0.01$ vs. NMDA 10 μM). NPFS alone did not affect the PSD-95 level ($100.0 \pm 5.7\%$). The insertion and removal of AMPA receptors are regulated by PSD-95 and PDZ-domain-containing protein SAP97, which binds to GluR1 and traffics GluR1 into the spine (Rumbaugh et al., 2003; Schluter et al., 2006). In the LA, the coupling of A-kinase-anchoring proteins (AKAPs) and protein kinase A (PKA) to GluR1 through SAP97 is essential for memory formation (Moita et al., 2002). We therefore examined whether SAP97 expression was involved in the extinction. Figure 6B

MOL #53728

shows that NMDA (20 μ M) application caused a reduction in the SAP97 level ($68.9 \pm 5.9\%$, $P < 0.001$ vs. control), whereas a lower concentration of NMDA (10 μ M) had no effect ($98.5 \pm 6.2\%$). In the presence of NFPS (1 μ M), 10 μ M NMDA induced a significant loss of SAP97 ($68.0 \pm 6.0\%$, $P < 0.01$ vs. NMDA 10 μ M). NFPS alone did not affect the SAP97 level ($95.4 \pm 6.0\%$).

We examined whether NFPS affected the levels of PSD-95 and SAP97 after extinction training. Rats were conditioned with 10 light–shock pairings and were randomly assigned to 4 groups, Paired, Extinction, Extinction+NFPS and Extinction+vehicle, as described in the Methods section. LA and BLA tissues were dissected out 24 h after extinction training and the levels of PSD-95 and SAP97 were determined. Extinction training did not significantly influence the conditioning-induced increase in PSD-95 ($P = 0.68$); however, the PSD-95 level was significantly lower in the NFPS-treated rats as compared with the levels in the vehicle-treated control and Ext groups ($F_{(2,15)} = 26.89$, $P < 0.001$) (Fig. 6C). Treatment with HA-966 (6 mg/kg, i.p.) 30 min before the administration of NFPS prevented the effect of NFPS, such that the level of PSD-95 in the HA966 plus NFPS-treated rats was not significantly different from that of rats that received extinction training only ($P > 0.5$). Furthermore, NFPS-treated rats that did not undergo extinction training had a PSD-95 level comparable to those of Paired or Ext rats ($P > 0.5$), suggesting that NFPS by itself did not influence the PSD-95 level. We also probed the level of SAP97. As shown in Figure 6D, conditioning increased the SAP97 expression level to $189.2 \pm 14.2\%$ that of the naïve controls. Light-alone trials 24 h after training did not significantly influence the increase in SAP97 expression ($185.7 \pm 12.1\%$, $P > 0.5$); however, infusion of NFPS (5 μ g per side) into the

MOL #53728

amygdala abolished the conditioning-induced increase in SAP97 ($101.9 \pm 11.2\%$, $P < 0.001$ vs. Extinction). Administration of NFPS without extinction training had no effect on the increase in the SAP97 level ($197.3 \pm 13.9\%$). Treatment with D-APV prevented the effect of NFPS, such that the level of SAP97 in the D-APV plus NFPS-treated rats was not significantly different to the levels in the rats that underwent extinction training only ($208.4 \pm 12.7\%$, $P > 0.1$).

Experiment 4: Effect of NFPS on the AMPA/NMDA ratio

To establish whether the decrease in surface receptors by extinction plus NFPS resulted in changes in excitatory synaptic transmission, we measured the relative contributions of AMPA receptors (AMPA receptors) and NMDA receptors (NMDARs) to the EPSCs, a method that has been shown to be a sensitive assay for detecting differences in glutamatergic synaptic strength (Ungless et al., 2001). Rats were conditioned and 24 h later were injected intraperitoneally with NFPS (1 or 10 mg/kg) or vehicle 30 min before light-alone trials. The rats were tested for fear-potentiated startle 24 h after extinction training. ANOVA for startle scores showed a significant effect for the group ($F_{(3,21)} = 51.15$, $P < 0.001$), and Newman-Keuls *post hoc* tests showed that the high-dose NFPS group differed from the vehicle group ($P < 0.05$) (Fig. 7A). This result was consistent with the facilitation of extinction by NFPS.

One hour after the test, amygdala slices were taken from rats of the four groups (Naïve, Paired, Extinction+vehicle, Extinction+NFPS). The AMPA/NMDA ratio was 1.26 ± 0.11 in slices from the naïve rats; this ratio was significantly higher in the Paired rats (3.35 ± 0.11 , $P < 0.001$, unpaired *t* test) suggesting that fear conditioning persistently increased

MOL #53728

AMPA-mediated synaptic transmission. There was no difference in the AMPA/NMDA ratio between the Paired and vehicle-treated extinction rats (3.56 ± 0.29 , $P > 0.1$), suggesting that extinction did not affect the conditioning-induced increase in the AMPA/NMDA ratio.

However, in the NFPS-treated extinction rats, the AMPA/NMDA ratio (1.17 ± 0.15) was significantly lower than that of the vehicle-treated extinction rats ($P < 0.001$) (Fig. 7B).

Importantly, there was no difference in the AMPA/NMDA ratio between the naïve and NFPS-treated extinction rats ($P > 0.5$). These results suggest that extinction failed to affect the conditioning-induced increase in the AMPA/NMDA ratio. Only in the presence of NFPS did extinction training reverse this increase.

GluR2_{3Y}, a synthetic peptide containing a short C-terminal sequence of GluR2 (₈₆₉YKEGYNVYG₈₇₇), which is critical for the expression of hippocampal CA1 LTD (Ahmadian et al., 2004), has been shown to block LTD, depotentiation and AMPA receptor endocytosis in the hippocampus, nucleus accumbens and amygdala (Ahmadian et al., 2004; Brebner et al., 2005; Kim et al., 2007). When GluR2_{3Y} was fused to the cell membrane transduction domain of the HIV-1 Tat protein (Tat-GluR2_{3Y}), it became membrane-permeable and was able to impair the extinction of fear memory (Kim et al., 2007; Dalton et al., 2008).

We performed behavioral assessment in order to determine whether Tat-GluR2_{3Y} influenced the effect of NFPS. Rats were divided into 4 groups: rats that received Tat-GluR2_{3Y} (15 pmol in 0.8 μ l saline per side) or Tat-GluR2_{3A}(₈₆₉AKEGANVAG₈₇₇), a control peptide that had no effect on AMPA receptor endocytosis, followed by NFPS (10 mg/kg, i.p.) or vehicle 30 min prior to extinction training. Two-way ANOVA revealed no group effect (Tat-GluR2_{3Y} vs.

Tat-GluR2_{3A}, $F_{(1,20)}=3.29$, $P=0.08$) or effects of drug treatment (NFPS vs. vehicle, $F_{(1,20)}=1.48$,

MOL #53728

$P=0.24$) but a significant effect of interaction ($F_{(1,20)}=4.75$, $P<0.05$). *Post hoc* comparisons revealed that the startle potentiation in the Tat-GluR2_{3A}-NFPS rats was significantly less than that in the Tat-GluR2_{3Y}-NFPS rats ($P<0.05$). Furthermore, the startle potentiation in the Tat-GluR2_{3Y}-vehicle rats was comparable with that in the Tat-GluR2_{3A}-vehicle rats ($P=0.81$), indicating that Tat-GluR2_{3Y} had no effect in the absence of NFPS. Interestingly, in the Tat-GluR2_{3Y} rats, the startle potentiation in the NFPS-treated rats was comparable with that in the vehicle-treated rats ($P=0.61$), indicating that Tat-GluR2_{3Y} specifically blocked the effect of NFPS on extinction.

MOL #53728

Discussion

From a classic viewpoint, extinction results from a positive cognitive enhancement of CS–no US learning concomitant with the increased inhibitory tone coming from ventromedial PFC projection neurons to the amygdala (Myers and Davis, 2002; Maren and Quirk, 2004; Quirk et al., 2003). Extinction can also be mediated by an increase in the inhibitory transmission within the amygdala, likely via modulation of the GABA_A receptor (Chhatwal et al., 2005). In the present study, we combined behavioral, biochemical and electrophysiological experiments to explore the effects of GlyT1 blockers on fear extinction. We showed for the first time that: (1) the GlyT1 blocker NFPS selectively enhanced the NMDA receptor-mediated synaptic response in the amygdala; (2) NFPS augmented extinction without exerting a general effect on the startle reflex; (3) the NFPS-treated rats exhibited partial reinstatement after the US presentations; (4) NMDA at a concentration that is normally ineffective reduced the surface expression of GluR1 and GluR2 and the levels of PSD-95 and SAP97 in the presence of NFPS; (5) NFPS plus extinction training reversed the conditioning-induced increases in PSD-95 and SAP97, and in the surface expression of GluR1 and GluR2; (6) NFPS treatment in conjunction with extinction training reversed the conditioning-induced AMPA/NMDA ratio; and (7) GluR2_{3Y}, a synthetic peptide that is critical for the expression of hippocampal CA1 LTD, specifically blocked the effect of NFPS on extinction. These results suggest that NFPS augments extinction via enhancement of NMDA-mediated AMPA receptor endocytosis in the amygdala. Intriguingly, DCS which binds at the glycine-binding site of the NMDA receptor produced similar facilitation of

MOL #53728

extinction, reversal of GluR1/2 expression and less reinstating effect (Ledgerwood et al., 2004). These results suggest that agents that augment NMDA responses may be useful in the treatment of anxiety disorders.

NFPS potentiates NMDA receptor-mediated synaptic response

Synaptic transmission in the IC-LA is mediated primarily by AMPA receptors. In this study, we showed that exogenous application of the GlyT1 inhibitor NFPS increased fEPSP, and that this effect was not seen in the presence of an NMDA receptor antagonist. In addition, the pharmacologically-isolated NMDA receptor-mediated response was enhanced by NFPS. These results demonstrate that the glycine concentration within the synaptic regions of the amygdala is strongly regulated by glycine transporters and under normal condition is not sufficient to saturate the glycine-binding site of the NMDA receptor. Thus, the blocking of GlyT1 selectively enhances the NMDA response without affecting the AMPA receptor-mediated synaptic response.

Mechanism of augmented extinction by NFPS

In hippocampal and cortical neurons, a brief application of NMDA induces long-term depression (LTD) that is associated with a reduction in surface GluR1 (Lee et al., 1998; Beattie et al., 2000). Using a similar procedure, we found that NMDA at a concentration that is normally unable to elicit a long-term effect reduced the surface expression of GluR1 and GluR2 in the presence of NFPS. NFPS *per se* had no effect on the conditioning-induced GluR1 expression; only in conjunction with extinction training did NFPS reverse the

MOL #53728

increased GluR1 expression. These results are consistent with the idea that a threshold activation of NMDA receptors and subsequent Ca^{++} influx must be reached in order to trigger endocytosis of AMPA receptors. In this way, NFPS specifically produced cue-induced extinction rather than exerting a general reduction of fear.

In behavioral tests, Tat-GluR2_{3Y}, a synthetic peptide that is critical for the expression of hippocampal CA1 LTD, blocked only the additional reduction caused by NFPS treatment, rather than returning the fear potentiation level to that of fear-conditioned animals that had not undergone extinction. These results suggest that extinction training-induced inhibitory learning is intact and NFPS only adds a bit of erasure. NFPS augments extinction via the facilitation of NMDA receptor activation, which induces Ca^{++} influx resulting in the activation of calcineurin. Calcineurin activation leading to local reorganization of F-actin causes displacement of PKA from the synapse, which may initiate endocytosis of receptors and trigger a loss of GluR1 and GluR2 from the surface of neurons. The present results together previous reports suggest that

PSD-95 is a prominent postsynaptic density molecule that interacts indirectly with AMPA receptors through the transmembrane protein stargazin (Chen et al., 2000) and regulates the trafficking and localization of AMPA receptors at synapses (EL-Husseini Ael et al., 2002; Schnell et al., 2002; Beique and Andrade, 2003). In the present study, we showed that the PSD-95 level was down-regulated in response to NMDA plus NFPS stimulation, which suggests that the degradation of GluR-interacting proteins leads to destabilization of the receptor and its subsequent endocytosis. In the LA, the coupling of A-kinase-anchoring proteins (AKAPs) and protein kinase A (PKA) to GluR1 through SAP97 is essential for

MOL #53728

memory formation (Moita et al., 2002). Here, we provide evidence suggesting that SAP97 is also involved in extinction, because the level of SAP97 decreased in response to NFPS stimulation, leading to augmentation of extinction.

Two recent reports showed that extinction reversed both the enhanced synaptic efficacy observed in conditioned rats and the conditioning-induced enhancement of surface expression of AMPA receptors (Kim et al., 2007; Dalton et al., 2008), implying the unlearning of extinction. In contrast to these reports, we found that extinction training failed to influence the conditioning-induced increase in the surface expression of GluR1 and GluR2, as well as the AMPA/NMDA ratio. Only in the presence of NFPS did extinction training reverse these changes. In addition, the NFPS-treated rats exhibited partial reinstatement after US alone presentations. The reasons for the difference are not known, but could be attributable to the different tests (freezing *vs.* fear-potentiated startle) employed. The extinction protocol in the recent reports consisted of 20 tone presentations, each 20–30 sec in duration, whereas we used 30 light presentations of 3.7 sec. The more extensive extinction protocol used in the previous studies may have reached the threshold for the activation of NMDA receptors and subsequent Ca^{++} influx to trigger endocytosis of AMPA receptors.

In summary, facilitation of extinction could result from a positive cognitive enhancement of CS-no US learning or an erasure of the original CS-US association. In the present study, we found that co-administration of NFPS with extinction training facilitated extinction and reversed the conditioning-induced increase in GluR1 and GluR2, as well as the AMPA/NMDA ratio. In this context, NFPS in combination with extinction training does erase the original excitatory association. As high-affinity glycine transporters are usually located

MOL #53728

around the synapses, glycine may normally exist at nonsaturating concentrations within the synaptic regions. Therefore, it would be beneficial to treat posttraumatic stress disorder patients with a glycine transporter blocker in conjunction with exposure therapy.

MOL #53728

Acknowledgements

We thank Dr Min-Der Lai for critical comments on the manuscript.

References

- Ahmadian G, Ju W, Liu L, Wyszynski M, Lee SH, and Dunah AW (2004) Tyrosine phosphorylation of GluR2 is required for insulin-stimulated AMPA receptor endocytosis and LTD. *Embo J* **23**:1040-1050.
- Aubrey KR and Vandenberg RJ (2001) N[3-(4'-Fluorophenyl)-3-(4'-phenylphenoxy)propyl]sarcosine (NFPS) is a selective persistent inhibitor of glycine transport. *Br J Pharmacol* **134**:1429-1436.
- Barad M (2006) Is extinction of fear erasure or inhibition? Why both, of course. *Learn Mem* **13**: 108-109.
- Beattie EC, Carroll RC, Yu X, Morishita W, Yasuda H, von Zastrow M, and Malenka RC (2000) Regulation of AMPA receptor endocytosis by a signaling mechanism shared with LTD. *Nat Neurosci* **3**: 1291-1300.
- Bergeron R, Meyer TM, Coyle JT, and Greene RW (1998) Modulation of N-methyl-D-aspartate receptor function by glycine transport. *Proc Natl. Acad Sci* **95**: 15730-15734.
- Brebner K, Wong TP, Liu L, Liu Y, Campsall P, and Gray S (2005) Nucleus accumbens long-term depression and the expression of behavioral sensitization. *Science* **310**: 1340-1343.
- Chen L, Chetkovich DM, Petralia RS, Sweeney NT, Kawasaki Y, Wenthold RJ, Brecht DS, and Nicoll RA (2000) Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature* **408**: 936-943.
- Chhatwal JP, Myers KM, Ressler KJ, and Davis M (2005) Regulation of gephyrin and GABAA receptor binding within the amygdala after fear acquisition and extinction. *J Neurosci* **25**: 502-506.
- Dalton GL, Wang YT, Floresco SB, and Phillips AG (2008) Disruption of AMPA receptor endocytosis impairs the extinction, but not acquisition of learned fear. *Neuropsychopharmacology* (In Press)
- Davis M, Ressler K, Rothbaum BO, and Richardson R (2006) Effects of D-cycloserine on extinction: translation from preclinical to clinical work. *Biol Psychiatry* **60**: 369-375.
- El-Husseini Ael D, Schnell E, Dakoji S, Sweeney N, Zhou Q, Prange O, Gauthier-Campbell C, Aguilera-Moreno A, Nicoll RA, and Brecht DS (2002) Synaptic strength regulated by palmitate cycling on PSD-95. *Cell* **108**: 849-863.
- Falls WA, Miserendino JD, and Davis M (1992) Extinction of fear-potentiated startle: blockade by infusion of an NMDA antagonist into the amygdala. *J Neurosci* **12**: 854-863.
- Gomez LL, Alam S, Smith KE, Horne E, and Dell'Acqua ML (2002) Regulation of A-kinase anchoring protein 79/150-cAMP-dependent protein kinase postsynaptic targeting by NMDA receptor activation of calcineurin and remodeling of dendritic actin. *J Neurosci* **22**: 7027-7044.
- Hermans D, Craske MG, Mineka S, and Lovibond PF (2006) Extinction in human fear conditioning. *Biol Psychiatry* **60**: 361-368.
- Johnson JW and Ascher P (1987) Glycine potentiates the NMDA response in cultured mouse brain neurons. *Nature* **325**: 529-531.
- Kim J, Lee S, Park K, Hong I, Song B, Son G, Park H, Kim WR, Park E, Choe HK, Kim H,

MOL #53728

- Lee C, Sun W, Kim K, Shin KS, and Choi S (2007) Amygdala depotentiation and fear extinction. *Proc Natl Acad Sci USA* **104**: 20955–20960.
- Ledgerwood L, Richardson R, and Cranney J (2003) Effects of D-cycloserine on the extinction of conditioned freezing. *Beh Neurosci* **117**: 341-349.
- Ledgerwood L, Richardson R, and Cranney J (2004) D-cycloserine and the facilitation of extinction of conditioned fear: consequences for reinstatement. *Behav Neurosci* **118**,505-513.
- Lee H and Kim JJ (1998) Amygdalar NMDA receptors are critical for new fear learning in previously fear-conditioned rats. *J Neurosci* **18**: 8444-8454.
- Lee HK, Kameyama K, Huganir RL, and Bear MF (1998) NMDA induces long-term synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in hippocampus. *Neuron* **21**: 1151-1162.
- Lin CH, Yeh SH, and Gean PW (2003) The similarities and diversities of signal pathways leading to consolidation and extinction of fear memory. *J Neurosci* **23**: 8310-8317.
- Man HY, Sekine-Aizawa Y, and Huganir RL (2007) Regulation of {alpha}-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit. *Proc Natl Aca Sci USA* **104**: 3579-3584.
- Maren S and Quirk GJ (2004) Neuronal signalling of fear memory. *Nat Rev Neurosci* **5**: 844-852.
- McBain CJ, Kleckner NW, Wyrick S, and Dingledine R (1989) Structural requirements for activation of the glycine coagonist site of N-methyl-D-aspartate receptors expressed in xenopus oocytes. *Mol Pharmacol* **36**: 556-565.
- Moita MA, Lamprecht R, Nader K, and LeDoux J (2002) A-kinase anchoring proteins in amygdala are involved in auditory fear memory. *Nature Neurosci* **5**: 837–838.
- Myers KM and Davis M (2002) Behavioral and neural analysis of extinction. *Neuron* **36**: 567-584.
- Quirk GJ, Likhtik E, Pelletier JG, and Paré D (2003) Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *J Neurosci* **23**: 8800-8807.
- Quirk GJ, Russo GK, Barron JL, and Lebron K (2000) The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *J Neurosci* **20**: 6225-6231.
- Ressler KJ, Rothbaum BO, Tannenbaum L, Anderson P, Graap K, and Zimand E (2004) Cognitive enhancers`as adjuncts to psychotherapy: use of D-cycloserine in phobic individuals to facilitate extinction of fear. *Arch Gen Psychiatry* **61**: 1136-1144.
- Rumbaugh G, Sia GM, Garner CC, and Huganir RL (2003) Synapse-associated protein-97 isoform-specific regulation of surface AMPA receptors and synaptic function in cultured neurons. *J Neurosci* **23**:4567–4576.
- Schluter OM, Xu W, and Malenka RC (2006) Alternative N-terminal domains of PSD-95 and SAP97 govern activity-dependent regulation of synaptic AMPA receptor function. *Neuron* **51**: 99-111.
- Schnell E, Sizemore M, Karimzadegan S, Chen L, Breidt DS, and Nicoll RA (2002) Direct interactions between PSD-95 and stargazin control synaptic AMPA receptor number. *Proc Natl Acad Sci USA* **99**: 13902-13907.

MOL #53728

- Snyder EM, Colledge M, Crozier RA, Chen WS, Scott JD, and Bear MF (2005) Role for A kinase-anchoring proteins (AKAPS) in glutamate receptor trafficking and long-term synaptic depression. *J Biol Chem* **280**: 16962-16968.
- Sotres-Bayon F, Bush DE, and LeDoux JE (2004) Emotional perseveration: an update on prefrontal-amygdala interactions in fear extinction. *Learn. Mem* **11**: 525-535.
- Taylor S, Thordarson DS, Maxfield L, Fedoroff IC, Lovell K, and Ogradniczuk J (2003) Comparative efficacy, speed, and adverse effects of three PTSD treatments: exposure therapy, EMDR, and relaxation training. *J Consult Psychol* **71**: 330-338.
- Thompson AM, Walker VE, and Flynn DM (1989) Glycine enhances NMDA receptor-mediated synaptic potentials in neocortical slices. *Nature* **338**: 422-424.
- Ungless MA, Whistler JL, Malenka RC, and Bonci A (2001) Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* **411**: 583-587.
- Walker DL, Ressler KJ, Lu KT, and Davis M (2002) Facilitation of conditioned fear extinction by systemic administration or intra-amygdala infusions of D-cycloserine as assessed with fear-potentiated startle in rats. *J Neurosci* **22**: 2343-2351.
- Westergren I, Nystrom B, Hamberger A, Nordborg AC, and Johansson BB (1994) Concentrations of amino acids in extracellular fluid after opening of the blood-brain barrier by intracarotid infusion of protamine sulfate. *J Neurochem* **62**: 159-165.
- Wright CI, Fischer H, Whalen PJ, McInemey SC, Shin LM, and Raugh SL (2001) Differential prefrontal cortex and amygdala habituation to repeatedly presented emotional stimuli. *NeuroReport* **12**: 379-383.
- Yehuda R (2002) Post-traumatic stress disorder. *N Eng J Med* **346**: 108-114.

MOL #53728

Footnotes

This study was supported by [Grants N08I97N] from the National Health Research Institute, [NSC94-2752-B-006-001-PAE] from the National Science Council, and a Landmark Project (A0031) of National Cheng-Kung University of Taiwan.

MOL #53728

Figure Legends

Figure 1. Blocking of GlyT1 selectively enhances the NMDA-mediated synaptic response.

A: The graph represents the mean \pm SE slope of fEPSPs plotted against time. Application of NFPS (1 μ M, n=8) reversibly increased the slope of fEPSP. B: The effect of NFPS was blocked in the presence of D-APV (50 μ M, n=6). C: NMDA receptor-mediated fEPSP (fEPSP_{NMDA}) was isolated pharmacologically by using Mg⁺⁺-free solution in the presence of CNQX (10 μ M). Application of NFPS (n=6) increased the amplitude of fEPSP_{NMDA}. At end of the experiment, D-APV (50 μ M) was applied to ensure that fEPSP_{NMDA} was mediated by the NMDA receptors. Calibration: 1 mV, 20 ms.

Figure 2. Augmentation of extinction by NFPS.

Rats were conditioned and 24 h later were administered light-alone trials. The percentage of fear-potentiated startle was measured 24 h after light-alone trials. NFPS (5 μ g per side, 0.8 μ l) or vehicle (50% DMSO, 0.8 μ l) was infused bilaterally into the amygdala 30 min before the light-alone trials. The effect of NFPS plus extinction training was completely abolished when NMDA receptor antagonist D-APV (12.5 nmol/side) was co-applied with NFPS (n=6). *** $P < 0.001$ vs. Paired; ## $P < 0.01$ vs. Ext+Veh.

Figure 3. Partial reinstatement after US alone presentations in rats treated with NFPS plus extinction training.

MOL #53728

A: Rats were administered 10 light–shock pairings and 24 h later were given intra-amygdala infusion of NFPS (5 μ g, n=8) or vehicle (n=8) 30 min before extinction training. Memory retention was tested 24 h later (day 3, Test 1). One hour later, the rats were administered 10 US-alone trials and retention of memory was tested 24 h later (day 4, Test 2). B: The cannulae tip placements in the rats infused with NFPS (●) or vehicle before extinction (▲).

Figure 4. GlyT1 blockers facilitate NMDA-induced reduction in the surface levels of GluR1 and GluR2.

Amygdala slices were treated with 20 μ M of NMDA, 10 μ M of NMDA, 10 μ M of NMDA plus 1 μ M NFPS (A), or 10 μ M of NMDA plus 1 mM sarcosine (B) for 3 min. One hour later, LA and BLA tissues were dissected out and the surface GluR1 (A and B) (n=7 per group) and GluR2 (C) (n=8 per group) levels were determined using biotin labeling. Control slices were incubated in ACSF for 1 hr. GluR1 and GluR2 levels in the NMDA- and NFPS-treated slices were expressed as a percentage of those in the control slices. *** p <0.001 vs. Control; ### P <0.001 vs. NMDA 10 μ M.

Figure 5. Extinction training plus NFPS reverses the conditioning-induced increase in the surface expression of GluR1 and GluR2.

Rats were conditioned and randomly assigned to 3 groups, Extinction, Extinction+vehicle and Extinction+NFPS on day 1. On day 2, the rats in the Extinction group were given light-alone trials (extinction training) while the Extinction+NFPS rats were infused with NFPS (5 μ g per side) bilaterally into the amygdala 30 min before extinction training. Twenty-four hours after

MOL #53728

extinction training, LA and BLA tissues were dissected out and the surface GluR1 (A) (n=6 per group) and GluR2 (B) (n=6 per group) levels were determined using biotin labeling. A separate group of rats was infused with HA-966 (6 mg/kg, i.p.) 30 min before NFPS; 30 min later, light-alone trials were administered. Co-treatment of NFPS with light-alone trials blocked the conditioning-induced increases in the surface expression of GluR1 and GluR2. The GluR1 and GluR2 levels in the conditioned animals were expressed as a percentage of those of the naïve controls that did not undergo light–shock pairings. *** $P < 0.001$ vs. Extinction.

Figure 6. NFPS facilitates NMDA-induced reduction in the cellular levels of PSD-95/SAP97 and reverses the conditioning-induced increase in PSD-95/SAP97 in conjunction with extinction training.

Amygdala slices were treated with 20 μ M of NMDA, 10 μ M of NMDA, or 10 μ M of NMDA plus 1 μ M NFPS for 3 min. One hour later, LA and BLA tissues were dissected out and PSD-95 (A) (n=6 per group) and SAP97 (B) (n=6 per group) levels were determined by Western blotting analysis. Control slices were incubated in ACSF for 1 hr. *** $P < 0.001$ vs. Control; ## $P < 0.01$ vs. NMDA 10 μ M. C and D: Rats were conditioned and randomly assigned to 3 groups, Extinction, Extinction+vehicle and Extinction+NFPS on day 1. On day 2, the rats in the Extinction group were given light-alone trials (extinction training) while the Extinction+NFPS rats were infused with NFPS (5 μ g per side) bilaterally into the amygdala 30 min before extinction training. Twenty-four hours after extinction training, LA and BLA tissues were dissected out and PSD-95 (C) (n=6 per group) and SAP97 (D) (n=6 per group)

MOL #53728

levels were determined. A separate group of rats was infused with HA-966 (6 mg/kg, i.p.) 30 min before NFPS; 30 min later light-alone trials were given. Co-treatment of NFPS with light-alone trials blocked the conditioning-induced increases in PSD-95 and SAP97. PSD-95 and SAP97 levels in the conditioned animals were expressed as a percentage of those in the naïve controls that did not undergo light–shock pairings. *** $P < 0.001$ vs. Extinction.

Figure 7. Extinction training plus NFPS reverses the conditioning-induced increase in the AMPA/NMDA ratio.

A: Rats received 10 light–shock pairings and 24 h later were injected intraperitoneally with NFPS (1 or 10 mg/kg) or vehicle 30 min before extinction training (3 blocks of 10 presentations of light-alone trials). Retention of memory was assessed 24 h after extinction training. *** $P < 0.001$ vs. Paired; # $P < 0.05$ vs. Ext+Veh. B: Plot of the AMPA/NMDA ratio in slices taken from Naïve, Paired, Ext+Veh and Ext+NFPS (10 mg/kg, ip) animals (n=6 per group). *** $P < 0.001$ vs. Naïve; ### $P < 0.001$ vs. Ext+Veh. Scale: 50 ms, 100 pA.

Figure 8. Tat-GluR_{23Y} specifically blocks the effect of NFPS on extinction.

A: Rats were divided into 4 groups: those that received Tat- GluR_{23Y} (15 pmol in 0.8 µl saline per side) or Tat- GluR_{23A}, followed by NFPS (10 mg/kg, i.p.) or vehicle administered bilaterally into the amygdala 30 min before extinction training. Retention of memory was assessed 24 h after extinction training. B: The cannulae tip placements in the rats infused with Tat-GluR_{23Y}+NFPS (●), Tat-GluR_{23Y}+vehicle (▲), Tat-GluR_{23A}+NFPS (■) or Tat-GluR_{23A}+vehicle (★) 30 min before extinction training.




Figure 1




Figure 2





Figure 3






Figure 4




Figure 5







Figure 6




Figure 7

