

MOL #58677

Title Page

Selective and Brain Penetrant Neuropeptide Y Y2 Receptor Antagonists Discovered by Whole-Cell High Throughput Screening

Shaun P. Brothers, S. Adrian Saldanha, Timothy P. Spicer, Michael Cameron, Becky A. Mercer, Peter Chase, Patricia McDonald, Claes Wahlestedt and Peter S. Hodder

¹ Department of Neuroscience (S.P.B., C.W.), ²Lead Identification Division, Scripps Research Institute Molecular Screening Center (S.A.S., T.P.S., B.A.M., P.C., P.S.H.), ³Drug Metabolism and Pharmacokinetics (M.C.), ⁴Discovery Biology (P.M.), and ⁵Department of Molecular Therapeutics (P.S.H.), The Scripps Research Institute, Jupiter, Florida; * these authors contributed equally to this work (S.P.B., S.A.S).

MOL #58677

Running Title Page

Running Title:

Novel Y2 Receptor Antagonists

Corresponding Author:

Peter S. Hodder, Ph.D.
Scripps Florida
130 Scripps Way #1A1
Jupiter, FL 33458
Ph. (561) 228-2100
Fax (561) 228-3054
Email: hodderp@scripps.edu

Number of text pages: 40

Number of tables: 5

Number of figures: 5

Number of supplemental figures: 1

Number of references: 40

Number of words in the *Abstract*: 204

Number of words in the *Introduction*: 667

Number of words in the *Discussion* section: 1626

List of nonstandard abbreviations:

Abbreviation	Description
Y2R	neuropeptide Y Y2 receptor
Y1R	NPY-Y1 receptor
¹²⁵ I-PPY	radiolabeled peptide YY
MLPCN	Molecular Probe Production Center Network
MLSMR	Molecular Libraries Small Molecule Repository
CNS	central nervous system
HTS	high throughput screening
GPCR	G protein coupled receptor
Z'	Z-factor

MOL #58677

Abstract

The role of neuropeptide Y Y2 receptor (Y2R) in human diseases such as obesity, mood disorders and alcoholism could be better resolved by use of small molecule chemical probes that are substantially different from the currently available Y2R antagonist, BIIE0246. Presented here are five potent, selective and publicly available Y2R antagonists identified by a high throughput screening (HTS) approach. These compounds belong to four chemical scaffolds that are structurally distinct from the peptidomimetic BIIE0246. In functional assays, IC_{50} values between 199 nM and 4400 nM against the Y2R were measured, with no appreciable activity against the related NPY-Y1 receptor (Y1R). Compounds also displaced radiolabeled peptide YY (^{125}I -PPY) from the Y2R with high affinity (K_i values between 1.55 nM and 60 nM), while not displacing the same ligand from the Y1R. In contrast to BIIE0246, Schild analysis with NPY suggests that two of the five compounds behave as competitive antagonists. Profiling against a panel of 40 receptors, ion channels, and transporters found in the central nervous system showed that the five Y2R antagonists demonstrate greater selectivity than BIIE0246. Furthermore, the ability of these antagonists to penetrate the blood-brain barrier makes them better suited for pharmacologic studies of Y2R function in both the brain and periphery.

MOL #58677

Introduction

Neuropeptide Y (NPY) is one of the most abundant neuropeptides in the mammalian brain (Catapano and Manji, 2007) and was originally identified in 1982 based on structural similarity to peptide YY (70% homology with PYY) and pancreatic polypeptide (50% homology with PP) (Tatemoto, 1982). Each of these 36 residue peptides is processed from a 94-95 residue prohormone (Balasubramaniam, 1997). Four related type 1 G protein coupled receptor (GPCR) family members (Y1, Y2, Y4 and Y5) mediate biological responses to NPY. These receptors couple to the $G_{\alpha i}$ signaling pathway and inhibit adenylate cyclase, preventing cAMP production.

While NPY receptors have mainly been of interest to the pharmaceutical industry as therapeutic targets for obesity and feeding disorders, NPY has numerous roles in the body including modulation of feeding behavior, circadian rhythm, learning, memory, vascular remodeling, cell proliferation, and angiogenesis (Tatemoto, 1982). Human studies suggest that NPY is involved in mood disorders as well as in alcoholism (Thorsell et al., 2006; Wahlestedt et al., 1989). In patients with major depression, several studies show that NPY levels in the central nervous system (CNS) are low and correlate inversely to anxiety (Heilig, 2004; Nikisch et al., 2005; Widerlov et al., 1988). In addition to major depressive disorders, human genetic studies show that polymorphisms in the NPY gene also associate with alcoholism (Mottagui-Tabar et al., 2005; Wahlestedt et al., 1986; Zhu et al., 2003).

The NPY-Y2 receptor (Y2R) is located presynaptically on neurons, and in part serves to negatively regulate neurotransmitter release in the brain (King et al., 1999). Antagonism of Y2R would thus be expected to increase NPY levels in the CNS and may prove useful in treating certain psychiatric diseases. Consistent with this, Y2R null mice demonstrate reduced anxiety-like behavior compared to wild type (Redrobe et al., 2003). In addition, administration of the

MOL #58677

Y2R antagonist BIIE0246 decreased ethanol consumption in rats and induced antidepressant-like effects in mice (Rimondini et al., 2005).

Few efforts have focused on developing Y2R antagonists, with only two selective Y2R antagonists being reported other than BIIE0246: the peptide ligand, T4-[NPY 33-36]₄ (Grouzmann et al., 1997) and the small molecule JNJ-5207787 (Bonaventure et al., 2004). Small molecule diamines have also been reported as Y2R ligands. However these compounds have not been shown to be functionally active nor selective for the Y2R (Andres et al., 2003).

The most widely used Y2R antagonist, BIIE0246, is selective with high receptor affinity and is efficacious both *in vitro* and *in vivo*, (Doods et al., 1999). However, there are significant drawbacks associated with this compound. For instance it has been suggested that BIIE0246 behaves as an insurmountable antagonist (Dautzenberg and Neysari, 2005), an undesirable trait for a probe or potential therapeutic agent, since this mode of inhibition might block the Y2R from further activation which would lead to chronic loss of Y2R function. As a consequence, studies with BIIE0246 are limited to *in vitro* binding and functional determinations of Y2R activity. Furthermore, this peptidomimetic is big (molecular weight of 896) with a large polar surface area (219 Å²), both factors contributing to poor brain penetration. For example, experiments performed *in vivo* require direct injection of BIIE0246 to the desired site (Abbott et al., 2005; Rimondini et al., 2005). Similar challenges exist for the studies involving peptides such as T4-[NPY 33-36]₄. In contrast, the small molecule, JNJ-5207787, has improved properties over T4-[NPY 33-36]₄ and BIIE0246. For example, *ex vivo* receptor autoradiography studies reveal that JNJ-5207787 is able to cross the blood brain barrier and occupy Y2R receptor binding sites (Bonaventure et al., 2004). However, the lack of commercial availability for JNJ-520778 prevents further investigation into its pharmacology.

MOL #58677

As a result of the drawbacks of current antagonists, there is a need to identify brain-penetrant Y2R antagonists to better define the role of the receptor in the CNS. This report presents five novel, selective, brain penetrant small molecule Y2R antagonists, discovered from a high throughput screening (HTS) approach, that appear to have a different mechanism of action from BIIE0246 and are better suited for neuropharmacology studies.

MOL #58677

Materials and Methods

Unless otherwise stated, reagents were purchased from commercial sources. The cAMP biosensor assay, ACTOne Membrane Potential assay kit (catalog # BD354663), was purchased from BD Biosciences (Rockville, MD USA). Puromycin, Ro 20-1724, and isoproterenol were from Sigma (St. Louis, MO USA). Neuropeptide Y was from American Peptide (Sunnyvale, CA USA). BIIE0246 and BIBP3226 were from Tocris Bioscience (Ellisville, MO USA). Human liver microsomes were from Xenotech (Lenexa, KS USA). Black, clear bottom 1536-well tissue culture-treated microtiter plates were from Greiner Bio-One (Longwood, FL USA). The NIH Molecular Libraries Small Molecule Repository (MLSMR, San Francisco, CA USA) provided compound collections (dissolved at concentrations up to 10 mM in DMSO) for the primary and follow-up assays. For dose-response assays, compounds were provided by the MLSMR or ordered as powders and prepared as 10 mM stock solutions in DMSO.

For studies involving powders, the compounds SF-11 (catalog #7354811), SF-21 (catalog # 7955494) and SF-22 (catalog # 7930722) were obtained from ChemBridge (San Diego, CA, USA). The compound SF-31 (catalog # K907-0250) and SF-41 (catalog # 41069539) were purchased from Chemical Diversity (San Diego, CA, USA) and Specs (Delft, Holland), respectively. Reference and ordering information for all compounds can be obtained from PubChem website (<http://pubchem.ncbi.nlm.nih.gov/>) by conducting queries with the following substance identifier (SID): SF-11 = SID 17507305, SF-12 = SID 17431723, SF-13 = SID 56365809, SF-14 = SID 17433143, SF-15 = SID 17505667, SF-16 = SID 56365810, SF-21 = SID 17413392, SF-22 = SID 17413034, SF-23 = SID 56365811, SF-24 = SID 17412946, SF-31 = SID 4242079, SF-32 = SID 4255014, SF-33 = SID 14736723, SF-34 = SID 56365812, SF-35 = SID 4241999, SF-36 = SID 56365813, SF-37 = SID 4246387, SF-38 = SID 4244380, SF-39 = SID 14731189, SF-41 = SID 22413249 and SF-42 = SID 22413387.

MOL #58677

International Union of Pure and Applied Chemistry (IUPAC) names for the compounds tested in this work are: BIIE0246 = N-[(1S)-4-[(Aminoiminomethyl)amino]-1-[[[2-(3,5-dioxo-1,2-diphenyl-1,2,4-triazolidin-4-yl)ethyl]amino]carbonyl]butyl]-1-[2-[4-(6,11-dihydro-6-oxo-5H-dibenz[b,e]azepin-11-yl)-1-piperazinyl]-2-oxoethyl]-cyclopentaneacetamide), BIBP3226 = (2R)-5-(diaminomethylideneamino)-2-[(2,2-diphenylacetyl)amino]-N-[(4-hydroxyphenyl)methyl]pentanamide, SF-11 = N-(4-ethoxyphenyl)-4-[hydroxy(diphenyl)methyl]piperidine-1-carbothioamide, SF-12 = 4-[hydroxy(diphenyl)methyl]-N-(4-methoxyphenyl)piperidine-1-carbothioamide, SF-13 = N-(4-chlorophenyl)-4-[hydroxy(diphenyl)methyl]piperidine-1-carbothioamide, SF-14 = N-(3,5-dimethoxyphenyl)-4-[hydroxy(diphenyl)methyl]piperidine-1-carbothioamide, SF-15 = 5-dimethoxyphenyl)-4-[hydroxy(diphenyl)methyl]piperidine-1-carbothioamide, SF-16 = N-(4-fluorophenyl)-4-[hydroxy(diphenyl)methyl]piperidine-1-carbothioamide, SF-21 = 4-chloro-3-[(2,5-dimethylphenyl)sulfamoyl]-N-(2-phenylphenyl)benzamide, SF-22 = N-2-biphenyl-3-[(2,5-dimethylphenyl)amino]sulfonyl)-4-methylbenzamide, SF-23 = 4-methyl-N-(2-phenylphenyl)-3-(phenylsulfamoyl)benzamide, SF-24 = 4-chloro-3-[(2-methylphenyl)sulfamoyl]-N-(2-phenylphenyl)benzamide, SF-31 = 2-(2-methoxyphenyl)-N-[4-[5-(3-methoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, SF-32 = N-[4-[5-(2-ethoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]-2-(2-methoxyphenyl)acetamide, SF-33 = 2-(4-methoxyphenyl)-N-[4-[5-(2-methoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, SF-34 = N-[4-[5-(4-fluorophenyl)-1,2,4-oxadiazol-3-yl]phenyl]-2-(2-methoxyphenyl)acetamide, SF-35 = N-[4-[5-(2-ethoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]-2-(4-methoxyphenyl)acetamide, SF-36 = N-[4-[5-(4-chlorophenyl)-1,2,4-oxadiazol-3-yl]phenyl]-2-(2-methoxyphenyl)acetamide, SF-37 = 2-(4-methoxyphenyl)-N-[4-[5-(3,4,5-trimethoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, SF-38 = 2-(4-methoxyphenyl)-N-[4-[5-(3-methoxyphenyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, SF-39 = N-[4-[5-(4-chlorophenyl)-1,2,4-oxadiazol-3-yl]phenyl]-2-(4-methoxyphenyl)acetamide, SF-41 = 3,5-dimethyl-4-[(4-

MOL #58677

methylphenyl)sulfonyl-phenylmethyl]-1,2-oxazole and SF-42 = 4-[benzenesulfonyl(phenyl)methyl]-3,5-dimethyl-1,2-oxazole.

Cell Culture. The cAMP biosensor assay cell lines were purchased from BD Biosciences (Rockville, MD USA) as HEK293 cells stably expressing a cyclic nucleotide-gated (CNG) channel and either Y2R (catalog # BD344870) or Y1R (catalog # BD344869). Cells were cultured in T-175 cm² flasks at 37 C and 95% relative humidity. Cells were plated and maintained in growth medium consisting of DMEM supplemented with 10% heat-inactivated fetal bovine serum, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 25 mM HEPES, 5 mM L-glutamine, 250 µg/mL geneticin, 1 µg /mL puromycin and 1% antibiotic mix containing penicillin, streptomycin and neomycin.

Primary HTS of the MLSMR Compound Collection. As part of the Molecular Libraries Probe Production Network (MLPCN, <http://mli.nih.gov/mli/>) program, the HTS campaign was executed on the automated Kalypsys robotic platform located at the Scripps Research Institute Molecular Screening Center (SRIMSC; Jupiter, FL). Screening results for the Y2R (PubChem AID 793) and Y1R (PubChem AID 1040) antagonist primary assays, and all subsequent assays within the HTS screening campaign have been deposited into the MLPCN database, PubChem (<http://pubchem.ncbi.nlm.nih.gov>). It is important to note that the MLSMR is constantly adding and removing compounds to the MLPCN screening library, and a net of ~50,000 compounds were added to the library between the Y2R and Y1R primary screens. Both primary assays were screened against the full MLPCN screening library, which consisted of either 140,094 compounds (Y2R assay) or 196,180 compounds (Y1R assay).

Y2R Primary Campaign Protocol. For Y2R HTS assays, HEK293-CNG cells were diluted in growth medium and dispensed into 1536-well black-wall, clear-bottom plates (final

MOL #58677

concentration: 3,750 cells/well) and allowed to incubate for 24 hours at 37 C. Next, 4.5x concentrated membrane potential dye, prepared according to manufacturer instructions, was dispensed into each well. After incubating for 3 hours at room temperature, an initial fluorescent measurement (T0) was performed (510-545 nm excitation and 565-625 nm emission) using a FLIPR^{TETRA} fluorescence reader (Molecular Devices, Sunnyvale, CA USA). Test compounds (2.7 μ M final concentration), DMSO alone (0.27% final concentration), or BIIE0246 (1 μ M final concentration) was then added to sample or appropriate control wells, respectively. The cells were then challenged with NPY (100 nM final concentration), isoproterenol (freshly prepared; 1 μ M final concentration) and the phosphodiesterase inhibitor Ro-20-1724 (25 μ M final concentration). The plates were incubated for 30 minutes at room temperature before the final fluorescence measurement (T30) was taken, as described above.

Y1R Primary Campaign Protocol. The Y1R HTS screen was performed using a similar protocol as used for Y2R, with the following exceptions: A total of 3,600 cells per well were seeded into assay plates and allowed to grow for 24 hours at 37 C. After the T0 measurement, NPY (25 nM final concentration) was added to all wells, followed by the addition of test compounds (3.6 μ M final concentration) or DMSO to respective sample and control wells. Plates were incubated for 1 hour at room temperature, followed by addition of isoproterenol (freshly prepared; 1 μ M final concentration) and Ro 20-1724 addition (25 μ M final concentration). After 45 minutes at room temperature, the final fluorescent measurement was performed (T45).

Calculation of Antagonist Activity. The data from each campaign were normalized with the T0 measurement by dividing the fluorescence at T30 (for the Y2R) or T45 (for the Y1R) by the initial basal fluorescence at T0. The percent inhibition for each compound was determined using calculated ratios on a per-plate basis as follows:

MOL #58677

$$\% \text{ Inhibition} = 100 * [1 - ((\text{Test Compound} - \text{High Control}) / (\text{Low Control} - \text{High Control}))]$$

Where *Test Compound* indicates the ratio of fluorescence of wells containing test compound, NPY and isoproterenol. *Low Control* indicates the average ratio of fluorescence of wells containing DMSO, NPY and isoproterenol. *High Control* indicates the average ratio of fluorescence of wells containing either Y2R antagonist (BIIE0246), NPY and isoproterenol for the Y2R assays, or no NPY and isoproterenol for the Y1R assays.

HTS Hit Selection and Activity Cutoff Criteria. To determine inhibitory compounds in each primary screen, two values were calculated: (1) the average percent inhibition of all compounds tested and (2) three times their standard deviation (Hodder et al., 2003). The sum of these two values was used as a cutoff parameter, i.e. any compound that exhibited greater % inhibition than the cutoff value was considered active. Z' values were determined to monitor assay quality and were calculated as previously described (Zhang et al., 1999). Assay plates were rejected and rerun if Z' was less than 0.5.

HTS Hit Confirmation and Counterscreening Assays. Test compounds active in the primary HTS campaign were subsequently confirmed using the same assay protocol and hit cutoff value as used in the primary campaign, except that compounds were tested in triplicate. All confirmation and counterscreen assay data has been uploaded to the PubChem website (AID 1257: Y2R hit confirmation; AID 1256: Y1R counterscreen).

Dose-response Assays. The potency of compounds that passed confirmation and counterscreening was determined using dose-response assays and IC₅₀ values were calculated from the resultant data. A 10-point dose-response curve with a 1:3 dilution series from 35 μM to 1.8 nM was used, with compounds being tested in triplicate. Compounds exhibiting an IC₅₀

MOL #58677

value less than 10 μM were considered active. Dose-response counterscreens were also performed in Y1R cells to confirm that compounds identified as Y2R antagonists were selective. (PubChem AID 1272: Y2R dose-response determinations; AID 1279: Y1R dose-response determinations).

Binding Assays. Whole cell binding assays were performed as previously described (Brothers et al., 2006). Briefly, Y2R HEK293-CNG cells Y1R HEK293-CNG cells were cultured and plated in growth medium as described above, except that 10^5 cells in 0.5 mL growth medium were seeded in 24-well poly-D-lysine coated cell culture plates. After 24 h, cells were washed twice with a wash buffer consisting of 0.5 mL DMEM/0.1% BSA/10 mM HEPES. Next a range of concentrations of each antagonist was added to the cells, followed by 20 nM ^{125}I -PYY (final concentration) in 0.25 mL in the same buffer. After incubation at room temperature for 60 minutes (Nikisch et al., 2005), the supernatant was removed, the cells were washed twice and solubilized, and radioactivity of the lysate was determined. K_i values from ^{125}I -PYY displacement were determined using a one-site competition binding model and the Cheng-Prusoff equation (Cheng and Prusoff, 1973). A K_d of 0.27 nM for ^{125}I -PYY as previously reported (Rose et al., 1995) and a concentration of 20 nM for ^{125}I -PYY in the competition binding experiments were used for K_i determination.

Selectivity binding studies were conducted by the National Institute of Mental Health's Psychoactive Drug Screening Program (NIMH PDSP), Contract # NO1MH32004. The NIMH PDSP is directed by Bryan L. Roth MD, PhD at the University of North Carolina at Chapel Hill and Project Officer Jamie Driscoll at NIMH, Bethesda MD, USA. For methodological details, which are not reported here due to length considerations, please refer to the PDSP web site <http://pdsp.med.unc.edu/> where comprehensive protocols are described which are similar to those reported previously (Armbruster et al., 2007).

MOL #58677

Schild Analysis. A 12-point dose-response curve was performed in quadruplet with the cAMP biosensor assay in 384-well plates for NPY (1:3 dilution series from 1 μ M to 6.6 pM) and in the presence of five different concentrations of BIIE0246 (1 nM, 3 nM, 10 nM, 30 nM, 50 nM) or the five HTS lead compounds (100 nM, 300 nM, 1000 nM, 3000 nM, 5000 nM). Identical to HTS assay protocols, agonist (NPY) and antagonists were administered simultaneously to Y2R HEK293-CNG cells. The EC₅₀ values for NPY at varying antagonist concentrations were used for Schild analysis.

Hepatic Microsomal Stability Assays. Microsome stability was evaluated by incubating 1 μ M compound with 2 mg/mL hepatic microsomes from either human, rat, or mouse in 100 mM potassium phosphate buffer, pH 7.4. The reactions were held at 37 C with continuous shaking. The reaction was initiated by adding NADPH (1 mM final concentration) and the final incubation volume was 300 μ L. Aliquots (40 μ L) were removed at 0, 3, 5, 10, 20, and 30 min and added to 160 μ L acetonitrile containing an internal standard (propranolol) to stop the reaction and precipitate the protein. At the end of the assay, the samples were centrifuged through a 0.45 micron filter plate (Millipore Solventer low binding hydrophilic plates, catalog # MSRLN0450) and analyzed by LC-MS/MS. The data was log transformed and results are reported as half life.

Plasma protein binding experiments. An ultracentrifugation method was employed for evaluation of plasma protein binding plasma (Nakai et al., 2004). Plasma sample (1 mL with 1 μ M of test compound) was prepared and 900 μ L transferred to a 2 mL polycarbonate ultracentrifuge tube. The sample was centrifuged at 400,000 x g for 2 hours using a Beckman Coulter Optima Max ultracentrifuge (130,000 RPM max) with a TLA 120.2 rotor held at 25 C. The middle layer (2-3 mm below the surface using the described conditions) was used to determine the amount of

MOL #58677

unbound drug. The fraction of compound unbound to plasma proteins was determined using LC-MS/MS by comparison to the total amount of drug in the uncentrifuged sample.

Hepatic Clearance Calculations. *In vitro* intrinsic hepatic clearance was calculated from the microsomal incubation data to determine using methods similar to those previously described (Lipscomb and Poet, 2008). Specifically, the plasma protein binding data was combined with the results from the microsome stability experiments to predict the theoretical hepatic clearance and hepatic extraction ratio.

Brain Penetration Assays. Plasma and brain levels of the compounds were assessed in C57Bl6 mice 30 minutes after dosing 10 mg/kg intraperitoneally. Samples were formulated at 2 mg/mL in 10/10/80 DMSO/tween/water. Blood was collected into EDTA containing tubes at 30 min and plasma was generated using standard centrifugation techniques. Brain samples were frozen upon collection and all samples were stored at -80 C until analyzed. Brain tissue was not perfused prior to freezing to prevent diffusion of the compound out of the tissue during the process. Plasma samples were analyzed by treating 25 μ L of plasma with 125 μ L of acetonitrile containing an internal standard (propranolol) and filtering through a Millipore Multiscreen Solvinter 0.45 μ m low binding PTFE hydrophilic filter. The filtrate was analyzed by LC-MS/MS using an API Sciex 4000. MRM methods were developed in positive ion mode and concentrations were determined using a standard curve between 2 to 2000 ng/mL. Samples with concentrations outside of the curve were diluted with blank plasma and reanalyzed. Similar conditions were used to determine brain levels except the samples were weighed and acetonitrile was added (10x, weight by volume). The samples were sonicated to extract the compound from the brain matrix and then filtered as described above. A density of 1 g/mL was used to convert compound per mg tissue into molar equivalents.

MOL #58677

Cell Viability Assays. Cytotoxicity assays were performed as previously described (Madoux et al., 2008). Y2R HEK293-CNG cells were seeded at 500 cells per well in 1536-well plates in 5 μ L growth medium. Compounds (in DMSO) prepared as 10-point, 1:3 serial dilutions and added to cells (highest final concentration: 99 μ M). Plates were then incubated for 72 h at 37 C. After incubation, 5 μ L of CellTiter-Glo (Promega, Madison, WI USA) were added to each well and the plates were allowed to incubate for 15 min at room temperature. Luminescence was then measured (ViewLux plate reader, PerkinElmer, Turku, Finland). Viability was calculated as a percentage relative to control cells treated with either DMSO alone (0% cytotoxicity) or 100 μ M Doxorubicin (100% cytotoxicity).

Data Analysis and Statistics. Data were analyzed using MDL Assay Explorer (version 3.1, Santa Clara, CA USA) or GraphPad Prism (version 5.01, San Diego, CA USA). Curve fitting and IC₅₀ determinations were performed using the variable slope sigmoidal dose-response analysis tool in Prism. K_i values were determined using a one-site competition binding model. Replicates of at least 3 data points for each treatment group within an experiment were analyzed by unpaired, two-tailed t-test or using one way ANOVA followed by Tukey's test. Schild non-linear regression analysis was performed in Graphpad Prism using the Gaddum/Schild EC50 shift analysis tool. Statistical significance was ascertained by F-test comparing the calculated Schild slope to a hypothetical Schild analysis with slope of unity.

MOL #58677

Results

High Throughput Screening Assay for NPY Receptor Antagonists. GPCRs that couple to the $G_{\alpha i}$ signaling pathway, such as Y1R and Y2R, modulate intracellular cAMP concentrations via adenylate cyclase. Several HTS-compatible assays may be used to measure $G_{\alpha i}$ -coupled receptor activity, including use of GPCR-fluorescent fusion proteins (Milligan et al., 2004), reporter genes (Doucette et al., 2009) or measurement of cellular cAMP concentrations directly (Eglen, 2005). For the research presented here, an HTS-compatible cell-based cAMP assay was employed (Visegrady et al., 2007). In this assay format, measurement of $G_{\alpha i}$ protein coupled receptor antagonism in mammalian cells is facilitated by the presence of a “cAMP biosensor”, *i.e.* a modified cyclic nucleotide-gated (CNG) channel. Activation of the CNG channel by cAMP results in membrane depolarization. Depolarization is detected using a fluorescent membrane potential dye.

In the NPY receptor cAMP biosensor assay protocols (**Figure 1**), isoproterenol, a β -adrenergic receptor agonist, was used to induce intracellular cAMP production and therefore membrane depolarization via the activated CNG channel. The addition of NPY receptor agonist, through inhibition of adenylate cyclase, inhibits the accumulation of cAMP induced by isoproterenol. As designed, an antagonist would be expected to block NPY-agonized receptor signaling, leading to increased cAMP levels, decreased membrane polarization, and consequently increased dye fluorescence.

Y2R Antagonist HTS Campaign. To identify selective Y2R antagonists via a high-throughput screening (HTS) approach, Y2R and Y1R cAMP biosensor assays were developed in 1536 well plates in a total assay volume of 9 μ L/well. The assays utilized human embryonic kidney (HEK)

MOL #58677

cells expressing either the Y2R or the Y1R and the CNG channel (necessary to affect a membrane potential change). Experimental conditions were optimized to give the best balance between assay performance (determined by Z' factor), reagent consumption, and suitability of the protocol for robotic-based screening. Several parameters were optimized independently for the Y2R and Y1R antagonist assays such as cell number, NPY concentration and assay incubation times (see materials and methods section for final conditions).

Parallel Counterscreen Outcomes and Hit Confirmation. As part of a Molecular Probe Production Center Network (MLPCN) research effort (Lazo et al., 2007) the Y2R cAMP biosensor assay was screened against a collection of 140,094 compounds from the Molecular Libraries Small Molecule Repository (MLSMR). All MLSMR compounds were tested for antagonism at a nominal concentration of 2.7 μ M. Compounds that antagonized Y2R activity greater than the 17.08% inhibition cutoff (determined from the mean response plus three standard deviations for the entire compound data set) were considered active ("hits"). A total of 1384 compounds met this cutoff for the Y2R antagonist campaign.

The Y1R cAMP biosensor assay was employed as a parallel counterscreen during the Y2R screening campaign with the goal to streamline Y2R hit validation (**Figure 2**). Specifically, all compounds tested in the Y2R antagonist HTS assay were also tested in the Y1R antagonist HTS assay in order to remove cAMP biosensor assay-specific artifact and non-selective antagonists from further consideration. Of the 1384 compounds found active in the Y2R screen, 624 were also active in the Y1R screen. Removal of compounds found active in both assays left 760 potentially Y2R-selective antagonists. From this subset of compounds active in the Y2R assay, 707 compounds were available for further testing from the MLSMR. Fresh aliquots of these 707 compounds were retested at a single concentration (2.7 μ M) in triplicate using the

MOL #58677

same assay conditions as the original screen. A total of 228 of these compounds confirmed activity (i.e. inhibited Y2R activity greater than 17.08%). From these 228 confirmed hits, 123 were found to have significant activity when tested in triplicate at a single concentration (3.6 μ M) in the Y1R assay; these 123 compounds were not considered further.

As a final step in the Y2R screening campaign, 119 out of 125 Y2R selective compounds (with six unavailable from the MLSMR) were freshly aliquoted as 10-point titrations (1:3 serial dilutions from 35 μ M to 1.8 nM). Antagonism was evaluated over this concentration range in both the Y2R and the Y1R cAMP biosensor assays. From this set, 72 compounds antagonized the Y2R with an IC_{50} value less than 10 μ M, while 74 compounds antagonized the Y1R with an IC_{50} value less than 10 μ M (**Figure 3**).

Identification of selective Y2R Antagonists. Any compound chosen for further follow-up studies was required to exhibit an IC_{50} value of less than 5 μ M at Y2R and greater than 35 μ M at Y1R. Five compounds met or exceeded these criteria: SF-11, SF-21, SF-22, SF-31 and SF-41 (**Figure 3**). They fall into four chemotypes, classified as SF-1 through SF-4, representing piperidinecarbothioamide, arylsulfamoylbenzamide, aryl-1,2,4-oxadiazole and arylsulfonylmethylisoxazole scaffolds, and are not structurally similar to the known Y2R antagonists BIIE0246 and JNJ-5207787. Molecular weight, partition coefficient (as Log P), and polar surface area (PSA) were calculated for the five HTS-derived compounds as well as BIIE0246 (**Table 1**). The molecular weight of all five HTS-derived antagonists was less than 500; taken together, the logP and PSA values suggest that they are lipophilic and likely to be cell membrane permeable.

MOL #58677

Fresh powders of all compounds were characterized by LC-MS and retested at the Y2R and the Y1R. All compounds confirmed antagonism in the Y2R assay with IC_{50} values of less than 5 μ M, and in the case of SF-11, SF-21 and SF-22, the IC_{50} values were less than 1 μ M (**Figure 4A**). For all compounds tested, the maximum inhibition observed was within 20% of that observed for the positive control (BIIE0246). Furthermore, four compounds (SF-22, SF-31 and SF-41) were inactive in the Y1R assay even at the highest concentration tested (35 μ M, **Figure 4B**), while SF-21 exhibited minor but significant inhibition (10% at 35 μ M).

Y2R antagonist SAR. As a first step in evaluating structure activity relationships (SAR) for the newly identified Y2R antagonists, 16 close analogs of the five antagonists were procured. All were then tested for IC_{50} values in both the Y2R and Y1R cAMP biosensor assays (**Table 2**). Although all 16 analogs demonstrated activity at the Y2R, none of the newly obtained compounds demonstrated improved potency compared to the antagonists derived from HTS efforts. Additionally, four analogs belonging to the aryl-1,2,4-oxadiazole scaffold (SF-33, SF-34, SF-35 and SF-36) exhibited greater than 30% inhibition in the Y1R assay, with SF-34 being equipotent in both assays.

Assessment of Compound Cytotoxicity. All five HTS-derived Y2R antagonists and their 16 structural analogs were tested as 10-point titrations (in the range 5 nM to 99 μ M) for their effect on the viability of Y2R-expressing HEK cells (**Table 2**). After a 72 hour incubation, the five compounds identified by HTS (SF-11, SF-21, SF-22, SF-31 and SF-41) were toxic (in terms of their CC_{50} values) at concentrations that were 10 to 100-fold higher than their respective IC_{50} values for the Y2R. In general the 16 analogs were also toxic at compound concentrations significantly higher than their respective Y2R IC_{50} values. However, nine analogs (SF-16, SF-33,

MOL #58677

SF-34, SF-35, SF-36, SF-37, SF-38, SF-39) demonstrated less than a 10-fold difference between the Y2R IC₅₀ values and their corresponding CC₅₀ values.

Binding of ¹²⁵I-PYY to Cells Treated with Y2R-selective antagonists. Radioligand displacement studies were used to measure the ability of SF-11, SF-21, SF-22, SF-31 or SF-41 to block agonist binding. These five compounds were titrated for 8-point dose-response curves (1:6 serial dilutions from 50 μM to 1.8 pM), and radioligand displacement was evaluated over this concentration range in HEK cells expressing the Y2R or the Y1R. Calculations of K_i values were performed using the Cheng-Prusoff model (Cheng and Prusoff, 1973). All five compounds competed ¹²⁵I-PYY binding in Y2R-expressing cells with K_i values of 1.55 ± 0.93 nM for SF-11, 1.93 ± 0.86 nM for SF-21, 2.25 ± 0.49 nM for SF-22, 6.0 ± 2.8 nM for SF-31 and 60.3 ± 36.2 nM for SF-41, respectively (**Figure 4C**). In contrast, none of the tested compounds significantly blocked ¹²⁵I-PYY binding in Y1R expressing cells (**Figure 4D**).

Schild Analysis of BIIE0246 and Y2R selective compounds. To investigate the molecular pharmacology of the Y2R antagonists, Schild analysis was performed on BIIE0246 and the five HTS leads using the Y2R cAMP biosensor assay. NPY was used as the challenge. From these experiments, the Schild slopes were 2.79 ± 0.09 for BIIE0246, 0.79 ± 0.03 for SF-11, 0.98 ± 0.05 for SF-21, 1.3 ± 0.1 for SF-22, 1.55 ± 0.12 for SF-31, and 1.23 ± 0.08 for SF-41 (see **supplementary Figure 1**).

Metabolic stability experiments. The metabolic stability of SF-11, SF-21, SF-22, SF-31 and SF-41 along with BIIE0246 were evaluated by incubation with human, mouse, and rat liver microsomes (**Table 3**). The concentration of microsomal protein was normalized to 2 mg/mL from each species and exogenous NADPH was added to initiate the reactions. BIIE0246 demonstrated the longest half-life of all compounds tested. Among the five compounds

MOL #58677

identified by HTS, SF-31 had the longest half-life in all species. The shortest half-life in rodent and human microsomes was observed with SF-21 and SF-22 with half-lives of approximately 1 minute.

Plasma protein binding experiments and Hepatic Clearance calculations. An ultracentrifugation method was employed for evaluation of plasma protein binding plasma (Nakai et al., 2004); the fraction of compound unbound to plasma proteins was determined using LC-MS/MS. As noted in **Table 3**, the fraction of compound unbound to plasma proteins (f_u) was high except for SF-11. Additionally, *in vitro* intrinsic hepatic clearance ($Cl_{hepatic}$) was calculated using methods similar to those previously described (Lipscomb and Poet, 2008). Overall, BIIE0246 had the lowest hepatic clearance; the hepatic clearance of the 5 HTS-derived antagonists was high except for SF-11 and SF-31, where it was below ten (**Table 3**). The hepatic extraction ratio (calculated as percent extracted per pass) for the five HTS derived compounds were also higher than those calculated for BIIE0246.

Blood-brain barrier penetration studies. To determine whether the five antagonists identified by HTS were brain-penetrant, each compound was injected intraperitoneally into adult mice at 10 mg/kg and levels in the brain tissue and plasma were measured after thirty minutes. All five newly identified antagonists were highly brain penetrant with brain to plasma concentration ratios ranging between 36% (SF-21) and 115% (SF-41) (**Table 4**). At 30 minutes, SF-11, SF-21, SF-22, and SF-31 had brain concentrations above their respective cell-based IC_{50} values. High concentrations of BIIE0246 were detected in the plasma after IP dosing, but the achieved brain levels were only 2% of plasma levels. All compounds appeared to be well-tolerated, with the exception of SF-11 which caused significant lethargy in the mice.

MOL #58677

Selectivity profiling using the NIMH Psychoactive Drug Screening Program. The selectivity of SF-11, SF-21, SF-22, SF-31 and SF-41 was further assessed by the National Institute of Mental Health Psychoactive Drug Screening Program (NIMH-PDSP), where the compounds were screened for binding to receptors (N=35), ion channels (N=2), and transporters (N=3) typically found in the CNS (**Figure 5**). Initial screens were conducted in quadruplicate at 10 μ M final concentration as previously detailed (Armbruster et al., 2007). For comparison, BIIE0246 was also tested; JNJ-5207787 could not be tested due to the lack of availability of this compound. In the initial screen, BIIE0246 had significant activity at 15 targets in the screening panel, while the Y2R antagonists discovered in this HTS effort were active at fewer targets. Namely, significant radioligand displacement was observed with two targets for SF-11 (the 5HT_{2B} serotonin receptor and the dopamine transporter), two targets for SF-21 (the 5HT_{2B} receptor and the dopamine transporter), three targets for SF-22 (the 5HT_{2B} and 5HT₆ serotonin receptors as well as the dopamine transporter), and three targets for SF-41 (5HT_{2C}, 5HT₆ and the κ opioid receptor).

Dose-response curves were generated where significant displacement was observed and K_i values were determined (**Table 5**). BIIE0246 was found to bind three members of the panel with high affinity (defined as $K_i < 1 \mu$ M). In contrast, only one of the HTS-derived compounds, SF-22, demonstrated comparable potency, binding the 5HT_{2B} receptor with a K_i value of 250 nM. SF-31 did not significantly displace radioligand from any of the 40 members in the screening panel.

MOL #58677

Discussion

In the Y2R screening campaign, all compounds tested in the Y2R cAMP biosensor primary assay were also counterscreened in the Y1R cAMP biosensor primary assay (**Figure 2**). We have found this approach useful for rapidly prioritizing selective and potent chemical probes (Madoux et al., 2008). Since both the Y2R and Y1R assays used the same detection technology, the counterscreen was effective at removing non-specific GPCR agonists as well as compounds that non-specifically modulate GPCR signaling. For example, the G_s-coupled adenosine receptor agonist NECA ((Baraldi et al., 2006) PubChem SID 7975554) was found active in both the Y1R and Y2R primary screening campaigns. Similarly, the laxative bisacodyl (SID 855868), a known activator of adenylate cyclase (Ratnaik and Jones, 1998), was removed from further consideration.

The elimination of compounds found active at the Y1R during each successive screening stage of the hit-to-lead process resulted in the rapid identification of five selective Y2R antagonists (SF-11, SF-21, SF-22, SF-31 and SF-41), with the most potent, SF-11, antagonizing the Y2R with an IC₅₀ value of 199 nM (**Figure 4A**). While the five HTS antagonists were found to be Y2R selective in the cAMP biosensor dose-response assays, it was necessary to determine whether their efficacy was a result of displacement of agonist from the NPY receptor. Through the use of competitive binding studies all five antagonists and BIIE0246 were shown to effectively block ¹²⁵I-PYY binding to the Y2R. Their rank order potency was comparable to that found in the Y2R cAMP biosensor functional assay (**Figure 4A, Figure 4C**). Interestingly, BIIE0246 blocked ¹²⁵I-PYY binding with a K_i of 0.02 nM, lending further support to the notion that it behaves as an insurmountable antagonist (Dautzenberg and Neysari, 2005). Conversely, all five novel antagonists failed to block ¹²⁵I-PYY binding to the Y1R as high as 50 μM, confirming functional assay results (**Figure 4D**).

MOL #58677

Mechanism of action for the five antagonists as well as for BIIE0246 was investigated via Schild Analysis (Arunlakshana and Schild, 1959)(see supplementary Figure 1). It has been reported that BIIE0246 effectively competes with agonist and functionally inhibits the Y2R (Dautzenberg and Neysari, 2005) (Ziemek et al., 2006). However, Schild analysis has not been reported for BIIE0246, and when performed in this study resulted in a steep slope (2.79 ± 0.09). Since BIIE0246 is known to be an insurmountable antagonist when administered prior to Y2R agonists, this high slope suggests a more sophisticated mechanism of action for BIIE0246, such as a binding site for the antagonist that is different from the orthosteric (NPY) site. In contrast, Schild slopes for the five HTS compounds were close to unity. One compound, SF-21 resulted in a slope that is not statistically different from unity (0.98 ± 0.05 , $P = 0.62$), suggesting that it behaves as a true competitive antagonist, and binds to the orthosteric site. Although the slope of compound SF-22 is significantly different from unity (1.3 ± 0.1 , $P < 0.0001$), SF-21 and SF-22 are closely related, making it unlikely that they exhibit substantially different mechanisms of action. The three other antagonists exhibited Schild slopes that are close to unity, but nonetheless statistically different (namely 0.79 ± 0.03 and $P < 0.0001$ for SF-11, 1.55 ± 0.12 and $P < 0.0001$ for SF-31, and 1.23 ± 0.08 and $P = 0.0001$ for SF-41).

When assessed *in silico*, all five HTS antagonists exhibit several physical properties desirable for CNS probes. Compared to drugs that act in the periphery, brain-penetrant drugs tend to be more lipophilic and rigid, have fewer hydrogen bonds, fewer formal charges, and lower polar surface area (Mahar Doan et al., 2002; van de Waterbeemd et al., 1998). Optimal molecular properties for brain penetration have been proposed (van de Waterbeemd et al., 1998), namely that desirable compounds exhibit a LogP between 1 and 4, molecular weight less than 450, and a polar surface area of less than 90 \AA^2 . All five Y2R antagonists display molecular properties

MOL #58677

close to these optimums, with the exception of a higher computed LogP value for all but SF-41 (**Table 1**).

Microsome stability studies were conducted for all five HTS antagonists as a first step towards evaluating the suitability of these compounds for CNS applications *in vivo*. To estimate the ability of the liver to remove compounds from the blood in the absence of confounding factors, *in vitro* intrinsic hepatic clearances were calculated from the microsome and protein plasma binding studies (Lipscomb and Poet, 2008). Studies in mouse, rat and human microsomes (**Table 3**) showed that across all species the antagonists exhibited moderate to low stability (in terms of $T_{1/2}$) when compared to BIIE0246. Among the identified Y2R antagonists, SF-21, SF-22, and SF-41 were rapidly metabolized in microsomal incubations and the predicted *in vivo* hepatic clearance is high. In contrast, SF-11 was rapidly metabolized by rat and mouse hepatic microsomes, but was highly bound to plasma protein (when 1 μ M SF-11 was added to mouse plasma, only 30 nM of the compound was unbound to plasma proteins). High plasma protein binding would be expected to decrease compound diffusion into hepatocytes and thus decrease the rate of elimination. SF-31 was the most stable of the identified antagonists in mouse microsomal incubations and had moderate plasma protein binding. Both SF-11 and SF-31 had predicted hepatic clearance values below ten and favorable hepatic extraction ratios.

Because the Y2R is implicated in neurological and behavioral disorders, mouse blood-brain barrier penetration experiments were conducted on all five HTS compounds and BIIE0246. In consideration of the predicted high hepatic extraction ratio of three of the antagonists, all compounds were dosed by intraperitoneal (IP) injection at 10 mg/kg and compound concentrations were evaluated in plasma and brain at 30 minutes. In general intraperitoneal dosing allows the compound to be absorbed over time and drug levels can be maintained over a short time period because metabolized compound is replenished by additional drug being

MOL #58677

absorbed from the IP space. As the IP reservoir is depleted and additional compound cannot be absorbed, drug concentrations can rapidly decrease. Therefore, the 30 minute time point was chosen due to the rapid hepatic metabolism predicted from the microsomal studies; continued absorption of compound from the intraperitoneal space maintained a higher plasma concentration and facilitated assessment of brain exposure.

Brain penetration for all five HTS antagonists was high, signifying their potential for *in vivo* neurological studies. Brain concentrations determined for all five compounds (0.4 μM to 4.5 μM) are comparable to levels achieved by the small molecule JNJ-5207787 (C_{max} of 2.6 μM at 30 min) (Bonaventure et al., 2004). Because of the potential for rapid hepatic metabolism, SF-21, SF-22, and SF-41 may not be suitable for rodent *in vivo* efficacy studies. Although a formal pharmacokinetic study was not conducted for the five compounds, plasma levels measured in the brain penetration experiment after intraperitoneal dosing in mice were consistent with the microsomal stability data.

The five antagonists identified by HTS were evaluated for their ability to displace radioligands from a panel of GPCRs (35), ion channels (2) and transporters (3) found in the CNS. Among these targets were dopamine, serotonin, muscarinic, histamine, adrenergic and opioid receptors (including subtypes), as well as dopamine, serotonin, and norepinephrine transporters, and rat (whole brain) GABA ion channels. BIIE0246, used as a control, displaced radioligand from 15 members of this panel at a test concentration of 10 μM , while each of the other compounds tested ≤ 3 panel members (**Figure 4**). In dose-response studies (**Table 5**), BIIE0246 displaced radioligand from three receptors with sub-micromolar potency: the $\alpha 1\text{A}$ adrenergic receptor ($K_i = 360 \pm 31$ nM) as well as the μ and κ opioid receptors ($K_i = 323 \pm 31$ nM and 948 ± 183 nM, respectively). In contrast, SF-22 was the only HTS compound that displaced radioligand with

MOL #58677

appreciable potency ($K_i = 255 \pm 26$ nM for the serotonin 5-HT_{2B} receptor). In summary, the results from selectivity profiling against this CNS panel provide further evidence for these newly discovered compounds as more selective Y₂R antagonists than the commonly used Y₂R antagonist, BIIE0246.

In an effort to better understand the structural requirements for antagonism and to ascertain whether the four identified chemotypes may be amenable to optimization by medicinal chemistry, 16 close structural analogs were tested for dose-dependent antagonism of Y₂R or Y₁R. While none of the analogs antagonized the Y₂R as well as the five original HTS compounds, all analogs did exhibit some significant Y₂R antagonism (**Table 2**), demonstrating that structural modification of the different chemotypes is possible without total loss of activity at the Y₂R. Off-target antagonism of the closely related Y₁R was either insignificant or low. In terms of CC₅₀ values, all 21 compounds exhibited relatively low cytotoxicity after 72 hours.

In summary, this report describes the use of cell-based high throughput screening (HTS) to discover selective Y₂R antagonists from a screening library available through the auspices of the NIH. The HTS compounds SF-11, SF-21, SF-22, SF-31, and SF-41 are small molecules that penetrate the brain and offer promising chemical probes for the studies of Y₂R antagonism both *in vitro* and *in vivo*. These compounds are structurally distinct from all previously reported NPY receptor ligands, and Schild analysis reveals a marked difference in mechanism of action between the peptidomimetic BIIE0246 and two of the five compounds. Further, these compounds have improved brain permeability over BIIE0246 and are comparable to JNJ-5207787. The antagonists display an improved selectivity profile compared to BIIE0246 when tested against a panel of CNS receptors, ion channels, and transporters found in the CNS. For these reasons, SF-11, SF-21, SF-22, SF-31 and SF-41 may prove to be valuable small molecule probes for further studies into the biology of the Y₂R in human diseases such as

MOL #58677

obesity, alcoholism, and psychiatric disorders. A continued medicinal chemistry effort is currently in progress to further optimize these compounds for CNS studies.

MOL #58677

Acknowledgements

Mr. Pierre Baillargeon and Mrs. Lina DeLuca (Lead Identification Division, Scripps Research Institute Molecular Screening Center, Scripps Florida) are thanked for their assistance with compound management. Dr. Louis Scampavia (Lead Identification Division, Scripps Research Institute Molecular Screening Center, Scripps Florida) is thanked for LC-MS analysis of the presented compounds. The helpful discussions with Bryan Roth (University of North Carolina at Chapel Hill) on the interpretation of PDSP binding data are appreciated. The support of Prof. Hugh Rosen (Scripps Research Institute, San Diego, CA) is acknowledged.

MOL #58677

References

- Abbott CR, Small CJ, Kennedy AR, Neary NM, Sajedi A, Ghatei MA and Bloom SR (2005) Blockade of the neuropeptide Y Y2 receptor with the specific antagonist BIIIE0246 attenuates the effect of endogenous and exogenous peptide YY(3-36) on food intake. *Brain Res* **1043**(1-2):139-144.
- Andres CJ, Antal Zimanyi I, Deshpande MS, Iben LG, Grant-Young K, Mattson GK and Zhai W (2003) Differentially functionalized diamines as novel ligands for the NPY2 receptor. *Bioorg Med Chem Lett* **13**(17):2883-2885.
- Armbruster BN, Li X, Pausch MH, Herlitz S and Roth BL (2007) Evolving the lock to fit the key to create a family of G protein-coupled receptors potentially activated by an inert ligand. *Proc Natl Acad Sci U S A* **104**(12):5163-5168.
- Arunlakshana O and Schild HO (1959) Some quantitative uses of drug antagonists. *Br J Pharmacol Chemother* **14**(1):48-58.
- Balasubramaniam AA (1997) Neuropeptide Y family of hormones: receptor subtypes and antagonists. *Peptides* **18**(3):445-457.
- Baraldi PG, Romagnoli R, Preti D, Fruttarolo F, Carrion MD and Tabrizi MA (2006) Ligands for A2B adenosine receptor subtype. *Curr Med Chem* **13**(28):3467-3482.
- Bonaventure P, Nepomuceno D, Mazur C, Lord B, Rudolph DA, Jablonowski JA, Carruthers NI and Lovenberg TW (2004) Characterization of N-(1-Acetyl-2,3-dihydro-1H-indol-6-yl)-3-(3-cyano-phenyl)-N-[1-(2-cyclopropyl-ethyl)-piperidin-4yl]acrylamide (JNJ-5207787), a small molecule antagonist of the neuropeptide Y Y2 receptor. *J Pharmacol Exp Ther* **308**(3):1130-1137.
- Brothers SP, Janovick JA and Conn PM (2006) Calnexin regulated gonadotropin-releasing hormone receptor plasma membrane expression. *J Mol Endocrinol* **37**(3):479-488.
- Catapano LA and Manji HK (2007) G protein-coupled receptors in major psychiatric disorders. *Biochim Biophys Acta* **1768**(4):976-993.
- Cheng Y and Prusoff WH (1973) Relationship between the inhibition constant (K₁) and the concentration of inhibitor which causes 50 per cent inhibition (I₅₀) of an enzymatic reaction. *Biochem Pharmacol* **22**(23):3099-3108.
- Dautzenberg FM and Neysari S (2005) Irreversible binding kinetics of neuropeptide Y ligands to Y2 but not to Y1 and Y5 receptors. *Pharmacology* **75**(1):21-29.
- Doods H, Gaida W, Wieland HA, Dollinger H, Schnorrenberg G, Esser F, Engel W, Eberlein W and Rudolf K (1999) BIIIE0246: a selective and high affinity neuropeptide Y Y(2) receptor antagonist. *Eur J Pharmacol* **384**(2-3):R3-5.
- Doucette C, Vedvik K, Koepnick E, Bergsma A, Thomson B and Turek-Etienne TC (2009) Kappa Opioid Receptor Screen with the Tango™ β -arrestin Recruitment Technology and Characterization of Hits with Second Messenger Assays. *Journal of Biomolecular Screening* **14**(4):381-394.
- Eglen RM (2005) Functional G protein-coupled receptor assays for primary and secondary screening. *Comb Chem High Throughput Screen* **8**(4):311-318.
- Grouzmann E, Buclin T, Martire M, Cannizzaro C, Dorner B, Razaname A and Mutter M (1997) Characterization of a selective antagonist of neuropeptide Y at the Y2 receptor. Synthesis and pharmacological evaluation of a Y2 antagonist. *J Biol Chem* **272**(12):7699-7706.
- Heilig M (2004) The NPY system in stress, anxiety and depression. *Neuropeptides* **38**(4):213-224.
- Hodder P, Cassaday J, Peltier R, Berry K, Inglese J, Feuston B, Culberson C, Bleicher L, Cosford ND, Bayly C, Suto C, Varney M and Strulovici B (2003) Identification of metabotropic glutamate receptor antagonists using an automated high-throughput screening system. *Anal Biochem* **313**(2):246-254.

MOL #58677

- King PJ, Widdowson PS, Doods HN and Williams G (1999) Regulation of neuropeptide Y release by neuropeptide Y receptor ligands and calcium channel antagonists in hypothalamic slices. *J Neurochem* **73**(2):641-646.
- Lazo JS, Brady LS and Dingledine R (2007) Building a pharmacological lexicon: small molecule discovery in academia. *Mol Pharmacol* **72**(1):1-7.
- Lipscomb JC and Poet TS (2008) In vitro measurements of metabolism for application in pharmacokinetic modeling. *Pharmacol Ther* **118**(1):82-103.
- Madoux F, Li X, Chase P, Zastrow G, Cameron MD, Conkright JJ, Griffin PR, Thacher S and Hodder P (2008) Potent, selective and cell penetrant inhibitors of SF-1 by functional ultra-high-throughput screening. *Mol Pharmacol* **73**(6):1776-1784.
- Mahar Doan KM, Humphreys JE, Webster LO, Wring SA, Shampine LJ, Serabjit-Singh CJ, Adkison KK and Polli JW (2002) Passive permeability and P-glycoprotein-mediated efflux differentiate central nervous system (CNS) and non-CNS marketed drugs. *J Pharmacol Exp Ther* **303**(3):1029-1037.
- Milligan G, Feng GJ, Ward RJ, Sartania N, Ramsay D, McLean AJ and Carrillo JJ (2004) G protein-coupled receptor fusion proteins in drug discovery. *Curr Pharm Des* **10**(17):1989-2001.
- Mottagui-Tabar S, Prince JA, Wahlestedt C, Zhu G, Goldman D and Heilig M (2005) A novel single nucleotide polymorphism of the neuropeptide Y (NPY) gene associated with alcohol dependence. *Alcohol Clin Exp Res* **29**(5):702-707.
- Nakai D, Kumamoto K, Sakikawa C, Kosaka T and Tokui T (2004) Evaluation of the protein binding ratio of drugs by a micro-scale ultracentrifugation method. *J Pharm Sci* **93**(4):847-854.
- Nikisch G, Agren H, Eap CB, Czernik A, Baumann P and Mathe AA (2005) Neuropeptide Y and corticotropin-releasing hormone in CSF mark response to antidepressive treatment with citalopram. *Int J Neuropsychopharmacol* **8**(3):403-410.
- Ratnaik RN and Jones TE (1998) Mechanisms of drug-induced diarrhoea in the elderly. *Drugs Aging* **13**(3):245-253.
- Redrobe JP, Dumont Y, Herzog H and Quirion R (2003) Neuropeptide Y (NPY) Y2 receptors mediate behaviour in two animal models of anxiety: evidence from Y2 receptor knockout mice. *Behav Brain Res* **141**(2):251-255.
- Rimondini R, Thorsell A and Heilig M (2005) Suppression of ethanol self-administration by the neuropeptide Y (NPY) Y2 receptor antagonist BIIIE0246: evidence for sensitization in rats with a history of dependence. *Neurosci Lett* **375**(2):129-133.
- Rose PM, Fernandes P, Lynch JS, Frazier ST, Fisher SM, Kodukula K, Kienzle B and Seethala R (1995) Cloning and functional expression of a cDNA encoding a human type 2 neuropeptide Y receptor. *J Biol Chem* **270**(39):22661-22664.
- Tatemoto K (1982) Neuropeptide Y: complete amino acid sequence of the brain peptide. *Proc Natl Acad Sci U S A* **79**(18):5485-5489.
- Thorsell A, Slawecki CJ, El Khoury A, Mathe AA and Ehlers CL (2006) The effects of social isolation on neuropeptide Y levels, exploratory and anxiety-related behaviors in rats. *Pharmacol Biochem Behav* **83**(1):28-34.
- van de Waterbeemd H, Camenisch G, Folkers G, Chretien JR and Raevsky OA (1998) Estimation of blood-brain barrier crossing of drugs using molecular size and shape, and H-bonding descriptors. *J Drug Target* **6**(2):151-165.
- Visegrady A, Boros A, Nemethy Z, Kiss B and Keseru GM (2007) Application of the BD ACTOne technology for the high-throughput screening of Gs-coupled receptor antagonists. *J Biomol Screen* **12**(8):1068-1073.
- Wahlestedt C, Ekman R and Widerlov E (1989) Neuropeptide Y (NPY) and the central nervous system: distribution effects and possible relationship to neurological and psychiatric disorders. *Prog Neuropsychopharmacol Biol Psychiatry* **13**(1-2):31-54.

MOL #58677

- Wahlestedt C, Yanaihara N and Hakanson R (1986) Evidence for different pre-and post-junctional receptors for neuropeptide Y and related peptides. *Regul Pept* **13**(3-4):307-318.
- Widerlov E, Lindstrom LH, Wahlestedt C and Ekman R (1988) Neuropeptide Y and peptide YY as possible cerebrospinal fluid markers for major depression and schizophrenia, respectively. *J Psychiatr Res* **22**(1):69-79.
- Zhang JH, Chung TD and Oldenburg KR (1999) A Simple Statistical Parameter for Use in Evaluation and Validation of High Throughput Screening Assays. *J Biomol Screen* **4**(2):67-73.
- Zhu G, Pollak L, Mottagui-Tabar S, Wahlestedt C, Taubman J, Virkkunen M, Goldman D and Heilig M (2003) NPY Leu7Pro and alcohol dependence in Finnish and Swedish populations. *Alcohol Clin Exp Res* **27**(1):19-24.
- Ziemek R, Brennauer A, Schneider E, Cabrele C, Beck-Sickinger AG, Bernhardt G and Buschauer A (2006) Fluorescence- and luminescence-based methods for the determination of affinity and activity of neuropeptide Y2 receptor ligands. *Eur J Pharmacol* **551**(1-3):10-18.

MOL #58677

Footnote

This work was supported by the National Institutes of Health [Grant R21NS056950-01S]
and [Grant U54MH084512].

MOL #58677

Legends for Figures

Figure 1. Principle of the cAMP biosensor assay used for the HTS campaign. (A) Addition of neuropeptide Y (NPY) receptor agonist stimulates the receptor, activates $G_{\alpha i}$ protein and decreases cAMP production via inhibition of adenylate cyclase (AC). Antagonism of the receptor increases production of cAMP by the same pathway, which serves to activate and open the cyclic nucleotide gated (CNG) ion channel. The relocation of extracellular Ca^{2+} and Na^+ from the extracellular space to the cytosol causes membrane depolarization. **(B)** A membrane potential fluorescent indicator (D) is used to measure the cell membrane polarization state. Prior to cellular depolarization, indicator dye molecules are mainly extracellular with minimal fluorescence. Upon depolarization, dye redistributes into the cytoplasm, becoming fluorescent.

Figure 2. Correlation plot of inhibition results from the Y2R and Y1R primary high-throughput screening (HTS) assays. Grey circles represent the response, expressed as percent inhibition, for each compound tested in the Y1R (y-axis) and Y2R (x-axis) antagonism-format cAMP biosensor assays. A total of 140,094 compounds were tested for inhibition in both the Y2R & Y1R primary HTS assays (details in text). The five HTS-derived Y2R antagonists presented in this manuscript are distributed in the lower quadrant of the plot, represented by the following symbols: ■ (SF-11), + (SF-21), ♦ (SF-22), • (SF-31), and ▲ (SF-41). The vertical and horizontal dashed lines represent the hit cutoff values of the Y2R (17.08%) and Y1R (20.11%) assays.

Figure 3. Correlation plot of results from the Y2R and Y1R dose-response assays. The IC_{50} values of 119 compounds are plotted for both the Y2R (x-axis) and Y1R (y-axis) cAMP biosensor assays. The five selective Y2R antagonists can be found in the upper part of the plot, represented by ■ (SF-11), + (SF-21), ♦ (SF-22), • (SF-31), and ▲ (SF-41). The five

MOL #58677

corresponding structures of antagonists, as well as previously described NPY receptor antagonists BIIE0246 & JNJ-5207787 (Y2R) and BIBP3226 (Y1R) are illustrated.

Figure 4. (A) Antagonist dose-response curves from the Y2R cAMP biosensor assay. Y2R antagonists SF-11 (■, $IC_{50}=199 \pm 10$ nM), SF-21 (+, $IC_{50}=440 \pm 30$ nM), SF-22 (◆, $IC_{50}=750 \pm 30$ nM), SF-31 (●, $IC_{50}=1200 \pm 100$ nM) and SF-41 (▲, $IC_{50}=4400 \pm 40$ nM) are graphed. For comparison BIIE0246's $IC_{50}=0.58 \pm 0.56$ nM. **(B) Antagonist dose-response curves from the Y1R cAMP biosensor assay.** Symbols for the different compounds are the same as those described in (A) above; For comparison, BIBP3226 (BIBP), a Y1R antagonist, is shown (x, $IC_{50}=3046 \pm 982$ nM). **(C) Results of ^{125}I -PYY displacement with Y2R-expressing cells.** Percent binding of symbols for the different compounds are the same as those described in (A). SF-11 ($K_i = 1.55 \pm 0.93$ nM), SF-21 ($K_i = 1.93 \pm 0.86$ nM), SF-22 ($K_i = 2.25 \pm 0.49$ nM), SF-31 ($K_i = 6.0 \pm 2.8$ nM) and SF-41 ($K_i = 60.3 \pm 36.2$ nM) versus concentration is graphed. As comparison $K_i = 0.02 \pm 0.01$ nM for BIIE0246. **(D) Results of ^{125}I -PYY displacement with Y1R-expressing cells.** Displacement was measured at a single high concentration (50 μ M) of Y1R selective antagonist (BIBP3226) or HTS-derived antagonist. Error bars on all graphs represent the standard deviation of at least three separate experiments.

Figure 5. Radioligand displacement screening assay results for 40 different receptors, ion channels and transporters found in the CNS. Measured percent activity vs. target name is graphed. All compounds were tested at 10 μ M. Inhibition greater than the 50% cutoff (dotted line) is considered significant. Negative binding values, obtained when non-specific binding was higher than specific binding, are reported as zero. Abbreviations are given for serotonin receptor subtypes (5HT1A-7); adrenergic receptor subtypes (Alpha1A-Beta3); rat brain GABA1 and GABA3 sites (BZP sites); dopamine receptor subtypes: (D2, D4) the dopamine (DAT),

MOL #58677

norepinephrine (NET) and serotonin (SERT) transporters; prostaglandin receptor subtypes (EP3, EP4); histamine receptor subtypes (H1, H2, H4); muscarinic receptor subtypes (M1-M5); opioid receptor subtypes (MOR, DOR, KOR) and the sigma 1 receptor.

MOL #58677

Table 1. Comparison of calculated physical properties for different Y2R antagonists.

ID	Molecular Weight	Log P	Polar Surface Area
SF-11	447	5.1	44.7
SF-21	471	6.5	75.3
SF-22	491	6.9	75.3
SF-31	415	4.7	86.5
SF-41	341	3.9	60.2
BIIE0246	896	4	219

MOL #58677

Table 2. Structure Activity Relationships for Y2R selective antagonists. SF numbers of compounds from HTS are underlined.

ID	R	IC ₅₀ (μM) (Max Inhibition)		CC ₅₀ (μM) (Max Toxicity)	ID	R1	R2	IC ₅₀ (μM) (Max Inhibition)		CC ₅₀ (μM) (Max Toxicity)
		Y2R	Y1R					Y2R	Y1R	
<u>SF-11</u>		0.199 ± 0.01	>35.4 (Inactive)	>33	<u>SF-21</u>	Cl		0.44 ± 0.03	>35.4 (10 ± 2%)	19.7 ± 7.2
SF-12		1.1 ± 0.1	>35.4 (5 ± 4%)	>99 (37 ± 8%)	<u>SF-22</u>	CH ₃		0.75 ± 0.03	>35.4 (Inactive)	61 ± 30
SF-13		1.3 ± 0.1	>35.4 (Inactive)	>33	SF-23	CH ₃		2.1 ± 0.2	>35.4 (Inactive)	>33
SF-14		2.99 ± 0.1	>35.4 (Inactive)	>99 (49 ± 6%)	SF-24	Cl		3.96 ± 0.3	>35.4 (Inactive)	>33
SF-15		4.9 ± 0.57	>35.4 (4 ± 3%)	>99 (41 ± 6%)						
SF-16		>35.4 (43 ± 5%)	>35.4 (Inactive)	>33						
					<u>SF-31</u>			1.2 ± 0.1	>35.4 (Inactive)	>99 (43 ± 7%)
<u>SF-41</u>		4.4 ± 0.04	>35.4 (Inactive)	>99 (35 ± 9%)	SF-32			4.9 ± 0.1	>35.4 (Inactive)	>99 (27 ± 11%)
SF-42		8.6 ± 0.5	>35.4 (Inactive)	>99 (9 ± 8%)	SF-33			5.5 ± 0.8	>35.4 (34 ± 4%)	30.8 ± 2.7
					SF-34			17.3 ± 6.7	17.3 ± 3.4	>33
					SF-35			>11.8	>11.8	17.2 ± 0.4
					SF-36			>35.4 (46 ± 4%)	>35.4 (36 ± 11%)	>11
					SF-37			>35.4 (45 ± 7%)	>35.4 (Inactive)	>33
					SF-38			>35.4 (43 ± 4%)	>35.4 (6 ± 3%)	>33
					SF-39			>35.4 (4 ± 2%)	>35.4 (Inactive)	22.9 ± 2.5

Downloaded from molpharm.aspetjournals.org at ASPET Journals on April 19, 2024

MOL #58677

Table 3. Results of microsome stability studies (expressed as $T_{1/2}$), protein binding experiments (expressed as fraction unbound, fu), calculated hepatic clearance ($Cl_{hepatic}$), and percent compound extracted per pass for five Y2R antagonists and BIIE0246.

ID	Species	$T_{1/2}$ (min)	fu	$Cl_{hepatic}$	% Extracted per pass
SF-11	Mouse	4	0.03	5.7	6.3
	Rat	3	0.02	6.1	11
	Human	10	0.01	0.3	1.4
SF-21	Mouse	1	0.82	90	100
	Rat	1	0.13	50	90
	Human	8	0.23	7	34
SF-22	Mouse	1	1	90	100
	Rat	1	0.19	53	96
	Human	6	0.28	10	49
SF-31	Mouse	14	0.12	6.4	7.2
	Rat	6	0.07	10	18
	Human	13	0.07	1.5	7.4
SF-41	Mouse	4	0.39	51	57
	Rat	1	0.19	53	96
	Human	3	0.19	12	60
BIIE0246	Mouse	166	0.079	0.4	0.4
	Rat	27	0.022	0.8	1.4
	Human	23	0.015	0.2	0.9

MOL #58677

Table 4. Results of blood-brain barrier penetration studies for five Y2R Antagonists and BIIE0246.

ID	Brain Penetration Studies		
	Plasma Concentration (μM)	Brain Concentration (μM)	Ratio [B]/[P] (%)
SF-11	8.5 \pm 3.3	3.7 \pm 1.7	44
SF-21	1.1 \pm 0.1	0.4 \pm 0.03	36
SF-22	1.6 \pm 0.6	1.7 \pm 0.6	106
SF-31	9.0 \pm 0.7	4.5 \pm 0.3	50
SF-41	1.3 \pm 0.3	1.5 \pm 0.2	115
BIIE0246	12 \pm 2	0.2 \pm 0.1	2

MOL #58677

Table 5. Radioligand displacement assay K_i values for BIIE0246 and five Y2R Antagonists. Target names correspond to those in Figure 5.

Target	K_i from PDSP (nM)					
	BIIE0246	SF-11	SF-21	SF-22	SF-31	SF-41
5ht2b	-	7093 ± 869	1212 ± 161	255 ± 26	-	-
5ht2c	-	-	-	-	-	9943 ± 1696
5ht5a	1361 ± 206	-	-	-	-	-
5ht6	-	-	-	2853 ± 419	-	2606 ± 359
Alpha1A	360 ± 31	-	-	-	-	-
Alpha1D	1294 ± 163	-	-	-	-	-
Beta3	1645 ± 160	-	-	-	-	-
D2	7294 ± 2509	-	-	-	-	-
D4	3151 ± 256	-	-	-	-	-
DAT	2222 ± 430	4989 ± 1047	5525 ± 987	4005 ± 676	-	-
H2	2979 ± 2979	-	-	-	-	-
M2	2452 ± 391	-	-	-	-	-
M3	1573 ± 280	-	-	-	-	-
M4	2232 ± 358	-	-	-	-	-
M5	1899 ± 231	-	-	-	-	-
MOR	323 ± 31	-	-	-	-	-
DOR	1132 ± 181	-	-	-	-	-
KOR	948 ± 183	-	-	-	-	1044 ± 57





Figure 1




Figure 2

Y2R Inhibition (%)








Figure 3







Figure 4




Figure 5