

MOL 64899

ROLE OF EPIGENETIC MECHANISMS IN DIFFERENTIAL REGULATION OF THE DIOXIN-INDUCIBLE HUMAN *CYP1A1* AND *CYP1B1* GENES.

**SUDHEER R. BEEDANAGARI, ROBERT T. TAYLOR, PETER BUI, FENG WANG, DEREK W.
NICKERSON, AND OLIVER HANKINSON**

Interdepartmental Molecular Toxicology Program, Department of Pathology and Laboratory
Medicine (S.R.B, P.B, F.W., O.H.), Jonsson Comprehensive Cancer Center, (O.H), Molecular
Biology Institute, (O.H.), and Department of Pediatrics, Hematology and Oncology, (D.W.N)
University of California, Los Angeles, California 90095, USA. Roche Pharmaceuticals, Nutly NJ
(R.T.T).

MOL 64899

Running title: Transcriptional regulation of CYP1A1 and CYP1B1 genes

To whom correspondence should be addressed:

Oliver Hankinson, Ph.D.

Departmental of Pathology and Laboratory Medicine, David Geffen School of Medicine,
University of California at Los Angeles.

650 Charles Young Drive, Los Angeles, CA. 90095

Tel: 310-825-2936 Fax: 310-794-9272 Email: ohank@mednet.ucla.edu

Text: 19 pages

Tables: 0

Figures: 10 Figures

Reference: 33 references

Abstract: 249 words

Introduction: 703 words

Discussion: 1138 words

Abbreviations: ChIP: Chromatin Immunoprecipitation, 5-AzaC : 5-aza-2'-deoxycytidine, AhR: Aryl
Hydrocarbon receptor, polII: RNA polymerase II, CBP: CREB binding protein, SRC-1, 2 and 3: Steroid
receptor coactivators 1,2, and 3,TBP: TATA binding protein.

ABSTRACT

The Aryl Hydrocarbon Receptor (Ahr) mediates induction of CYP1A1 and CYP1B1 by 2,3,7,8-tetrachlorodibenzo-p-dioxin (dioxin) via binding to Xenobiotic Responsive Elements (XREs) in their enhancer regions. CYP1A1 and CYP1B1 were both inducible by dioxin in human MCF-7 cells. However, only CYP1A1 was inducible in human HepG2 cells. Further experiments focused on providing an explanation for this last observation. Dioxin induced recruitment of AHR and the transcriptional coactivators p300 and PCAF to the *CYP1B1* enhancer in HepG2 cells, but failed to induce recruitment of RNA polymerase II (polII) or the TATA binding protein (TBP) and acetylations of histones 3 and 4 or methylation of histone 3 at the promoter. Since p300 was required for dioxin induction of the aforementioned histone modifications at the *CYP1B1* promoter and for induction of CYP1B1 transcription (in MCF-7 cells), the recruitments of p300 and AhR, although necessary, are not sufficient for eliciting the above responses to dioxin. Cytosine residues within CpG dinucleotides at the enhancer, including those within the XREs, were partially methylated, whereas those at the promoter were fully methylated. Treatment of HepG2 cells with 5-aza-2'-deoxycytidine led to partial demethylation of the promoter, and restored polII and TBP binding, and CYP1B1 inducibility. Thus the deficiency of CYP1B1 induction in HepG2 cells is ascribable to cytosine methylation at the promoter which prevents recruitment of TBP and polII. Importantly, our data indicate that stable recruitment of p300 and PCAF to the *CYP1B1* gene does not require their tethering to the promoter as well as to the enhancer.

INTRODUCTION

Dioxin (2,3,7,8-tetrachlorodibenzo-p-dioxin or TCDD), is a common persistent trace environmental pollutant and the most potent carcinogen known (Huff et al., 1994; Mandal, 2005). Dioxin induces a broad range of biological responses, including robust induction of the *CYP1* family of genes, via its binding to the aryl hydrocarbon receptor (AhR). Besides dioxin, AhR binds to a wide spectrum of environmental pollutants and xenobiotics, including polycyclic aromatic hydrocarbons (PAHs) and certain polychlorinated biphenyls (Hankinson, 1995). Unliganded AhR resides in the cytosol, complexed with p23, Hsp90 and XAP2/ARA9 (Meyer et al., 1998). Upon ligand binding AhR translocates to the nucleus, discards its associated proteins, and forms a heterodimer with the aryl hydrocarbon nuclear translocator (ARNT) to form the aryl hydrocarbon receptor complex (AHRC) (Pollenz et al., 1994; Reyes et al., 1992). The AHRC binds to xenobiotic responsive elements (XREs) located in the regulatory regions of AhR responsive genes, leading to the activation of their transcription.

The CYP1 family includes three genes, *CYP1A1*, *CYP1A2*, and *CYP1B1*, all of which are inducible by AhR agonists. A variety of environmental toxicants and xenobiotics are metabolized by these three enzymes. *CYP1A1* and *CYP1B1* are particularly effective at metabolically activating polycyclic aromatic hydrocarbons, some of which represent important carcinogenic components of cigarette smoke, smog and cooked foods (Nebert et al., 1993). Both the *CYP1A1* and *CYP1B1* genes contain a cluster of XREs about 1kb upstream of their transcription start sites, within regions that represent the enhancers of these genes. Although there are many similarities in the modes of dioxin induction of these two genes, there are interesting differences. *CYP1A1* is usually expressed at very low levels in relevant tissues or cells, and dioxin typically induces its expression hundreds to thousands of fold. In contrast, the constitutive levels of *CYP1B1* are usually substantial, and it is generally induced to a lesser degree than *CYP1A1* (Tang et al., 1996). Upon dioxin induction, AhR, and the histone acetylase (HAT) coactivators, p300, and Steroid Receptor Coactivator 2 (SRC-2) are recruited to the enhancers of both the *CYP1A1* and *CYP1B1* genes, and they are required for their induction by dioxin. However, although the ATPase-dependent nucleosome remodeling factor Brahma/ Switch 2-related gene 1 (BRG-1) is recruited to the *CYP1A1* gene

MOL 64899

upon dioxin treatment and is required for its induction, neither is the case for the *CYP1B1* gene (Taylor et al., 2009). Both the *CYP1A1* and *CYP1B1* genes are expressed in a large number of tissues although there are differences in the tissue-specific expression of the two genes (Bieche et al., 2007). Our understanding of the precise mechanisms involved in the transcriptional regulation of the *CYP1A1* and *CYP1B1* genes is far from complete and a fuller understanding of the molecular mechanisms underlying the regulation of these genes will be of great importance to delineate the toxic effects mediated by these enzymes.

Epigenetic modifications play a significant role in transcriptional regulation of genes (Kouzarides, 2007; Li et al., 2007). Two of the most important modifications in the context of transcription regulation are covalent chromatin/ histone modifications and DNA methylation. Operationally, covalent chromatin modifications function either by altering the nucleosomal architecture and/ or by affecting the recruitment of nonhistone proteins to chromatin (Kouzarides, 2007; Li et al., 2007). DNA methylation suppresses gene expression either directly by interfering with the binding of transcription factors, or indirectly by attracting methylated DNA binding factors that recruit histone deacetylases to generate an inactive heterochromatin structure (Guo et al., 2002; Singal and Ginder, 1999). CpG islands have been identified in the enhancer and in the promoter regions of the human *CYP1A1* and *CYP1B1* genes, and alterations in the DNA methylation status of both the *CYP1A1* and *CYP1B1* genes occur in various types of cancer (Habano et al., 2009; Kang et al., 2008; Okino et al., 2006; Tokizane et al., 2005).

In the current study, we studied the potential role of epigenetic mechanisms in dioxin-induced transcriptional regulation of the *CYP1B1* gene. We demonstrate that the *CYP1B1* gene is silenced in HepG2 cells due to hypermethylation of its promoter, which affects some, but not all, of the relevant dioxin-induced changes that normally occur at the gene. Our studies provide important insights both into the mechanism of *CYP1B1* induction by dioxin and into the mechanisms of gene regulation in mammalian cells in general.

MATERIALS AND METHODS

Cell culture and Reagents: The human breast cancer cell line MCF-7, and human hepatic cancer cell line, HepG2, were obtained from the American type Culture Collection (ATCC; Manassas, VA), and were grown as monolayers in α -minimal essential media and DMEM media respectively, containing 10% fetal bovine serum, 5% fungizone, 5% Penicillin-Streptomycin (Invitrogen, Carlsbad, CA) at 37 ° C and 5 % CO₂. The tissue culture dishes used to grow HepG2 cells were coated with 5 ml of 50 μ g/ml poly-L-lysine and dried before plating the cells. Cells were treated with 100 nM dioxin (Wellington Laboratories, Guelph, Ontario, Canada) dissolved in dimethyl sulphoxide (DMSO), at a final concentration of 0.1% DMSO in the medium. 5-aza-2'-deoxycytidine was purchased from Sigma (5-AzaC; St. Louis, MO). The antibodies used for ChIP analysis are indicated in the supplementary material (Supplementary Table1)

Reverse Transcription and Real-Time PCR: The levels of the mRNAs for CYP1A1, CYP1B1 and the constitutively expressed ribosomal subunit 36B4 were determined by SYBR green real-time PCR. Total RNA was isolated using RNeasy mini columns (Qiagen, Valencia, CA) according to the manufacturer's protocol and quantified on a SmartSpec 3000 spectrophotometer (BioRad, Hercules, CA). Five micrograms of total RNA was used for complementary DNA (cDNA) synthesis in a 20 μ l reaction using Superscript III reverse transcriptase (Invitrogen, Carlsbad, CA) and primed with random hexamers (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. cDNA synthesis was performed using incubations at 25 ° C for 10 minutes, 48 ° C for 30 minutes, and 95 ° C for 5 minutes, using an Icycler thermalcycler (BioRad, Hercules, CA). cDNAs were diluted 10-fold in autoclaved water. Standard curves were generated using the 72 hr dioxin treated MCF-7 cDNA sample and performing a 10-fold dilution series. The primers for real time PCR were designed using Primer Express software (Applied Biosystems, CA) and were synthesized by Integrated DNA Technologies, Inc (Coralville, IA) (Supplementary Table 2). Real time PCR assays were performed using an Applied Biosystems 7500 machine. Real-time PCR reaction parameters were 50 ° C for 2 minutes, and 40 cycles of 95 ° C for 10

MOL 64899

minutes, 92 ° C for 15 seconds, 60 ° C for 1 minute. CYP1A1 and CYP1B1 mRNA quantities were normalized to the amount of 36B4 mRNA. The relative expression levels of all genes measured were reported using standard curve generated from MCF-7 cDNA treated for 72 hrs with dioxin, allowing us to compare HepG2 mRNA expression levels relative to MCF-7 expression levels. In all real-time PCR analyses, three replicates were analyzed for each biological sample, and the standard deviations from those three replicates are reported.

Chromatin Immunoprecipitation assay: MCF-7 and HepG2 cells were treated with 100 nM dioxin for the indicated amount of time. ChIP analyses were carried out as described previously (Beedanagari et al., 2009). DNA was extracted using DNeasy mini columns (Qiagen), and finally eluted in a volume of 50 µl using PCR grade water. The primers used for the housekeeping gene, 36B4, were reported previously (Beedanagari et al., 2009; Hsu et al., 2007). All the ChIP analyses were carried out at least three times, and the data from representative experiments are reported in the manuscript. The antibodies used in ChIP analyses are listed in Supplementary Table 1.

Sodium Bisulfite modification and sequencing: MCF-7 and HepG2 cells were plated on day 0 and treated with 5 µM 5-aza-2'-deoxycytidine (5-Aza-dC; Sigma, St. Louis, MO) starting on day 1 for 72 hrs. Cells were also treated with 100 nM dioxin on day 3 for 24 hrs. Control samples were treated with the vehicle, DMSO. The genomic DNA was extracted using the DNeasy kit, (Qiagen, Valencia, CA USA). Bisulfite modification of DNA was performed using the EZ DNA methylation kit according to the manufacturer's protocol (Zymo Research, Orange, CA USA). The sense strand of bisulfite-modified genomic DNA of the *CYP1B1* enhancer and promoter regions were amplified using two rounds of PCR with specific primers reported previously (Han et al., 2006), followed by quantification of CpG methylation by tag-modified bisulfite genomic sequencing (tBGS) as described previously by (Han et al., 2006). The amplification products were confirmed and purified from 2% agarose gels and sequenced directly by an outside vendor using the same primer sequences (Genewiz, San Diego, CA). The sequences

MOL 64899

were analyzed using Sequencher 4.8 version software purchased from Gene Codes Inc (Anna Arbor, MI USA)

Statistical Analysis:

All the real-time PCR and ChIP analyses were performed at least twice and a representative data set is presented in the figures. All the real-time PCR samples were run in triplicate and the standard deviation from these three replicates are presented as standard error bars in the representative graphs.

RESULTS

Effect of dioxin on the expression of the CYP1A1, AhR and ARNT genes in the MCF-7 and HepG2

cell lines: The human breast cancer cell line MCF-7, and human hepatic cancer cell line HepG2, were treated with 100 nM dioxin for 0, 1, 3, 6, 12, 24, 48, and 72 hrs. This concentration of dioxin maximally induces *CYP1A1* and *CYP1B1* (Taylor et al., 2009) and was not toxic to the cell lines. The *CYP1A1*, *CYP1B1*, AhR and ARNT mRNA levels were quantified using SYBR green real-time PCR. The standard curves for real-time PCR were generated from the corresponding cDNAs derived from MCF-7 cDNAs treated with dioxin for 72 hrs, allowing us to report the mRNA expression levels of these genes in the two cell lines relative to MCF-7 expression levels. Dioxin treatment had no effect on AhR and ARNT mRNA levels at the different time points tested. However, dioxin markedly induced *CYP1A1* mRNA levels in both cell lines. Interestingly, dioxin induced *CYP1B1* mRNA expression in MCF-7 cells, but not in HepG2 cells, in which *CYP1B1* mRNA was undetectable both in the presence and absence of dioxin (Fig. 1). Subsequent experiments were directed at determining the basis for the lack of *CYP1B1* induction in HepG2 cells.

Effect of dioxin on recruitment of AhR and PolII to the enhancer and promoter regions, respectively,

of the *CYP1A1* and *CYP1B1* Genes: We previously reported that, in MCF-7 cells, dioxin induces rapid recruitment of AhR to the enhancer regions of the *CYP1A1* and *CYP1B1* genes, located in each case approximately 1kb 5' to the transcriptional start site of the corresponding gene, and also induced rapid recruitment of RNA polymerase II (polII) to the proximal promoter regions of both genes (Taylor et al., 2009). Here, we investigated the kinetics of AhR recruitment to the enhancers and polII to the promoters of the same genes in HepG2 cells after dioxin treatment, utilizing ChIP analysis. Dioxin rapidly induced AhR recruitment to the enhancers of both the *CYP1A1* and *CYP1B1* genes in HepG2 cells (Fig. 2A & 2C). Interestingly, AhR binding to the *CYP1B1* enhancer followed a cyclical pattern. Such cycling of AhR binding has previously been reported in some but not all studies on the human *CYP1A1* and *CYP1B1* enhancers after AhR ligand treatment (Hestermann and Brown, 2003; Matthews et al., 2007;

MOL 64899

Taylor et al., 2009; Wihlen et al., 2009). The basis for this cycling and its potential significance are not clear. Dioxin induced polII recruitment to the *CYP1A1* promoter (Fig 2B). However, no polII recruitment to the *CYP1B1* promoter was observed in HepG2 cells (Fig. 2D), consistent with the lack of induction of this gene in HepG2 cells. From these studies, we conclude that the loss of dioxin inducibility of *CYP1B1* in HepG2 cells is ascribable to a defect(s) in the induction pathway subsequent to AhR recruitment to the enhancer.

Chromatin modifications at the promoter regions of the CYP1A1 and CYP1B1 genes correlate with dioxin induction:

We performed time course ChIP analyses to investigate whether dioxin induces any changes in the extent of acetylation of histone H3 at lysines 9 and 14 (AcH3K9 and AcH3K14), acetylation of histone H4 (AcH4), and trimethylation of histone H3 at lysine 4 (me3H3K4) at the *CYP1A1* and *CYP1B1* promoters in MCF-7 and HepG2 cells. Dioxin induced increases in these modifications at both the *CYP1A1* and *CYP1B1* promoters in MCF-7 cells, and these increases reached maximal levels between 60 and 120 min after dioxin treatment (Figs. 3A and 3C). The extent of each histone modification at the *CYP1A1* promoter was generally somewhat greater in HepG2 cells than in MCF-7 cells (compare Figures 3B and 3A). In contrast, the extent of the modifications at the *CYP1B1* promoter were much less in HepG2 cells than in MCF-7 cells both before and after dioxin treatment (Figs. 3D and 3C). Thus dioxin induction of these chromatin modifications correlates with dioxin inducibility of the *CYP1A1* and *CYP1B1* genes in the two cell lines.

Coactivator recruitment to the enhancers and promoters of the CYP1B1 and CYP1A1 genes in HepG2 cells: We then studied the recruitment of p300 and the transcriptional coactivator p300/CBP-associated factor (PCAF) to the enhancer and promoter regions of the *CYP1A1* and *CYP1B1* genes in HepG2 cells. Dioxin induced recruitment of p300 and PCAF (and AhR) to the enhancers, but not the promoters, of the *CYP1A1* and *CYP1B1* genes in HepG2 cells (Figs. 4A and 4C). Furthermore, dioxin induced the

MOL 64899

recruitment of these proteins to the *CYP1B1* enhancer as efficiently as it induced their recruitment to the *CYP1A1* enhancer. Thus, we conclude that lack of CYP1B1 induction in HepG2 cells is not due to deficiencies in recruitment of either p300 or PCAF to the corresponding enhancer. Furthermore, we conclude that although p300 is required, it is not sufficient for dioxin induction of the above chromatin modifications at the *CYP1B1* enhancer, since these modifications are not induced by dioxin in HepG2 cells. Interestingly, the recruitment of coactivators at the promoters of both genes in HepG2 cells was negligible. We previously made similar observations with regard to AhR and p300 recruitment at both genes in MCF-7 cells, and furthermore found that polII was recruited to the promoter but not to the enhancer regions of the genes. Thus under our ChIP protocol conditions, recruitment to the enhancers was distinguishable from recruitment to the promoters.

Knockdown of p300 inhibits dioxin induction of chromatin modifications at the CYP1B1 promoter in MCF-7 Cells: We previously showed that dioxin induces recruitment of p300 to the enhancer regions of the *CYP1A1* and *CYP1B1* genes in MCF-7 cells. Furthermore, we previously demonstrated, that p300 is required for maximal dioxin induction of the *CYP1A1* and *CYP1B1* mRNAs in this cell line (Taylor et al., 2009). In the current study we investigated whether p300 is required for dioxin induction of the above chromatin modifications at the *CYP1B1* enhancer. In these experiments we utilized MCF-7 cells stably infected with a vector expressing a shRNA for p300 (shp300 MCF-7 cells) that were previously generated in our laboratory. MCF-7 cells stably infected with the empty vector, RVGP, were used as a control (Taylor et al., 2009). Dioxin treatment increased the levels of the AcH3K9, AcH3K14, AcH4, and me3H3K4 histone modifications at the *CYP1B1* promoter in RVGP MCF-7 cells, but failed to induce these modification in shp300 MCF-7 cells (Figure 5A and 5B). We therefore conclude that p300 is required for dioxin induction of these histone modifications at the *CYP1B1* promoter.

Reactivation of dioxin inducibility of CYP1B1 in HepG2 cells by 5-AzadC: To examine the role of DNA methylation in the regulation of the *CYP1A1* and *CYP1B1* genes, MCF-7 and HepG2 cells were

MOL 64899

treated with the DNA methyltransferase (DNMT) inhibitor, 5-AzaC, for 72 hrs. 100nM dioxin or the vehicle (DMSO) was included in the medium for the last 24 hrs before harvest. In HepG2 cells, 5 μ M 5-AzaC reactivated dioxin-induced CYP1B1 mRNA expression to a level comparable to that in MCF-7 cells. However, the mRNA expression levels of CYP1A1 in both cell lines and of CYP1B1 in MCF-7 cells were not affected by 5-AzaC treatment (Fig. 6). These results indicate that silencing of the *CYP1B1* gene in HepG2 cells is most likely due to hypermethylation of the *CYP1B1* gene, or possibly of a gene encoding a negative regulator of CYP1B1 expression.

DNA methylation at the *CYP1B1* enhancer and promoter: High frequencies of CpG dinucleotides occur at both the enhancer and promoter of the human *CYP1B1* gene, and these regions represent “CpG islands.” Thus there are 29 CpG sites within a 360 bp sequence (-560 bp to -920 bp) encompassing the enhancer, and 24 CpG sites within a 280 bp sequence (-260 to +20) encompassing the promoter. To study the cytosine methylation status of the *CYP1B1* promoter and enhancer regions, we utilized the tag-modified bisulfite genomic sequencing (tBGS) procedure as described by Han and coworkers (Han et al., 2006). Direct bisulfite sequencing was performed on the amplified PCR products of the *CYP1B1* regulatory regions, without cloning. Bisulfite treatment of DNA converts unmethylated cytosines (C) to uracils, which will be ultimately converted to thymines (T) during the amplification process, whereas the methylated cytosines remain unaffected. Representative bisulfite sequencing chromatograms are presented in the Figs. 7A, B, and C. In these figures, red peaks in positions corresponding to cytosines in genomic DNA represent unmethylated cytosines, and blue peaks represent methylated cytosines. Bisulfite sequencing demonstrated that 22 of the 24 CpG sites are fully methylated, 1 is partially methylated, and 1 is unmethylated in the *CYP1B1* promoter in HepG2 cells, while only one of these sites is methylated in MCF-7 cells (Fig. 7D). 22 of the 29 CpG sites were found to be partially methylated in the *CYP1B1* enhancer in HepG2 cells. In contrast, only two of these sites were partially methylated in MCF-7 cells, the remainder being totally unmethylated (Fig. 7E). The XRE sequence harbors a CpG site. Methylation of the CpG dinucleotide within the XRE prevents binding of the AhR/ARNT dimer *in vitro*, and inhibits the

MOL 64899

ability of the XRE to mediate dioxin induction of gene transcription (Shen and Whitlock, 1989). We were therefore particularly interested in the methylation status of the CpG dinucleotides in the XRE2 and XRE3 sequences, which are located in the enhancer and which play the most significant role in the dioxin-mediated induction of *CYP1B1* (Shehin et al., 2000; Tsuchiya et al., 2003). Bisulfite sequencing revealed that both XREs were approximately 30% methylated in HepG2 cells (Fig 7E). Thus either or both of XRE2 and XRE3 are unmethylated in a majority of chromosomes, consistent with the observation that AhR is recruited to the *CYP1B1* enhancer of HepG2 cells after dioxin treatment. In summary, these studies indicate that the loss of dioxin induction of CYP1B1 in HepG2 cells is due to methylation of cytosines at CpG dinucleotides at the gene's promoter. We did not directly analyze the methylation status of the *CYP1A1* gene promoter, as its dioxin induction was unaltered after treatment with 5-AzadC in both MCF-7 and HepG2 cells (Fig. 6), suggesting that DNA methylation does not play an important role in the dioxin induced transcriptional regulation of this gene in these cell lines.

Partial demethylation of the CYP1B1 promoter by 5'-AzadC treatment in HepG2 cells, and restoration of polIII binding at the promoter. In 5-AzadC treated HepG2 cells, DNA methylation was partially reversed at the CYP1B1 promoter (Fig. 7C). Furthermore, ChIP analysis demonstrated that dioxin-induced polIII recruitment to the *CYP1B1* promoter was restored in these cells (Fig. 8). Both these observations are consistent with our observation that 5'-AzadC restores dioxin inducibility to the *CYP1B1* gene in HepG2 cells.

Binding of the TATA binding protein (TBP) to the CYP1B1 promoter in MCF-7 and HepG2 cells.

Since dioxin failed to induce polIII recruitment to the *CYP1B1* promoter, we also studied the recruitment of the TATA binding protein (TBP) to this promoter in MCF-7 and HepG2 cells by ChIP analysis. TBP binding generally precedes the binding of other general transcription factors and polIII at the promoter and "seeds" their association with the promoter (Kornberg, 2007). TBP presumably binds to the functional TATA-like box sequence located 35 bp upstream of the transcription start site of *CYP1B1* (Wo et al.,

MOL 64899

1997). Dioxin was found to induce TBP recruitment to the *CYP1B1* promoter in MCF-7 cells but not in HepG2 cells (Fig 9A). TBP recruitment was restored at the *CYP1B1* promoter, after treatment with 5-AzadC. However, the TBP recruitments levels at the *CYP1B1* promoter remain unaltered in MCF-7 cells after treatment with 5-AzadC (Fig 9B). Thus, the loss of dioxin induction of CYP1B1 in HepG2 cells attributable to defect(s) occurring prior to TBP recruitment to the *CYP1B1* promoter.

DISCUSSION

We report a detailed analysis of the mechanism responsible for the lack of induction of CYP1B1 by dioxin in HepG2 cells. Our observations provide insights both into mechanisms of gene regulation in general, and into the regulation of human CYP1B1 in particular.

Whereas CYP1A1 was induced in both MCF-7 and HepG2 cells, and CYP1B1 was induced in MCF-7 cells, CYP1B1 mRNA was neither expressed in the absence of dioxin nor induced by dioxin in HepG2 cells. However, dioxin did induce recruitment of AhR to the enhancer of the *CYP1B1* gene in HepG2 cells, thereby confirming that binding of AhR does not necessarily equate with dioxin inducibility. This conclusion was previously also made by our laboratory and by others based on studies on other cell lines (Beedanagari et al., 2010; Yang et al., 2008). Dioxin induced four different types of histone modifications at the *CYP1A1* promoter in HepG2 cells and at the *CYP1A1* and *CYP1B1* promoters in MCF-7 cells. Other investigators have shown that dioxin induces these same modifications at the promoter of the *Cyp1a1* gene in mouse hepatoma cells (Schnekenburger et al., 2007). These modifications are generally associated with actively transcribed genes (Li et al., 2007). Of interest, these modifications were not induced by dioxin at the *CYP1B1* promoter in HepG2 cells, demonstrating that these cells are blocked in the CYP1B1 induction pathway at a step prior to the generation of these modifications. We previously showed that the HAT coactivator p300 is required for dioxin induction of CYP1B1 (Taylor et al., 2009), and we now show p300 is also required for dioxin induction of the above histone modifications at the *CYP1B1* promoter (in MCF-7 cells). The three acetylations we studied, AcH3K9, AcH3K14 and AcH4, are known to be catalyzed by p300 (as well as certain other HAT coactivators) (Kouzarides, 2007) and they may therefore represent direct targets of p300 in the *CYP1B1* gene. We show that dioxin induces recruitment of p300 and the HAT coactivator PCAF to the *CYP1B1* enhancer in HepG2 cells. Thus the block in the induction pathway in HepG2 cells occurs at a step beyond these recruitments, but prior to the generation of the above chromatin modifications at the promoter.

We show that CpG dinucleotides are fully methylated within the CpG island encompassing the promoter of the *CYP1B1* gene in HepG2 cells, but are only partially methylated within the CpG island

MOL 64899

encompassing the enhancer. Importantly, the XREs in the enhancer are only partially methylated and are therefore available for binding of the AhR/ARNT dimer. Treatment of HepG2 cells with the DNA methyltransferase inhibitor, 5-AzadC, partially demethylated the CpG dinucleotides in the *CYP1B1* promoter, restored dioxin induction of the recruitment of RNA polymerase II to the *CYP1B1* promoter, and restored induction of *CYP1B1* expression. These results indicate that DNA methylation at the *CYP1B1* promoter is directly responsible for silencing the gene's transcriptional response to dioxin. Interestingly, Shehin and coworkers also previously observed no induction of *CYP1B1* in HepG2 cells by dioxin. (Shehin et al., 2000). Since Shehin and coworkers used 5-Azacytidine, rather than its deoxy analogue 5-AzadC that we used in our study, we also tested the former compound in our studies and demonstrated that it is equally effective in reactivating *CYP1B1* mRNA as 5-AzadC (data not shown). We and Shehin and coworkers obtained HepG2 cells from the same source, and we have no explanation for the differences in our results.

Binding of the TATA binding protein (TBP) to gene promoters usually occurs subsequent to the binding of sequence specific transcription factors enhancers, and prior to association of the general transcription factors and polII to the promoter (Kornberg, 2007). Our observation that dioxin fails to induce TBP binding to the *CYP1B1* promoter in HepG2 cells, suggests that one of the critical effects of DNA methylation may be to preclude association of this factor with the promoter. However, since the TATA-like box sequence in the *CYP1B1* promoter does not contain any CpG dinucleotides, DNA methylation may not directly prevent binding of this protein, but may inhibit binding of another protein or protein(s) at the promoter required for TBP recruitment. This inhibition could be due to the presence of methylated DNA-binding proteins, such as MeCP2, MBD1, MBD2 and/or MBD3 (Szyf, 2009), with the *CYP1B1* promoter. Our conclusions concerning the alterations to the regulatory processes occurring at the *CYP1B1* gene in HepG2 cells are presented in Figure 10.

It is recognized that transcriptional coactivators are incorporated in large multi-protein complexes straddling both the enhancer and the promoter of responsive genes (O'Malley, 2007). We observed much greater levels of p300 and PCAF recruitment to the *CYP1A1* and *CYP1B1* enhancers than to the corresponding promoters, after dioxin treatment. This observation is consistent with the notion that these

MOL 64899

proteins are in closer proximity to the AhR/ARNT dimer at the *CYP1B1* enhancer than to TBP and the general transcription factors located at the promoter. The lack of detectable binding at the promoter is most likely due to the inefficiency of formaldehyde at cross-linking proteins under the conditions of our ChIP procedure, coupled with the presence of multiple proteins spanning the distance between these coactivators and the promoter. It is therefore of considerable interest that p300 and PCAF are recruited efficiently at the *CYP1B1* enhancer after dioxin treatment in HepG2 cells, despite the fact that TBP and pol II are not recruited to the corresponding promoter. These observations strongly imply that p300 and PCAF can be recruited to the enhancer even when they are not incorporated into a multiprotein complex spanning the enhancer and the promoter. The lack of a requirement for coactivators to be tethered at both promoter and enhancer can probably be generalized to other coactivators and other genes and this represents an important area for future research. It should be noted that several uncommon features of our experimental system, including (i) our ability to distinguish the *in vivo* binding of proteins to the enhancer from their binding to the promoter, and vice versa, and most importantly (ii) the fact that the promoter is fully methylated but the enhancer is not, provided us with the opportunity to draw the above conclusion, and therefore to provide important insight into the general mechanism of gene transcription in mammalian cells. Altogether, the *CYP1B1* gene in HepG2 cells provides a valuable experimental system for further experiments focusing on potential regulatory mechanisms affecting the relationship between transcription and epigenetic modifications. Since *CYP1B1* has been observed to be inducible by dioxin in other human hepatic cancer cell lines (MacPherson et al., 2009), the mechanism of inhibition of *CYP1B1* expression we describe is not universal to such cell lines.

Of considerable importance are the observations that the *CYP1B1* promoter exhibits DNA methylation in a portion of colorectal (Habano et al., 2009) and gastric (Kang et al., 2008) cancers. Interestingly, in a portion of colorectal cancers, CpG dinucleotides at the *CYP1B1* promoter are fully methylated, while CpG dinucleotides at the enhancer are only partially methylated, just as we observed with HepG2 cells (18). Our studies are therefore potentially relevant to the development of these cancers,

MOL 64899

as well as providing insights into the mechanism of CYP1B1 induction, and into the regulation of gene expression in mammalian cells in general.

ACKNOWLEDGEMENTS

We thank Kelly Joiner for her help in formatting the manuscript.

REFERENCES

- Beedanagari SR, Bebenek I, Bui P and Hankinson O (2009) Resveratrol inhibits dioxin-induced expression of human CYP1A1 and CYP1B1 by inhibiting recruitment of the aryl hydrocarbon receptor complex and RNA polymerase II to the regulatory regions of the corresponding genes. *Toxicol Sci* **110**(1):61-67.
- Beedanagari SR, Taylor RT and Hankinson O (2010) Differential regulation of the dioxin-induced Cyp1a1 and Cyp1b1 genes in mouse hepatoma and fibroblast cell lines. *Toxicol Lett*.
- Bieche I, Narjoz C, Asselah T, Vacher S, Marcellin P, Lidereau R, Beaune P and de Waziers I (2007) Reverse transcriptase-PCR quantification of mRNA levels from cytochrome (CYP)1, CYP2 and CYP3 families in 22 different human tissues. *Pharmacogenet Genomics* **17**(9):731-742.
- Guo Y, Pakneshan P, Gladu J, Slack A, Szyf M and Rabbani SA (2002) Regulation of DNA methylation in human breast cancer. Effect on the urokinase-type plasminogen activator gene production and tumor invasion. *J Biol Chem* **277**(44):41571-41579.
- Habano W, Gamo T, Sugai T, Otsuka K, Wakabayashi G and Ozawa S (2009) CYP1B1, but not CYP1A1, is downregulated by promoter methylation in colorectal cancers. *Int J Oncol* **34**(4):1085-1091.
- Han W, Cauchi S, Herman JG and Spivack SD (2006) DNA methylation mapping by tag-modified bisulfite genomic sequencing. *Anal Biochem* **355**(1):50-61.
- Hankinson O (1995) The aryl hydrocarbon receptor complex. *Annu Rev Pharmacol Toxicol* **35**:307-340.
- Hestermann EV and Brown M (2003) Agonist and chemopreventative ligands induce differential transcriptional cofactor recruitment by aryl hydrocarbon receptor. *Mol Cell Biol* **23**(21):7920-7925.
- Hsu EL, Yoon D, Choi HH, Wang F, Taylor RT, Chen N, Zhang R and Hankinson O (2007) A proposed mechanism for the protective effect of dioxin against breast cancer. *Toxicol Sci* **98**(2):436-444.
- Huff J, Lucier G and Tritscher A (1994) Carcinogenicity of TCDD: experimental, mechanistic, and epidemiologic evidence. *Annu Rev Pharmacol Toxicol* **34**:343-372.
- Kang GH, Lee S, Cho NY, Gandamihardja T, Long TI, Weisenberger DJ, Campan M and Laird PW (2008) DNA methylation profiles of gastric carcinoma characterized by quantitative DNA methylation analysis. *Lab Invest* **88**(2):161-170.
- Kornberg RD (2007) The molecular basis of eukaryotic transcription. *Proc Natl Acad Sci U S A* **104**(32):12955-12961.
- Kouzarides T (2007) Chromatin modifications and their function. *Cell* **128**(4):693-705.
- Li B, Carey M and Workman JL (2007) The role of chromatin during transcription. *Cell* **128**(4):707-719.
- MacPherson L, Lo R, Ahmed S, Pansoy A and Matthews J (2009) Activation function 2 mediates dioxin-induced recruitment of estrogen receptor alpha to CYP1A1 and CYP1B1. *Biochem Biophys Res Commun* **385**(2):263-268.
- Mandal PK (2005) Dioxin: a review of its environmental effects and its aryl hydrocarbon receptor biology. *J Comp Physiol B* **175**(4):221-230.
- Matthews J, Wihlen B, Heldring N, MacPherson L, Helguero L, Treuter E, Haldosen LA and Gustafsson JA (2007) Co-planar 3,3',4,4',5-pentachlorinated biphenyl and non-co-planar

MOL 64899

- 2,2',4,6,6'-pentachlorinated biphenyl differentially induce recruitment of oestrogen receptor alpha to aryl hydrocarbon receptor target genes. *Biochem J* **406**(2):343-353.
- Meyer BK, Pray-Grant MG, Vanden Heuvel JP and Perdew GH (1998) Hepatitis B virus X-associated protein 2 is a subunit of the unliganded aryl hydrocarbon receptor core complex and exhibits transcriptional enhancer activity. *Mol Cell Biol* **18**(2):978-988.
- Nebert DW, Puga A and Vasiliou V (1993) Role of the Ah receptor and the dioxin-inducible [Ah] gene battery in toxicity, cancer, and signal transduction. *Ann N Y Acad Sci* **685**:624-640.
- O'Malley BW (2007) Coregulators: from whence came these "master genes". *Mol Endocrinol* **21**(5):1009-1013.
- Okino ST, Pookot D, Li LC, Zhao H, Urakami S, Shiina H, Igawa M and Dahiya R (2006) Epigenetic inactivation of the dioxin-responsive cytochrome P4501A1 gene in human prostate cancer. *Cancer Res* **66**(15):7420-7428.
- Pollenz RS, Sattler CA and Poland A (1994) The aryl hydrocarbon receptor and aryl hydrocarbon receptor nuclear translocator protein show distinct subcellular localizations in Hepa 1c1c7 cells by immunofluorescence microscopy. *Mol Pharmacol* **45**(3):428-438.
- Reyes H, Reisz-Porszasz S and Hankinson O (1992) Identification of the Ah receptor nuclear translocator protein (Arnt) as a component of the DNA binding form of the Ah receptor. *Science* **256**(5060):1193-1195.
- Schnekenburger M, Talaska G and Puga A (2007) Chromium cross-links histone deacetylase 1-DNA methyltransferase 1 complexes to chromatin, inhibiting histone-remodeling marks critical for transcriptional activation. *Mol Cell Biol* **27**(20):7089-7101.
- Shehin SE, Stephenson RO and Greenlee WF (2000) Transcriptional regulation of the human CYP1B1 gene. Evidence for involvement of an aryl hydrocarbon receptor response element in constitutive expression. *J Biol Chem* **275**(10):6770-6776.
- Shen ES and Whitlock JP, Jr. (1989) The potential role of DNA methylation in the response to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Biol Chem* **264**(30):17754-17758.
- Singal R and Ginder GD (1999) DNA methylation. *Blood* **93**(12):4059-4070.
- Szyf M (2009) Epigenetics, DNA methylation, and chromatin modifying drugs. *Annu Rev Pharmacol Toxicol* **49**:243-263.
- Tang YM, Wo YY, Stewart J, Hawkins AL, Griffin CA, Sutter TR and Greenlee WF (1996) Isolation and characterization of the human cytochrome P450 CYP1B1 gene. *J Biol Chem* **271**(45):28324-28330.
- Taylor RT, Wang F, Hsu EL and Hankinson O (2009) Roles of coactivator proteins in dioxin induction of CYP1A1 and CYP1B1 in human breast cancer cells. *Toxicol Sci* **107**(1):1-8.
- Tokizane T, Shiina H, Igawa M, Enokida H, Urakami S, Kawakami T, Ogishima T, Okino ST, Li LC, Tanaka Y, Nonomura N, Okuyama A and Dahiya R (2005) Cytochrome P450 1B1 is overexpressed and regulated by hypomethylation in prostate cancer. *Clin Cancer Res* **11**(16):5793-5801.
- Tsuchiya Y, Nakajima M and Yokoi T (2003) Critical enhancer region to which AhR/ARNT and Sp1 bind in the human CYP1B1 gene. *J Biochem* **133**(5):583-592.
- Wihlen B, Ahmed S, Inzunza J and Matthews J (2009) Estrogen receptor subtype- and promoter-specific modulation of aryl hydrocarbon receptor-dependent transcription. *Mol Cancer Res* **7**(6):977-986.
- Wo YY, Stewart J and Greenlee WF (1997) Functional analysis of the promoter for the human CYP1B1 gene. *J Biol Chem* **272**(42):26702-26707.

MOL 64899

Yang X, Solomon S, Fraser LR, Trombino AF, Liu D, Sonenshein GE, Hestermann EV and Sherr DH (2008) Constitutive regulation of CYP1B1 by the aryl hydrocarbon receptor (AhR) in pre-malignant and malignant mammary tissue. *J Cell Biochem* **104**(2):402-417.

MOL 64899

FOOTNOTES

The research was supported by a grant from the National Institute of Health [R01CA28868] and was partially supported by an underrepresented minority supplement to R01CA28868, and by a National Institute of Environmental Health Sciences training grant [T32 ES015457].

FIGURE LEGENDS

Figure1: Effect of dioxin on expression of CYP1A1, CYP1B1, AhR and ARNT mRNAs. MCF-7 and HepG2 cells were treated with 100nM dioxin for the indicated times. RNA was isolated and mRNA levels were assayed by real time PCR. The relative amounts of AhR, ARNT, CYP1B1, and CYP1A1 mRNAs were corrected against the levels of mRNA expression for the constitutively expressed ribosomal subunit, 36B4. All mRNA expression levels were reported using a standard curve generated from a standard cDNA sample obtained from MCF-7 cells treated with dioxin for 72 hrs.

Figure 2: AhR and polII recruitment to the *CYP1A1* and *CYP1B1* genes in HepG2 cells. HepG2 cells were treated with 100 nM dioxin for the indicated times and subjected to ChIP analysis. AhR recruitment to the enhancers (Fig 2A & 2C) and polII to the promoters (Fig 2B & 2D) of the *CYP1A1* and *CYP1B1* genes were measured by real-time PCR and reported relative to total inputs.

Figure 3: Characterization of chromatin modifications at the *CYP1A1* and *CYP1B1* promoter regions in MCF-7 and HepG2 Cells. ChIP analyses were carried out using antibodies targeted to four chromatin modifications on MCF-7 and HepG2 cells treated with 100 nM dioxin for the indicated amounts of time at the *CYP1A1* promoter (Fig. 3A& 3B) and *CYP1B1* promoter (Fig. 3C and 3D) . The relative levels of the chromatin modifications were measured by real-time PCR and are reported relative to total inputs.

Figure 4: Coactivator recruitment to the *CYP1B1* and *CYP1A1* enhancer and promoter regions in HepG2 cells ChIP analyses were carried out using antibodies targeted to AhR, PCAF and p300 in HepG2 cells treated with 100 nM dioxin (black bars), or with vehicle (DMSO, gray bars) for 60 mins. AhR, PCAF and p300 recruitment to the *CYP1B1* and *CYP1A1* enhancer (Fig. 4A & 4C) and promoter (Fig. 4B & 4D) regions were measured by real-time PCR and are reported relative to total inputs.

Figure 5: Effect of p300 knock-down on chromatin modifications at the *CYP1B1* promoter in MCF-7 cells. MCF-7 cells stably knocked down for shp300, or treated with control vector (RVGP MCF-7) were treated with 100 nM dioxin (black bars), or with vehicle (DMSO, gray bars) for 60 mins and subjected to ChIP analysis. The chromatin modifications at the *CYP1B1* promoter were measured and reported as described in Figure 4.

Figure 6: Effect of 5'-AzadC treatment on CYP1A1 and CYP1B1 expression in MCF-7 and HepG2 cells. MCF-7 and HepG2 cells were treated with 5 μ M 5-AzadC or DMSO (vehicle) for 72 hrs. The cells were also treated with 24 hrs of either DMSO or dioxin, overlapping with the last 24 hrs of 5-AzadC treatment. Expression levels of the mRNAs for CYP1B1 and CYP1A1 in HepG2 cells and MCF-7 cells were quantified using real-time PCR and reported relative to the mRNA expression levels of the housekeeping gene, 36B4. All mRNA expression levels were calculated using a standard curve generated from cDNA sample obtained from MCF-7 cells treated with dioxin for 72 hrs.

Figure 7: Bisulfite DNA sequencing of the *CYP1B1* enhancer and promoter regions. The MCF-7 and HepG2 enhancer and promoter regions were amplified using bisulfite converted genomic DNA templates from cells treated or untreated with 5'-AzadC. The bisulfite converted DNA was amplified using bisulfite sequencing primers and the amplicons were separated by electrophoresis on 2% agarose gels. The PCR products were excised from the gel, purified and both the sense and antisense strands were directly sequenced for quantitative analysis of DNA methylation. A representative bisulfite sequencing chromatograms from the *CYP1B1* promoter (Fig. 7A), enhancer (Fig. 7B) and 5-AzadC treated promoter (Fig. 7C) are shown. All the CpG dinucleotides are underlined. The red peaks of the bisulfite sequenced chromatogram at CpG dinucleotides represent unmethylated cytosines, and the blue peaks represent methylated cytosines. Overlapping both red and blue peaks indicate a partial methylation status of

MOL 64899

cytosine at that particular CpG site. The relative heights of the blue vs red peaks from bisulfite sequenced chromatograms were used to calculate the methylation status of cytosines and was represented in terms of percentage of methylation. The summary of the methylation patterns and percentage of cytosine methylation at each CpG sites of *CYP1B1* enhancer (Fig. 7D) and promoter (Fig. 7E) are represented pictorially.

Figure 8: polII recruitment to the *CYP1B1* gene in 5'-AzadC treated HepG2 cells. HepG2 cells were treated with 5 μ M 5'-AzadC for 3 days, followed with 100 nM dioxin treatment for the indicated times, and subjected to ChIP analysis. AhR recruitment to the enhancer and polII to the promoter of the *CYP1B1* gene were measured by real-time PCR and reported relative to total inputs.


Figure 9: TBP recruitment to the *CYP1B1* promoter in MCF-7 and HepG2 cells.

ChIP analyses were carried out using a TBP antibody in HepG2 and MCF-7 cells treated with a 100 nM dioxin or vehicle (DMSO) for 60 mins. TBP recruitment to the *CYP1B1* promoter with or without 5'-AzadC treatment in MCF-7 and HepG2 cells was measured by real-time PCR and reported relative to that of total inputs.


Figure 10: Proposed silencing model for the *CYP1B1* gene in HepG2 Cells: Proposed changes in regulatory process occurring at the *CYP1B1* promoter in HepG2 cells (compared with MCF-7 cells) after dioxin treatment.

Figure 1


AHR



ARNT



CYP1B1



CYP1A1




Figure 2




Figure 2A




Figure 2B




Figure 2C




Figure 2D

Figure 3




Figure 3A




Figure 3B




Figure 3C




Figure 3D

Figure 4




Figure 4A




Figure 4B




Figure 4C




Figure 4D

Figure 5




Figure 5A




Figure 5B

Figure 6







Figure 7




Figure 7A




Figure 7B




Figure 7C

Figure 7




Figure 7D: Bisulfite sequenced CpG sites in the *CYP1B1* Promoter region




Figure 7E: Bisulfite sequenced CpG sites in the *CYP1B1* Enhancer region

Figure 8




Figure 9




Figure 9A





Figure 9B

Figure 10

CYP1B1 Silencing Model in HepG2 cells

MCF-7 Cells



HepG2 Cells

