

MOL #97220

## **Molecular insights into metabotropic glutamate receptor allosteric modulation**

Karen J. Gregory and P. Jeffrey Conn

Drug Discovery Biology, Monash Institute of Pharmaceutical Sciences and Department of Pharmacology, Monash University, Parkville, VIC, Australia (KJG).

Vanderbilt Center for Neuroscience Drug Discovery & Department of Pharmacology, Vanderbilt University Medical Center, Nashville, TN, USA (PJC).

MOL #97220

Running Title: Molecular insights into mGlu allosteric modulation

Corresponding Author:

P. Jeffrey Conn

1215 Light Hall

2215-B Garland Ave

Nashville, TN, USA

37232

[jeff.conn@vanderbilt.edu](mailto:jeff.conn@vanderbilt.edu)

Number of text pages: 47

Number of tables: 4

Number of figures: 4

Number of references: 150

Number of words in Abstract: 136

Number of words in Introduction: 471

Number of words in Discussion: 6089

**Non-standard abbreviations:** 7TMs, seven transmembrane spanning  $\alpha$ -helical domains; ACPD, 1-Amino-1,3-dicarboxycyclopentane; AMN082, N,N'-dibenzhydriethane-1,2-diamine dihydrochloride; BAY36-7620: (3a*S*,6a*S*)-Hexahydro-5-methylene-6a-(2-naphthalenylmethyl)-1*H*-cyclopenta[*c*]furan-1-one; BINA, biphenyl-indanone A; cAMP: cyclic adenosine-monophosphate; CBiPES, *N*-[4'-cyano-biphenyl-3-yl]-*N*-(3-pyridinylmethyl)-ethanesulfonamide hydrochloride; CPCCOEt, 7-(hydroxyimino)cyclopropa[*b*]chromen-1a-carboxylate ethyl ester; CPPHA, *N*-(4-chloro-2-[(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)methyl]phenyl)-2-hydroxybenzamide; DCG-IV, (2*S*,2'*R*,3'*R*)-2-(2',3'-dicarboxycyclopropyl)glycine; DFB, ((3-Fluorophenyl)methylene)hydrazone-3-fluorobenzaldehyde; DHPG, 3,5-dihydroxyphenylglycine; DPFE, 1-(4-(2,4-difluorophenyl)piperazin-1-yl)-2-((4-fluorobenzyl)oxy)ethanone; FITM, 4-fluoro-*N*-(4-(6-(isopropylamino)pyrimidin-4-yl)thiazole-2-yl)-*N*-methylbenzamide; GIRK, G protein-coupled inwardly-rectifying potassium channel; GPCR, G protein-coupled receptor;  $iCa^{2+}$ , intracellular  $Ca^{2+}$  mobilization; IP<sub>1</sub>, inositol phosphate accumulation; HYDIA, (1*S*,2*R*,3*R*,5*R*,6*S*)-2-amino-3-Hydroxy-bicyclo[3.1.0]hexane-2,6-dicarboxylic Acid; JNJ-40068782, 3-cyano-1-cyclo-propylmethyl-4-(4-phenyl-piperidin-1-yl)-pyridine-2(1*H*)-one; L-AP4, L-2-amino-4-phosphonobutyrate; LSP4-2022, (2*S*)-2-amino-4-([(4-(carboxymethoxy)phenyl](hydroxy)methyl](hydroxy)phosphoryl)butanoic acid; LTD, long-term depression; LTP, long-term potentiation; LY341495, (2*S*)-2-Amino-2-[(1*S*,2*S*)-2-carboxycycloprop-1-yl]-3-(xanth-9-yl) propanoic acid; LY354740, (1*S*,2*R*,5*R*,6*S*)-2-amino-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid; LY379268, (-)-2-oxa-4-aminobicyclo[3.1.0]hexane-4,6-dicarboxylate; LY541850, (1*S*,2*S*,4*R*,5*R*,6*S*)-2-amino-4-methylbicyclo[3.1.0]hexane-2,6-dicarboxylic acid; M-5MPEP, 2-(2-(3-methoxyphenyl)ethynyl)-5-methylpyridine; MPEP, 2-Methyl-6-(phenylethynyl)pyridine; mPEPy, 3-methoxy-5-(2-pyridinylethynyl)pyridine; MMPIP, 6-(4-Methoxyphenyl)-5-methyl-3-(4-pyridinyl)-isoxazolo [4,5-*c*]pyridine-4(5*H*)-one hydrochloride; NAM, negative allosteric modulator; NCFP, *N*-(4-chloro-2-[(1,3-dioxoisindolin-2-yl)methyl]phenyl)picolinamide; PAM, positive allosteric modulator; pERK1/2, phosphorylated extracellular-signal regulated kinases 1 and 2; PET, positron emission tomography; PHCCC, *N*-phenyl-7-(hydroxyimino)cyclopropa[*b*] chromen-1a-carboxamide; TM, transmembrane;

MOL #97220

R214127, 1-(3,4-dihydro-2H-pyrano[2,3-b]quinolin-7-yl)-2-phenylethanone; RO4988546, 5-[7-Trifluoromethyl-5-(4-trifluoromethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-3-ylethynyl]-pyridine-3-sulphonic acid; RO5488608, 3'-(8-Methyl-4-oxo-7-trifluoromethyl-4,5-dihydro-3H-benzo[*b*][1,4]diazepin-2-yl)-biphenyl-3-sulphonic acid; Ro 67-4853, butyl (9H-xanthene-9-carbonyl)carbamate; Ro 67-7476, (S)-2-(4-fluorophenyl)-1-(toluene-4-sulfonyl)pyrrolidine; VFT: venus flytrap domain; VU0092273, 3'-(8-Methyl-4-oxo-7-trifluoromethyl-4,5-dihydro-3H-benzo[*b*][1,4]diazepin-2-yl)-biphenyl-3-sulphonic acid; VU0155094, methyl 4-(3-(2-((4-acetamidophenyl)thio)acetyl)-2,5-dimethyl-1*H*-pyrrol-1-yl)benzoate; VU0366248, *N*-(3-chloro-2-fluorophenyl)-3-cyano-5-fluorobenzamide; VU0409551, 5-[(4-Fluorophenyl)carbonyl]-2-(phenoxymethyl)-4,5,6,7-tetrahydro[1,3]oxazolo[5,4-*c*]pyridine; VU0422288, *N*-(3-chloro-4-((5-chloropyridin-2-yl)oxy)phenyl)picolinamide.

MOL #97220

## **Abstract**

The metabotropic glutamate (mGlu) receptors are a group of eight family C G protein-coupled receptors that are expressed throughout the CNS and periphery. Within the CNS the different subtypes are found in neurons, both pre- and/or post-synaptically where they mediate modulatory roles and in glial cells. The mGlu receptor family provides attractive targets for numerous psychiatric and neurological disorders, with the majority of discovery programs focussed on targeting allosteric sites, with allosteric ligands now available for all mGlu receptor subtypes. However, the development of allosteric ligands remains challenging. Biased modulation, probe dependence and molecular switches all contribute to the complex molecular pharmacology exhibited by mGlu receptor allosteric ligands. In recent years we have made significant progress in our understanding of this molecular complexity coupled with an increased understanding of the structural basis of mGlu allosteric modulation.

MOL #97220

## Introduction

Glutamate is the major excitatory neurotransmitter in the central nervous system (CNS), eliciting its effects via two distinct receptor classes. The ionotropic glutamate receptors are ligand-gated ion channels that mediate fast synaptic responses. The metabotropic glutamate (mGlu) receptors are G protein-coupled receptors (GPCRs) that play a modulatory role in the CNS. Ubiquitously expressed throughout the CNS in glia and neurons, found both pre and post synaptically, the mGlu receptors are involved in a broad range of physiological functions in the brain. Comprised of eight members, the mGlu receptor family is often sub-classified into three groups based on homology, pharmacology and G protein coupling. Group I includes mGlu<sub>1</sub> and mGlu<sub>5</sub> that preferentially couple to G<sub>q/11</sub> and are typically found postsynaptically. Group II, mGlu<sub>2</sub> and mGlu<sub>3</sub>; and group III, mGlu<sub>4</sub>, mGlu<sub>6</sub>, mGlu<sub>7</sub> and mGlu<sub>8</sub>, members couple to G<sub>i/o</sub> and with the exception of mGlu<sub>6</sub>, which is exclusively found in the retina, are predominantly located pre-synaptically. Multiple recent reviews have extensively discussed the roles of mGlu receptors in different neuronal circuits, modulating synaptic transmission, neuronal excitability and other cellular functions within the CNS (Anwyl, 1999; Bellone et al., 2008; Conn and Pin, 1997; Coutinho and Knopfel, 2002; Pinheiro and Mulle, 2008; Valenti et al., 2002) as well as physiology in the periphery (Julio-Pieper et al., 2011). Based on this understanding, targeting activation or inhibition of specific mGlu receptor subtypes may provide a therapeutic benefit for a range of psychiatric and neurological disorders. For neuropathic pain, inhibition of mGlu<sub>1</sub> is being pursued as a therapeutic intervention with success reported in pre-clinical models (Bennett et al., 2012; Mabire et al., 2005). Agonists of group II receptors have reached clinical trials for schizophrenia and anxiety with varying success (Dunayevich et al., 2008; Kinon et al., 2011; Patil et al., 2007; Schoepp et al., 2003). Group II mGlu receptors are also attractive targets for

MOL #97220

addiction and depression (Chaki et al., 2013; Ma et al., 2007; Moussawi and Kalivas, 2010). Inhibitors of mGlu<sub>5</sub> have demonstrated efficacy in pre-clinical studies for anxiety, depression, and Parkinson's disease Levodopa-induced dyskinesias (Porter et al., 2005; Li et al., 2006; Belozertseva et al., 2007; Rylander et al., 2010; Hughes et al., 2012). Promisingly, multiple mGlu<sub>5</sub> inhibitors have reached (Berry-Kravis et al., 2009; Keywood et al., 2009; Zerbib et al., 2010) or are currently in phase II clinical trials ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)) with varying successes and failures for fragile X syndrome, Parkinson's disease Levodopa-induced dyskinesias, gastroesophageal reflux disorder and migraine. On the other hand, activators and/or potentiators of mGlu<sub>5</sub> have efficacy in preclinical models for schizophrenia and cognition (Moghaddam, 2004; Noetzel et al., 2012a). Activation or potentiation of group III mGlu receptors, in particular mGlu<sub>4</sub>, has therapeutic potential for Parkinson's disease (Jones et al., 2012; Marino et al., 2003), with mGlu<sub>4</sub> also a potential target for pain (Vilar et al., 2013) and autism (Becker et al., 2014).

### **Therapeutic targeting of mGlu's with allosteric modulators**

The prevailing strategy for GPCR-based drug discovery has been to target the endogenous ligand binding site, referred to as an orthosteric site, to competitively block or mimic the actions of the endogenous ligand. However, multiple receptor families and transporters recognize glutamate, and as such, glutamate binding pockets are highly conserved such that orthosteric ligands often lack sufficient selectivity. Based on this and difficulties in optimizing glutamate analogs as drug candidates, mGlu receptor-based drug discovery programs have turned their attention to alternative strategies to develop more selective agents. One such approach is to target allosteric binding sites that are topographically distinct from the orthosteric site, such that the receptor can be simultaneously bound by more than one ligand. These ligands are referred to as allosteric

MOL #97220

modulators and have the capacity to perturb the affinity and/or efficacy of an orthosteric ligand. Since allosteric sites are often located in regions of the receptor that show greater diversity between subtypes, allosteric modulators offer the potential for greater subtype selectivity than their orthosteric counterparts. Allosteric modulators may potentiate or inhibit the affinity and/or efficacy of an orthosteric ligand a property referred to as “cooperativity”. A potentiator that has positive cooperativity with the orthosteric ligand is a positive allosteric modulator (PAM); while an inhibitor is a negative allosteric modulator (NAM). Cooperativity is saturable, therefore allosteric modulators have the theoretical advantage of being safer in the case of overdose, as there is a limit to the modulatory effect. A third class of ligands that interact with allosteric sites but have neutral cooperativity with the orthosteric ligand are called neutral allosteric ligands (NAL) and represent excellent pharmacological tools for dissecting allosteric modulator pharmacology (Christopoulos et al., 2014). A third advantage of allosteric modulators is the potential to maintain spatial and temporal aspects of receptor activation, i.e. modulation will only occur when and where the orthosteric agonist is present. However, it is important to note that allosteric ligands may possess intrinsic efficacy (either positive or inverse) in addition to, or exclusive of, cooperativity with orthosteric ligands (Annoura et al., 1996; Duvoisin et al., 2010; Gregory et al., 2013a; Gregory et al., 2012; Lavreysen et al., 2013; Litschig et al., 1999; Niswender et al., 2008; Noetzel et al., 2012b; Rook et al., 2013). The first allosteric modulator discovered for the mGlu family was CPCCOEt, an mGlu<sub>1</sub> NAM of multiple orthosteric agonists, including glutamate (Annoura et al., 1996; Litschig et al., 1999). The past fifteen years has seen the discovery of selective allosteric modulators for most of the mGlu family and for many subtypes multiple chemotypes and different classes of allosteric ligands have been disclosed (Tables 1-3).

MOL #97220

### **Complexities of mGlu receptor allosteric modulator pharmacology**

**Stimulus-bias.** Despite definite success in the discovery of mGlu allosteric ligands, the pharmacology of allosteric modulators is inherently more complex than their orthosteric counterparts. The overall pharmacological effect of an allosteric ligand includes affinity and efficacy as well as the cooperativity between the allosteric and orthosteric ligands. With respect to ligand efficacy it has become increasingly accepted that occupancy and efficacy are not necessarily linearly linked. The concept that orthosteric ligands may differentially impact the G-protein coupling and other functional responses mediated by activation of the receptor has come to the fore in recent years (Figure 1). This phenomenon referred to as ‘stimulus-bias’ (although has had many other monikers including ‘agonist-directed trafficking of receptor stimulus’, ‘functional selectivity’ and ‘biased agonism’) has now been observed for many GPCRs, including mGlu<sub>1</sub>, mGlu<sub>4</sub>, mGlu<sub>7</sub> and mGlu<sub>8</sub> (Emery et al., 2012; Gregory et al., 2012; Jalan-Sakrikar et al., 2014). For example stimulus-bias is evident when comparing orthosteric agonists L-AP4 and LSP4-2022 with glutamate between Ca<sup>2+</sup> mobilization and GIRK channel activation at a single group III subtype (Jalan-Sakrikar et al., 2014). Interestingly, when these orthosteric agonist bias profiles are compared between mGlu<sub>7</sub> and either mGlu<sub>4</sub> or mGlu<sub>8</sub> there is a complete reversal in the direction of bias for the same ligands across the same two pathways. This finding highlights the important concept that the endogenous ligand itself will have a ‘natural bias’, revealed in this study by differences between different group III mGlu subtypes. Stimulus-bias is thought to arise from ligands stabilizing different complements of receptor conformations that favour one suite of receptor activation outcomes over others. It is perhaps not surprising then that there is preliminary evidence for stimulus-bias when comparing allosteric agonist activity to that



MOL #97220

with glutamate. For example, VU29 and CDPPB are PAM-agonists of mGlu<sub>5</sub> for both intracellular Ca<sup>2+</sup> mobilization and pERK1/2 and notably have higher intrinsic efficacy for pERK1/2 relative to Ca<sup>2+</sup> signalling (Gregory et al., 2012). Glutamate, however, shows the reverse, with higher efficacy for Ca<sup>2+</sup> signalling over pERK1/2. Similar observations have been reported for mGlu<sub>1</sub> PAM-agonists relative to glutamate between Ca<sup>2+</sup> mobilization and cAMP accumulation (Sheffler and Conn, 2008). Moreover, many of the mGlu<sub>5</sub> PAM-agonists for pERK1/2 are super-agonists, capable to achieving higher maximal responses than glutamate (Gregory et al., 2012). Although it should be noted that both ligand and system kinetics may contribute to these observations of biased agonism. Negative allosteric modulators are also often reported to be inverse agonists (Tables 1-3), however, these observations are assay dependent, likely a result of receptor constitutive activity and assay resolution. Comprehensive investigation into stimulus-bias that factors in these contributing mechanisms and confounds is sorely needed. An additional mechanism that may also contribute to observations of stimulus-bias relates to receptor compartmentalisation. For example, a large pool of intracellular mGlu<sub>5</sub> is located on the nuclear membrane (Jong et al., 2005; Kumar et al., 2008). Activation of mGlu<sub>5</sub> at the cell surface versus the nuclear membrane results in the activation of different signalling cascades, as demonstrated through the use of permeable/transported and impermeable/non-transported orthosteric and allosteric ligands of mGlu<sub>5</sub> (Jong et al., 2009; Kumar et al., 2012). Furthermore, activation of cell surface mGlu<sub>5</sub> leads to long-term potentiation (LTP) and long-term depression (LTD) in the hippocampus, whereas activation of intracellular mGlu<sub>5</sub> only mediates LTD (Purgert et al., 2014). These findings highlight that stimulus-bias may also arise from a differential ability of different chemotypes to access distinct subcellular compartments where

MOL #97220

receptors may be located. Therefore the lipophilicity or transportability of an allosteric ligand may impact the overall functional outcome observed.

**Biased modulation.** A related concept to biased agonism is the phenomenon of biased modulation (Figure 1). If we consider that the conformations engendered by the simultaneous occupation of a receptor with an orthosteric and allosteric ligand that give rise to cooperativity differ from those engendered by binding of orthosteric ligand alone. Therefore this raises the possibility that the ‘natural bias’ of an endogenous ligand may be influenced by an allosteric modulator. Conceptually, this may manifest as a change in the magnitude and/or direction of the cooperativity factors that describe the allosteric interaction depending on the receptor activation outcome measured. Alternatively, allosteric modulators may also show differential apparent affinity estimates depending upon the receptor activation outcome measured. In recent years, there have been a growing number of reports of biased mGlu allosteric modulation. For example at mGlu<sub>5</sub>, CPPHA and DFB are both PAMs for glutamate-mediated Ca<sup>2+</sup> mobilization in rat cortical astrocytes; however, DFB potentiates whereas CPPHA inhibits DHPG-stimulated pERK1/2 (Zhang et al., 2005). Further, multiple mGlu<sub>5</sub> modulators have lower positive cooperativity for glutamate-mediated pERK1/2 versus intracellular Ca<sup>2+</sup> mobilization (Gregory et al., 2012). Recent studies suggest that biased modulation has relevance in more physiological systems, raising the possibility that biased modulation may have clinical implications. For example, a novel mGlu<sub>5</sub> selective modulator from the same chemotype as CPPHA, NCFP, potentiated DHPG-induced depolarization of subthalamic nucleus neurons but lacked the ability to potentiate induction of long-term potentiation or long-term depression in the hippocampus (Noetzel et al., 2013). Importantly, another mGlu<sub>5</sub> allosteric modulator, VU0092273 that shared

MOL #97220

a similar pharmacological mGlu<sub>5</sub> PAM profile in recombinant cells, potentiated all three slice electrophysiological responses (Noetzel et al., 2013), suggesting that NCFP is showing true biased modulation in a native environment. Another novel mGlu<sub>5</sub> allosteric modulator, VU0409551, was recently disclosed that potentiates multiple glutamate-induced mGlu<sub>5</sub> signalling responses in cell-based assays, but has no efficacy in modulating hippocampal LTP or NMDA receptor currents (Rook et al., 2015). However, VU0409551 has efficacy in preclinical rodent models of psychosis and cognition (Rook et al., 2015), suggesting that biased allosteric modulators of mGlu<sub>5</sub> can possess preclinical efficacy.

Biased modulation is not limited to mGlu<sub>5</sub> PAMs, both positive and negative modulators of group III receptors have been suggested to induce biased receptor signaling (Niswender et al., 2010a; Jalan-Sakrikar et al., 2014). The extent and prevalence of biased modulators is likely to be highly underappreciated. This is due to the fact that the vast majority of mGlu allosteric modulators are characterized in a single cell-based assay, generally Ca<sup>2+</sup> mobilization, to classify and rank compounds on the basis of potency. Further, the therapeutic relevance of biased pharmacology has yet to be realized, however, it is tempting to speculate that unappreciated bias may contribute to differential efficacy of allosteric modulators in behavioural models. For example DPFE, an mGlu<sub>5</sub> PAM of glutamate, requires far lower doses for efficacy in cognitive models compared with reversing amphetamine-induced hyperlocomotion, a model for anti-psychotic efficacy (Gregory et al., 2013a). Moreover, there is the potential that biased modulation may be exploited to develop therapeutics that are not only subtype selective but also pathway selective; modulating receptor responses that yield therapeutic effects and avoiding those that give rise to on-target adverse effects.

MOL #97220

**Probe dependence.** A related phenomenon to biased modulation is that of probe dependence, where allosteric interactions are dependent upon the chemical nature of the two ligands under investigation. The result being that the direction and/or magnitude of cooperativity may be significantly different. There are multiple examples of probe dependence at the group II mGlu receptors. BINA potentiates  $\text{Ca}^{2+}$  oscillations in cortical neurons induced by the mGlu<sub>2</sub> selective orthosteric agonist, LY541850, but not the mixed group II orthosteric agonist, LY379268 (Sanger et al., 2012; Sanger et al., 2013). JNJ-40068782, a PAM-agonist of glutamate-stimulation of [<sup>35</sup>S]GTP $\gamma$ S binding and  $\text{Ca}^{2+}$  mobilization at mGlu<sub>2</sub>, has neutral cooperativity with the orthosteric antagonist radioligand [<sup>3</sup>H]LY341495 and positive cooperativity with the orthosteric agonist radioligand [<sup>3</sup>H]DCG-IV (Lavreysen et al., 2013). For group I and III mGlu receptors there are also select examples of probe dependence. Affinity modulation occurs between the orthosteric agonist quisqualate and mGlu<sub>5</sub> PAMs (Bradley et al., 2011); however, a separate study found no evidence for affinity modulation (i.e. neutral cooperativity) between glutamate and mGlu<sub>5</sub> PAMs of glutamate in functional assays (Gregory et al., 2012). At mGlu<sub>4</sub> and mGlu<sub>7</sub>, VU0155094 and VU0422288 have different degrees of cooperativity depending upon both the orthosteric agonist and the assay endpoint used (Jalan-Sakrikar et al., 2014). For example, at mGlu<sub>7</sub> VU0155094 is neutral with respect to the apparent affinity of LSP4-2022 and L-AP4 but a positive modulator of glutamate apparent affinity (Jalan-Sakrikar et al., 2014). Clearly, probe dependence is evident for the mGlu family and should be taken into consideration when designing screening paradigms and characterizing mGlu allosteric modulators. The choice of assay and/or probe can result in a very different pharmacological profile. This is of particular importance for mGlu<sub>7</sub> where L-AP4 is most commonly used as the orthosteric agonist given the low affinity of this subtype for glutamate (in the millimolar range). Moreover, extracellular  $\text{Ca}^{2+}$

MOL #97220

ions can activate mGlu<sub>1</sub> influencing the responses to orthosteric agonists, indicating that two endogenous orthosteric agonists need to be considered with respect to probe dependence but also that different endogenous ligands can allosterically interact at mGlu<sub>1</sub> (Jiang et al., 2014). Importantly, extracellular Ca<sup>2+</sup> activation of mGlu<sub>1</sub> can be modulated by small molecule positive and negative allosteric modulators of glutamate (Ro67-4853 and CPCCOEt) (Jiang et al., 2014). Probe dependence can also be an important consideration when assessing allosteric modulators in native systems (brain slice electrophysiology or primary neuronal cultures) where surrogate agonists are typically used to selectively activate a particular mGlu subtype and avoid the confounds of glutamate activating ionotropic glutamate receptors and being subject to endogenous transporters.

**Protein-protein interactions.** Another important consideration in assessing allosteric modulator pharmacology in native systems is the fact that the mGlu family are obligate dimers (El-Moustaine et al., 2012) that also have the capacity to form heteromers with other subtypes (Doumazane et al., 2011; Kammermeier, 2012; Sevastyanova and Kammermeier, 2014; Yin et al., 2014) as well as with other unrelated GPCRs (Cabello et al., 2009; Gama et al., 2001; Gonzalez-Maeso et al., 2008). Unsurprisingly, such complexity in receptor configurations can influence the pharmacology of small molecules. Heterodimerization of mGlu<sub>4</sub> and mGlu<sub>2</sub> has differential effects on the pharmacology of different allosteric ligands (Kammermeier, 2012; Yin et al., 2014). For example at the mGlu<sub>2/4</sub> heteromer PHCCC has lower cooperativity with L-AP4, whereas Lu AF21934 and VU0155041 have decreased affinity but higher cooperativity, compared with mGlu<sub>4</sub> homodimers (Yin et al., 2014). This perturbation in the pharmacology of allosteric ligands has consequences in vivo at synapses where mGlu<sub>2</sub> and mGlu<sub>4</sub> are co-

MOL #97220

expressed. VU0155041 and Lu AF21934 are both effective at potentiating the actions of L-AP4 at corticostriatal synapses (Gubellini et al., 2001; Yin et al., 2014). However, PHCCC does not potentiate the actions of L-AP4 at corticostriatal synapses (Yin et al., 2014), despite being an efficacious mGlu<sub>4</sub> PAM of L-AP4-induced responses at other synapses (Jones et al., 2008; Marino et al., 2003; Valenti et al., 2005). Direct protein-protein interactions are not necessarily required for a co-expressed receptor to impact the pharmacological effect of an mGlu ligand. Co-expression and co-activation of G<sub>q</sub>-coupled receptors enhances mGlu<sub>4</sub> stimulus-response coupling to orthosteric and allosteric agonists in intracellular Ca<sup>2+</sup> mobilization but not cAMP accumulation assays (Yin et al., 2012). It is apparent that cellular context, measure of receptor function and co-expression of other GPCRs can impact the pharmacology of small molecule allosteric ligands, leading to unanticipated consequences in native systems. However, this complexity extends further to the structure-activity relationships of allosteric modulators themselves.

**Molecular switches.** Given the complexities of allosteric modulation surrounding stimulus-bias and probe dependence it is perhaps not surprising that structure-activity relationships (SAR) for mGlu receptor allosteric modulators are notoriously challenging. SAR is often reported to be ‘steep’ with minimal changes resulting in a complete loss of activity (Conn et al., 2014; Lindsley, 2014). In addition, mGlu allosteric chemotypes often show ‘molecular switches’ where the same scaffold can give rise to modulators with negative, positive and neutral cooperativity with the same orthosteric agonist (Wood et al., 2011). The phenomenon was first observed during the development of DFB, an mGlu<sub>5</sub> PAM for glutamate-mediated intracellular Ca<sup>2+</sup> mobilization (O'Brien et al., 2003). Recently, this propensity for molecular switching was

MOL #97220

exploited with the development of a photoswitchable allosteric modulator for mGlu<sub>5</sub>, where negative cooperativity between with glutamate could be controlled by exposure to different light wavelengths (Pittolo et al., 2014). However, unexpected changes in cooperativity continue to be a challenge for medicinal chemistry programs and interpretation of SAR (Hammer et al., 1980; Lamb et al., 2011; Rodriguez et al., 2010; Sams et al., 2011; Schann et al., 2010; Sharma et al., 2009; Sheffler et al., 2012; Zhou et al., 2010). These difficulties in SAR interpretation can in part be attributed to a paucity in our understanding of the structural basis of allosterism. The majority of discovery programs rely on modulator potency curves in the presence of a single concentration of agonist in order to classify allosteric ligands as active/inactive and ascribe either positive or negative cooperativity. However, potency represents a composite measure of efficacy, affinity and cooperativity (Gregory et al., 2012). As such, modifications to a chemotype may influence any one of these three parameters. Therefore, more quantitative analysis of allosteric chemotypes is required to truly understand how SAR relates to efficacy, affinity and cooperativity. Going forward a key challenge will be coupling SAR interpretation as we improve our understanding of the structural basis of allosterism.

### **Structural basis of allosteric modulation**

The mGlu's are family C GPCRs typified by their large extracellular N-terminal Venus flytrap domain (VFT), linked via a cysteine-rich domain to the seven transmembrane spanning  $\alpha$ -helical domains (7TMs) that are the hallmark of the GPCR superfamily. The orthosteric site is located in the N-terminal domain and to date all small molecule allosteric modulators are thought to bind within the 7TMs. Historically, receptor chimeras proved valuable to 1) definitively demonstrate an allosteric mechanism of action for small molecule ligands and 2) to localise allosteric binding

MOL #97220

sites to the 7TM domains. Chimeric constructs swapping the VFT of one subtype onto another, or even exchange with non-mGlu family C GPCRs have been utilised. Chimeras between mGlu<sub>1</sub> and mGlu<sub>5</sub>, mGlu<sub>4</sub> and the Calcium-sensing receptor were used to validate the allosteric mechanism of action of mGlu<sub>1</sub> selective allosteric modulator CPCCOEt (Brauner-Osborne et al., 1999; Gasparini et al., 2001; Litschig et al., 1999). Subsequent studies used mGlu<sub>1/5</sub> chimeras to demonstrate MPEP (an mGlu<sub>5</sub> selective NAM) and multiple mGlu<sub>1</sub> PAMs recognised an overlapping binding site with that of CPCCOEt (Knoflach et al., 2001; Pagano et al., 2000). Chimeric constructs were also used to validate an allosteric mechanism of action for PHCCC, AMN082 and BAY36-7620 at mGlu<sub>4</sub>, mGlu<sub>7</sub> and mGlu<sub>1</sub> respectively (Carroll et al., 2001; Maj et al., 2003; Mitsukawa et al., 2005). Another approach to achieve gross localisation of allosteric binding sites is through the generation of N-terminal truncated receptors. The extracellular VFT can be removed to generate a “headless” mGlu, yielding a receptor construct that does not respond to glutamate, but can couple to intracellular signalling pathways (Goudet et al., 2004). Headless mGlu<sub>1</sub> and mGlu<sub>5</sub> are constitutively active such that negative allosteric modulators behave as inverse agonists (Chen et al., 2007; Goudet et al., 2004; Suzuki et al., 2007). Further, headless mGlu<sub>5</sub> can be stimulated by small molecule positive allosteric modulators that do not necessarily have intrinsic efficacy at the full-length receptor (Chen et al., 2007; Goudet et al., 2004; Noetzel et al., 2013). Interestingly, NCFP had little or no efficacy for stimulating headless mGlu<sub>5</sub>, despite having a similar magnitude of positive cooperativity as VU0092273, which can activate headless mGlu<sub>5</sub> (Noetzel et al., 2013). The headless mGlu 7TM is thought to behave as a monomer (Goudet et al., 2004), so perhaps this discrepancy relates to receptor dimerization in the full-length receptor being important either for NCFP binding or cooperativity.



MOL #97220

Two recent x-ray crystal structures were solved for the 7TMs of group I mGlu's with allosteric modulators bound (PDB ID's: 4OO9 & 4OR2; (Dore et al., 2014; Wu et al., 2014)). Prior to this, research relied on family A 7TM templates to perform homology modelling and interpret site-directed mutagenesis data. Despite low sequence homology (<20%), these approaches proved fruitful, localising allosteric binding pockets of mGlu<sub>1</sub>, mGlu<sub>2</sub>, mGlu<sub>4</sub> and mGlu<sub>5</sub> to a region analogous to that of the biogenic amine binding sites in family A GPCRs (Gregory et al., 2014; Gregory et al., 2013b; Lundstrom et al., 2011; Malherbe et al., 2003a; Malherbe et al., 2003b; Molck et al., 2012; Ott et al., 2000; Pagano et al., 2000; Rovira et al., 2015). In light of the new crystal structures, while alignments and secondary structure prediction of TM5-7 deviated from that observed in the mGlu<sub>1</sub> and mGlu<sub>5</sub> structures, the general localisation of the pocket was consistent with earlier reports. Many residues that had previously been identified in mutagenesis-based studies were recapitulated in the crystal structures (discussed in detail below). Ongoing research continues to refine our understanding of binding pocket/s within the 7TM bundle in particular in relation to determinants that govern selectivity, affinity, cooperativity and efficacy of allosteric modulators.

### **The common mGlu allosteric site**

The concept for a common allosteric site between mGlu subtypes arises from three lines of evidence: 1) allosteric modulators that lack subtype selectivity; 2) competitive interactions between diverse chemotypes and allosteric radioligands; 3) overlapping binding pockets based on mutagenesis studies and observed in x-ray crystal structures. While attaining selectivity with allosteric modulators is considerably more effective than for orthosteric ligands, there are a number of examples of allosteric modulators that have efficacy at multiple subtypes. MPEP is

MOL #97220

both a negative allosteric modulator of glutamate at mGlu<sub>5</sub> and a positive allosteric modulator of L-AP4 at mGlu<sub>4</sub> (Mathiesen et al., 2003). Conversely, DFB and CPPHA, which potentiate glutamate signalling at mGlu<sub>5</sub> are weak negative allosteric modulators of glutamate at mGlu<sub>4</sub> and mGlu<sub>8</sub> (O'Brien et al., 2003; O'Brien et al., 2004). In addition, PHCCC, an mGlu<sub>4</sub> PAM for glutamate and L-AP4, negatively modulates glutamate at mGlu<sub>1</sub> (Annoura et al., 1996; Maj et al., 2003). The recent report of a molecular switch to mGlu<sub>3</sub> NAM from an mGlu<sub>5</sub> PAM scaffold (Sheffler et al., 2012), reveals that the subtleties of interactions within this common pocket that dictate subtype selectivity are not limited to cross-over between group I and group III mGlu receptors alone. This lack of selectivity within the common binding pocket is also associated with differential cooperativity between subtypes, highlighting that the small molecule interactions within the common allosteric site can elicit either positive or negative cooperativity.

Identification of allosteric ligands that interact with this common mGlu allosteric site has been facilitated by the development of radiolabelled mGlu allosteric modulators. Multiple selective tritiated and positron emission tomography (PET) allosteric ligands have been developed for mGlu<sub>1</sub>, mGlu<sub>2</sub> and mGlu<sub>5</sub> (Ametamey et al., 2007; Anderson et al., 2003; Anderson et al., 2002; Baumann et al., 2010; Cosford et al., 2003b; Gasparini et al., 2002; Kohara et al., 2005; Lavreysen et al., 2003; Malherbe et al., 2006; Treyer et al., 2007). Through the use of inhibition binding assays with these allosteric ligands a direct assessment of competitive versus non-competitive binding modes can be performed (Chen et al., 2007; Hemstapat et al., 2006; Kinney et al., 2005; Lavreysen et al., 2003; Lavreysen et al., 2004; Lundstrom et al., 2011; Malherbe et al., 2006). Such experiments are often the first step towards identifying allosteric modulators that bind to alternative allosteric sites ((Chen et al., 2008; Hammond et al., 2010) discussed in further

MOL #97220

detail later). However, competition binding experiments may also be used to identify neutral allosteric ligands that interact with the common allosteric site but have neutral cooperativity with agonist/s in functional screening assays (Rodriguez et al., 2005; Sams et al., 2011).

Early mutational studies investigating group I mGlu receptor allosteric binding pockets exploited non-conserved residues within the 7TMs. Substitution of mGlu<sub>1</sub> non-conserved amino acids onto the equivalent positions in mGlu<sub>5</sub>, and vice versa, resulted in gain-of-function of selective allosteric modulators (Goudet et al., 2005; Knoflach et al., 2001; Litschig et al., 1999; Pagano et al., 2000; Surin et al., 2007). These mutagenesis studies pointed to a common location for allosteric binding pockets for group I mGlu receptors. Additional studies have shown that residues at positions 3.40 and 7.38 are important for allosteric modulation of both mGlu<sub>1</sub> and mGlu<sub>5</sub> (Ballesteros-Weinstein numbering (Ballesteros and Weinstein, 1995) adopted from mGlu<sub>1</sub> structural alignment to family A GPCRs in Wu et al., 2014; (Fukuda et al., 2009; Gregory et al., 2014; Gregory et al., 2013b; Gregory et al., 2012; Malherbe et al., 2006; Malherbe et al., 2003a; Malherbe et al., 2003b; Muhlemann et al., 2006; Suzuki et al., 2007; Turlington et al., 2014; Wu et al., 2014). Subsequently, key residues for the potency and/or binding of mGlu<sub>1</sub>, mGlu<sub>2</sub>, mGlu<sub>4</sub> and mGlu<sub>5</sub> allosteric modulators from diverse chemical scaffolds have been mapped to TMs 3-7. Across the four subtypes multiple residues have repeatedly been reported as contributing to allosteric modulation. For all four subtypes, residues in positions 3.36, 5.43 and 6.48 are crucial for both positive and negative allosteric modulators (Fukuda et al., 2009; Gregory et al., 2014; Gregory et al., 2013b; Gregory et al., 2012; Lundstrom et al., 2011; Malherbe et al., 2003a; Malherbe et al., 2006; Molck et al., 2012; Muhlemann et al., 2006; Pagano et al., 2000; Rovira et al., 2015; Turlington et al., 2014; Wu et al., 2014). Position 7.45 has also been implicated in

MOL #97220

mGlu<sub>4</sub> and mGlu<sub>5</sub> allosteric modulation (Gregory et al., 2014; Gregory et al., 2013b; Molck et al., 2012; Rovira et al., 2015; Turlington et al., 2014); there are no reports investigating the influence of this amino acid on selective allosteric modulators of mGlu<sub>1</sub> or mGlu<sub>2</sub>. Between group I mGlu receptors and mGlu<sub>2</sub>, three residues are common: 5.44, 5.47 and 6.55 (Fukuda et al., 2009; Gregory et al., 2014; Gregory et al., 2013b; Gregory et al., 2012; Hemstapat et al., 2006; Knoflach et al., 2001; Lundstrom et al., 2011; Malherbe et al., 2006; Malherbe et al., 2003b; Molck et al., 2012; Muhlemann et al., 2006; Rowe et al., 2008; Schaffhauser et al., 2003). Group 1 receptors and mGlu<sub>4</sub> share a common determinant in F6.51 (Fukuda et al., 2009; Gregory et al., 2014; Malherbe et al., 2003a; Malherbe et al., 2006; Malherbe et al., 2003b; Muhlemann et al., 2006; Rovira et al., 2015; Suzuki et al., 2007). Two residues at the top of TM3 (3.28 and 3.29) are important for allosteric modulation of both mGlu<sub>2</sub> and mGlu<sub>4</sub> (Lundstrom et al., 2011; Rovira et al., 2015), with position 3.29 also implicated in the activity of some mGlu<sub>5</sub> allosteric modulators (Malherbe et al., 2006; Malherbe et al., 2003b; Molck et al., 2012). Therefore these mutagenesis data reinforce the notion of overlapping binding pockets across the mGlu family.

Comparison of the recently solved mGlu<sub>1</sub> and mGlu<sub>5</sub> crystal structures reveals a similar architecture for the backbone of the transmembrane-spanning helices (Figure 2A). Overlay of the 7TM binding pockets for FITM and mavoglurant in the mGlu<sub>1</sub> and mGlu<sub>5</sub> respectively confirmed that these subtypes have overlapping allosteric binding pockets within the 7TMs (Dore et al., 2014; Wu et al., 2014). Examination of binding pockets for mavoglurant and FITM reveals that mavoglurant binds deeper within the 7TM bundle (Figure 2B). Side chain conformations of two key non-conserved residues in TM3 of mGlu<sub>5</sub>: P655<sup>3.36</sup> and S658<sup>3.39</sup> (mGlu<sub>1</sub> equivalents are: S668 and C671 respectively) create greater space within the pocket that

MOL #97220

allows mavoglurant to bind deeper within the 7TM bundle (Figure 2C). These crystal structures of closely related mGlu receptors have yielded structural insights with respect to how selectivity can be achieved in the common mGlu allosteric site.

### **Delineating affinity versus cooperativity determinants**

The vast majority of research on allosteric modulator SAR and structure-function has relied on measures of allosteric modulator potency; however, potency values represent a composite of affinity and cooperativity. In recent years the application of rigorous analytical pharmacological methods has facilitated delineation of the impact of mutations on allosteric modulator affinity, cooperativity and efficacy based on functional interaction data alone. By combining this research with earlier studies using radiolabelled allosteric modulators where affinity was determined or a loss of binding was observed we can start to map affinity determinants in the binding pocket. Residues known to be crucial affinity determinants are summarised in Table 4 (also see Supplementary Table 1) where the most well-characterised representatives from individual allosteric modulator scaffolds were included. These affinity determinants are also mapped onto the mGlu<sub>1</sub> and mGlu<sub>5</sub> structures (Figure 2A, B), Significant overlap is observed between mGlu<sub>1</sub> and mGlu<sub>5</sub>, although there is limited affinity data for mGlu<sub>1</sub>. For mGlu<sub>5</sub>, six residues have been implicated in the binding of most PAMs and NAMs: P655<sup>3,36</sup>, Y659<sup>3,40</sup>, T781<sup>6,44</sup>, W785<sup>6,48</sup>, S809<sup>7,45</sup> and A810<sup>7,46</sup> within the common allosteric site. Mapping the known affinity determinants for MPEP, the most well studied mGlu<sub>5</sub> allosteric ligand with respect to structure-function, onto the mGlu<sub>5</sub> crystal structure yields additional insights (Figure 4; (Gregory et al., 2014; Gregory et al., 2013b; Gregory et al., 2012; Malherbe et al., 2003b; Pagano et al., 2000)). Coloring residues on the basis of the impact of mutations reveals that substitutions that cause a

MOL #97220

moderate (<10-fold) reduction in affinity are located on the fringes of the mavoglurant binding pocket or are immediately adjacent to key residues (Figure 4, Supplementary Table 1).

In addition to effects on affinity, molecular determinants for a large set of allosteric modulators spanning multiple scaffolds with both positive and negative cooperativity have been explored at mGlu<sub>5</sub>. Some residues that are important for modulator affinity also contribute to cooperativity; however, there are select amino acids that only influence cooperativity. Negative cooperativity with glutamate of multiple allosteric modulators at mGlu<sub>5</sub> is decreased at P655<sup>3,36</sup> mutants (Gregory et al., 2014; Gregory et al., 2013b). Additionally mutations in TM3-7 (Y659<sup>3,40</sup>, N747<sup>5,47</sup>, T781<sup>6,44</sup>, W785<sup>6,48</sup>, S809<sup>7,45</sup>; (Gregory et al., 2014; Molck et al., 2012)) reduced cooperativity for negative modulators of glutamate at mGlu<sub>5</sub>. Notably, in mGlu<sub>1</sub> mutations of T794<sup>6,44</sup> and S822<sup>7,45</sup> had no effect on the interaction between glutamate and FITM (Wu 2014), suggesting that the structural determinants that govern cooperativity can be subtype selective even for conserved amino acids.

Negative cooperativity with glutamate at mGlu<sub>5</sub> was increased for M-5MPEP and/or VU0366248 when mutations were introduced to S658<sup>3,39</sup>, P743<sup>5,43</sup>, G748<sup>5,48</sup> and V789<sup>6,52</sup> (Gregory et al., 2014). Both M-5MPEP and VU0366248 have limited negative cooperativity at the wild type receptor, with saturating concentrations unable to completely abolish the glutamate response (Felts et al., 2010; Rodriguez et al., 2005). The cooperativity of other mGlu<sub>5</sub> negative allosteric modulators tested at these mutations was not discernibly different to wild-type (Gregory et al., 2014). However, given that these all have strong negative cooperativity at wild-type, fully abolishing the maximal glutamate response, further increases in negative cooperativity were likely beyond the limits of detection. For P743<sup>5,43</sup> this increase is opposite to

MOL #97220

that observed for the equivalent mGlu<sub>1</sub> mutation (Wu et al., 2014), again highlighting the subtype selective nature of cooperativity. Interestingly, two point mutations (N747<sup>5.47</sup> and S809<sup>7.45</sup>) had differential effects on allosteric ligands, actually increasing negative cooperativity of M-5MPEP and VU0366248 respectively (Figure 3C; Gregory et al., 2014). It is clear that the amino acids within the common allosteric site can differentially contribute to affinity and the manifestation of negative cooperativity in a manner that is very much ligand dependent.

With respect to positive allosteric modulators, no mutations have yet been demonstrated to unequivocally reduce cooperativity. However, there are multiple examples of mutations that increase the magnitude of positive cooperativity with glutamate (Figure 3C). Shared with the low cooperativity negative modulators, mutation of P743<sup>5.43</sup> and V789<sup>6.52</sup> also increased positive cooperativity of diverse allosteric modulators with glutamate at mGlu<sub>5</sub> (Gregory et al., 2014; Gregory et al., 2013b). Four residues (L744<sup>5.44</sup>, Y792<sup>6.55</sup>, A810<sup>7.46</sup> and C816<sup>7.52</sup>) are implicated in positive but not negative cooperativity (Gregory et al., 2014; Gregory et al., 2013b). Interestingly, compared to negative allosteric modulators, mutation of T781<sup>6.44</sup> and W785<sup>6.48</sup> had differential effects, increasing positive cooperativity with glutamate at mGlu<sub>5</sub> (Gregory et al., 2014; Gregory et al., 2013b). The complicated nature of cooperativity determinants is further highlighted through evidence of mutations that engender ‘molecular switches’ in allosteric modulator cooperativity (Figure 3D). Substitution of Y6.51 at mGlu<sub>5</sub> switches DFB from a PAM of glutamate to a NAM, whereas the reverse is true for YM298198 at mGlu<sub>1</sub> (Fukuda et al., 2009; Muhlemann et al., 2006). Multiple additional point mutations in TMs 3, 6 and 7 can also switch acetylenic PAMs to have either neutral or negative cooperativity with glutamate at mGlu<sub>5</sub> (Gregory et al., 2013b; Turlington et al., 2014). Mutation of conserved W785<sup>6.48</sup> in mGlu<sub>5</sub> can

MOL #97220

switch the cooperativity of select NAMs of glutamate to positive (Gregory et al., 2014). Collectively, these data highlight the subtleties and complexities of interactions within the common allosteric site that dictate whether or not the binding of a ligand will result in enhancement or inhibition of mGlu receptor function. Further, developing a better understanding of the dynamic ligand-receptor interactions that give rise to cooperativity will likely improve interpretation of allosteric modulator SAR and perhaps inform as to why ‘molecular switches’ are prevalent for some chemical scaffolds.

### **Additional mGlu allosteric sites**

As mentioned earlier, the development of radiolabelled allosteric ligands has identified allosteric ligands that are non-competitive. The first such ligand identified was CPPHA, a group I mGlu selective positive allosteric modulator of glutamate. CPPHA is unable to fully displace the mGlu<sub>5</sub> radioligands [<sup>3</sup>H]mPEPy and [<sup>3</sup>H]MPEP or compete with [<sup>3</sup>H]R214127 for binding to mGlu<sub>1</sub> (Bradley et al., 2011; Chen et al., 2008; Noetzel et al., 2013; O'Brien et al., 2004). Another structurally related selective mGlu<sub>5</sub> PAM of glutamate, NCFP, is also thought to bind to this alternate allosteric site (Noetzel et al., 2013). Potentiation of glutamate by both NCFP and CPPHA is non-competitively inhibited by the neutral allosteric ligand, 5MPEP (Chen et al., 2008; Noetzel et al., 2013). This indicates that the second allosteric site is conformationally linked to the common allosteric site such that ligand binding to these sites can allosterically influence one another. Isoleucine substitution of F1.42 reduces potentiation by CPPHA (both mGlu<sub>1</sub> and mGlu<sub>5</sub>) and NCFP (mGlu<sub>5</sub> only) (Chen et al., 2008; Noetzel et al., 2013). Interestingly, NCFP also loses either affinity or the capacity to allosterically modulate [<sup>3</sup>H]mPEPy binding and has very little efficacy as an agonist at an N-terminus truncated mGlu<sub>5</sub>



MOL #97220

(Noetzel et al., 2013). This suggests that the intact full-length receptor is required for this alternate allosteric site and/or receptor dimerization is also necessary. Also for mGlu<sub>5</sub>, VU0357121 and VU0400100 do not completely displace [<sup>3</sup>H]mPEPy binding (Gregory et al., 2012; Hammond et al., 2010; Rodriguez et al., 2012). Multiple mGlu<sub>1</sub> PAM scaffolds have also been flagged as binding to alternate allosteric site/s based on no or incomplete displacement of the allosteric radioligand [<sup>3</sup>H]R214127 (Hemstapat et al., 2006). Overall, these data strongly support the hypothesis that multiple allosteric sites are present on the group I mGlu receptors.

In the absence of an allosteric radioligand, multiple allosteric sites have also been proposed for mGlu<sub>4</sub>. The concentration-response curve for VU0155041 as an allosteric agonist at mGlu<sub>4</sub> was unaffected by co-application of PHCCC, an mGlu<sub>4</sub> PAM of glutamate and L-AP4 (Niswender et al., 2008). Recent mutational analysis of mGlu<sub>4</sub> allosteric modulators suggests that these two allosteric ligands interact with the common allosteric pocket made up of TM3-7 but in a non-overlapping manner (Rovira et al., 2015). Based on homology modelling and mutational data, PHCCC is proposed to bind deep within the allosteric site, analogous to the binding pose of mavoglurant in the mGlu<sub>5</sub> structure (Rovira et al., 2015). On the other hand, VU0155041 binds higher in the pocket, interacting with residues at the very top of the TM domains as well as extracellular loops 1 and 2 (Rovira et al., 2015). This suggests that two allosteric ligands can be accommodated simultaneously by mGlu<sub>4</sub>; however, at least for PHCCC and VU0155041, there is neutral cooperativity between the binding of these ligands. Interestingly, VU0415374 appears to be bitopic, with homology modelling proposing it can span both binding sites. It is tempting to speculate that the reason VU0415374 has ten-fold (or more) higher apparent affinity than other mGlu<sub>4</sub> positive allosteric modulators is due to its ability to engage more residues within the pocket. Further, it is interesting to consider that CPPHA has weak activity at mGlu<sub>4</sub> (Chen et al.,

MOL #97220

2008), as such there is the possibility that allosteric ligands may interact with mGlu<sub>4</sub> in three distinct binding modes. The extensive literature surrounding orthosteric and allosteric binding pockets of class A GPCRs may offer some insights into the location of alternative allosteric sites (Christopoulos, 2014). As confirmed in the recent co-crystal structure of the M2 muscarinic acetylcholine receptor, class A GPCR allosteric modulators recognise a binding pocket defined by the top of the TMs and the second and third extracellular loops (Kruse et al., 2013). Given the parallels between the common allosteric site of class C GPCRs and the orthosteric site in class A GPCRs, there is the possibility that alternate allosteric sites in class C GPCRs may also involve the extracellular surfaces of the receptor. Allosteric modulators that interact with different binding pockets are likely to engender a different complement of receptor conformational states than those that interact with the common allosteric site. Therefore the functional consequences of allosteric modulation through different binding pockets are likely to be divergent. Indeed, CPPHA and related compounds have biased pharmacology at mGlu<sub>5</sub> when compared to common site modulators (Noetzel et al., 2013; Zhang et al., 2005). We are just beginning to scratch the surface with respect to our understanding of what drives efficacy, cooperativity and affinity through the different allosteric sites across the entire mGlu family.

### **Concluding remarks**

Since the discovery of the first mGlu receptor allosteric modulator in 1996, the mGlu allosteric modulator pharmacological toolbox has continued to expand with selective allosteric ligands now available for the majority of mGlu family members. This toolbox includes chemically diverse allosteric ligands that may possess positive, negative or neutral cooperativity and/or positive or inverse agonism. At least two allosteric sites have been proposed, one of which is

MOL #97220

shared across mGlu receptor subtypes, but can still be selectively targeted. As to which allosteric site is best targeted for therapeutic efficacy remains unknown. Given the complexity introduced by the phenomenon of biased modulation, the optimal allosteric ligand pharmacological profile and the relative contribution of affinity, efficacy and cooperativity will likely be both pathophysiology and mGlu subtype dependent. With two x-ray crystal structures of the 7TMs for group I mGlu receptors now available, the field is now entering an exciting new era in our understanding of the dynamic interactions that govern allosteric ligand affinity, cooperativity and efficacy.

MOL #97220

### **Authorship Contributions**

Wrote or contributed to writing of manuscript: Gregory and Conn.

MOL #97220

## References

- Ametamey SM, Treyer V, Streffer J, Wyss MT, Schmidt M, Blagoev M, Hintermann S, Auberson Y, Gasparini F, Fischer UC and Buck A (2007) Human PET studies of metabotropic glutamate receptor subtype 5 with <sup>11</sup>C-ABP688. *J Nucl Med* **48**(2): 247-252.
- Anderson JJ, Bradbury MJ, Giracello DR, Chapman DF, Holtz G, Roppe J, King C, Cosford ND and Varney MA (2003) In vivo receptor occupancy of mGlu5 receptor antagonists using the novel radioligand [<sup>3</sup>H]3-methoxy-5-(pyridin-2-ylethynyl)pyridine). *Eur J Pharmacol* **473**(1): 35-40.
- Anderson JJ, Rao SP, Rowe B, Giracello DR, Holtz G, Chapman DF, Tehrani L, Bradbury MJ, Cosford ND and Varney MA (2002) [<sup>3</sup>H]Methoxymethyl-3-[(2-methyl-1,3-thiazol-4-yl)ethynyl]pyridine binding to metabotropic glutamate receptor subtype 5 in rodent brain: in vitro and in vivo characterization. *J Pharmacol Exp Ther* **303**(3): 1044-1051.
- Annoura H, Fukunaga A, Uesugi M, Tatsuoka T and Horikawa Y (1996) A novel class of antagonists for metabotropic glutamate receptors, 7-(hydroxyimino)cyclopropa[b]chromen-1a-carboxylates. *Bioorg Med Chem Lett* **6**(7): 763-766.
- Anwyl R (1999) Metabotropic glutamate receptors: electrophysiological properties and role in plasticity. *Brain Res Brain Res Rev* **29**(1): 83-120.
- Ayala JE, Chen Y, Banko JL, Sheffler DJ, Williams R, Telk AN, Watson NL, Xiang Z, Zhang Y, Jones PJ, Lindsley CW, Olive MF and Conn PJ (2009) mGluR5 positive allosteric modulators facilitate both hippocampal LTP and LTD and enhance spatial learning. *Neuropsychopharmacology* **34**(9): 2057-2071.
- Ayala JE, Niswender CM, Luo Q, Banko JL and Conn PJ (2008) Group III mGluR regulation of synaptic transmission at the SC-CA1 synapse is developmentally regulated. *Neuropharmacology* **54**(5): 804-814.
- Ballesteros J and Weinstein H (1995) Integrated methods for the construction of three dimensional models and computational probing of structure-function relations in G-protein coupled receptors. *Methods Neuroscience* **25**: 366-428.
- Baumann CA, Mu L, Wertli N, Kramer SD, Honer M, Schubiger PA and Ametamey SM (2010) Syntheses and pharmacological characterization of novel thiazole derivatives as potential mGluR5 PET ligands. *Bioorg Med Chem* **18**(16): 6044-6054.
- Becker JA, Cleese D, Spiegelhalter C, Schwab Y, Le Merrer J and Kieffer BL (2014) Autistic-like syndrome in mu opioid receptor null mice is relieved by facilitated mGluR4 activity. *Neuropsychopharmacology* **39**(9): 2049-2060.
- Bellone C, Luscher C and Mameli M (2008) Mechanisms of synaptic depression triggered by metabotropic glutamate receptors. *Cell Mol Life Sci* **65**(18): 2913-2923.
- Belozertseva IV, Kos T, Popik P, Danysz W, and Bessalov AY (2007) Antidepressant-like effects of mGluR1 and mGluR5 antagonists in the rat forced swim and the mouse tail suspension tests. *Eur Neuropsychopharmacol* **17**(3): 172-179.
- Bennett CE, Burnett DA, Greenlee WJ, Knutson CE, Korakas P, Li C, Tulshian D, Wu WL, Bertorelli R, Fredduzzi S, Grilli M, Lozza G, Reggiani A and Veltri A (2012) Fused tricyclic mGluR1 antagonists for the treatment of neuropathic pain. *Bioorg Med Chem Lett* **22**(4): 1575-1578.

MOL #97220

- Beqollari D and Kammermeier PJ (2008) The mGlu(4) receptor allosteric modulator N-phenyl-7-(hydroxyimino)cyclopropa[b]chromen-1a-carboxamide acts as a direct agonist at mGlu(6) receptors. *Eur J Pharmacol* **589**(1-3): 49-52.
- Berry-Kravis E, Hessel D, Coffey S, Hervey C, Schneider A, Yuhas J, Hutchison J, Snape M, Tranfaglia M, Nguyen DV and Hagerman R (2009) A pilot open label, single dose trial of fenobam in adults with fragile X syndrome. *J Med Genet* **46**(4): 266-271.
- Bradley SJ, Langmead CJ, Watson JM and Challiss RA (2011) Quantitative analysis reveals multiple mechanisms of allosteric modulation of the mGlu5 receptor in rat astroglia. *Mol Pharmacol* **79**(5): 874-885.
- Brauner-Osborne H, Jensen AA and Krosgaard-Larsen P (1999) Interaction of CPCCOEt with a chimeric mGlu1b and calcium sensing receptor. *Neuroreport* **10**(18): 3923-3925.
- Cabello N, Gandia J, Bertarelli DC, Watanabe M, Lluís C, Franco R, Ferre S, Lujan R and Ciruela F (2009) Metabotropic glutamate type 5, dopamine D2 and adenosine A2a receptors form higher-order oligomers in living cells. *J Neurochem* **109**(5): 1497-1507.
- Campo B, Kalinichev M, Lambeng N, El Yacoubi M, Royer-Urios I, Schneider M, Legrand C, Parron D, Girard F, Bessif A, Poli S, Vaugeois JM, Le Poul E and Celanire S (2011) Characterization of an mGluR2/3 negative allosteric modulator in rodent models of depression. *J Neurogenet* **25**(4): 152-166.
- Carroll FY, Stolle A, Beart PM, Voerste A, Brabet I, Mauler F, Joly C, Antonicek H, Bockaert J, Muller T, Pin JP and Prezeau L (2001) BAY36-7620: a potent non-competitive mGlu1 receptor antagonist with inverse agonist activity. *Mol Pharmacol* **59**(5): 965-973.
- Chaki S, Ago Y, Palucha-Paniewiera A, Matrisciano F and Pilc A (2013) mGlu2/3 and mGlu5 receptors: potential targets for novel antidepressants. *Neuropharmacology* **66**: 40-52.
- Chen Y, Goudet C, Pin JP and Conn PJ (2008) N-{4-Chloro-2-[(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)methyl]phenyl}-2-hydroxybenzamide (CPPHA) acts through a novel site as a positive allosteric modulator of group 1 metabotropic glutamate receptors. *Mol Pharmacol* **73**(3): 909-918.
- Chen Y, Nong Y, Goudet C, Hemstapat K, de Paulis T, Pin JP and Conn PJ (2007) Interaction of novel positive allosteric modulators of metabotropic glutamate receptor 5 with the negative allosteric antagonist site is required for potentiation of receptor responses. *Mol Pharmacol* **71**(5): 1389-1398.
- Christopoulos A (2014) Advances in G Protein-Coupled Receptor Allosterism: From Function to Structure. *Mol Pharmacol* **86**(5): 463-478.
- Christopoulos A, Changeux JP, Catterall WA, Fabbro D, Burriss TP, Cidlowski JA, Olsen RW, Peters JA, Neubig RR, Pin JP, Sexton PM, Kenakin TP, Ehlert FJ, Spedding M and Langmead CJ (2014) International union of basic and clinical pharmacology. XC. multisite pharmacology: recommendations for the nomenclature of receptor allosterism and allosteric ligands. *Pharmacol Rev* **66**(4): 918-947.
- Conn PJ, Lindsley CW, Meiler J and Niswender CM (2014) Opportunities and challenges in the discovery of allosteric modulators of GPCRs for treating CNS disorders. *Nat Rev Drug Discov* **13**(9): 692-708.
- Conn PJ and Pin JP (1997) Pharmacology and functions of metabotropic glutamate receptors. *Annu Rev Pharmacol Toxicol* **37**: 205-237.
- Cosford ND, Roppe J, Tehrani L, Schweiger EJ, Seiders TJ, Chaudary A, Rao S and Varney MA (2003b) [3H]-methoxymethyl-MTEP and [3H]-methoxy-PEPy: potent and selective

MOL #97220

- radioligands for the metabotropic glutamate subtype 5 (mGlu5) receptor. *Bioorg Med Chem Lett* **13**(3): 351-354.
- Coutinho V and Knopfel T (2002) Metabotropic glutamate receptors: electrical and chemical signaling properties. *Neuroscientist* **8**(6): 551-561.
- Dore AS, Okrasa K, Patel JC, Serrano-Vega M, Bennett K, Cooke RM, Errey JC, Jazayeri A, Khan S, Tehan B, Weir M, Wiggin GR and Marshall FH (2014) Structure of class C GPCR metabotropic glutamate receptor 5 transmembrane domain. *Nature* **511**(7511): 557-562.
- Doria JG, Silva FR, de Souza JM, Vieira LB, Carvalho TG, Reis HJ, Pereira GS, Dobransky T and Ribeiro FM (2013) Metabotropic glutamate receptor 5 positive allosteric modulators are neuroprotective in a mouse model of Huntington's disease. *Br J Pharmacol* **169**(4): 909-921.
- Doumazane E, Scholler P, Zwier JM, Trinquet E, Rondard P and Pin JP (2011) A new approach to analyze cell surface protein complexes reveals specific heterodimeric metabotropic glutamate receptors. *FASEB J* **25**(1): 66-77.
- Dunayevich E, Erickson J, Levine L, Landbloom R, Schoepp DD and Tollefson GD (2008) Efficacy and tolerability of an mGlu2/3 agonist in the treatment of generalized anxiety disorder. *Neuropsychopharmacology* **33**(7): 1603-1610.
- Duvoisin RM, Pfankuch T, Wilson JM, Grabell J, Chhajlani V, Brown DG, Johnson E and Raber J (2010) Acute pharmacological modulation of mGluR8 reduces measures of anxiety. *Behavioural brain research* **212**(2): 168-173.
- El Moustaine D, Granier S, Doumazane E, Scholler P, Rahmeh R, Bron P, Mouillac B, Baneres JL, Rondard P and Pin JP (2012) Distinct roles of metabotropic glutamate receptor dimerization in agonist activation and G-protein coupling. *Proc Natl Acad Sci USA* **109**(40): 16342-16347.
- Emery AC, Diraddo JO, Miller E, Hathaway HA, Pshenichkin S, Takoudjou GR, Grajkowska E, Yasuda RP, Wolfe BB and Wroblewski J (2012) Ligand Bias at Metabotropic Glutamate 1a Receptor: Molecular Determinants that Distinguish beta-arrestin from G Protein Mediated Signaling. *Mol Pharmacol* **82**(2): 95-105.
- Fell MJ, Witkin JM, Falcone JF, Katner JS, Perry KW, Hart J, Rorick-Kehn L, Overshiner CD, Rasmussen K, Chaney SF, Benvenga MJ, Li X, Marlow DL, Thompson LK, Luecke SK, Wafford KA, Seidel WF, Edgar DM, Quets AT, Felder CC, Wang X, Heinz BA, Nikolayev A, Kuo MS, Mayhugh D, Khilevich A, Zhang D, Ebert PJ, Eckstein JA, Ackermann BL, Swanson SP, Catlow JT, Dean RA, Jackson K, Tauscher-Wisniewski S, Marek GJ, Schkeryantz JM and Svensson KA (2011) N-(4-((2-(trifluoromethyl)-3-hydroxy-4-(isobutyl)phenoxy)methyl)benzyl)-1-methyl-1H-imidazole-4-carboxamide (THIIC), a novel metabotropic glutamate 2 potentiator with potential anxiolytic/antidepressant properties: in vivo profiling suggests a link between behavioral and central nervous system neurochemical changes. *J Pharmacol Exp Ther* **336**(1): 165-177.
- Felts AS, Lindsley SR, Lamb JP, Rodriguez AL, Menon UN, Jadhav S, Jones CK, Conn PJ, Lindsley CW and Emmitte KA (2010) 3-Cyano-5-fluoro-N-arylbenzamides as negative allosteric modulators of mGlu(5): Identification of easily prepared tool compounds with CNS exposure in rats. *Bioorg Med Chem Lett* **20**(15): 4390-4394.
- Fukuda J, Suzuki G, Kimura T, Nagatomi Y, Ito S, Kawamoto H, Ozaki S and Ohta H (2009) Identification of a novel transmembrane domain involved in the negative modulation of

MOL #97220

- mGluR1 using a newly discovered allosteric mGluR1 antagonist, 3-cyclohexyl-5-fluoro-6-methyl-7-(2-morpholin-4-ylethoxy)-4H-chromen-4-one. *Neuropharmacology* **57**(4): 438-445.
- Galici R, Jones CK, Hemstapat K, Nong Y, Echemendia NG, Williams LC, de Paulis T and Conn PJ (2006) Biphenyl-indanone A, a positive allosteric modulator of the metabotropic glutamate receptor subtype 2, has antipsychotic- and anxiolytic-like effects in mice. *J Pharmacol Exp Ther* **318**(1): 173-185.
- Gama L, Wilt SG and Breitwieser GE (2001) Heterodimerization of calcium sensing receptors with metabotropic glutamate receptors in neurons. *J Biol Chem* **276**(42): 39053-39059.
- Gasparini F, Andres H, Flor PJ, Heinrich M, Inderbitzin W, Lingenhohl K, Muller H, Munk VC, Omilusik K, Stierlin C, Stoehr N, Vranesic I and Kuhn R (2002) [(3)H]-M-MPEP, a potent, subtype-selective radioligand for the metabotropic glutamate receptor subtype 5. *Bioorg Med Chem Lett* **12**(3): 407-409.
- Gasparini F, Floersheim P, Flor PJ, Heinrich M, Inderbitzin W, Ott D, Pagano A, Stierlin C, Stoehr N, Vranesic I and Kuhn R (2001) Discovery and characterization of non-competitive antagonists of group I metabotropic glutamate receptors. *Farmacologia* **56**(1-2): 95-99.
- Gonzalez-Maeso J, Ang RL, Yuen T, Chan P, Weisstaub NV, Lopez-Gimenez JF, Zhou M, Okawa Y, Callado LF, Milligan G, Gingrich JA, Filizola M, Meana JJ and Sealfon SC (2008) Identification of a serotonin/glutamate receptor complex implicated in psychosis. *Nature* **452**(7183): 93-97.
- Goudet C, Gaven F, Kniazeff J, Vol C, Liu J, Cohen-Gonsaud M, Acher F, Prezeau L and Pin JP (2004) Heptahelical domain of metabotropic glutamate receptor 5 behaves like rhodopsin-like receptors. *Proc Natl Acad Sci USA* **101**(1): 378-383.
- Goudet C, Kniazeff J, Hlavackova V, Malhaire F, Maurel D, Acher F, Blahos J, Prezeau L and Pin JP (2005) Asymmetric functioning of dimeric metabotropic glutamate receptors disclosed by positive allosteric modulators. *J Biol Chem* **280**(26): 24380-24385.
- Gregory KJ, Herman EJ, Ramsey AJ, Hammond AS, Byun NE, Stauffer SR, Manka JT, Jadhav S, Bridges TM, Weaver CD, Niswender CM, Steckler T, Drinkenburg WH, Ahnaou A, Lavreysen H, Macdonald GJ, Bartolome JM, Mackie C, Hrupka BJ, Caron MG, Daigle TL, Lindsley CW, Conn PJ and Jones CK (2013a) N-aryl piperazine metabotropic glutamate receptor 5 positive allosteric modulators possess efficacy in pre-clinical models of NMDA hypofunction and cognitive enhancement. *J Pharmacol Exp Ther* **347**:438-457.
- Gregory KJ, Nguyen ED, Malosh C, Mendenhall JL, Zic JZ, Bates BS, Noetzel MJ, Squire EF, Turner EM, Rook JM, Emmitte KA, Stauffer SR, Lindsley CW, Meiler J and Conn PJ (2014) Identification of specific ligand-receptor interactions that govern binding and cooperativity of diverse modulators to a common metabotropic glutamate receptor 5 allosteric site. *ACS Chem Neurosci* **5**(4): 282-295.
- Gregory KJ, Nguyen ED, Reiff SD, Squire EF, Stauffer SR, Lindsley CW, Meiler J and Conn PJ (2013b) Probing the metabotropic glutamate receptor 5 (mGlu(5)) positive allosteric modulator (PAM) binding pocket: discovery of point mutations that engender a "molecular switch" in PAM pharmacology. *Mol Pharmacol* **83**(5): 991-1006.
- Gregory KJ, Noetzel MJ, Rook JM, Vinson PN, Stauffer SR, Rodriguez AL, Emmitte KA, Zhou Y, Chun AC, Felts AS, Chauder BA, Lindsley CW, Niswender CM and Conn PJ (2012) Investigating metabotropic glutamate receptor 5 allosteric modulator cooperativity,



MOL #97220

- affinity, and agonism: enriching structure-function studies and structure-activity relationships. *Mol Pharmacol* **82**(5): 860-875.
- Gubellini P, Saulle E, Centonze D, Bonsi P, Pisani A, Bernardi G, Conquet F and Calabresi P (2001) Selective involvement of mGlu1 receptors in corticostriatal LTD. *Neuropharmacology* **40**(7): 839-846.
- Hammer R, Berrie CP, Birdsall NJ, Burgen AS and Hulme EC (1980) Pirenzepine distinguishes between different subclasses of muscarinic receptors. *Nature* **283**(5742): 90-92.
- Hammond AS, Rodriguez AL, Townsend SD, Niswender CM, Gregory KJ, Lindsley CW and Conn PJ (2010) Discovery of a Novel Chemical Class of mGlu(5) Allosteric Ligands with Distinct Modes of Pharmacology. *ACS Chem Neurosci* **1**(10): 702-716.
- Hemstapat K, Da Costa H, Nong Y, Brady AE, Luo Q, Niswender CM, Tamagnan GD and Conn PJ (2007) A novel family of potent negative allosteric modulators of group II metabotropic glutamate receptors. *J Pharmacol Exp Ther* **322**(1): 254-264.
- Hemstapat K, de Paulis T, Chen Y, Brady AE, Grover VK, Alagille D, Tamagnan GD and Conn PJ (2006) A novel class of positive allosteric modulators of metabotropic glutamate receptor subtype 1 interact with a site distinct from that of negative allosteric modulators. *Mol Pharmacol* **70**(2): 616-626.
- Hughes ZA, Neal SJ, Smith DL, Sukoff Rizzo SJ, Pulicicchio CM, Lotarski S, Lu S, Dwyer JM, Brennan J, Olsen M, Bender CN, Kouranova E, Andree TH, Harrison JE, Whiteside GT, Springer D, O'Neil SV, Leonard SK, Schechter LE, Dunlop J, Rosenzweig-Lipson S, and Ring RH (2013) Negative allosteric modulation of metabotropic glutamate receptor 5 results in broad spectrum activity relevant to treatment resistant depression. *Neuropharmacology* **66**: 202-214.
- Iacovelli L, Felicioni M, Nistico R, Nicoletti F and De Blasi A (2014) Selective regulation of recombinantly expressed mGlu7 metabotropic glutamate receptors by G protein-coupled receptor kinases and arrestins. *Neuropharmacology* **77**: 303-312.
- Jalan-Sakrikar N, Field JR, Klar R, Mattmann ME, Gregory KJ, Zamorano R, Engers DW, Bollinger SR, Weaver CD, Days EL, Lewis LM, Utley TJ, Hurtado M, Rigault D, Acher F, Walker AG, Melancon BJ, Wood MR, Lindsley CW, Conn PJ, Xiang Z, Hopkins CR and Niswender CM (2014) Identification of Positive Allosteric Modulators VU0155094 (ML397) and VU0422288 (ML396) Reveals New Insights into the Biology of Metabotropic Glutamate Receptor 7. *ACS Chem Neurosci*.
- Jiang JY, Nagaraju M, Meyer RC, Zhang L, Hamelberg D, Hall RA, Brown EM, Conn PJ and Yang JJ (2014) Extracellular calcium modulates actions of orthosteric and allosteric ligands on metabotropic glutamate receptor 1alpha. *J Biol Chem* **289**(3): 1649-1661.
- Johnson MP, Barda D, Britton TC, Emkey R, Hornback WJ, Jagdmann GE, McKinzie DL, Nisenbaum ES, Tizzano JP and Schoepp DD (2005) Metabotropic glutamate 2 receptor potentiators: receptor modulation, frequency-dependent synaptic activity, and efficacy in preclinical anxiety and psychosis model(s). *Psychopharmacology (Berl)* **179**(1): 271-283.
- Jones CK, Bubser M, Thompson AD, Dickerson JW, Turle-Lorenzo N, Amalric M, Blobaum AL, Bridges TM, Morrison RD, Jadhav S, Engers DW, Italiano K, Bode J, Daniels JS, Lindsley CW, Hopkins CR, Conn PJ and Niswender CM (2012) The metabotropic glutamate receptor 4-positive allosteric modulator VU0364770 produces efficacy alone and in combination with L-DOPA or an adenosine 2A antagonist in preclinical rodent models of Parkinson's disease. *J Pharmacol Exp Ther* **340**(2): 404-421.

MOL #97220

- Jones PJ, Xiang Z and Conn PJ (2008) Metabotropic glutamate receptors mGluR4 and mGluR8 regulate transmission in the lateral olfactory tract-piriform cortex synapse. *Neuropharmacology* **55**(4): 440-446.
- Jong YJ, Kumar V, Kingston AE, Romano C and O'Malley KL (2005) Functional metabotropic glutamate receptors on nuclei from brain and primary cultured striatal neurons. Role of transporters in delivering ligand. *J Biol Chem* **280**(34): 30469-30480.
- Jong YJ, Kumar V and O'Malley KL (2009) Intracellular metabotropic glutamate receptor 5 (mGluR5) activates signaling cascades distinct from cell surface counterparts. *J Biol Chem* **284**(51): 35827-35838.
- Julio-Pieper M, Flor PJ, Dinan TG and Cryan JF (2011) Exciting times beyond the brain: metabotropic glutamate receptors in peripheral and non-neural tissues. *Pharmacol Rev* **63**(1): 35-58.
- Kalinichev M, Rouillier M, Girard F, Royer-Urios I, Bournique B, Finn T, Charvin D, Campo B, Le Poul E, Mutel V, Poli S, Neale SA, Salt TE and Lutjens R (2013) ADX71743, a potent and selective negative allosteric modulator of metabotropic glutamate receptor 7: in vitro and in vivo characterization. *J Pharmacol Exp Ther* **344**(3): 624-636.
- Kammermeier PJ (2012) Functional and pharmacological characteristics of metabotropic glutamate receptors 2/4 heterodimers. *Mol Pharmacol* **82**(3): 438-447.
- Keyword C, Wakefield M and Tack J (2009) A proof-of-concept study evaluating the effect of ADX10059, a metabotropic glutamate receptor-5 negative allosteric modulator, on acid exposure and symptoms in gastro-oesophageal reflux disease. *Gut* **58**(9): 1192-1199.
- Kinney GG, O'Brien JA, Lemaire W, Burno M, Bickel DJ, Clements MK, Chen TB, Wisnoski DD, Lindsley CW, Tiller PR, Smith S, Jacobson MA, Sur C, Duggan ME, Pettibone DJ, Conn PJ and Williams DL, Jr. (2005) A novel selective positive allosteric modulator of metabotropic glutamate receptor subtype 5 has in vivo activity and antipsychotic-like effects in rat behavioral models. *J Pharmacol Exp Ther* **313**(1): 199-206.
- Kinon BJ, Zhang L, Millen BA, Osuntokun OO, Williams JE, Kollack-Walker S, Jackson K, Kryzhanovskaya L and Jarkova N (2011) A multicenter, inpatient, phase 2, double-blind, placebo-controlled dose-ranging study of LY2140023 monohydrate in patients with DSM-IV schizophrenia. *J Clin Psychopharmacol* **31**(3): 349-355.
- Knoflach F, Mutel V, Jolidon S, Kew JN, Malherbe P, Vieira E, Wichmann J and Kemp JA (2001) Positive allosteric modulators of metabotropic glutamate 1 receptor: characterization, mechanism of action, and binding site. *Proc Natl Acad Sci USA* **98**(23): 13402-13407.
- Kohara A, Toya T, Tamura S, Watabiki T, Nagakura Y, Shitaka Y, Hayashibe S, Kawabata S and Okada M (2005) Radioligand binding properties and pharmacological characterization of 6-amino-N-cyclohexyl-N,3-dimethylthiazolo[3,2-a]benzimidazole-2-carboxamide (YM-298198), a high-affinity, selective, and noncompetitive antagonist of metabotropic glutamate receptor type 1. *J Pharmacol Exp Ther* **315**(1): 163-169.
- Kruse AC, Ring AM, Manglik A, Hu J, Eitel K, Hubner H, Pardon E, Valant C, Sexton PM, Christopoulos A, Felder CC, Gmeiner P, Steyaert J, Weis WI, Garcia KC, Wess J, Kobilka BK (2013) Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature* **504**(7478): 101-106.
- Kumar V, Fahey PG, Jong YJ, Ramanan N and O'Malley KL (2012) Activation of intracellular metabotropic glutamate receptor 5 in striatal neurons leads to up-regulation of genes

MOL #97220

- associated with sustained synaptic transmission including Arc/Arg3.1 protein. *J Biol Chem* **287**(8): 5412-5425.
- Kumar V, Jong YJ and O'Malley KL (2008) Activated nuclear metabotropic glutamate receptor mGlu5 couples to nuclear Gq/11 proteins to generate inositol 1,4,5-trisphosphate-mediated nuclear Ca<sup>2+</sup> release. *J Biol Chem* **283**(20): 14072-14083.
- Lamb JP, Engers DW, Niswender CM, Rodriguez AL, Venable DF, Conn PJ and Lindsley CW (2011) Discovery of molecular switches within the ADX-47273 mGlu5 PAM scaffold that modulate modes of pharmacology to afford potent mGlu5 NAMs, PAMs and partial antagonists. *Bioorg Med Chem Lett* **21**(9): 2711-2714.
- Lavreysen H, Janssen C, Bischoff F, Langlois X, Leysen JE and Lesage AS (2003) [3H]R214127: a novel high-affinity radioligand for the mGlu1 receptor reveals a common binding site shared by multiple allosteric antagonists. *Mol Pharmacol* **63**(5): 1082-1093.
- Lavreysen H, Langlois X, Ahnaou A, Drinkenburg W, te Riele P, Biesmans I, Van der Linden I, Peeters L, Megens A, Wintmolders C, Cid JM, Trabanco AA, Andres JI, Dautzenberg FM, Lutjens R, Macdonald G and Attack JR (2013) Pharmacological characterization of JNJ-40068782, a new potent, selective, and systemically active positive allosteric modulator of the mGlu2 receptor and its radioligand [3H]JNJ-40068782. *J Pharmacol Exp Ther* **346**(3): 514-527.
- Lavreysen H, Wouters R, Bischoff F, Nobrega Pereira S, Langlois X, Blokland S, Somers M, Dillen L and Lesage AS (2004) JNJ16259685, a highly potent, selective and systemically active mGlu1 receptor antagonist. *Neuropharmacology* **47**(7): 961-972.
- Li X, Need AB, Baez M, and Witkin JM (2006) Metabotropic glutamate 5 receptor antagonism is associated with antidepressant-like effects in mice *J Pharmacol Exp Ther* **319**(1): 254-259.
- Lindsley CW (2014) 2013 Philip S. Portoghese Medicinal Chemistry Lectureship: drug discovery targeting allosteric sites. *J Med Chem* **57**(18): 7485-7498.
- Litschig S, Gasparini F, Rueegg D, Stoehr N, Flor PJ, Vranesic I, Prezeau L, Pin JP, Thomsen C and Kuhn R (1999) CPCCOEt, a noncompetitive metabotropic glutamate receptor 1 antagonist, inhibits receptor signaling without affecting glutamate binding. *Mol Pharmacol* **55**(3): 453-461.
- Lundstrom L, Bissantz C, Beck J, Wettstein JG, Woltering TJ, Wichmann J and Gatti S (2011) Structural determinants of allosteric antagonism at metabotropic glutamate receptor 2: mechanistic studies with new potent negative allosteric modulators. *Br J Pharmacol* **164**(2b): 521-537.
- Ma AWS, Redka DS, Pisterzi LF, Angers S and Wells JW (2007) Recovery of Oligomers and Cooperativity When Monomers of the M2 Muscarinic Cholinergic Receptor Are Reconstituted into Phospholipid Vesicles. *Biochemistry* **46**(26): 7907-7927.
- Mabire D, Coupa S, Adelinet C, Poncelet A, Simonnet Y, Venet M, Wouters R, Lesage AS, Van Beijsterveldt L and Bischoff F (2005) Synthesis, structure-activity relationship, and receptor pharmacology of a new series of quinoline derivatives acting as selective, noncompetitive mGlu1 antagonists. *J Med Chem* **48**(6): 2134-2153.
- Maj M, Bruno V, Dragic Z, Yamamoto R, Battaglia G, Inderbitzin W, Stoehr N, Stein T, Gasparini F, Vranesic I, Kuhn R, Nicoletti F and Flor PJ (2003) (-)-PHCCC, a positive allosteric modulator of mGluR4: characterization, mechanism of action, and neuroprotection. *Neuropharmacology* **45**(7): 895-906.

MOL #97220

- Malherbe P, Kratochwil N, Knoflach F, Zenner MT, Kew JN, Kratzeisen C, Maerki HP, Adam G and Mutel V (2003a) Mutational analysis and molecular modeling of the allosteric binding site of a novel, selective, noncompetitive antagonist of the metabotropic glutamate 1 receptor. *J Biol Chem* **278**(10): 8340-8347.
- Malherbe P, Kratochwil N, Muhlemann A, Zenner M-T, Fischer C, Stahl M, Gerber PR, Jaeschke G and Porter RHP (2006) Comparison of the binding pockets of two chemically unrelated allosteric antagonists of the mGlu5 receptor and identification of crucial residues involved in the inverse agonism of MPEP. *J Neurochem* **98**(2): 601-615.
- Malherbe P, Kratochwil N, Zenner MT, Piussi J, Diener C, Kratzeisen C, Fischer C and Porter RH (2003b) Mutational analysis and molecular modeling of the binding pocket of the metabotropic glutamate 5 receptor negative modulator 2-methyl-6-(phenylethynyl)-pyridine. *Mol Pharmacol* **64**(4): 823-832.
- Marino MJ, Williams DL, Jr., O'Brien JA, Valenti O, McDonald TP, Clements MK, Wang R, DiLella AG, Hess JF, Kinney GG and Conn PJ (2003) Allosteric modulation of group III metabotropic glutamate receptor 4: a potential approach to Parkinson's disease treatment. *Proc Nat Acad Sci USA* **100**(23): 13668-13673.
- Mathiesen JM, Svendsen N, Brauner-Osborne H, Thomsen C and Ramirez MT (2003) Positive allosteric modulation of the human metabotropic glutamate receptor 4 (hmGluR4) by SIB-1893 and MPEP. *Br J Pharmacol* **138**(6): 1026-1030.
- Mitsukawa K, Yamamoto R, Ofner S, Nozulak J, Pescott O, Lukic S, Stoehr N, Mombereau C, Kuhn R, McAllister KH, van der Putten H, Cryan JF and Flor PJ (2005) A selective metabotropic glutamate receptor 7 agonist: activation of receptor signaling via an allosteric site modulates stress parameters in vivo. *Proc Nat Acad Sci USA* **102**(51): 18712-18717.
- Moghaddam B (2004) Targeting metabotropic glutamate receptors for treatment of the cognitive symptoms of schizophrenia. *Psychopharmacology* **174**(1): 39-44.
- Molck C, Harpsoe K, Gloriam DE, Clausen RP, Madsen U, Pedersen LO, Jimenez HN, Nielsen SM, Mathiesen JM and Brauner-Osborne H (2012) Pharmacological Characterization and Modeling of the Binding Sites of Novel 1,3-bis(pyridinylethynyl)benzenes as Metabotropic Glutamate Receptor 5-selective Negative Allosteric Modulators. *Mol Pharmacol* **82**:929-937.
- Moussawi K and Kalivas PW (2010) Group II metabotropic glutamate receptors (mGlu2/3) in drug addiction. *Eur J Pharmacol* **639**(1-3): 115-122.
- Muhlemann A, Ward NA, Kratochwil N, Diener C, Fischer C, Stucki A, Jaeschke G, Malherbe P and Porter RH (2006) Determination of key amino acids implicated in the actions of allosteric modulation by 3,3'-difluorobenzaldazine on rat mGlu5 receptors. *Eur J Pharmacol* **529**(1-3): 95-104.
- Niswender CM, Johnson KA, Miller NR, Ayala JE, Luo Q, Williams R, Saleh S, Orton D, Weaver CD and Conn PJ (2010a) Context-dependent pharmacology exhibited by negative allosteric modulators of metabotropic glutamate receptor 7. *Mol Pharmacol* **77**(3): 459-468.
- Niswender CM, Johnson KA, Miller NR, Ayala JE, Luo Q, Williams R, Saleh S, Orton D, Weaver CD and Conn PJ (2010b) Context-dependent pharmacology exhibited by negative allosteric modulators of metabotropic glutamate receptor 7. *Mol Pharmacol* **77**(3): 459-468.

MOL #97220

- Niswender CM, Johnson KA, Weaver CD, Jones CK, Xiang Z, Luo Q, Rodriguez AL, Marlo JE, de Paulis T, Thompson AD, Days EL, Nalywajko T, Austin CA, Williams MB, Ayala JE, Williams R, Lindsley CW and Conn PJ (2008) Discovery, characterization, and antiparkinsonian effect of novel positive allosteric modulators of metabotropic glutamate receptor 4. *Mol Pharmacol* **74**(5): 1345-1358.
- Noetzel MJ, Gregory KJ, Vinson PN, Manka JT, Stauffer SR, Lindsley CW, Niswender CM, Xiang Z and Conn PJ (2013) A Novel Metabotropic Glutamate Receptor 5 Positive Allosteric Modulator Acts at a Unique Site and Confers Stimulus Bias to mGlu5 Signaling. *Mol Pharmacol* **83**:835-847.
- Noetzel MJ, Jones CK and Conn PJ (2012a) Emerging approaches for treatment of schizophrenia: modulation of glutamatergic signaling. *Discov Med* **14**(78): 335-343.
- Noetzel MJ, Rook JM, Vinson PN, Cho HP, Days E, Zhou Y, Rodriguez AL, Lavreysen H, Stauffer SR, Niswender CM, Xiang Z, Daniels JS, Jones CK, Lindsley CW, Weaver CD and Conn PJ (2012b) Functional impact of allosteric agonist activity of selective positive allosteric modulators of metabotropic glutamate receptor subtype 5 in regulating central nervous system function. *Mol Pharmacol* **81**(2): 120-133.
- O'Brien JA, Lemaire W, Chen TB, Chang RS, Jacobson MA, Ha SN, Lindsley CW, Schaffhauser HJ, Sur C, Pettibone DJ, Conn PJ and Williams DL, Jr. (2003) A family of highly selective allosteric modulators of the metabotropic glutamate receptor subtype 5. *Mol Pharmacol* **64**(3): 731-740.
- O'Brien JA, Lemaire W, Wittmann M, Jacobson MA, Ha SN, Wisnoski DD, Lindsley CW, Schaffhauser HJ, Rowe B, Sur C, Duggan ME, Pettibone DJ, Conn PJ and Williams DL, Jr. (2004) A novel selective allosteric modulator potentiates the activity of native metabotropic glutamate receptor subtype 5 in rat forebrain. *J Pharmacol Exp Ther* **309**(2): 568-577.
- Odagaki Y, Kinoshita M and Toyoshima R (2011) Functional coupling between metabotropic glutamate receptors and G-proteins in rat cerebral cortex assessed by guanosine-5'-O-(3-[(35)S]thio)triphosphate binding assay. *Basic Clin Pharmacol Toxicol* **109**(3): 175-185.
- Odagaki Y, Kinoshita M and Toyoshima R (2013) Group II metabotropic glutamate receptor-mediated activation of G-proteins in rat hippocampal and striatal membranes. *Neuroscience letters* **539**: 22-26.
- Ott D, Floersheim P, Inderbitzin W, Stoehr N, Francotte E, Lecis G, Richert P, Rihs G, Flor PJ, Kuhn R and Gasparini F (2000) Chiral resolution, pharmacological characterization, and receptor docking of the noncompetitive mGlu1 receptor antagonist (+/-)-2-hydroxyimino-1a, 2-dihydro-1H-7-oxacyclopropa[b]naphthalene-7a-carboxylic acid ethyl ester. *J Med Chem* **43**(23): 4428-4436.
- Pagano A, Ruegg D, Litschig S, Stoehr N, Stierlin C, Heinrich M, Floersheim P, Prezeau L, Carroll F, Pin JP, Cambria A, Vranesic I, Flor PJ, Gasparini F and Kuhn R (2000) The non-competitive antagonists 2-methyl-6-(phenylethynyl)pyridine and 7-hydroxyiminocyclopropan[b]chromen-1a-carboxylic acid ethyl ester interact with overlapping binding pockets in the transmembrane region of group I metabotropic glutamate receptors. *J Biol Chem* **275**(43): 33750-33758.
- Parmentier-Batteur S, Hutson PH, Menzel K, Uslander JM, Mattson BA, O'Brien JA, Magliaro BC, Forest T, Stump CA, Tynebor RM, Anthony NJ, Tucker TJ, Zhang XF, Gomez R, Huszar SL, Lambeng N, Faure H, Le Poul E, Poli S, Rosahl TW, Rocher JP, Hargreaves R and Williams TM (2014) Mechanism based neurotoxicity of mGlu5 positive allosteric

MOL #97220

- modulators-development challenges for a promising novel antipsychotic target. *Neuropharmacology* **82**: 161-173.
- Patil ST, Zhang L, Martenyi F, Lowe SL, Jackson KA, Andreev BV, Avedisova AS, Bardenstein LM, Gurovich IY, Morozova MA, Mosolov SN, Neznanov NG, Reznik AM, Smulevich AB, Tochilov VA, Johnson BG, Monn JA and Schoepp DD (2007) Activation of mGlu2/3 receptors as a new approach to treat schizophrenia: a randomized Phase 2 clinical trial. *Nat Med* **13**(9): 1102-1107.
- Pelkey KA, Yuan X, Lavezzari G, Roche KW and McBain CJ (2007) mGluR7 undergoes rapid internalization in response to activation by the allosteric agonist AMN082. *Neuropharmacology* **52**(1): 108-117.
- Pinheiro PS and Mulle C (2008) Presynaptic glutamate receptors: physiological functions and mechanisms of action. *Nat Rev Neurosci* **9**(6): 423-436.
- Pittolo S, Gomez-Santacana X, Eckelt K, Rovira X, Dalton J, Goudet C, Pin JP, Llobet A, Giraldo J, Llebaria A and Gorostiza P (2014) An allosteric modulator to control endogenous G protein-coupled receptors with light. *Nat Chem Biol* **10**(10): 813-815.
- Porter RH, Jaeschke G, Spooren W, Ballard TM, Buttleman B, Kolczewski S, Peter JU, Prinssen E, Wichmann J, Vieira E, Muhlemann A, Gatti S, Mutel V and Malherbe P (2005) Fenobam: a clinically validated nonbenzodiazepine anxiolytic is a potent, selective, and noncompetitive mGlu5 receptor antagonist with inverse agonist activity. *J Pharmacol Exp Ther* **315**(2): 711-721.
- Purgert CA, Izumi Y, Jong YJ, Kumar V, Zorumski CF and O'Malley KL (2014) Intracellular mGluR5 can mediate synaptic plasticity in the hippocampus. *J Neurosci* **34**(13): 4589-4598.
- Rodriguez AL, Grier MD, Jones CK, Herman EJ, Kane AS, Smith RL, Williams R, Zhou Y, Marlo JE, Days EL, Blatt TN, Jadhav S, Menon UN, Vinson PN, Rook JM, Stauffer SR, Niswender CM, Lindsley CW, Weaver CD and Conn PJ (2010) Discovery of novel allosteric modulators of metabotropic glutamate receptor subtype 5 reveals chemical and functional diversity and in vivo activity in rat behavioral models of anxiolytic and antipsychotic activity. *Mol Pharmacol* **78**(6): 1105-1123.
- Rodriguez AL, Nong Y, Sekaran NK, Alagille D, Tamagnan GD and Conn PJ (2005) A close structural analog of 2-methyl-6-(phenylethynyl)-pyridine acts as a neutral allosteric site ligand on metabotropic glutamate receptor subtype 5 and blocks the effects of multiple allosteric modulators. *Mol Pharmacol* **68**(6): 1793-1802.
- Rodriguez AL, Zhou Y, Williams R, Weaver CD, Vinson PN, Dawson ES, Steckler T, Lavreysen H, Mackie C, Bartolome JM, Macdonald GJ, Daniels JS, Niswender CM, Jones CK, Conn PJ, Lindsley CW and Stauffer SR (2012) Discovery and SAR of a novel series of non-MPEP site mGlu(5) PAMs based on an aryl glycine sulfonamide scaffold. *Bioorg Med Chem Lett* **22**(24): 7388-7392.
- Rook JM, Noetzel MJ, Pouliot WA, Bridges TM, Vinson PN, Cho HP, Zhou Y, Gogliotti RD, Manka JT, Gregory KJ, Stauffer SR, Dudek FE, Xiang Z, Niswender CM, Daniels JS, Jones CK, Lindsley CW and Conn PJ (2013) Unique signaling profiles of positive allosteric modulators of metabotropic glutamate receptor subtype 5 determine differences in in vivo activity. *Biol Psychiatry* **73**(6): 501-509.
- Rook JM, Xiang Z, Lv X, Ghoshal A, Dickerson JW, Bridges TM, Johnson KA, Foster DJ, Gregory KJ, Vinson PN, Thompson AD, Byun N, Collier RL, Bubser M, Nedelcovych MT, Gould RW, Stauffer SR, Daniels JS, Niswender CM, Lavreysen H, Mackie C,

MOL #97220

- Conde-Ceide S, Alcazar J, Bartolome JM, Macdonald GJ, Steckler T, Jones CK, Lindsley CW and Conn PJ (2015) Biased mGlu<sub>5</sub> positive allosteric modulators provide *in vivo* efficacy without potentiating mGlu<sub>5</sub> modulation of NMDAR currents. *Neuron* (in press).
- Rovira X, Malhaire F, Scholler P, Rodrigo J, Gonzalez-Bulnes P, Llebaria A, Pin JP, Giraldo J and Goudet C (2015) Overlapping binding sites drive allosteric agonism and positive cooperativity in type 4 metabotropic glutamate receptors. *FASEB J* **29**(1): 116-130.
- Rowe BA, Schaffhauser H, Morales S, Lubbers LS, Bonnefous C, Kamenecka TM, McQuiston J and Daggett LP (2008) Transposition of three amino acids transforms the human metabotropic glutamate receptor (mGluR)-3-positive allosteric modulation site to mGluR2, and additional characterization of the mGluR2-positive allosteric modulation site. *J Pharmacol Exp Ther* **326**(1): 240-251.
- Rylander D, Iderberg H, Li Q, Dekundy A, Zhang J, Li H, Baishen R, Danysz W, Bezard E and Cenci MA (2010) A mGluR5 antagonist under clinical development improves L-DOPA-induced dyskinesia in parkinsonian rats and monkeys. *Neurobiol Dis* **39**(3): 352-361.
- Sams AG, Mikkelsen GK, Brodbeck RM, Pu X and Ritzen A (2011) Efficacy switching SAR of mGluR5 allosteric modulators: highly potent positive and negative modulators from one chemotype. *Bioorg Med Chem Lett* **21**(11): 3407-3410.
- Sanger H, Hanna L, Colvin EM, Grubisha O, Ursu D, Heinz BA, Findlay JD, Vivier RG, Sher E, Lodge D, Monn JA and Broad LM (2013) Pharmacological profiling of native group II metabotropic glutamate receptors in primary cortical neuronal cultures using a FLIPR. *Neuropharmacology* **66**: 264-273.
- Scandroglio P, Brusa R, Lozza G, Mancini I, Petro R, Reggiani A and Beltramo M (2010) Evaluation of cannabinoid receptor 2 and metabotropic glutamate receptor 1 functional responses using a cell impedance-based technology. *J Biomol Screen* **15**(10): 1238-1247.
- Schaffhauser H, Rowe BA, Morales S, Chavez-Noriega LE, Yin R, Jachec C, Rao SP, Bain G, Pinkerton AB, Vernier JM, Bristow LJ, Varney MA and Daggett LP (2003) Pharmacological characterization and identification of amino acids involved in the positive modulation of metabotropic glutamate receptor subtype 2. *Mol Pharmacol* **64**(4): 798-810.
- Schann S, Mayer S, Franchet C, Frauli M, Steinberg E, Thomas M, Baron L and Neuville P (2010) Chemical switch of a metabotropic glutamate receptor 2 silent allosteric modulator into dual metabotropic glutamate receptor 2/3 negative/positive allosteric modulators. *J Med Chem* **53**(24): 8775-8779.
- Schoepp DD, Wright RA, Levine LR, Gaydos B and Potter WZ (2003) LY354740, an mGlu2/3 receptor agonist as a novel approach to treat anxiety/stress. *Stress* **6**(3): 189-197.
- Sevastyanova TN and Kammermeier PJ (2014) Cooperative signaling between homodimers of metabotropic glutamate receptors 1 and 5. *Mol Pharmacol* **86**(5): 492-504.
- Sharma S, Kedrowski J, Rook JM, Smith RL, Jones CK, Rodriguez AL, Conn PJ and Lindsley CW (2009) Discovery of molecular switches that modulate modes of metabotropic glutamate receptor subtype 5 (mGlu5) pharmacology in vitro and in vivo within a series of functionalized, regioisomeric 2- and 5-(phenylethynyl)pyrimidines. *J Med Chem* **52**(14): 4103-4106.
- Sheffler DJ and Conn PJ (2008) Allosteric potentiators of metabotropic glutamate receptor subtype 1a differentially modulate independent signaling pathways in baby hamster kidney cells. *Neuropharmacology* **55**(4): 419-427.

MOL #97220

- Sheffler DJ, Wenthur CJ, Bruner JA, Carrington SJS, Vinson PN, Gogi KK, Blobaum AL, Morrison RD, Vamos M, Cosford NDP, Stauffer SR, Scott Daniels J, Niswender CM, Jeffrey Conn P and Lindsley CW (2012) Development of a novel, CNS-penetrant, metabotropic glutamate receptor 3 (mGlu3) NAM probe (ML289) derived from a closely related mGlu5 PAM. *Bioorg Med Chem Lett* **22**(12): 3921-3925.
- Sukoff Rizzo SJ, Leonard SK, Gilbert A, Dollings P, Smith DL, Zhang MY, Di L, Platt BJ, Neal S, Dwyer JM, Bender CN, Zhang J, Lock T, Kowal D, Kramer A, Randall A, Huselton C, Vishwanathan K, Tse SY, Butera J, Ring RH, Rosenzweig-Lipson S, Hughes ZA and Dunlop J (2011) The metabotropic glutamate receptor 7 allosteric modulator AMN082: a monoaminergic agent in disguise? *J Pharmacol Exp Ther* **338**(1): 345-352.
- Surin A, Pshenichkin S, Grajkowska E, Surina E and Wroblewski JT (2007) Cyclothiazide selectively inhibits mGluR1 receptors interacting with a common allosteric site for non-competitive antagonists. *Neuropharmacology* **52**(3): 744-754.
- Suzuki G, Kimura T, Satow A, Kaneko N, Fukuda J, Hikichi H, Sakai N, Maehara S, Kawagoe-Takaki H, Hata M, Azuma T, Ito S, Kawamoto H and Ohta H (2007) Pharmacological characterization of a new, orally active and potent allosteric metabotropic glutamate receptor 1 antagonist, 4-[1-(2-fluoropyridin-3-yl)-5-methyl-1H-1,2,3-triazol-4-yl]-N-isopropyl-N-methyl-3,6-dihydropyridine-1(2H)-carboxamide (FTIDC). *J Pharmacol Exp Ther* **321**(3): 1144-1153.
- Treyer V, Streffer J, Wyss MT, Bettio A, Ametamey SM, Fischer U, Schmidt M, Gasparini F, Hock C and Buck A (2007) Evaluation of the metabotropic glutamate receptor subtype 5 using PET and 11C-ABP688: assessment of methods. *J Nucl Med* **48**(7): 1207-1215.
- Turlington M, Noetzel MJ, Bridges TM, Vinson PN, Steckler T, Lavreysen H, Mackie C, Bartolome-Nebreda JM, Conde-Ceide S, Tong HM, Macdonald GJ, Daniels JS, Jones CK, Niswender CM, Conn PJ, Lindsley CW and Stauffer SR (2014) Discovery and SAR of a novel series of metabotropic glutamate receptor 5 positive allosteric modulators with high ligand efficiency. *Bioorg Med Chem Lett* **24**(15): 3641-3646.
- Valenti O, Conn PJ and Marino MJ (2002) Distinct physiological roles of the Gq-coupled metabotropic glutamate receptors Co-expressed in the same neuronal populations. *J Cell Physiol* **191**(2): 125-137.
- Valenti O, Mannaioni G, Seabrook GR, Conn PJ and Marino MJ (2005) Group III metabotropic glutamate-receptor-mediated modulation of excitatory transmission in rodent substantia nigra pars compacta dopamine neurons. *J Pharmacol Exp Ther* **313**(3): 1296-1304.
- Vieira E, Huwyler J, Jolidon S, Knoflach F, Mutel V and Wichmann J (2009) Fluorinated 9H-xanthene-9-carboxylic acid oxazol-2-yl-amides as potent, orally available mGlu1 receptor enhancers. *Bioorg Med Chem Lett* **19**(6): 1666-1669.
- Vilar B, Busserolles J, Ling B, Laffray S, Ulmann L, Malhaire F, Chapuy E, Aissouni Y, Etienne M, Bourinet E, Acher F, Pin JP, Eschalier A and Goudet C (2013) Alleviating pain hypersensitivity through activation of type 4 metabotropic glutamate receptor. *J Neurosci* **33**(48): 18951-18965.
- Woltering TJ, Wichmann J, Goetschi E, Knoflach F, Ballard TM, Huwyler J and Gatti S (2010) Synthesis and characterization of 1,3-dihydro-benzo[b][1,4]diazepin-2-one derivatives: Part 4. In vivo active potent and selective non-competitive metabotropic glutamate receptor 2/3 antagonists. *Bioorg Med Chem Lett* **20**(23): 6969-6974.



MOL #97220

- Wood MR, Hopkins CR, Brogan JT, Conn PJ and Lindsley CW (2011) "Molecular switches" on mGluR allosteric ligands that modulate modes of pharmacology. *Biochemistry* **50**(13): 2403-2410.
- Wu H, Wang C, Gregory KJ, Han GW, Cho HP, Xia Y, Niswender CM, Katritch V, Meiler J, Cherezov V, Conn PJ and Stevens RC (2014) Structure of a class C GPCR metabotropic glutamate receptor 1 bound to an allosteric modulator. *Science* **344**(6179): 58-64.
- Yin S, Noetzel MJ, Johnson KA, Zamorano R, Jalan-Sakrikar N, Gregory KJ, Conn PJ and Niswender CM (2014) Selective actions of novel allosteric modulators reveal functional heteromers of metabotropic glutamate receptors in the CNS. *J Neurosci* **34**(1): 79-94.
- Yin S, Zamorano R, Conn PJ and Niswender CM (2012) Functional selectivity induced by mGlu(4) receptor positive allosteric modulation and concomitant activation of G(q) coupled receptors. *Neuropharmacology*.
- Zerbib F, Keywood C and Strabach G (2010) Efficacy, tolerability and pharmacokinetics of a modified release formulation of ADX10059, a negative allosteric modulator of metabotropic glutamate receptor 5: an esophageal pH-impedance study in healthy subjects. *Neurogastroenterol Motil* **22**(8): 859-865, e231.
- Zhang Y, Rodriguez AL and Conn PJ (2005) Allosteric potentiators of metabotropic glutamate receptor subtype 5 have differential effects on different signaling pathways in cortical astrocytes. *J Pharmacol Exp Ther* **315**(3): 1212-1219.
- Zhou Y, Manka JT, Rodriguez AL, Weaver CD, Days EL, Vinson PN, Jadhav S, Hermann EJ, Jones CK, Conn PJ, Lindsley CW and Stauffer SR (2010) Discovery of N-Aryl Piperazines as Selective mGluR5 Potentiators with Improved In Vivo Utility. *ACS Med Chem Lett* **1**(8): 433-438.

MOL #97220

### Footnotes

a) Karen J. Gregory is a recipient of a National Health and Medical Research Council of Australia (NHMRC) Overseas Biomedical Postdoctoral Training Fellowship, work in her laboratory is supported by a NHMRC project grant [1084775]. Research on metabotropic glutamate receptors in the Conn laboratory is supported by the National Institutes of Health [Grants MH062646-13; NS031373-16A2; DA023947].

b) Reprint requests should be addressed to P. Jeffrey Conn, Department of Pharmacology & Vanderbilt Center for Neuroscience Drug Discovery, Vanderbilt University Medical Center, 1215 Light Hall, 2215-B Garland Ave, Nashville, TN, USA, 37232-0697; jeff.conn@vanderbilt.edu

MOL #97220

## Legends for Figures

### Figure 1: Schematic depicting the concepts of biased agonism and modulation.

**A)** The endogenous ligand (blue triangle) activates the receptor resulting in intracellular signal transduction. However, not all pathways will be activated to the same extent, this is the natural bias of the receptor/system. **B)** A surrogate orthosteric agonist (red triangle) may engender a different suite of active receptor conformations resulting in biased signalling when compared to the endogenous ligand, in this case the strength of signalling to response 1 versus 2 is reversed. **C)** The binding of an allosteric modulator (orange star) potentiates the response to the endogenous agonist for all signal pathways to the same extent. **D)** In contrast, a biased allosteric modulator (purple star) may have different magnitude or direction on cooperativity depending on the response measured, in this case, response 1 is potentiated while response 2 is inhibited.

### Figure 2: Comparison of allosteric pockets within the 7TM domains of mGlu<sub>1</sub> and mGlu<sub>5</sub>.

**A)** X-ray crystal structures of 7TMs of mGlu<sub>5</sub> (grey, PDB ID: 4OO9) and mGlu<sub>1</sub> (teal, PDB ID: 4OR2) were aligned using MacPyMOL (The PyMOL Molecular Graphics System, Version 1.3, Schrödinger, LLC). **B)** Inside surface rendering of the allosteric binding pockets of mavoglurant (black) in mGlu<sub>5</sub> (grey) and FITM (blue) in mGlu<sub>1</sub> (teal). **C)** Zoomed in image of the bottom of the allosteric pockets with mavoglurant shown. Two amino acids differ between mGlu<sub>1</sub> and mGlu<sub>5</sub> in this region, the differences in the side chains (depicted as sticks) create more space in mGlu<sub>5</sub>, allowing mavoglurant to bind deeper within the 7TM bundle. All images created using MacPyMOL.

### Figure 3: Affinity and cooperativity determinants for allosteric modulation of metabotropic glutamate receptors.

**A)** X-ray crystal structure of mGlu<sub>1</sub> 7TMs with FITM (rendered in blue sticks) bound (PDB ID: 4OR2) with known mGlu<sub>1</sub> allosteric ligand affinity determinants highlighted in dark blue. **B)** X-ray crystal structure of mGlu<sub>5</sub> 7TMs with mavoglurant (rendered in black sticks) bound (PDB ID: 4OO9), affinity determinants for mGlu<sub>5</sub> allosteric ligands are highlighted in red. **C)** X-ray crystal structure of mGlu<sub>5</sub> and mGlu<sub>1</sub> 7TM domains were aligned, only the mGlu<sub>5</sub> structure is represented in cartoon. The binding poses of FITM and mavoglurant show the overlapping allosteric pockets of mGlu<sub>1</sub> and mGlu<sub>5</sub>. Highlighted are residues that when mutated change allosteric ligand cooperativity: dark red: reduce mGlu<sub>5</sub> NAM cooperativity; light red: increase NAM cooperativity; orange: differential effects on NAM cooperativity; dark blue: increase PAM cooperativity; light blue: increase cooperativity of NAMs and PAMs; purple: differential effects on cooperativity of NAMs versus PAMs. **D)** Structures of mGlu<sub>1</sub> and mGlu<sub>5</sub> were aligned as per panel C, highlighted are residues that when mutated switch allosteric modulator cooperativity: blue: PAM to either NAM or neutral switch; red: NAM to PAM switch; purple: PAM to NAM and vice versa for NAMs. All images created using MacPyMOL.

### Figure 4: Relative contribution of 7TM residues to the binding affinity of prototypical mGlu<sub>5</sub> allosteric modulator, MPEP.

Known affinity determinants for MPEP are mapped onto the mGlu<sub>5</sub>-7TM x-ray crystal structure with mavoglurant bound (PDB ID: 4OO9). The relative contribution of pocket residues are colour-coded with side chains rendered as sticks based on mutational data as indicated in the key. All images created using MacPyMOL.

**Table 1: Molecular pharmacology of selected examples of allosteric ligands for group I metabotropic glutamate receptors.**

Ligand/selectivity		Allosteric pharmacology	Ref.
CPCCOEt	1	<b>NAM:</b> glutamate, DHPG, quisqualate and ACPD mediated inositol phosphate accumulation (IP <sub>1</sub> ); glutamate mediated intracellular Ca <sup>2+</sup> mobilization (iCa <sup>2+</sup> ); quisqualate induced changes in cell impedance (xCelligence); extracellular Ca <sup>2+</sup> induced changes in iCa <sup>2+</sup> . <b>NAL:</b> [ <sup>3</sup> H]glutamate binding.	1-4
JNJ-16259685	1>>5	<b>NAM:</b> glutamate mediated iCa <sup>2+</sup> and IP <sub>1</sub> ; quisqualate induced changes in cell impedance (xCelligence). <b>Inverse agonist:</b> IP <sub>1</sub> . <b>NAL:</b> [ <sup>3</sup> H]quisqualate binding.	5-6
R214127	1/5	<b>NAM:</b> glutamate mediated iCa <sup>2+</sup> . <b>Inverse agonist:</b> IP <sub>1</sub> .	7
Ro 67-7476	1	<b>PAM:</b> glutamate efficacy for mediating iCa <sup>2+</sup> and cAMP accumulation. <b>Agonist:</b> pERK1/2.	8-9
Ro0711401	1	<b>Agonist:</b> iCa <sup>2+</sup> . <b>PAM:</b> glutamate mediated K <sup>+</sup> currents via GIRK activation.	10
YM-298198	1	<b>NAM:</b> glutamate induced IP <sub>1</sub> ; quisqualate induced changes in cell impedance (xCelligence). <b>NAL:</b> orthosteric agonists and affinity of radiolabelled ligand.	3, 11
5MPEP	5	<b>NAL:</b> glutamate efficacy for iCa <sup>2+</sup> ; glutamate and DHPG efficacy for IP <sub>1</sub> ; quisqualate affinity.	12-17
5PAM523	5	<b>PAM:</b> glutamate efficacy for iCa <sup>2+</sup> . <b>NAL:</b> [ <sup>3</sup> H]quisqualate binding.	18
ADX47273	5	<b>PAM:</b> quisqualate affinity; quisqualate efficacy for IP <sub>1</sub> .	17
CDPPB	5	<b>PAM:</b> glutamate efficacy for iCa <sup>2+</sup> ; quisqualate affinity; quisqualate efficacy for IP <sub>1</sub> . <b>PAM-agonist:</b> glutamate efficacy for iCa <sup>2+</sup> , pERK1/2 and p-Akt.	13, 17-20
CPPHA	5>1 >4/8	<b>mGlu<sub>5</sub> PAM:</b> glutamate efficacy for iCa <sup>2+</sup> ; quisqualate efficacy for IP <sub>1</sub> and iCa <sup>2+</sup> . <b>mGlu<sub>5</sub> NAL:</b> quisqualate and glutamate affinity. <b>mGlu<sub>5</sub> PAM-agonist:</b> glutamate efficacy for pERK1/2. <b>NAM-agonist:</b> DHPG efficacy for pERK1/2 (cortical astrocytes). <b>mGlu<sub>4</sub> and mGlu<sub>8</sub> NAM:</b> glutamate-mediated Ca <sup>2+</sup> signalling.	14, 17, 19, 21, 22
DPFE	5	<b>PAM-agonist:</b> glutamate efficacy for iCa <sup>2+</sup> & pERK1/2.	23
M-5MPEP	5	<b>NAM (low cooperativity):</b> glutamate efficacy for iCa <sup>2+</sup> ; quisqualate & DHPG efficacy for IP <sub>1</sub> . <b>NAM:</b> glutamate, DHPG and quisqualate mediated Ca <sup>2+</sup> oscillations; glutamate efficacy for pERK1/2. <b>NAL:</b> quisqualate affinity.	12, 17, 19
MPEP	5>>4	<b>mGlu<sub>5</sub> NAM:</b> glutamate efficacy for iCa <sup>2+</sup> and pERK1/2; quisqualate and DHPG stimulated IP <sub>1</sub> ; glutamate, DHPG and quisqualate induced Ca <sup>2+</sup> oscillations. <b>mGlu<sub>5</sub> inverse agonist:</b> IP <sub>1</sub> . <b>mGlu<sub>5</sub> NAL:</b> quisqualate and glutamate affinity. <b>mGlu<sub>4</sub> PAM-agonist:</b> L-AP4 mediated iCa <sup>2+</sup> . <b>mGlu<sub>4</sub> PAM:</b> L-AP4 induced changes in VFT.	13, 17, 19, 24, 25
VU0357121	5	<b>NAL:</b> glutamate affinity. <b>PAM:</b> glutamate efficacy for iCa <sup>2+</sup> . <b>PAM-agonist:</b> glutamate efficacy for pERK1/2	19, 16
VU0360172	5	<b>PAM:</b> glutamate efficacy for iCa <sup>2+</sup> . <b>NAL:</b> glutamate affinity. <b>PAM-agonist:</b> glutamate efficacy for pERK1/2	19

References: 1, Annoura et al., 1996; 2, Litschig et al., 1999; 3, Scandroglio et al., 2010; 4, Jiang et al., 2014; 5, Lavreysen et al., 2004; 6, Scandroglio et al., 2010; 7, Lavreysen et al., 2003; 8, Hemstapat et al., 2006; 9, Sheffler et al., 2008; 10, Vieira et al., 2009; 11, Kohara et al., 2005; 12, Rodriguez et al., 2005; 13, Chen et al., 2007; 14, Chen et al., 2008; 15, Ayala et al., 2009; 16, Hammond et al., 2010; 17, Bradley et al., 2011; 18, Parmentier-Batteur et al., 2014; 19, Gregory et al., 2012; 20, Doria et al., 2013; 21, O'Brien et al., 2004; 22, Zhang et al., 2005; 23, Gregory et al., 2013b; 24, Gasparini et al., 1999; 24, Rovira et al., 2015.

**Table 2: Molecular pharmacology of selected allosteric ligands for group II metabotropic glutamate receptors.**

Ligand/selectivity		Allosteric pharmacology	Ref.
BINA	2	<b>PAM-agonist:</b> glutamate stimulated [ <sup>35</sup> S]GTPγS binding at human mGlu <sub>2</sub> . <b>PAM:</b> glutamate stimulated [ <sup>35</sup> S]GTPγS binding at rat mGlu <sub>2</sub> and iCa <sup>2+</sup> at human and rat mGlu <sub>2</sub> ; [ <sup>3</sup> H]DCG-IV binding to human mGlu <sub>2</sub> ; LY541850 mediated decreases in Ca <sup>2+</sup> oscillation frequency in primary neuronal cultures. <b>NAL:</b> [ <sup>3</sup> H]LY341495 binding to human mGlu <sub>2</sub> , LY379268 mediated decreases in Ca <sup>2+</sup> oscillation frequency in primary neuronal cultures.	1, 2, 3, 4
CBiPES		<b>PAM:</b> glutamate mediated iCa <sup>2+</sup> ; LY379268 and LY541850 mediated decreases in Ca <sup>2+</sup> oscillation frequency in primary neuronal cultures.	3, 5
JNJ-40068782	2	<b>PAM-agonist:</b> glutamate stimulated [ <sup>35</sup> S]GTPγS binding and iCa <sup>2+</sup> at human mGlu <sub>2</sub> . <b>PAM:</b> glutamate stimulated [ <sup>35</sup> S]GTPγS binding at rat mGlu <sub>2</sub> ; [ <sup>3</sup> H]DCG-IV binding to human mGlu <sub>2</sub> ; glutamate affinity (displacement of [ <sup>3</sup> H]DCG-IV). <b>NAL:</b> [ <sup>3</sup> H]LY341495 binding	1
LY2607540 (THIC)	2	<b>PAM-agonist:</b> glutamate stimulation of iCa <sup>2+</sup> and [ <sup>35</sup> S]GTPγS binding at human receptor. <b>PAM:</b> glutamate stimulation of iCa <sup>2+</sup> and [ <sup>35</sup> S]GTPγS binding at rat receptor	1, 6
LY487379	2	<b>PAM-agonist:</b> glutamate stimulation of [ <sup>35</sup> S]GTPγS binding to rat cerebral cortical membranes. <b>PAM:</b> glutamate, DCG-IV, LCCG-I and LY379268 stimulated [ <sup>35</sup> S]GTPγS binding and glutamate mediated iCa <sup>2+</sup> ; [ <sup>3</sup> H]DCG-IV binding to human mGlu <sub>2</sub> . <b>NAL:</b> [ <sup>3</sup> H]LY341495 binding to human mGlu <sub>2</sub> .	1, 2, 7, 8, 9
MNI-137	2/3	<b>mGlu<sub>2</sub> NAM:</b> glutamate and DCG-IV mediated iCa <sup>2+</sup> and glutamate stimulated [ <sup>35</sup> S]GTPγS binding. <b>mGlu<sub>3</sub> NAM:</b> glutamate mediated iCa <sup>2+</sup> and [ <sup>35</sup> S]GTPγS binding. <b>mGlu<sub>2</sub> NAL:</b> [ <sup>3</sup> H]LY341495 binding and glutamate affinity.	2, 10
RO4491533	2/3	<b>mGlu<sub>2</sub> NAM:</b> [ <sup>3</sup> H]LY354740 affinity; ACPD inhibition of cAMP accumulation; glutamate stimulation of GIRK channels and [ <sup>35</sup> S]GTPγS binding. <b>mGlu<sub>2</sub> NAL:</b> [ <sup>3</sup> H]LY341495 binding to human mGlu <sub>2</sub> . <b>mGlu<sub>3</sub> NAM:</b> glutamate stimulation of GIRK channels.	11, 12
RO4988546	2/3	<b>mGlu<sub>2</sub> NAM:</b> [ <sup>3</sup> H]LY354740 binding; LY354740 stimulation of [ <sup>35</sup> S]GTPγS binding and iCa <sup>2+</sup> . <b>mGlu<sub>2</sub> NAL:</b> [ <sup>3</sup> H]HYDIA binding.	13
RO5488608	2/3	<b>mGlu<sub>2</sub> NAM:</b> [ <sup>3</sup> H]LY354740 binding; LY354740 stimulation of [ <sup>35</sup> S]GTPγS binding and iCa <sup>2+</sup> . <b>mGlu<sub>2</sub> NAL:</b> [ <sup>3</sup> H]HYDIA binding.	13

References: 1, Lavreysen et al., 2013; 2, Hemstapat et al., 2007; 3, Sanger et al., 2013; 4, Galici et al., 2006; 5, Johnson et al., 2005; 6, Fell et al., 2011; 7, Odagaki et al., 2013; 8, Schaffhauser et al., 2003; 9, Odagaki et al., 2011; 10, Yin et al., 2014; 11, Woltering et al., 2010; 12, Campo et al., 2011; 13, Lundstrom et al., 2011.

**Table 3: Molecular pharmacology of selected allosteric ligands for group III metabotropic glutamate receptors.**

Ligand/selectivity		Allosteric pharmacology	Ref.
ADX71743	7	<b>NAM:</b> L-AP4 efficacy for $iCa^{2+}$ and inhibition of cAMP accumulation; glutamate in mediating inhibition of cAMP accumulation.	1
AMN082	7 <sup>#</sup>	<b>NAL-Agonist:</b> inhibition of cAMP accumulation, stimulates pERK1/2 and receptor internalization. <b>NAL:</b> MSOP (does not block AMN082 internalization).	2-5
AZ12216052	8	<b>PAM-agonist:</b> glutamate stimulation of [ <sup>35</sup> S]GTP $\gamma$ S binding	6
MMPIP	7	<b>NAM:</b> L-AP4 efficacy for $iCa^{2+}$ and changes in cell mass distribution; L-CCG-I and CPPG efficacy for $iCa^{2+}$ ; glutamate stimulation of GIRK channels. <b>Inverse/agonist:</b> cAMP accumulation and Epic label-free assay. <b>NAL:</b> [ <sup>3</sup> H]LY341495 binding	4, 7
PHCCC	4/1 6	<b>PAM:</b> glutamate for $iCa^{2+}$ and GIRK channel activation; L-AP4 stimulated $iCa^{2+}$ and changes in VFT. <b>PAM-agonist:</b> L-AP4 and glutamate stimulated [ <sup>35</sup> S]GTP $\gamma$ S binding. <b>mGlu<sub>1</sub> NAM:</b> glutamate mediated $iCa^{2+}$ . <b>mGlu<sub>6</sub> agonist:</b> glutamate mediated calcium current inhibition.	10-15
VU0155041	4	<b>PAM-agonist:</b> glutamate mediated $iCa^{2+}$ and GIRK channel activation. <b>PAM:</b> L-AP4 mediated $iCa^{2+}$ and changes in VFT.	13, 14, 16
VU0155094	4-8	<b>PAM:</b> glutamate apparent affinity (mGlu <sub>4,7,8</sub> ); glutamate efficacy for stimulation of GIRK channels (mGlu <sub>8</sub> ) and $iCa^{2+}$ (mGlu <sub>4,7</sub> ); L-AP4 apparent affinity (mGlu <sub>7,8</sub> ); L-AP4 efficacy for $iCa^{2+}$ (mGlu <sub>4</sub> ); LSP4-2022 apparent affinity (mGlu <sub>4,7,8</sub> ); LSP4-2022 efficacy for $iCa^{2+}$ (mGlu <sub>4</sub> ). <b>NAL/NAM:</b> L-AP4 apparent affinity (mGlu <sub>4</sub> ); L-AP4 and LSP4-2022 efficacy for $iCa^{2+}$ (mGlu <sub>7</sub> ). <b>NAL:</b> glutamate, L-AP4, LSP4-2022 efficacy for $iCa^{2+}$ (mGlu <sub>8</sub> ).	17
VU0422288	4-8	<b>PAM:</b> glutamate, L-AP4 and LSP4-2022 efficacy for $iCa^{2+}$ and GIRK channel activation (mGlu <sub>4,7,8</sub> ); glutamate, L-AP4, LSP4-2022 apparent affinity (mGlu <sub>7,8</sub> ). <b>NAL:</b> glutamate apparent affinity (mGlu <sub>4</sub> ). <b>NAM:</b> L-AP4 and LSP4-2022 apparent affinity (mGlu <sub>4</sub> ).	17

<sup>#</sup> compound and/or its metabolite hits other non-mGlu targets (Ayala et al., 2008; Sukoff Rizzo et al., 2011).

References: 1, Kalinichev et al., 2013; 2, Mitsukawa et al., 2005; 3, Pelkey et al., 2007; 4, Suzuki et al., 2007; 5, Iacovelli et al., 2014; 6, Duvoisin et al., 2010; 7, Niswender et al., 2010b; 10, Beqollari and Kammermeier, 2008; 11, Maj et al., 2003; 12, Marino et al., 2003; 13, Niswender et al., 2008; 14, Rovira et al., 2015; 15, Yin et al., 2014; 16, Yin et al., 2012; 17, Jalan-Sakrikar et al., 2014.

**Table 4: Known affinity determinants for mGlu allosteric modulators interacting with the common allosteric site.**

Ligand/subtype	Amino acid residue position																		
	3.29	3.36	3.39	3.40	5.43	5.44	5.47	5.48	6.44	6.48	6.51	6.52	6.55	6.56	7.38	7.45	7.46		
EM-TBPC	1			Y		V/L				W+	F	--	Y		T				
FITM	1	s			P				--						T	--			
MPEP	5	R	P	S	Y	P	L	--	G	T	W	F	V	Y	f+	--	S	A	
fenobam	5	R	P	S	Y					T	W	F		Y				A	
VU0366248	5		P	S	Y			N	--	t	W	f <sup>#</sup>	--	Y	--		S	S+	A
VU0285683	5		p <sup>#</sup>		Y	--		N	--	T	W	F	--	Y			S	A	
VU0366058	5		p <sup>#</sup>		Y			--	g+		W	F+	v+	Y+			s	A	
VU0409106	5		p <sup>#</sup>		Y			N	--	T	W	F	--	Y	f		S	A	
DPFE	5	--	P		Y	P	L+		--	T	W	--	V+	y+	f+		s	A	
VU29	5	--	p <sup>#</sup>		--	p	l		G	T	w+	f	v	y	f+		-	A	
VU0403602	5	--	P	s	Y <sup>#</sup>	p	--	--		T	W		--				S	A	

Most well-characterised representatives from individual allosteric modulator scaffolds were selected for inclusion. Alignment of amino acids and assignment of Ballesteros-Weinstein numbering was adopted from the structural alignment to class A GPCRs in Wu et al., 2014. A colorised version of this table (Supplementary Table 1) is included in the online supplementary materials.

lower case: mutations caused a change (2-10 fold) in affinity estimates that did not reach significance (determined from either radioligand binding or from functional interactions with orthosteric agonists).

upper case: mutations caused a significant change in affinity (determined from either radioligand binding or from functional interactions with orthosteric agonists; not included: T6.43 for MPEP, C7.52 for VU0403602)

<sup>#</sup> mutation caused a loss in allosteric modulation of orthosteric agonist functional response; impact on affinity inferred from effects across other scaffolds.

-- mutation/s had no effect on modulator affinity (determined from either radioligand binding or from functional interactions with orthosteric agonists).

+ mutation increases modulator affinity, in all other cases mutations cause a decrease (determined from either radioligand binding or from functional interactions with orthosteric agonists).

Grey scale: white: 3-10 fold change; light grey: 10-30 fold change; mid grey: 30-100 fold decrease; dark grey: >100 fold decrease or complete loss of radiolabelled allosteric modulator binding (Data summarised from: Gregory et al., 2012; Gregory et al., 2013b; Gregory et al., 2014; Malherbe et al., 2003a; Malherbe et al., 2003b; Malherbe et al., 2006; Pagano et al., 2000; Turlington et al., 2013; Wu et al., 2014).




Figure 1






Figure 2





Figure 3



- Mutations reduce affinity 3-10 fold
- Mutations reduce affinity 10-30 fold
- Mutations reduce affinity 30-100 fold
- Mutations reduce affinity >100 fold