

## **The phosphodiesterase inhibitor IBMX blocks the potassium channel THIK-1 from the extracellular side**

**Xinle Zou<sup>1</sup>, Linus J. Conrad<sup>1</sup>, Kristin Koschinsky, Günter Schlichthörl, Regina Preisig-Müller, Eugen Netz, Jens Krüger, Jürgen Daut<sup>2</sup> and Vijay Renigunta<sup>2</sup>**

<sup>1</sup>*Contributed equally*

<sup>2</sup>*Corresponding authors*

*Institute of Physiology and Pathophysiology, Marburg University,  
35037 Marburg, Germany (X.Z., L.J.C., K.K, G.S., R.P.M., J.D., V.R.)*

*Biomolecular Interactions, Max Planck Institute for Developmental Biology,  
Tübingen, Germany (E.N.)*

*High Performance and Cloud Computing Group, IT Center, University of  
Tübingen, Germany (J.K.)*

## Running title: IBMX blocks THIK-1

*Please send editorial correspondence to:*

Prof. Jürgen Daut  
Institute of Physiology and Pathophysiology  
Marburg University  
Deutschhausstr. 2  
35037 Marburg, Germany

Tel.: 0049-6421-2866494

Fax: 0049-6421-2866495

Email: [jdaut@staff.uni-marburg.de](mailto:jdaut@staff.uni-marburg.de).

Text pages:	16
Tables:	0
Figures:	9
Abstract:	215 words
Introduction:	618 words
Discussion:	1105 words

### Non-standard abbreviations

CHO cells:	Chinese hamster ovary cells
IBMX:	3-isobutyl-1-methyl-xanthine
THIK-1:	Two-pore-domain halothane-inhibited K <sup>+</sup> channel 1
THIK-2:	Two-pore-domain halothane-inhibited K <sup>+</sup> channel 2
TRAAK:	TWIK-related arachidonic acid stimulated K <sup>+</sup> channel
TREK-1:	TWIK-related K <sup>+</sup> channel
TWIK:	tandem of P domains in a weakly inwardly rectifying K <sup>+</sup> channel

## Abstract

The two-pore-domain potassium channel ( $K_{2P}$  channel) THIK-1 has several predicted protein kinase A (PKA) phosphorylation sites. In trying to elucidate whether THIK-1 is regulated via PKA we expressed THIK-1 channels in a mammalian cell line (CHO cells) and used the phosphodiesterase inhibitor 3-isobutyl-1-methyl-xanthine (IBMX) as a pharmacological tool to induce activation of PKA. Using the whole-cell patch-clamp recording we found that THIK-1 currents were inhibited by application of IBMX with an  $IC_{50}$  of 120  $\mu$ M. Surprisingly, intracellular application of IBMX or of the second messenger cAMP via the patch pipette had no effect on THIK-1 currents. In contrast, extracellular application of IBMX produced a rapid and reversible inhibition of THIK-1. In patch-clamp experiments with outside-out patches, THIK-1 currents were also inhibited by extracellular application of IBMX. Expression of THIK-1 channels in *Xenopus* oocytes was used to compare wild-type channels with mutated channels. Mutation of the putative PKA phosphorylation sites did not change the inhibitory effect of IBMX on THIK-1 currents. Mutational analysis of all residues of the (extracellular) helical cap of THIK-1 showed that mutation of the arginine residue at position 92, which is in the C2-P1 linker, markedly reduced the inhibitory effect of IBMX. This flexible linker region, which is unique for each  $K_{2P}$  channel subtype, may be a possible target of channel-specific blockers.

## Significance statement

The potassium channel THIK-1 is strongly expressed in the central nervous system. We studied the effect of 3-isobutyl-1-methyl-xanthine (IBMX) on THIK-1 currents. IBMX inhibits breakdown of cAMP and thus activates protein kinase A (PKA). Surprisingly, THIK-1 current was inhibited when IBMX was applied from the extracellular side of the membrane, but not from the intracellular side. Our results suggest that IBMX binds directly to the channel and that the inhibition of THIK-1 current was not related to activation of PKA.

## Introduction

There are about 80 potassium channel subunits in the human genome (Yu and Catterall, 2004). These channels regulate the function of many cell types by three basic mechanisms: (i)  $K^+$  channels are responsible for the repolarisation process of action potentials. (ii)  $K^+$  channels modulate the excitability by setting the resting membrane potential. (iii) By modulating the membrane potential,  $K^+$  channels indirectly control the influx of calcium ions through voltage-activated  $Ca^{2+}$  channels and TRP (transient receptor potential) channels, which influences many cellular processes. So far, it has been difficult to develop blockers that specifically block only one  $K^+$  channel or only a particular subfamily of  $K^+$  channels. This can be explained by the fact that most channel blockers target the central cavity of the channel, which is accessible from the inside of the cell. Because the dimensions and the inner walls of the channels are quite well conserved most drugs inhibit several different potassium channels to some extent.

The family of two-pore-domain  $K^+$  channels ( $K_{2P}$ -channels) is unique among potassium channels in that their transmembrane pore is a dimer consisting of two identical or different subunits.  $K_{2P}$ -channels also possess a unique extracellular 'helical cap' structure which allows the efflux of potassium ions into the extracellular space through two lateral openings (Brohawn et al., 2012; Miller and Long, 2012). This structural peculiarity may allow the development of drugs that specifically target  $K_{2P}$ -channels. Since the helical caps of different  $K_{2P}$ -channels show only moderate homology, it may even be possible to develop tailored drugs that are specific for one particular  $K_{2P}$ -channel.

In humans, the family of  $K_{2P}$ -channels has fifteen members (Enyedi and Czirjak, 2010). Most of the  $K_{2P}$ -channels are characterized by a very complex regulation of open probability (Feliciangeli et al., 2014; Renigunta et al., 2015), and the function of various  $K_{2P}$ -channel subtypes in the brain, the heart and in endocrine glands has been characterised in some detail (Bayliss and Barrett, 2008; Bichet et al., 2014; Duprat et al., 2007; Lesage and Barhanin, 2011). THIK-1 and THIK-2 (Tandem-pore-domain Halothane Inhibited  $K^+$  channels) represent a  $K_{2P}$ -channel subfamily that has received relatively little attention so far (Chatelain et al., 2013; Rajan et al., 2001; Renigunta et al., 2014). A distinguishing feature of these channels is their inhibition by the inhalational anaesthetics halothane and isoflurane (Chatelain et al., 2013; Kang et al., 2014; Lazarenko et al., 2010; Rajan et al., 2001; Renigunta et al., 2014). RT-PCR analysis showed that THIK-1 is expressed ubiquitously both in the adult and during embryogenesis (Aller and Wisden, 2008; Rajan et al., 2001). In situ hybridization experiments showed that THIK-1 is robustly expressed in rat brain, especially in some thalamic nuclei, the dentate gyrus, the ventromedial hypothalamic nucleus and the

olfactory bulb (Girard et al., 2001; Rajan et al., 2001). The function of THIK-1 and THIK-2 channels in the brain is still unclear. Interestingly, THIK-1 channels have recently been shown to be strongly expressed in microglial cells (Butovsky et al., 2014; Zhang et al., 2014) and to play an essential role in controlling the morphology and the immunological function of these cells (Madry et al., 2018).

The present study was initiated by a serendipitous finding. When studying the signal transduction between G-protein coupled receptors (GPCRs) and THIK-1 channels in heterologous expression systems we found that application of the phosphodiesterase inhibitor 3-isobutyl-1-methyl-xanthine (IBMX) rapidly and reversibly inhibited THIK-1 currents. Investigation of the underlying mechanism showed that THIK-1 channels were insensitive to intracellular application of IBMX and that mutation of a residue of the extracellular cap (R92) reduced the sensitivity to extracellular application of IBMX. Taken together, our results suggest that the effect of IBMX was not caused by inhibition of an enzyme but by direct binding to the extracellular cap of THIK-1.

## Methods

### Ethical approval

For experiments involving *Xenopus* oocytes, adult female African clawed frogs (*Xenopus laevis*) were used. The frogs were anaesthetised by putting them in water containing 1g/l tricaine. Stage-V oocytes were obtained from ovarian lobes. Anaesthesia and operation were carried out in accordance with the principles of German legislation with approval of the animal welfare officer of the Medical Faculty of Marburg University under the governance of the Regierungspräsidium Giessen (the regional veterinary health authority).

### Molecular cloning and mutagenesis

Full length rat THIK-1 (accession no. NM\_022293), human TREK-1c (accession no. AY552980) and human TRAAK (accession no. AF247042) channels were cloned into the pSGEM oocyte expression vector and into the mammalian expression vector pCDNA3.1 (Rajan et al., 2001; Renigunta et al., 2014; Rinné et al., 2014). The QuikChange site-directed mutagenesis kit (Agilent Technologies) was used to introduce point mutations. All constructs were verified by sequencing.

### Patch-clamp measurements of THIK currents in transfected CHO cells

CHO cells were cultured in MEM alpha with glutamine (Thermo Fisher), 10% FCS and 1% penicillin/streptomycin (PAA). The cells were seeded in 35 mm glass-bottom dishes (ibidi, Martinsried, Germany) and transfected with either rat THIK-1, human TREK-1c, hTRAAK or hTREK-1c<sup>S344A</sup> cloned into the pCDNA3.1 mammalian expression vector (Invitrogen) using jetPRIME reagent (Polyplus, Illkirch, France), according to the manufacturers' protocols. Patch-clamp measurements were performed 24 h after transfection. For whole-cell recording, the cells were superfused with a bath solution containing (mM): 150 NaCl, 5 KCl, 1.6 CaCl<sub>2</sub>·6 H<sub>2</sub>O, 1.13 MgCl<sub>2</sub>·6 H<sub>2</sub>O; 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 10 glucose, 2 Na pyruvate and 10 HEPES; the pH was adjusted to 7.4 with NaOH. The patch pipettes (resistance, 3-6 MΩ) were filled with an 'intracellular' solution containing (mM): 60 KCl, 65 K-glutamate, 5 EGTA, 3.5 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 3 K<sub>2</sub>ATP, 0.2 Na<sub>2</sub>GTP, 5 HEPES; the pH was adjusted to 7.2 with KOH. Steady-state current-voltage relations were obtained by applying slow voltage ramps (40 mV s<sup>-1</sup>) between -90 and +30 mV. The liquid junction potential between the patch electrode and the bath solution (~ -8 mV) was not compensated.

For cell-attached or outside-out recording the patch pipettes (resistance, 3 to 6 MΩ) were coated with the elastomer R-6101 (Dow Corning) to reduce capacitive noise; the pipettes were filled with an 'intracellular' solution containing (mM): 140 KCl, 5 HEPES and 1 MgCl<sub>2</sub>·6H<sub>2</sub>O, and the cells were superfused with a solution containing (mM): 150 NaCl, 4.5 KCl, 1.6 CaCl<sub>2</sub>·6 H<sub>2</sub>O, 1.13 MgCl<sub>2</sub>·6 H<sub>2</sub>O; 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 10 glucose, 2 Na pyruvate and 10 HEPES; the pH was adjusted to 7.4 with NaOH. The liquid junction potential between the patch electrode and the bath solution (~ -8 mV) was not compensated. The sampling rate was 10 kHz; the filter had a cut-off frequency of 2 kHz. All electrophysiological experiments were carried out at room temperature (20-23°C).

### **Voltage-clamp measurements with *Xenopus* oocytes**

For *Xenopus* oocyte expression studies, complementary RNA (cRNA) coding for THIK-1, TRAAK, TREK-1c or TREK-1c<sup>S344A</sup> was transcribed *in vitro* using T7 RNA polymerase (mMessage mMachine T7 Kit, Ambion). cRNA quality was determined by gel electrophoresis and UV spectroscopy. Defolliculated *Xenopus* oocytes (stage V) were injected with nuclease-free water containing cRNA of THIK-1 (6 ng/oocyte), TRAAK (2.5 ng/oocyte) or TREK-1c (2.5 ng/oocyte). The oocytes were incubated at 19°C for 24-48 h in ND96 solution containing (mM) 96 NaCl, 2 KCl, 1 MgCl<sub>2</sub>, 1.8 CaCl<sub>2</sub> and 5 HEPES (pH 7.5), supplemented with 100 µg/ml gentamycin, 2.5 mM sodium pyruvate. For voltage-clamp recording the oocytes were placed in a small-volume perfusion chamber and superfused with ND96 solution. Two-microelectrode voltage-clamp measurements with ramp-shaped voltage commands were performed with a TurboTec-10 C amplifier (npi electronic, Tamm, Germany). Data acquisition and analysis were performed with a software developed in our laboratory (PC.DAQ1.2); data were recorded at a sampling rate of 120 Hz. For quantification of current amplitude under different experimental conditions the currents were measured at 0 mV (2 mM K<sup>+</sup>). All electrophysiological experiments were carried out at room temperature (20-23 °C).

### **Homology modelling of THIK-1 channels**

A homology model of rat THIK-1 was constructed based on the structure of human TRAAK channels (Brohawn et al., 2013; Brohawn et al., 2012) using the Schrödinger Maestro 10.1 software package (Release 2015-1), which includes, among others, the tools 'Small Molecule Drug Discovery Suite', 'Biologics Suite', 'Multiple Sequence Viewer', 'Prime 3.9' and 'Structure Prediction wizard'. rTHIK-1, hTHIK-1 and hTRAAK

were aligned using the 'Multiple Sequence Viewer'. In the original study of Brohawn et al. (Brohawn et al., 2012), the structure of hTRAACK could be resolved from position 28 to 286 in chain A and from position 28 to 284 in chain B, except for a flexible region unresolved for position 104 to 109 for chain A and 105 to 109 for chain B, which corresponds to the C2-P1 linker (shown in red in Figure 1A). The structure coordinates of hTRAACK and the part of the alignment of rTHIK-1 that covers the amino acids represented in the structure were fed into 'Structure Prediction Wizard' and the final homology model was constructed using the energy-based building method of Prime 3.9.

## Statistics

Data are reported as means  $\pm$  S.D. Statistical significance was determined using Student's two-tailed t-test. Usually, the individual measurements are also shown as scatter dot plots. In the Figures, statistically significant differences to control values are marked by asterisks (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ ).

## Results

### THIK-1 currents are inhibited by IBMX

The rat ortholog of the  $K_{2P}$ -channel THIK-1 has four plausible protein kinase A (PKA) phosphorylation sites, as predicted by linear motif analysis (NetPhosK1.0, scansite3, PhosphoELM): S7, T170, S343 and S356 (Fig. 1A). To test whether the open probability of THIK-1 channels may be modulated by PKA we studied the effect of the phosphodiesterase inhibitor 3-isobutyl-1-methyl-xanthine (IBMX) on THIK-1 current. Application of IBMX is expected to cause a rise in intracellular cAMP and a subsequent activation of endogenous protein kinase A.

THIK-1 channels were expressed in a mammalian cell line (CHO cells) and whole-cell currents were measured using the patch-clamp technique. The cells were superfused with a physiological salt solution containing 5 mM extracellular potassium and current-voltage relations were obtained using slow voltage ramps (40 mV/s). Application of 1 mM IBMX caused a block of inward and outward currents with a reversal potential at about -80 mV, consistent with a potassium-selective channel (Fig. 1B). The steady-state dose-response curve for the effect of IBMX on THIK-1 currents could be fitted with a Hill coefficient of 1 and an  $IC_{50}$  of 120  $\mu$ M (Fig. 1C).

The time course of the effect of IBMX on THIK-1 currents was determined by measuring the outward current at 0 mV at intervals of 1 s. The inhibition of outward current elicited by application of 1 mM IBMX reached a steady state within 15 s; it was 90 % complete within 10 s (Fig. 1D; red curve). The effect of IBMX was readily reversible, the increase in current during wash-out reached a steady-state within 40 s (Fig. 1D). To assess the time course of the solution change in the recording chamber we changed the extracellular potassium concentration from 5 to 24 mM and monitored the resulting change in the inward current at -80 mV (Fig. 1D; black curve). We found that the solution change was 90 % complete within 12 s. These results suggest that the time constant ( $\tau_{on}$ ) of the binding of 1 mM IBMX to its target was too fast to be measured adequately in our recording chamber, whereas the dissociation of IBMX was somewhat slower than the solution change; the dissociation time constant of the drug ( $\tau_{off}$ ) is estimated to be 7 s or less. We conclude from these findings that the onset and wash-out of effects of IBMX were faster than would be expected for an intracellular site of action of the drug.

We then tried to inhibit the effect of IBMX on THIK-1 currents by pre-incubating the CHO cells for 10 min with the PKA inhibitor H89. We found that application of 5  $\mu$ M H89 by itself had no effect on THIK-1 current (Fig. 2A). After pre-incubation of the cells with H89 for at least 10 min (Figure 2B), the effect of IBMX was the same as without

preincubation (Fig. 2C). Stimulation of adenylate cyclase by application of 20  $\mu$ M forskolin (which is expected to lead to activation of PKA) also had no effect on THIK-1 current (Fig. 2D,F). As a control for the action forskolin under our experimental conditions, we repeated these experiments with CHO cells transfected with the  $K_{2P}$ -channel TREK-1c (Rinné et al., 2014). Application of 20  $\mu$ M forskolin caused a complete and reversible inhibition of TREK-1c (Fig. 2E,F), consistent with previous results (Fink et al., 1996; Terrenoire et al., 2001). In contrast, a mutant of TREK-1c in which the PKA phosphorylation site was removed (TREK-1c<sup>S344A</sup>) was insensitive to the action of forskolin.

### **IBMX blocks THIK-1 channels only from the extracellular side**

The results presented above suggest that TREK-1, but not THIK-1 channels can be inhibited via activation of PKA. To confirm this, we tried to apply cAMP directly to the cytosol of CHO cells via the pipette solution. Whole-cell recordings were carried out in CHO cells transfected with THIK-1. To induce activation of PKA, 100  $\mu$ M cAMP was included in the pipette solution. Currents were monitored by continuous voltage clamp at 0 mV. After formation of the Gigaseal the patch membrane was ruptured by applying negative pressure. Immediately after rupture of the patch, an outward current was observed which remained nearly constant as long as the cell was viable (Fig. 3A,B), despite the fact that cAMP was allowed to diffuse into the cytosol from the patch pipette. When we repeated this experiment with CHO cells transfected with TREK-1c, a rapid decay of the outward current (recorded at 0 mV) was observed (Fig. 3C). A mutant of TREK-1c in which the PKA phosphorylation site at position 344 was mutated (TREK-1c<sup>S344A</sup>) was unaffected by inclusion of 100  $\mu$ M cAMP in the pipette solution (Fig. 3D). These experiments show that endogenous PKA in CHO cells could be activated within less than 20 s by dialysis of the cytoplasm with cAMP. Wild-type TREK-1 channels (but not the S344A mutant) could be inhibited by dialysis of the cells with cAMP. In contrast, THIK-1 channels were not affected by activation of PKA via cAMP. Thus, the effect of extracellular application of IBMX on THIK-1 currents shown in Figures 1 and 2 must be mediated by other mechanisms.

To further characterise the effect of IBMX on THIK-1 channels we applied the drug via the pipette solution. CHO cells were transfected with THIK-1, a Gigaseal was formed and after rupture of the patch the cells were dialysed with a pipette solution containing 1 mM IBMX (Fig. 4A). The outward current at 0 mV was constant for several minutes. We then superfused the same cell with a bath solution containing 1 mM IBMX (Fig. 4A), and we found that the outward current was inhibited in the same way as shown in

Figs. 1D and 2B. When we repeated the IBMX dialysis experiments with CHO cells transfected with TREK-1c, the results were quite different: the TREK-1 current was inhibited completely within 10 s after rupture of the patch (Fig. 4B). This effect was most likely due to diffusion of IBMX into the cell, inhibition of phosphodiesterases and subsequent activation of PKA. The fractional residual current ( $I_{\text{IBMX}}/I_0$ ) measured after dialysis of the cells with 1 mM IBMX for 20 s is shown in Figure 4C. It can be seen that dialysis of IBMX caused a complete block of outward current in cells transfected with TREK-1 but had no effect in cells transfected with THIK-1. Taken together, these data suggest that intracellular application of IBMX has no effect on THIK-1 channels, whereas extracellular application of IBMX causes rapid inhibition of THIK-1 channels. Our results also imply that the permeability of the cell membrane to IBMX was not large enough to cause binding of intracellularly applied IBMX to an extracellular binding site; in other words, any IBMX appearing outside the cell was rapidly diluted by diffusion and by superfusion of the cell.

As a positive control of our experimental approach, we dialysed the cells the  $K^+$  channel blocker tetrapentylammonium (TPA). We found that THIK-1 channels could be inhibited by intracellular dialysis with 20  $\mu\text{M}$  TPA, as illustrated in Fig. 4D. TREK-1c channels could also be inhibited by intracellular application of 20  $\mu\text{M}$  TPA (Fig. 4E), in agreement with previous results (Piechotta et al., 2011).

### **THIK-1 channels in cell-attached and outside-out patches**

The results presented above suggest that IBMX blocks THIK-1 channels from the extracellular side but not from the intracellular side of the membrane. To obtain further information on the localization of the binding site for IBMX (intracellular or extracellular) we performed cell-attached and outside-out patch-clamp measurements in CHO cells transfected with THIK-1 channels. As a first step we characterised the properties of THIK-1 channels in cell-attached patches (Fig. 5A). The cells were superfused with physiological salt solution containing 5 mM  $K^+$ . The (extracellular) pipette solution contained 145 mM  $K^+$ . The membrane potential of the cell measured after subsequent rupture of the patch had a value of -60 mV. The current noise in the cell-attached patch was minimal at pipette potentials of about -80 mV (in the range -60 to -100 mV), which is consistent with the estimated  $K^+$  equilibrium potential across the patch membrane. We then tried to inhibit the noise attributable to  $K^+$  channels by applying tetrapentylammonium, which blocks THIK-1 channels from the inside (see Figure 4). Figure 5B shows that the noise at -80 to +40 mV was markedly reduced after

application of 300  $\mu$ M TPA in the bath solution. The concentration of TPA in the cytosol was probably much lower than 300  $\mu$ M.

These results suggest that we can indeed observe an increased noise level related to THIK-1 channel activity in cell-attached patch-clamp recordings. However, the single-channel amplitude could not be resolved, despite the fact that the noise level of our patch-clamp set-up was quite low (< 200 fA RMS noise with a sampling rate of 10 kHz and a low-pass filter with a cut-off frequency of 2 kHz). The difficulty of recording THIK-1 single-channel currents is probably related to the fact that (i) the channels have low conductance, (ii) the gating kinetics of the channels is extremely fast, and (iii) there were several channels in a single patch. Similar characteristics of THIK-1 channels have been reported previously (Kang et al., 2014).

Next we carried out measurements of THIK-1 channels in outside-out patches. The aim was to demonstrate directly the effect of extracellular IBMX in the absence of any kinase activity and in the absence of any intracellular second messengers. For these experiments we also used an extracellular (bath) solution containing 5 mM  $K^+$ ; the intracellular (pipette) solution contained 145 mM  $K^+$ . We found that in outside-out patches application of 1 mM IBMX caused a rapid reduction in outward current and in current noise (Figure 5C). This finding suggests that the patch contained several THIK-1 channels with a relatively high open probability. Like in the whole-cells recordings with a rapid extracellular solution change (Figure 1D), the binding of IBMX to the channels appeared to be faster than the dissociation. The average current-voltage relations of the outside-out patches before and after application of IBMX (Fig. 5D) crossed at about -75 mV (which corresponds to about -83 mV if the liquid junction potential of -8 mV is taken into account). This reversal potential is consistent with the calculated potassium equilibrium potential. Thus, our results obtained with cell free membrane patches confirm the idea that IBMX blocks THIK-1 by binding to an extracellular domain of the channel.

### **The block of THIK-1 currents by IBMX is not modified by mutation of putative PKA phosphorylation sites**

As a final test for the mechanism underlying the effects of IBMX we turned to the predicted PKA phosphorylation sites (S7, T170, S343 and S356; see Figure 1A). If mutation of one of these sites diminished the effects of IBMX on the channels this would suggest that the inhibitory action of the drug on THIK-1 was - at least to some extent - mediated by phosphorylation of the channel protein. THIK-1 channels were expressed in *Xenopus* oocytes and THIK-1 currents were studied with the two-

microelectrode voltage-clamp technique. When the oocytes were superfused with a bath solution containing 2 mM K<sup>+</sup> (ND96 solution, see Methods) the current-voltage relation of THIK-1 was outwardly rectifying (Fig. 6A), as described previously (Rajan et al., 2001). The inward and outward currents carried by THIK-1 channels were inhibited by application of 1 mM IBMX (Fig. 6A,B). The current change produced by IBMX reversed its polarity at approximately -100 mV, as expected for block of a K<sup>+</sup>-selective channel. We then mutated one by one all the predicted PKA phosphorylation sites: their serine or threonine residues were changed to alanine. As can be seen in 6B, the inhibition of THIK-1 by IBMX was unchanged after mutation of any of these putative phosphorylation sites. These findings suggest that the effect of IBMX on THIK-1 current was not mediated by phosphorylation of the channel protein. There are two possible explanations for the result illustrated in Figure 6B: (i) the rise in intracellular cAMP elicited by the phosphodiesterase inhibitor IBMX and the subsequent activation of protein kinase A did not cause phosphorylation of THIK-1 channels, or (ii) phosphorylation of THIK-1 via PKA did not produce any measurable change the open probability of the channels.

### **Identification of the IBMX binding site by mutagenesis of the helical cap of THIK-1**

To identify the binding site of IBMX on THIK-1 we mutated one-by-one all amino acids of the helical cap and compared the effect of IBMX on wild-type THIK-1 with that on the mutants (Figure 7A). Alanine residues were mutated to serine, all other residues were mutated to alanine. For each mutant we calculated the fractional residual THIK-1 current during application of 500  $\mu$ M IBMX ( $I_{\text{IBMX}}/I_0$ ), i.e. the ratio of the residual current measured during application of IBMX in the steady state ( $I_{\text{IBMX}}$ ) to the current measured before application of the drug ( $I_0$ ). Overall, the effect of 500  $\mu$ M IBMX on THIK-1 current was tested in  $n = 422$  oocytes ( $n = 5$  to 21 for different mutants). During application of IBMX, most mutants showed a fractional residual current ( $I_{\text{IBMX}}/I_0$ ) of ~40 %, similar to wild-type THIK-1 ( $n = 26$ ; black bar in Figure 7A), whereas one mutant (R92A) was clearly less sensitive to inhibition by IBMX (red bar in Fig. 7A).

Figure 7A also shows that there was some scatter in the residual THIK-1 current measured during application of IBMX with different mutants. This may have been partially due to the fact that different batches of oocytes were used for different mutants; different batches showed different absolute current magnitudes. To check for the possibility of a systematic error due to variable expression levels in different oocyte batches, we plotted the absolute amplitude of the THIK-1 current against the fractional

residual current (Figure 7B). The correlation coefficient (calculated including the mutant R92A) was -0.10, which suggests that there was practically no linear correlation between absolute THIK-1 current and the sensitivity of the channel to inhibition by IBMX. The grey lines in Figure 7B indicate a distance of three standard deviations from the mean fractional residual current observed in the presence of IBMX. It can be seen that only one mutant, R92A, was outside this range (red circle in Figure 7B). This makes it very unlikely that this result was obtained by chance, and it raises the possibility that this residue was somehow involved in the action of IBMX on THIK-1.

To further confirm this hypothesis, we mutated the residue R92 also to glutamate and to glutamine. We found that the fractional residual current of THIK-1 current observed after application of 1 mM IBMX was even larger in the R92E and R92Q mutants than in the mutant R92A (Figure 7C).

The two simplest explanations for the effect of the three R92 mutations on the sensitivity of THIK-1 channels to IBMX are (i) residue R92 is involved in the binding of IBMX to the extracellular cap of the channel, or that (ii) residue R92 is somehow involved in the gating of the channel. The latter effect would be expected to give rise to a change in the shape or position of the current-voltage relation. In Figure 7D the current-voltage relations of wild-type THIK-1 and the mutants R92A, R92E and R92Q are superimposed. No change in the current voltage relation could be detected, which makes it less likely that residue R92 is involved in the gating of the channel (see Discussion).

Figure 7E shows an alignment of the helical cap region of the  $K_{2P}$  channels TRAAK, TWIK-1 and TREK-2 (which have all been crystallized), together with rat THIK-1, the highly homologous rat THIK-2, and human THIK-1. It can be seen that the cap helices 1 and 2 are highly conserved in all of these channels, except for the linker region (indicated by a red dashed line) between cap helix 2 (C2) and the pore helix 1 (P1). Part of the linker region was found to be disordered (Brohawn et al., 2012; Miller and Long, 2012), it was not resolved in the crystal structure; the residues missing in the crystal structure are highlighted in red. It can be seen that the arginine residue at position 92 is at the end of the unstructured region, three residues proximal to the pore helix.

We also tested the effect of IBMX on the  $K_{2P}$ -channel TRAAK (long AT1 variant), which has a different C2-P1 linker (see Fig. 7E). We found that the current produced by TRAAK channels expressed in *Xenopus* oocytes was completely insensitive to extracellular application of 1 mM IBMX ( $n = 8$  oocytes; not illustrated).

Taken together, our mutagenesis experiments suggest that the 'disordered' C2-P1 linker region (Brohawn et al., 2014; Brohawn et al., 2012; Dong et al., 2015; Miller and Long, 2012) may play a role in the block of THIK-1 by IBMX. The mutations R92A, R92E and R92Q were the only ones in the entire helical cap that markedly attenuated the effect of IBMX on THIK-1 channels. This observation is consistent with the idea that residue R92 – located near the end of the C2-P1 linker – may form part of the extracellular binding site of IBMX. To visualize the localisation of R92 we constructed a homology model of THIK-1 (see methods) using the TRAAK-fab structure (Brohawn et al., 2013) as template. Figure 8 shows that in the homology model residue R92 is in the wall of the outer vestibule, not far from the outer opening of the selectivity filter.

### **The effects of other xanthine derivatives on THIK-1 currents**

To find out which molecular features of IBMX are involved in the binding to THIK-1 we studied the effects of some other xanthine derivatives on THIK-1 channels expressed in *Xenopus* oocytes and CHO cells. In both expression systems, 1 mM theophylline had no effect and 1 mM caffeine had only a very weak effect (Fig. 9 A-D). Xanthine, 1-methyl-xanthine and 1,3-diethyl-7-methyl-xanthine were applied at a concentration of 100  $\mu$ M because of their low water solubility. These substances also had no effect on THIK-1 currents in *Xenopus* oocytes, whereas 100  $\mu$ M IBMX reduced THIK-1 currents by approximately 20 % (Fig. 9E). Taken together, our results with various xanthine derivatives suggest that the 3-isobutyl group of IBMX is important for its inhibitory effect on THIK-1 channels; a methyl group at this position, as in theophylline, or a hydrogen atom, as in 1-methyl-xanthine, does not appear to be sufficient.

## Discussion

### *IBMX blocks THIK-1 channels from the extracellular side*

Our experimental results suggest that the phosphodiesterase inhibitor IBMX acts on THIK-1 channels exclusively from the outside and that it inhibits THIK-1 currents by binding to the helical cap of the channel. This inference is based on the following findings: (i) Extracellular application of IBMX caused a rapid and reversible block of THIK-1 currents in *Xenopus* oocytes (two-electrode voltage clamp) and in a mammalian cell line (whole-cell recording). The time course of the onset and the wash-out of the inhibitory effect on the channel was limited only by the speed of the solution change in the recording chamber. (ii) Intracellular application of IBMX through the patch pipette in the whole-cell recording mode had no effect on THIK-1 currents. Intracellular application of TPA, which was used as a positive control, inhibited both THIK-1 and TREK-1 currents. (iii) In outside-out patches, application of IBMX rapidly and reversibly blocked potassium-selective channels. (iv) Mutation of an arginine residue in the outer cavity of THIK-1 to alanine (R92A) markedly reduced the effect of IBMX on THIK-1 current.

Our discovery of the direct block of the channel from the extracellular side was fortuitous because our experiments were originally aimed at studying the modulation of THIK-1 currents via PKA using IBMX as a blocker of cAMP phosphodiesterase. Since THIK-1 has four predicted PKA phosphorylation sites (Figure 1A) it is important to show that the effects of IBMX on THIK-1 currents described here are not in any way related to a rise in intracellular cAMP and subsequent activation of PKA. The evidence for this can be summarized as follows: (i) The effect of IBMX on THIK-1 current was unchanged after mutation of any of the four predicted PKA phosphorylation sites. (ii) The effect of IBMX persisted when the cells were pre-incubated with the PKA inhibitor H89. (iii) Intracellular application of cAMP through the patch pipette had no effect on THIK-1 current, but inhibited TREK-1 current, which was used as a positive control. (iv) intracellular application of tetrapentylammonium blocked both THIK-1 and TREK-1 channels. (v) Application of the PKA activator forskolin had no effect on THIK-1 currents, but inhibited TREK-1 currents. (v) The inhibitory effect of IBMX on THIK-1 current persisted in outside-out patches, where phosphorylation via PKA should be eliminated due to absence of ATP in the pipette solution.

Caffeine (1 mM) had a much smaller inhibitory effect on THIK-1 channels than IBMX, whereas theophylline (1 mM), 1-methyl-xanthine (1 mM) and 1,3,-diethyl-7-methylxanthine (1 mM) had no effect at all, indicating that perhaps the isobutyl group of IBMX may be relevant for the binding to THIK-1. Now that we have identified a seed

molecule that blocks THIK-1 channels from the extracellular side of the membrane, it appears possible to start a structure-based search of specific THIK-1 modulators that do not have unwanted 'side effects'.

THIK-1 channels are very unusual potassium channels. In cell-attached and outside-out recordings the (voltage dependent) noise attributable to the channel openings could be clearly seen. However, the single-channel conductance of THIK-1 could not be resolved using sampling rate of 10 kHz and a bandwidth of 2 kHz. Since the RMS noise in cell-attached or outside-out patches at the  $K^+$  equilibrium potential was <200 fA and the currents in the patches were between 10 and 40 pA, it may be concluded that each patch contained several channels with very short mean open times.

Furthermore it should be noted that THIK-1 channels are robustly expressed in microglial cells and in neurons of many nuclei in the brain. Thus, experiments on these cells using IBMX or caffeine as pharmacological tools should take into account possible direct effects of IBMX on THIK-1.

#### *Localization of the binding site on the extracellular cap*

Our extensive mutagenesis experiments analysing the effects of IBMX on THIK-1 currents point to an important role of residue R92 in the C2-P1 linker, which is localised to the outer cavity of the channel near the entry of the selectivity filter (Fig. 8). Of all the residues localised to the extracellular cap, only mutation of R92 produced a major change in the effect of IBMX on THIK-1 currents. How can we interpret the observation that mutations in R92 attenuated the inhibitory effect of IBMX? In principle, there are three possibilities: (i) The simplest mechanistic interpretation of our results is that R92 may form part of the IBMX binding site and the binding of the drug to the outer cavity of the channel near the selectivity filter or to the side portal occludes the permeation of potassium ions. (ii) Another possible interpretation is that binding of IBMX may have a purely allosteric effect, i.e. IBMX does not bind to R92 but elsewhere, and this may affect the gating of the channel by inducing a conformational change. The modulatory effect of the R92 mutations would then be explained by the fact that these mutations affect the gating of the channel (considering that R92 is not too far from the selectivity filter). However, it is then difficult to see where the IBMX binding site could be because none of the other mutations of the extracellular cap modulated the effect of IBMX. (iii) Finally, it is conceivable that residue R92 is involved in both the binding of IBMX and in the gating at the selectivity filter, and both are affected by the mutations. In this case, the binding of IBMX would not necessarily occlude the permeation of potassium ions. At present, none of these interpretations can be excluded. The observation that

the mutations did not change the shape of the current voltage relation is consistent with the idea that R92 is involved in the binding of IBMX rather than in the gating of the channel. Structural studies of THIK-1 are required to clarify the most likely orientations of R92.

Taken together, our results strongly suggest that some xanthine derivatives may act as extracellular blockers of THIK-1 channels. The outer cavity and/or the side portals of the extracellular cap were also found to harbour binding sites for water soluble drugs in some other  $K_{2P}$ -channels. The cationic dye ruthenium red was found to block TASK-3, TREK-1 and TREK-2 channels by binding to a site within the extracellular cavity (Braun et al., 2014; Braun et al., 2015; Czirjak and Enyedi, 2003). Another small-molecule drug, the quinoline derivative TKDC, has recently been suggested to block TREK-1 and TREK-2 channels by binding to their extracellular caps (Luo et al., 2017). Since the extracellular cap is quite unique among ion channels, and the C2-P1 linker of  $K_{2P}$ -channels is quite variable, it may be possible to design drugs specific for extracellular binding sites of particular  $K_{2P}$ -channels.

## **Acknowledgements**

We thank Kirsten Ramlow and Doris Wagner for excellent technical support. The support of the High Performance and Cloud Computing Group at the Zentrum für Datenverarbeitung of the University of Tübingen is gratefully acknowledged. None of the authors has any competing interests.

## **Authorship contributions**

Participated in research design: Zou, Conrad, Schlichthörl, Krüger, Daut, Renigunta

Conducted experiments: Zou, Conrad, Schlichthörl, Netz, Renigunta

Contributed new reagents or analytic tools: Koschinsky, Preisig-Müller, Krüger

Performed data analysis: Zou, Conrad, Schlichthörl, Netz, Krüger, Renigunta, Daut

Wrote or contributed to the writing of the manuscript: Daut, Renigunta, Conrad

## REFERENCES

- Aller MI and Wisden W (2008) Changes in expression of some two-pore domain potassium channel genes (KCNK) in selected brain regions of developing mice. *Neuroscience* **151**(4):1154-1172.
- Bayliss DA and Barrett PQ (2008) Emerging roles for two-pore-domain potassium channels and their potential therapeutic impact. *Trends Pharmacol Sci* **29**(11):566-575.
- Bichet D, Blin S, Feliciangeli S, Chatelain FC, Bobak N and Lesage F (2014) Silent but not dumb: how cellular trafficking and pore gating modulate expression of TWIK1 and THIK2. *Pflugers Arch*.
- Braun G, Lengyel M, Enyedi P and Czirjak G (2014) Differential sensitivity of TREK-1, TREK-2 and TRAAK background potassium channels to the polycationic dye ruthenium red. *Br J Pharmacol*.
- Braun G, Lengyel M, Enyedi P and Czirjak G (2015) Differential sensitivity of TREK-1, TREK-2 and TRAAK background potassium channels to the polycationic dye ruthenium red. *Br J Pharmacol* **172**(7):1728-1738.
- Brohawn SG, Campbell EB and MacKinnon R (2013) Domain-swapped chain connectivity and gated membrane access in a Fab-mediated crystal of the human TRAAK K<sup>+</sup> channel. *Proc Natl Acad Sci U S A* **110**(6):2129-2134.
- Brohawn SG, Campbell EB and MacKinnon R (2014) Physical mechanism for gating and mechanosensitivity of the human TRAAK K<sup>+</sup> channel. *Nature* **516**(7529):126-130.
- Brohawn SG, del Marmol J and MacKinnon R (2012) Crystal structure of the human K<sub>2P</sub> TRAAK, a lipid- and mechano-sensitive K<sup>+</sup> ion channel. *Science* **335**(6067):436-441.
- Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglsperger T, Dake B, Wu PM, Doykan CE, Fanek Z, Liu L, Chen Z, Rothstein JD, Ransohoff RM, Gygi SP, Antel JP and Weiner HL (2014) Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat Neurosci* **17**(1):131-143.
- Chatelain FC, Bichet D, Feliciangeli S, Larroque MM, Braud VM, Douguet D and Lesage F (2013) Silencing of the tandem pore domain halothane-inhibited K<sup>+</sup> channel 2 (THIK2) relies on combined intracellular retention and low intrinsic activity at the plasma membrane. *J Biol Chem* **288**(49):35081-35092.
- Czirjak G and Enyedi P (2003) Ruthenium Red Inhibits TASK-3 Potassium Channel by Interconnecting Glutamate 70 of the Two Subunits. *Mol Pharmacol* **63**(3):646-652.
- Dong YY, Pike AC, Mackenzie A, McClenaghan C, Aryal P, Dong L, Quigley A, Grieben M, Goubin S, Mukhopadhyay S, Ruda GF, Clausen MV, Cao L, Brennan PE, Burgess-Brown NA, Sansom MS, Tucker SJ and Carpenter EP (2015) K2P channel gating mechanisms revealed by structures of TREK-2 and a complex with Prozac. *Science* **347**(6227):1256-1259.

- Duprat F, Lauritzen I, Patel A and Honore E (2007) The TASK background K<sub>2</sub>P channels: chemo- and nutrient sensors. *Trends Neurosci* **30**(11):573-580.
- Enyedi P and Czirjak G (2010) Molecular background of leak K<sup>+</sup> currents: two-pore domain potassium channels. *Physiol Rev* **90**(2):559-605.
- Feliciangeli S, Chatelain FC, Bichet D and Lesage F (2014) The family of K<sub>2</sub>P channels: salient structural and functional properties. *J Physiol*.
- Fink M, Duprat F, Lesage F, Reyes R, Romey G, Heurteaux C and Lazdunski M (1996) Cloning, functional expression and brain localization of a novel unconventional outward rectifier K<sup>+</sup> channel. *EMBO J* **15**(24):6854-6862.
- Girard C, Duprat F, Terrenoire C, Tinel N, Fosset M, Romey G, Lazdunski M and Lesage F (2001) Genomic and functional characteristics of novel human pancreatic 2P domain K<sup>+</sup> channels. *Biochem Biophys Res Commun* **282**(1):249-256.
- Kang D, Hogan JO and Kim D (2014) THIK-1 (K2P13.1) is a small-conductance background K<sup>+</sup> channel in rat trigeminal ganglion neurons. *Pflugers Arch* **466**(7):1289-1300.
- Lazarenko RM, Fortuna MG, Shi Y, Mulkey DK, Takakura AC, Moreira TS, Guyenet PG and Bayliss DA (2010) Anesthetic activation of central respiratory chemoreceptor neurons involves inhibition of a THIK-1-like background K<sup>+</sup> current. *J Neurosci* **30**(27):9324-9334.
- Lesage F and Barhanin J (2011) Molecular physiology of pH-sensitive background K<sub>2</sub>P channels. *Physiology (Bethesda)* **26**(6):424-437.
- Luo Q, Chen L, Cheng X, Ma Y, Li X, Zhang B, Li L, Zhang S, Guo F, Li Y and Yang H (2017) An allosteric ligand-binding site in the extracellular cap of K<sub>2</sub>P channels. *Nature communications* **8**(1):378.
- Madry C, Kyrargyri V, Arancibia-Carcamo IL, Jolivet R, Kohsaka S, Bryan RM and Attwell D (2018) Microglial Ramification, Surveillance, and Interleukin-1 $\beta$  Release Are Regulated by the Two-Pore Domain K(+) Channel THIK-1. *Neuron* **97**(2):299-312 e296.
- Miller AN and Long SB (2012) Crystal structure of the human two-pore domain potassium channel K2P1. *Science* **335**(6067):432-436.
- Piechotta PL, Rapedius M, Stansfeld PJ, Bollepalli MK, Ehrlich G, Andres-Enguix I, Fritzenschaft H, Decher N, Sansom MS, Tucker SJ and Baukrowitz T (2011) The pore structure and gating mechanism of K<sub>2</sub>P channels. *EMBO J* **30**(17):3607-3619.
- Rajan S, Wischmeyer E, Karschin C, Preisig-Müller R, Grzeschik KH, Daut J, Karschin A and Derst C (2001) THIK-1 and THIK-2, a novel subfamily of tandem pore domain K<sup>+</sup> channels. *J Biol Chem* **276**(10):7302-7311.
- Renigunta V, Schlichthörl G and Daut J (2015) Much more than a leak: structure and function of K<sub>2</sub>P-channels. *Pflügers Arch* **467**(5):867-894.
- Renigunta V, Zou X, Kling S, Schlichthörl G and Daut J (2014) Breaking the silence: functional expression of the two-pore-domain potassium channel THIK-2. *Pflügers Arch* **466**(9):1735-1745.
- Rinné S, Renigunta V, Schlichthörl G, Zuzarte M, Bittner S, Meuth SG, Decher N, Daut J and Preisig-Müller R (2014) A splice variant of the two-pore

domain potassium channel TREK-1 with only one pore domain reduces the surface expression of full-length TREK-1 channels. *Pflugers Arch* **466**(8):1559-1570.

Terrenoire C, Lauritzen I, Lesage F, Romey G and Lazdunski M (2001) A TREK-1-like potassium channel in atrial cells inhibited by beta-adrenergic stimulation and activated by volatile anesthetics. *Circ Res* **89**(4):336-342.

Yu FH and Catterall WA (2004) The VGL-kanome: a protein superfamily specialized for electrical signaling and ionic homeostasis. *Sci STKE* **2004**(253):re15.

Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, Phatnani HP, Guarnieri P, Caneda C, Ruderisch N, Deng S, Liddelow SA, Zhang C, Daneman R, Maniatis T, Barres BA and Wu JQ (2014) An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**(36):11929-11947.

## Footnotes

This study was Supported by grants of the Deutsche Forschungsgemeinschaft (DFG) to J.D. [FOR 1086 TP7; SFB 593, TP4] and to J.K. [INST 37/935-1 FUGG].

<sup>1</sup> X.Z. and L.J.C. contributed equally to this work.

<sup>2</sup> V.R. and J.D. are corresponding authors

Please send reprint requests either to

Dr. Vijay Renigunta  
Institute of Physiology and Pathophysiology  
Marburg University  
Deutschhausstr. 2  
35037 Marburg, Germany  
Email: vijay@staff.uni-marburg.de

or to

Professor Jürgen Daut  
Institute of Physiology and Pathophysiology  
Marburg University,  
Deutschhausstr. 2  
35037 Marburg, Germany  
Email: jdaut@staff.uni-marburg.de

## Figure legends

**Fig. 1.** The effect of IBMX on THIK-1 currents measured in CHO cells. (A) Topology of THIK-1 and localization of predicted PKA phosphorylation sites. (B) Typical current-voltage relations in CHO cells transfected with THIK-1, measured before (black line) and after (red line) application of 1 mM IBMX. (C) Dose-response curve for the fractional block of THIK-1 current by IBMX; the number of CHO cells from which the data was obtained is indicated in brackets. The error bars indicate the standards deviation. (D) Red curve: Typical recording of the time course of the effect of IBMX on THIK-1 outward currents measured at 0 mV. Black curve: the effect of changing the extracellular potassium concentration from 5 to 24 mM on THIK-1 inward currents measured at -80 mV. The two time courses (red and black curve) were recorded sequentially in the same CHO cell.

**Fig. 2.** The effects of the PKA inhibitor H89 and the PKA activator forskolin on THIK-1 currents were measured in CHO cells at a holding potential of 0 mV. (A) Application of 5  $\mu$ M H89 had no effect on whole-cell currents in CHO cells transfected with THIK-1. (B) The effect of 200  $\mu$ M IBMX on THIK-1 currents persisted after pre-incubation with

H89. The residual current is defined as  $I_{\text{IBMX}}/I_0$ , where  $I_0$  is the current measured before application of IBMX and  $I_{\text{IBMX}}$  is the current measured after application of 200  $\mu\text{M}$  IBMX. (C) Scatter dot plot (mean  $\pm$  S.D.) of the inhibitory effect of IBMX on THIK-1 currents with and without pre-incubation with H89. Open circles represent individual measurements (individual CHO cells). The fractional residual current measured in the presence of IBMX is shown; it is defined as  $I_{\text{IBMX}}/I_0$ , where  $I_0$  is the current measured before application of IBMX and  $I_{\text{IBMX}}$  is the current measured after application of 200  $\mu\text{M}$  IBMX. (D) Application of 20  $\mu\text{M}$  forskolin had no effect on THIK-1 currents in CHO cells. (E) Application of 20 mM forskolin strongly inhibited TREK-1 currents in CHO cells. (F) Scatter dot plot (mean  $\pm$  S.D.) of the effect of 20  $\mu\text{M}$  forskolin on THIK-1 and TREK-1 currents. The fractional residual current is shown; it is defined as  $I_{\text{for}}/I_0$ , where  $I_0$  is the current measured before application of forskolin and  $I_{\text{for}}$  is the current measured after application of 20  $\mu\text{M}$  forskolin.

**Fig. 3.** The effect of intracellular application of 100  $\mu$ M cAMP through the patch pipette on THIK-1 and TREK-1 channels in CHO cells. (A) Typical measurement of the effect of intracellular cAMP on THIK-1 outward current in a CHO cell. The holding potential was set to 0 mV, a cell-attached patch was formed and ruptured by application of negative pressure at the time indicated by the red arrow, allowing cAMP to diffuse into the cell. (B) Scatter dot plot (mean  $\pm$  S.D.) of the relative outward current ( $I_1/I_0$ ) measured after dialysis of the cell with cAMP;  $I_1$  is the current measured 1 min after rupture of the patch membrane; ( $I_0$ ) is the current recorded immediately after rupture of the patch. The cells were transfected with THIK-1, TREK-1 or TREK-1S344A. (C) Typical measurement of the effect of intracellular cAMP on TREK-1 current. (D) Typical measurement of the effect of intracellular cAMP on TREK-1S344A current.

**Fig. 4.** The effects of intracellular and extracellular application of IBMX on THIK-1 in CHO cells, with TREK-1 channels being used as controls. (A) Application of 1 mM IBMX via the patch pipette (left) and via the bath solution (right) in a CHO cell transfected with THIK-1 (continuous recording); please note that the time scale varies between left and right part of the graph. The experimental procedure for intracellular drug application was the same as described in Figure 3; the patch was ruptured at the time indicated by the arrow. (B) Application of 1 mM IBMX via the patch pipette in a CHO cell transfected with TREK-1. (C) Scatter dot plot of fractional residual current ( $I_{\text{IBMX}}/I_0$ ) measured 1 min after intracellular application of 1 mM IBMX in CHO cells transfected with THIK-1 or TREK-1. (D) Application of 20  $\mu$ M TPA via the patch pipette in a CHO cell transfected with THIK-1. (E) Application of 20  $\mu$ M TPA via the patch pipette in a CHO cell transfected with TREK-1.

**Fig. 5.** THIK-1 currents in cell-attached and outside-out patches from CHO cells. The cells were superfused with physiological salt solution containing 5 mM  $K^+$ . (A) Typical cell-attached patch recordings at different pipette potentials (indicated on the right); the sampling rate was 15 kHz, the low-pass filter had a cut-off frequency of 2 kHz. (B) Recordings from the same patch after application of 300  $\mu$ M TPA to the bath solution. (C) Typical recording from an outside-out patch with 5 mM  $K^+$  in the bath solution; the time course of the effect of 1 mM IBMX, was measured at a pipette potential of 0 mV. (D) Average current-voltage relation of outside-out patches measured before (black) and after (red) application of 1 mM IBMX; the 95 % confidence interval is indicated by the grey shading; green: the average difference curve ( $n = 4$  cells). The voltage on the x-axis corresponds to the applied pipette potential (which was not corrected for the liquid junction potential).

**Fig. 6.** The effects of IBMX on THIK-1 channels expressed in *Xenopus* oocytes. (A) Typical example of the current-voltage relation of THIK-1 channels in *Xenopus* oocytes before and after application of 1 mM IBMX. (B) Fractional residual current ( $I_{\text{IBMX}} / I_0$ ) measured at 0 mV after application of 1 mM IBMX. The oocytes were injected with cRNA coding for wild-type THIK-1 or for the THIK-1 mutants indicated.

**Fig. 7.** Effect of mutations in the extracellular cap of rat THIK-1 on the magnitude of the inhibitory effect of IBMX on THIK-1 current in *Xenopus* oocytes. (A) The fractional residual THIK-1 current ( $I_{\text{IBMX}}/I_0$ ) measured at 0 mV in the steady state during application of 500  $\mu\text{M}$  IBMX in wt THIK-1 and in various mutants. The residues shown on the abscissa represent the amino acids of the extracellular cap; the wild-type channel is shown in black (the bar is denoted X). The number of oocytes from which the data were obtained were 5 to 21 for the different mutants; the error bars represent s.d. Only one mutant (R92A, indicated in red;  $n = 18$ ) showed a much larger residual current (i.e. less inhibition by IBMX). Two mutants (indicated in blue) did not produce any current. (B) Plot of the absolute current measured for each mutant against the fractional residual THIK-1 current. The dark grey horizontal line represents the mean value of  $I_{\text{IBMX}}/I_0$  for all mutants (including wild-type and R92A), the light grey lines are three standard deviations above or below the mean. The red circle indicates the mutant R92A. (C) The inhibitory effect of 1 mM IBMX on three THIK-1 mutants with different substitutions for residue R92. (D) The shape of the current-voltage relation in different R92 mutants. Normalized current-voltage relations of wt THIK-1 (grey) and mutants R92A (green), R92E (blue) and R92Q (red); the curves were set to 1 at 0 mV. (E) Alignment of three  $K_{2P}$ -channels whose crystal structure has been determined (TRAAK, TWIK-1 and TREK-2) as well as rat THIK-2, rat THIK-1 and human THIK-1. The residues not resolved in the crystal structure are shown in red; the cysteine residue forming the disulfide bridge between the subunits is highlighted in green; the beginning of the pore helix is highlighted in blue; the unstructured region of the C2-P1 linker is indicated on top by a dotted red line; conserved residues are shown in bold; residue R92 is highlighted in red.

**Fig. 8.** Homology model of THIK-1 based on the TRAAK (4I9W) crystal structure. (A) Ribbon representation viewed from within the membrane; the C2-P1 linker is shown in yellow; residue R92 is shown as spheres (C atoms, yellow; N atoms, blue). (B) View of the helical cap rotated by 90° and enlarged. R92 is localized in the connecting loop between cap domain and the first pore helix. This region of the channel has not been resolved in any crystal structure of K<sub>2P</sub> channels published so far and was modeled *de novo* here. (C) View from the outside into the channel cavity. All residues of the extracellular cap beyond residue R92 have been removed; R92 is shown as spheres.

**Fig. 9.** Effects of various xanthine derivatives on THIK-1 currents. Typical effects of 1 mM caffeine (A) and 1 mM theophylline (B) on THIK-1 currents in *Xenopus* oocytes. (C) and (D) Fractional residual THIK-1 current measured during application of 1 mM theophylline, 1 mM caffeine and 1 mM IBMX in *Xenopus* oocytes (C) and in CHO cells (D). (E) Fractional residual THIK-1 current measured during application of 100 μM IBMX, 100 μM xanthine, 100 μM 1-methylxanthine and 100 μM 1,3-diethyl-7-methylxanthine in *Xenopus* oocytes (there was a slight 'run-up' of current, which explains the increase in fractional residual current).




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9