

MOL #61861

## **cAMP Analogs and Their Metabolites Enhance TREK-1 mRNA and K<sup>+</sup> Current Expression in Adrenocortical Cells**

*Judith A. Enyeart, Haiyan Liu, and John J. Enyeart*

**The Department of Neuroscience, The Ohio State University  
College of Medicine and Public Health (JAE, HL, JJE),  
Columbus, OH 43210-1239**

**Running Title:** cAMP analog metabolites enhance TREK-1 K<sup>+</sup> Current

**Address Correspondence to:** John J. Enyeart, Ph.D. Department of Neuroscience, OSU College of Medicine and Public Health, 5196 Graves Hall, 333 W.10th Avenue, Columbus, OH 43210-1239, Tel: (614) 292-3511, E-Mail: [enyeart.1@osu.edu](mailto:enyeart.1@osu.edu)

**Number of text pages:** 39

**Number of Tables:** 1 (supplemental)

**Number of Figures:** 11 (1 supplemental)

**Number of References:** 47

**Number of words in the Abstract:** 250

**Number of words in the Introduction:** 661

**Number of words in the Discussion:** 1498

**List of non-standard abbreviations:** ACTH, adrenocorticotrophic hormone; AZF, adrenal zona fasciculata; Epac, exchange protein directly activated by cAMP; ESCA, Epac-selective cAMP analog; PKA, cyclic AMP-dependent protein kinase; PBS, phosphate-buffered saline; FBS, fetal bovine serum; DMEM, Dulbecco's Modified Eagle Medium; BAPTA,1,2 *bis*-(2-aminophenoxy)ethane-*N,N,N',N''*-tetraacetic acid; DMSO, dimethyl sulfoxide; 8-Br-cAMP, 8-bromoadenosine- 3', 5'- cyclic monophosphate; 6-Bnz-cAMP, N<sup>6</sup>- benzoyladenosine- 3', 5'- cyclic monophosphate; 8CPT-2'-OMe-cAMP, 8-(4-chlorophenylthio)-2'-O-methyl-cAMP; Sp-8CPT-2'-OMe-cAMP, hydrolysis-resistant 8-(4-chlorophenylthio)-2'-O-methyl-cAMP; 8CPT-2'-OMe-5' AMP, 8-(4-chlorophenylthio)-2'-O-methyladenosine-5'-O-monophosphate; 8CPT-2'-OMe-Ado, 8-(4-chlorophenylthio)-2'-O-methyladenosine; 8CPT-cAMP, 8-(4chlorophenylthio)-cAMP; Sp-8CPT-cAMP, hydrolysis-resistant 8-(4-chlorophenylthio)-cAMP; 8CPT-Ado, 8-(4-chlorophenylthio)-adenosine; 8CPT-Ade, 8-(4-chlorophenylthio) adenine.

## **ABSTRACT**

bTREK-1  $K^+$  channels set the resting membrane potential of bovine adrenal zona fasciculata (AZF) cells and function pivotally in the physiology of cortisol secretion. Adrenocorticotrophic hormone (ACTH) controls the function and expression of bTREK-1 channels through signaling mechanisms that may involve cAMP and downstream effectors including PKA and Epac2. Using patch-clamp and Northern blot analysis, we explored the regulation of bTREK-1 mRNA and  $K^+$  current expression by cAMP-analogs and several of their putative metabolites in bovine AZF cells. At concentrations sufficient to activate both PKA and Epac2, 8-Br-cAMP enhanced the expression of both bTREK-1 mRNA and  $K^+$  current. 6-Bnz-cAMP, which activates PKA but not Epac, also enhanced the expression of bTREK-1 mRNA and  $K^+$  current measured at times from 24-96 h. An Epac-selective cAMP analog 8CPT-2'-OMe-cAMP potently stimulated bTREK-1 mRNA and  $K^+$  current expression, while the non-hydrolyzable Epac-activator Sp-8CPT-2'-OMe-cAMP was ineffective. Metabolites of 8CPT-2'-OMe-cAMP including 8CPT-2'-OMe-5'AMP and 8CPT-2'-OMe-adenosine promoted the expression of bTREK-1 transcripts and ion current with a temporal pattern, potency, and effectiveness resembling that of the parent compound. Similarly, at low concentrations, 8CPT-cAMP (30  $\mu$ M), but not its non-hydrolyzable analog Sp-8CPT-cAMP, enhanced the expression of bTREK-1 mRNA and current. 8CPT-cAMP metabolites, including 8CPT-adenosine and 8CPT-adenine, also increased bTREK-1 expression. These results indicate that cAMP increases the expression of bTREK-1 mRNA and  $K^+$  current through a cAMP-dependent, but Epac2-independent mechanism. They further demonstrate that one or more metabolites of 8-(4-chlorophenylthio)-cAMP analogs potently stimulate bTREK-1 expression by activation of a novel cAMP-independent mechanism. These findings raise significant questions regarding the specificity of 8-(4-chlorophenylthio)-cAMP-analogs as cAMP-mimetics.

## **INTRODUCTION**

Bovine adrenal zona fasciculata (AZF) cells express bTREK-1 (or KCNK2) K<sup>+</sup> leak-type channels that set the resting membrane potential and function pivotally in the physiology of cortisol secretion (Mlinar et al., 1993; Enyeart et al., 2002; Enyeart et al., 1993). Cortisol synthesis is stimulated by the pituitary peptide adrenocorticotrophic hormone (ACTH) (Simpson and Waterman, 1988). ACTH exerts rapid and long-term control over the electrical and secretory properties of AZF cells by regulating both the activity of pre-existing ion channels and the expression of genes coding for these same channel proteins (Mlinar et al., 1993; Enyeart et al., 2000; Enyeart et al., 2003; Enyeart et al., 1996; Liu et al., 2008). In particular, in whole cell recordings, ACTH rapidly (within seconds to minutes) inhibits the activity of bTREK-1 K<sup>+</sup> channels by a cAMP-dependent mechanism (Mlinar et al., 1993; Enyeart et al., 2000; Enyeart et al., 2003; Enyeart et al., 1996). ACTH also induces, with a delay of several hours, an increase in bTREK-1 mRNA and maintains the expression of the associated K<sup>+</sup> current (Enyeart et al., 2003).

The signaling pathways by which ACTH regulates the expression of bTREK-1 K<sup>+</sup> channel mRNA and current are only partially understood. Early studies of cortisol synthesis established cAMP as the principal intracellular messenger for ACTH in AZF cells (Haynes and Berthet, 1957; Grahame-Smith et al., 1967; Richardson and Schulster, 1973; Sala et al., 1979). Accordingly, bovine AZF cells express a high affinity MC2R melanocortin receptor coupled to adenylate cyclase through Gs (Penhoat et al., 1989; Raikhin et al., 1994). Until recently, all of the cAMP-dependent actions of ACTH were thought to be mediated by PKA. However, alternative signaling pathways for cAMP-mediated responses are present in these cells. Specifically, two cAMP-activated guanine nucleotide exchange factors Epac1 and Epac2 (also known as cAMP-GEFI and

cAMP-GEFII) have been identified and implicated in the regulation of cellular processes, including gene expression (de Rooij et al., 1998;Kawasaki et al., 1998;Holz et al., 2006). While Epac1 is expressed in many tissues, Epac2 is robustly expressed in selected areas of the brain and the adrenal glands of rats and humans (de Rooij et al., 1998). Recently, we discovered that Epac2 is strongly expressed in bovine AZF cells, raising the possibility that cAMP could produce responses through this protein and PKA (Liu et al., 2008).

Differentiating between PKA- and Epac-dependent signaling pathways in cells has been hampered by the absence of specific agents that selectively activate each of these two proteins. Exploiting differences in the cAMP-binding domains of these proteins, rational drug design has been employed to synthesize cAMP derivatives that, at appropriate concentrations, specifically activate Epacs or PKA (Enserink et al., 2002;Christensen et al., 2003;Holz et al., 2008). In a patch clamp study, we found that in addition to the well-documented PKA-dependent inhibition of bTREK-1 channel activity, the Epac-selective cAMP analog (ESCA) 8CPT-2'-OMe-cAMP potently inhibited these channels (Liu et al., 2008). Thus, cAMP appears to inhibit bTREK-1 channel function by activation of both PKA and Epac2.

In a more recent study, we found that 8CPT-2'-OMe-cAMP stimulated a delayed increase in cortisol synthesis by inducing the expression of genes coding for steroidogenic proteins, including several steroid hydroxylases (Enyeart and Enyeart, 2009). However, the 8CPT-2'-OMe-cAMP-stimulated increases in cortisol synthesis were not mediated through activation of Epac2. Rather, the effect was produced by one or more metabolites of this 8-(4-chlorophenylthio)-cAMP derivative.

With this knowledge in hand, the present study was done to characterize the signaling pathways by which ACTH and cAMP regulate the expression of bTREK-1 mRNA and

corresponding ion current. It was discovered that several cAMP analogs, including those that selectively activate PKA or Epac2, and those that activate both of these proteins, enhanced the expression of bTREK-1 mRNA and membrane current. Consequently, treatment of AZF cells with any of these cAMP analogs suppresses the time-dependent disappearance of bTREK-1 that typically occurs in culture. However, the increases in bTREK-1 transcripts and current induced by low concentrations of 8CPT-2'-OMe-cAMP and 8CPT-cAMP were mediated indirectly through one or more metabolites of these compounds by activation of an unknown signaling pathway.

## **MATERIALS AND METHODS**

*Materials* - Tissue culture media, antibiotics, fibronectin, and fetal bovine sera (FBS) were obtained from Invitrogen (Carlsbad, CA). Phosphate-buffered saline (PBS), 1,2 bis-(2-aminophenoxy)ethane-*N,N,N',N''*-tetraacetic acid (BAPTA), MgATP, collagenase, DNase, H-89, and ACTH (1-24) were obtained from Sigma (St. Louis, MO). 8- Bromoadenosine- 3', 5'- cyclic monophosphate (8-Br-cAMP, Biolog #B007), N<sup>6</sup>- Benzoyladenosine- 3', 5'- cyclic monophosphate (6-Bnz-cAMP, Biolog #B009), 8-(4-chlorophenylthio)-2'-O-methyl-cAMP (8CPT-2'-OMe-cAMP -Biolog #C041), hydrolysis-resistant 8-(4-chlorophenylthio)-2'-O-methyl-cAMP, Sp-isomer (Sp-8CPT-2'-OMe-cAMP - Biolog #C052), 8-(4-chlorophenylthio)-2'-O-methyladenosine-5'-O-monophosphate (8CPT-2'-OMe-5'AMP – Biolog # C078), 8-(4-chlorophenylthio)-2'-O-methyladenosine (8CPT-2'-OMe-Ado – Biolog #C070), 8-(4chlorophenylthio)-cAMP (8CPT-cAMP- Biolog #C010), hydrolysis-resistant 8-(4-chlorophenylthio)-cAMP (Sp-8CPT-cAMP – Biolog #C012), 8-(4-chlorophenyl-thio)-adenosine (8CPT-Ado- Biolog #C086), and 8-(4-chlorophenylthio) adenine (8CPT-Ade – Biolog #C069) were purchased from Axxora, LLC (San Diego, CA). [ $\alpha$ -<sup>32</sup>P] dCTP was purchased from Perkin Elmer (Waltham, MA). Ultrahyb was

purchased from Ambion (Austin, TX). RNeasy columns for total RNA isolation were obtained from QIAGEN (Valencia, CA). bTREK-1 probe was labeled with [<sup>32</sup>P]dCTP by random priming (Prime-It II kit; Stratagene, La Jolla, CA). Full-length bTREK-1 cDNA (1414 base pairs) was obtained as described previously (Enyeart et al., 2002).

*Isolation and Culture of AZF Cells* - Bovine adrenal glands were obtained from steers (age 2-3 yr) at a local slaughterhouse. Isolated AZF cells were obtained and prepared as previously described (Enyeart et al., 1997). After isolation, cells were either resuspended in DMEM/F12 (1:1) with 10 % FBS, 100 U/ml penicillin, 0.1 mg/ml streptomycin, and the antioxidants  $\alpha$ -tocopherol (1  $\mu$ M), 20 nM selenite and 100  $\mu$ M ascorbic acid (DMEM/F12+) and plated for immediate use, or resuspended in FBS/5% DMSO, divided into 1 ml aliquots, and stored in liquid nitrogen for future use. To ensure cell attachment, dishes were treated with fibronectin (10  $\mu$ g/ml) at 37°C for 30 minutes then rinsed with warm, sterile PBS immediately before adding cells. For patch clamp experiments, cells were plated in DMEM/F12+ in 35 mm dishes containing 9 mm<sup>2</sup> glass coverslips (Bellco, Vineland NJ). . Coverslips were treated with fibronectin (10  $\mu$ g/ml) as described above. Cells were maintained at 37°C in a humidified atmosphere of 95% air-5% CO<sub>2</sub>.

*Measurement of bTREK-1 mRNA*- RNeasy columns (Qiagen, Valencia, CA), treated with RNase-free DNase (Qiagen, Valencia, CA) to remove genomic contamination, were used to extract total RNA from AZF cells. Total mRNA (10  $\mu$ g/lane) was separated on denaturing 8% formaldehyde, 1.0% agarose gels, and transferred to nylon membranes (Gene Screen Plus, NEN). RNA was fixed to the membrane by UV-crosslinking (Stratalinker, Stratagene La Jolla, CA), prehybridized for 2 hrs at 42°C in ULTRAhyb (Ambion, Austin, TX), then hybridized with a [ $\alpha$  <sup>32</sup>P]dCTP-labelled bTREK-1

full-length cDNA as previously described (Enyeart et al., 2003). Northern autoradiograms were imaged using a Typhoon 9200 variable mode phosphorimager and quantitated using ImageQuant TL v2003.3 software (GE Healthcare Life Sciences, Piscataway, NJ). mRNA values are presented as mean  $\pm$  SEM of at least 3 independent determinations. For the figures, a representative Northern blot of at least 3 independent experiments is shown. Statistically significant differences were determined by unpaired *t*-test analysis (GraphPad Software, Inc., La Jolla, CA). *P* values  $<0.05$  were considered statistically significant.

*Patch Clamp Experiments*- Patch clamp recordings of K<sup>+</sup> channel currents were made in the whole cell configuration from bovine AZF cells. The standard external solution consisted of 140 mM NaCl, 5 mM KCl, 2 mM CaCl<sub>2</sub>, 2 mM MgCl<sub>2</sub>, 10 mM HEPES, and 5 mM glucose, with pH adjusted to 7.3 using NaOH. The standard pipette solution consisted of 120 mM KCl, 1 mM CaCl<sub>2</sub>, 2 mM MgCl<sub>2</sub>, 11 mM BAPTA, 10 mM HEPES, 5 mM ATP, and 200  $\mu$ M GTP, with pH titrated to 6.8 using KOH.

*Recording Conditions and Electronics*- AZF cells were used for patch clamp experiments 2–12 h after plating. Typically, cells with diameters  $<15\ \mu$ m and capacitances of 10–15 pF were selected. Coverslips were transferred from 35-mm culture dishes to the recording chamber (volume: 1.5 ml) that was continuously perfused by gravity at a rate of 3–5 ml/min. For whole cell recordings, patch electrodes with resistances of 1.0–2.0 M $\Omega$  were fabricated from Corning 0010 glass (World Precision Instruments, Sarasota, FL). These electrodes routinely yielded access resistances of 1.5–4.0 M $\Omega$  and voltage-clamp time constants of  $<100\ \mu$ s. K<sup>+</sup> currents were recorded at room



temperature (22–25 °C) according to the procedure of Hamill *et al.* (Hamill et al., 1981) using a List EPC-7 patch clamp amplifier.

Pulse generation and data acquisition were done using a personal computer and pCLAMP software with a Digidata 1200 interface (Molecular Devices, Sunnyvale, CA). Currents were digitized at 2–10 KHz after filtering with an 8-pole Bessel filter (Frequency Devices, Haverhill, MA). Linear leak and capacity currents were subtracted from current records using summed scaled hyperpolarizing steps of ½ to ¼ pulse amplitude. Data were analyzed using CLAMPFIT 9.2 (Molecular Devices, Sunnyvale, CA) and SigmaPlot (version 10.0) software. Drugs were applied by bath perfusion, controlled manually by a six-way rotary valve.

## **RESULTS**

### *ACTH and 8-Br-cAMP Promote the Expression of bTREK-1 K<sup>+</sup> Current*

Bovine AZF cells express two types of K<sup>+</sup> channels – a voltage-gated, rapidly inactivating Kv1.4 (or KCNA4) channel, and a two-pore domain, four transmembrane-spanning segment (2P/4TMS) bTREK-1 (or KNCK2) background K<sup>+</sup> channel (Mlinar et al., 1993; Mlinar and Enyeart, 1993; Enyeart et al., 2002). In whole cell patch clamp recordings bTREK-1 amplitude spontaneously increases over a period of 10-20 minutes to a stable maximum. The absence of time- and voltage-dependent inactivation allows bTREK-1 K<sup>+</sup> currents to be isolated in whole cell recordings using either of two voltage clamp protocols. When voltage steps of several hundred milliseconds duration are applied from a holding potential of -80 mV, bTREK-1 current can be measured near the end of a voltage step when the Kv1.4 current has completely inactivated (Figure 1A-D, left traces). Alternatively, bTREK-1 current can be selectively activated by an identical

voltage step, applied immediately after a 10 s prepulse to -20 mV has fully inactivated Kv1.4 channels (Figure 1A-D, right traces).

When bovine AZF cells were cultured in serum-supplemented media for periods of 24 h or more, the bTREK-1 K<sup>+</sup> current was markedly diminished compared to currents recorded 1-3 h after plating (Figure 1A,B,E). In contrast, when these cells were continuously exposed to ACTH (2 nM) for 48 h after plating, bTREK-1 current was well maintained, compared to time-matched controls (Figure 1C,E). Overall, when AZF cells were cultured in serum-supplemented (control) media for 48 h, bTREK-1 current density decreased from its original density of  $47.1 \pm 6.5$  pA/pF (n=7) to  $5.37 \pm 1.35$  pA/pF (n=17). In contrast, bTREK-1 current density in ACTH-treated cells decreased to only  $32.5 \pm 12.3$  pA/pF (n=7) (Figure 1D).

cAMP analogs substituted at the 8-position of the adenine ring activate both PKA and Epac2 (Christensen et al., 2003;Poppe et al., 2008). Although 8-substituted cAMP analogs bind to PKA and Epac proteins with K<sub>d</sub>s in the low micromolar range, limited membrane permeability dictates that higher concentrations are required to produce effects in intact cells (Liu et al., 2008;Christensen et al., 2003). 8-Br-cAMP stimulates large increases in cortisol secretion by bovine AZF cells only when applied at concentrations above 100  $\mu$ M (Supplementary Figure 1A).

We tested the effect of 8-Br-cAMP (300  $\mu$ M) on the expression of bTREK-1 in AZF cells. As illustrated in Figure 1D and E, 8-Br-cAMP mimicked ACTH in promoting the expression of bTREK-1 current. After a 48 h exposure to 8-Br-cAMP, bTREK-1 current density was 4.6 times greater than the time-matched control. Similar to ACTH, exposing cells to 8-Br-cAMP for prolonged periods suppressed the expression of Kv1.4 (Figure 1C, D). This response was not further explored in the present study.

### *6-Bnz-cAMP Enhances bTREK-1 Expression*

The increases in bTREK-1 current density induced by ACTH and 8-Br-cAMP could have occurred through activation of Epac2, PKA or perhaps an unidentified signaling pathway. cAMP derivatives with substitutions at the 6-position of the adenine ring selectively activate PKA over Epac proteins (Christensen et al., 2003). Similar to 8-Br-cAMP, limited membrane permeability likely diminishes the potency of 6-Bnz-cAMP when it is applied to intact cells (Liu et al., 2009). 6-Bnz-cAMP stimulates large increases in cortisol secretion from AZF cells only at concentrations above 100  $\mu$ M (Supplemental Figure 1B). Accordingly, we found that prolonged incubation of AZF cells with 6-Bnz-cAMP (200  $\mu$ M), but not 30  $\mu$ M, markedly enhanced the expression of bTREK-1 current measured at times from 48 to 96 h compared to time-matched controls (Figure 2C,D). After 48 h in the presence of 6-Bnz-cAMP (200  $\mu$ M), bTREK-1 current density was 9-fold greater than that of untreated cells (Figure 2D). These results suggest that cAMP can enhance the expression of bTREK-1 by activating PKA alone.

### *Epac-selective cAMP Analogs and Metabolites Promote bTREK-1 Expression*

Experiments with 6-Bnz-cAMP indicated that activation of PKA was sufficient to stimulate bTREK-1 current expression. Additional studies were done to determine the role of Epac2 in the expression of this current. The ESCA 8CPT-2'-OMe-cAMP has been shown to activate Epac proteins when applied to intact cells at concentrations from 10-100  $\mu$ M (Enserink et al., 2002; Holz et al., 2008). We found that 8CPT-2'-OMe-cAMP (30  $\mu$ M) effectively stimulated bTREK-1 current expression measured at times from 48 to 96 h (Figure 3A,C). As illustrated in Figure 3C, in control media, bTREK-1 current density decreased spontaneously with time, reaching an apparent minimum of  $5.27 \pm 1.35$  pA/pF (n=17) by 48 h, after which no further decay occurred. However, in

the presence of 8CPT-2'-OMe-cAMP, bTREK-1 current density was significantly increased at each time point. By 48 h, bTREK-1 current density reached a nearly constant value that was approximately 4.5 fold greater than that of the time-matched controls at 48, 72, and 96 h (Figure 3C).

The marked stimulation of bTREK-1 expression by 8CPT-2'-OMe-cAMP suggested that cAMP could induce the expression of bTREK-1 channels by activation of Epac2. However, at the same concentration, Sp-8CPT-2'-OMe-cAMP, a membrane-permeable ESCA that is resistant to hydrolysis by cyclic nucleotide phosphodiesterase, failed to stimulate the expression of bTREK-1 current at 48 and 72 h. In fact, in the presence of this ESCA, bTREK-1 current was significantly reduced compared to controls (Figure 3B,C).

The effective stimulation of bTREK-1 current expression by 8CPT-2'-OMe-cAMP combined with the failure of its non-hydrolyzable analog to mimic this action indicated that, at low concentrations, 8CPT-2'-OMe-cAMP acts independently of Epac2. In this regard, it has become apparent that, in some cells, significantly higher concentrations are required for effective activation of Epac2 (Chepurny et al., 2009; Holz et al., 2008). Previously, we showed that in AZF cells, Sp-8CPT-2'-OMe-cAMP (100  $\mu$ M) activated Rap1, a downstream effector of Epac2 (Enyeart and Enyeart, 2009). However, we now report that treating AZF cells for 48 h with Sp-8CPT-2'-OMe-cAMP (100  $\mu$ M) did not enhance the expression of bTREK-1 currents (Figure 4A,C).

The results of experiments with the ESCAs indicated that activation of Epac2 does not induce the expression of bTREK-1 current. They further suggested that 8CPT-2'-OMe-cAMP increased bTREK-1  $K^+$  current expression indirectly by generation of one or more active metabolites.

8CPT-2'-OMe-cAMP can be sequentially converted to 8CPT-2'-OMe-5' AMP and 8CPT-2'-OMe-adenosine (8CPT-2'-OMe-Ado) by cyclic nucleotide phosphodiesterase and 5' nucleotidases, respectively (Price and Stevens, 1999; Holz et al., 2008) (Figure 5). We discovered that 8CPT-2'-OMe-Ado potently and effectively stimulated the expression of bTREK-1 current. After exposing cells for 48 h to this metabolite at a concentration of 30  $\mu$ M, bTREK-1 current density reached a value 6 times that of its time-matched control (Figure 4C). This large increase in bTREK-1 current persisted after 72 h.

#### *Effect of cAMP Analogs and Metabolites on bTREK-1 mRNA*

Previously, we showed that ACTH-induced increases in bTREK-1 expression were associated with increases in bTREK-1 mRNA (Enyeart et al., 2003). Experiments were done to determine if the increases in bTREK-1 current induced by cAMP analogs and their presumptive metabolites were mediated through the enhancement of bTREK-1 gene expression. In Northern blots using RNA isolated from AZF cells, a 1414 base pair cDNA probe that included the entire coding sequence of bTREK-1 hybridized to separate mRNA transcripts of ~4.9, 3.6, and 2.8 Kb (Figure 6A) (Enyeart et al., 2002). When bovine AZF cells were cultured in serum-supplemented media for 5 - 48 h, the quantity of bTREK-1 mRNA decreased over time (Figure 6A).

Similar to ACTH, 8-Br-cAMP (300  $\mu$ M) markedly increased the expression of bTREK-1 mRNA in AZF cells. In the experiment illustrated in Figure 6B, AZF cells were cultured overnight in serum-supplemented media before exposing them for 48 h to ACTH or 8-Br-cAMP at the indicated concentrations. ACTH (2nM) and 8-Br-cAMP (300  $\mu$ M) increased mRNA expression by  $2.0 \pm 0.1$  and  $2.3 \pm 0.3$  fold, respectively. Although each of the three bTREK-1 transcripts were induced by 8-Br-cAMP and ACTH, the smaller 3.6 and 2.8 Kb transcripts were preferentially

increased. At a 10 fold lower concentration, 30  $\mu$ M 8-Br-cAMP failed to stimulate bTREK-1 mRNA expression (Figure 6B).

In addition to 8-Br-cAMP, we found that 6-Bnz-cAMP (200  $\mu$ M) also enhanced the expression of bTREK-1 mRNA (Figure 6C). However, the PKA antagonist H-89 (10  $\mu$ M) only partially inhibited 6-Bnz-cAMP-stimulated increases in bTREK-1 expression. In 3 similar, independent experiments, H-89 reduced 6-Bnz-cAMP-stimulated bTREK-1 expression by  $56 \pm 2\%$ . These results indicate that 6-Bnz-cAMP-stimulated increases in bTREK-1 expression are mediated only in part by PKA.

Patch clamp experiments indicated that the ESCA 8CPT-2'-OMe-cAMP induced bTREK-1 current indirectly after conversion to one or more active metabolites (Figures 3 and 4). Accordingly, we found that 8CPT-2'-OMe-cAMP potently and effectively induced the expression of bTREK-1 mRNA, but this effect was again mediated through hydrolysis products of this cAMP analog, rather than by activation of Epac2.

8CPT-2'-OMe-cAMP stimulated the expression of bTREK-1 mRNA at the same concentrations that enhanced the expression of bTREK-1 current. In the experiment illustrated in Figure 7A, bTREK-1 mRNA was measured after exposing AZF cells to 8CPT-2'-OMe-cAMP at concentrations ranging from 10 to 50  $\mu$ M for 48 h. A significant increase was observed at 20  $\mu$ M, while 50  $\mu$ M 8CPT-2'-OMe-cAMP induced a maximum  $2.8 \pm 0.4$ -fold increase in this mRNA. The 8CPT-2'-OMe-cAMP-induced increases in bTREK-1 transcripts could be observed after a delay of 5 to 10 h, and bTREK-1 mRNA continued to increase for at least 48 h (Figure 7B). The temporal pattern of these increases in mRNA was therefore well correlated with the corresponding enhancement in bTREK-1 current.

In contrast to 8CPT-2'-OMe-cAMP, Sp-8CPT-2'-OMe-cAMP failed to promote the expression of bTREK-1 mRNA when applied to cells at 30  $\mu$ M, or at higher concentrations where this non-hydrolyzable ESCA produces significant activation of Rap1 (Figure 7C, left and right panels)(Enyeart and Enyeart, 2009). Accordingly, we found that the two putative metabolites 8CPT-2'-OMe-5' AMP and 8CPT-2'-OMe-Ado increased the expression of bTREK-1 mRNA with potency similar to that of the parent compound (50  $\mu$ M 8CPT-2'-OMe-5' AMP,  $2.4 \pm 0.1$  fold; 50  $\mu$ M 8CPT-2'-OMe-Ado,  $2.1 \pm 0.6$  fold, Figure 7D). In contrast, adenosine (50  $\mu$ M) failed to stimulate any increase in bTREK-1 expression ( $0.9 \pm 0.1$  fold, Figure 7D, right panel).

Overall, these results indicate that cAMP induces bTREK-1 mRNA and  $K^+$  current through activation of PKA but not Epac2. Remarkably, increases in bTREK-1 mRNA and  $K^+$  current induced by the ESCA 8CPT-2'-OMe-cAMP appear to be produced indirectly by one or more metabolites of this compound.

#### *Effect of 8CPT-cAMP and Metabolites on bTREK-1 Expression*

In a previous study, we showed that 8CPT-cAMP (250  $\mu$ M) increased the expression of bTREK-1 mRNA (Enyeart et al., 2003). Although it likely activates both PKA and Epac2 at this concentration, this 8-(4-chlorophenylthio)-derivative of cAMP can be metabolized by the same enzymes that hydrolyze 8CPT-2'-OMe-cAMP (Price and Stevens, 1999; Holz et al., 2008). This raised the possibility that metabolites of 8CPT-cAMP contributed to the observed increases in bTREK-1 mRNA.

To answer this question, we compared 8CPT-cAMP and its membrane-permeable non-hydrolyzable analog Sp-8CPT-cAMP with respect to their potency as enhancers of bTREK-1 current and mRNA (Dostmann et al., 1990). Exposing cells for 48 h to 30  $\mu$ M 8CPT-cAMP

markedly increased bTREK-1 current density from  $5.85 \pm 1.22$  pA/pF (n=20) to  $35.2 \pm 7.6$  (n=7) (Figure 8A). In contrast, treating cells with Sp-8CPT-cAMP (30  $\mu$ M) failed to increase bTREK-1 expression (Figure 8B, D). These results indicated that, at this low concentration, 8CPT-cAMP increased bTREK-1 expression by a mechanism independent of cAMP. At a 10-fold higher concentration, the poorly hydrolyzable Sp-8CPT-cAMP (300  $\mu$ M) increased bTREK-1 current density 4.2-fold over the time-matched control to  $24.5 \pm 2.9$  pA/pF (n=7) (Figure 8C,D).

These results suggested that, at low concentrations where 8CPT-cAMP does not effectively activate PKA, it stimulates bTREK-1 current expression only after it is converted to one or more active metabolites. In contrast, at higher concentrations, the non-hydrolyzable Sp-8CPT-cAMP increases bTREK-1 current expression by activation of PKA.

8CPT-cAMP can be metabolized to 8CPT-5'AMP and 8CPT-adenosine (8CPT-Ado) by cyclic nucleotide phosphodiesterase and 5'nucleotidase, respectively (Price and Stevens, 1999). 8CPT-Ado can then be converted to 8CPT-adenine (8CPT-Ade) by purine nucleoside phosphorylase (Price and Stevens, 1999). Interestingly, 8CPT-Ade may be synthesized from both 8CPT-2'-OMe-cAMP and 8CPT-cAMP in AZF cells (Figure 5). Therefore, we examined the effects of 8CPT-Ado and 8CPT-Ade on bTREK-1 expression.

bTREK-1 current was markedly enhanced in response to a 48 h incubation with both 8CPT-Ado (30  $\mu$ M) and 8CPT-Ade (30  $\mu$ M) (Figure 9A-C). The 4.6- and 7.2-fold increases in current density induced by these metabolites compare with the 6.0 fold increase produced by 8CPT-cAMP at the same concentration. Thus, patch clamp experiments indicated that, at low concentrations, 8CPT-cAMP enhanced the expression of bTREK-1 K<sup>+</sup> current indirectly after conversion to one or more active metabolites.



*Effect of 8CPT-cAMP, Sp-8CPT-cAMP, and 8CPT-adenine on bTREK-1 mRNA*

*Expression*

In Northern blot analysis, we found that the effects of 8CPT-cAMP and Sp-8CPT-cAMP on bTREK-1 mRNA expression were consistent with their effects on bTREK-1 K<sup>+</sup> current. Similar to results seen in experiments with 8CPT-2'-OMe-cAMP and Sp-8CPT-2'-OMe-cAMP (Figure 7C), exposing AZF cells to 30  $\mu$ M 8CPT-cAMP for 48 h significantly increased bTREK-1 mRNA ( $1.8 \pm 0.2$  fold), while non-hydrolyzable Sp-8CPT-cAMP was ineffective in this regard (Figure 10A).

8CPT-Ade stimulated a concentration-dependent increase in bTREK-1 mRNA transcripts with a potency, temporal pattern, and effectiveness similar to that of other cAMP metabolites used in this study (Figure 10B). In the experiment illustrated in Figure 10B, 8CPT-Ade increased bTREK-1 mRNA at concentrations from 10 to 50  $\mu$ M. With respect to temporal pattern, ACTH induced the expression of bTREK-1 mRNA more rapidly than 8CPT-Ade, but by 48 h, these two agents were similarly effective. In the experiment illustrated in Figure 10C, at 48 h, 8CPT-Ade and ACTH induced  $2.9 \pm 0.1$ - and  $2.4 \pm 0.1$ -fold increases in bTREK-1 transcript relative to the time-matched control.

## **DISCUSSION**

In this study, we presented evidence indicating that cAMP enhanced the expression of bTREK-1 mRNA and K<sup>+</sup> current and prevented their disappearance by activating PKA, but not Epac2. Further, hydrolysis products of 8-(4-chlorophenylthio)-cAMP analogs enhanced bTREK-1 mRNA transcripts and corresponding membrane current. The hydrolyzable 8-(4-chlorophenylthio)-cAMP analogs and their putative metabolites were significantly more potent than other cAMP analogs at promoting bTREK-1 expression. These metabolites all induced bTREK-1 with a

potency, temporal pattern, and effectiveness similar to the parent compound, but by an unknown cAMP-independent pathway.

Overall, we have identified 10 separate cAMP analogs and metabolites that enhanced the expression of bTREK-1 mRNA and associated K<sup>+</sup> current (Supplemental Table 1). For each of these, the effects on TREK-1 mRNA and current were well correlated. These results are consistent with the hypothesis that each of these agents increases the rate of transcription of the bTREK-1 gene. However, our results do not exclude the possibility that post-transcriptional mechanisms are involved, including bTREK-1 mRNA stabilization as well as translational or post-translational control by these compounds. It is unlikely that the cAMP-dependent and independent actions of these on bTREK-1 are produced by a single common mechanism.

#### *cAMP, PKA, Epac2, and bTREK-1 Expression*

Experiments with 8-Br-cAMP and Sp-8CPT-cAMP clearly established that cAMP stimulates the expression of bTREK-1 mRNA and associated K<sup>+</sup> current. These experiments were necessary because our previous study demonstrating that 8CPT-cAMP induced bTREK-1 mRNA was completed without the knowledge that hydrolyzable 8-(4-chlorophenylthio) derivatives of cAMP induce bTREK-1 mRNA indirectly through metabolites (Enyeart et al., 2003). Consequently, it was important to determine whether cAMP could induce bTREK-1 expression, because ACTH appears to produce effects in bovine AZF cells by cAMP-dependent and independent mechanisms (Yamazaki et al., 1998; Yamazaki et al., 2006; Moyle et al., 1973).

The robust stimulation of bTREK-1 mRNA and K<sup>+</sup> current expression by 6-Bnz-cAMP suggests that the expression of this K<sup>+</sup> channel is controlled at the pre-translational level by cAMP

through the activation of PKA. However, several questions remain regarding the molecular mechanisms and signaling pathways by which 6-Bnz-cAMP functions.

First, with regard to molecular mechanism, ACTH- and cAMP induce increases in steroid hydroxylase mRNAs by accelerating the rate of gene transcription (John et al., 1986; Waterman, 1994). Our results are consistent with a similar effect of 6-Bnz-cAMP on bTREK-1 gene transcription. However, they do not rule out an effect of this cAMP analog on bTREK-1 mRNA stability.

With respect to signaling mechanism, cAMP analogs substituted in the 6-position of the adenine ring selectively activate PKA over Epac proteins (Christensen et al., 2003). Therefore, our observation that 6-Bnz-cAMP robustly induced the expression of bTREK-1 mRNA and K<sup>+</sup> current argue that cAMP-stimulated increases in the expression of this K<sup>+</sup> channel were mediated through PKA rather than Epac2. However, several lines of evidence suggest that 6-Bnz-cAMP could function through a third unidentified cAMP receptor. First, when applied at concentrations that suppress PKA activity, H-89 only partially suppressed increases in bTREK-1 mRNA expression (Davies et al., 2000). Further, in patch clamp experiments, we have found that, when included in the patch electrode at very low concentrations (< 10  $\mu$ M), 6-Bnz-cAMP completely blocks this activity of bTREK-1 channels, even in the presence of several PKA antagonists (Liu et al., 2009).

The steroidogenic actions of cAMP in AZF cells also suggest the presence of additional cAMP binding proteins. PKA-stimulated transcription typically occurs within minutes and does not require *de novo* protein synthesis (Parker and Schimmer, 1995). In contrast, cAMP-induced increases in steroid hydroxylase-specific mRNAs are cycloheximide-sensitive and require periods of up to several hours before they can be observed (Waterman, 1994; Simpson and Waterman, 1988; Waterman and Simpson, 1988). Further, although the steroid hydroxylase genes are all

induced by cAMP, they lack consensus cAMP response elements in their 5' flanking regions and therefore do not bind PKA phosphorylated transcription factors (Parker and Schimmer, 1995; Payne and Hales, 2004; Simpson and Waterman, 1988; Lund et al., 1990; Kagawa and Waterman, 1990; Ahlgren et al., 1990).

In a number of other cells, cAMP synthesized in response to activation of G protein-coupled receptors produces effects that are independent of PKA or Epac proteins (Fujita et al., 2002; Ivins et al., 2004; Iocovelli et al., 2001; Stork and Schmitt, 2002; Busca et al., 2000). Overall, these results suggest that 6-Bnz-cAMP may induce the expression of bTREK-1 through a cAMP-dependent pathway in addition to PKA.

Experiments with the ESCAs 8CPT-2'-OMe-cAMP and its non-hydrolyzable analog Sp-8CPT-2'-OMe-cAMP clearly demonstrated that cAMP does not enhance the expression of bTREK-1 K<sup>+</sup> channels by activation of Epac2. Specifically, 8CPT-2'-OMe-cAMP induced bTREK-1 mRNA and K<sup>+</sup> current at concentrations that produce little or no activation of the Epac2 effector Rap1, while Sp-8CPT-2'-OMe-cAMP failed to increase bTREK-1 expression at higher concentrations previously shown to activate Rap1 in these cells (Enyeart and Enyeart, 2009). Thus, ACTH- and cAMP-stimulated expression of bTREK-1 appears to be mediated by in part by PKA, but independently of Epac2.

#### *8-(4-chlorophenylthio)-cAMP Metabolites and bTREK-1 Expression*

Results of experiments with hydrolyzable and non-hydrolyzable 8-(4-chlorophenylthio)-cAMP analogs strongly suggest that the hydrolyzable cAMP analogs stimulated bTREK-1 expression indirectly through metabolites. Accordingly, we found that a total of 5 potential metabolites of 8CPT-cAMP and 8CPT-2'-OMe-cAMP each induced the expression of bTREK-1 mRNA transcripts and ion current with potency and effectiveness similar to the parent compounds.

Enzymes that could catalyze the conversion of these two cAMP derivatives to each of these putative metabolites are expressed in mammalian cells. These include cyclic nucleotide phosphodiesterases, 5' nucleotidases, hypoxanthine phosphoribosyltransferase, and nucleotide phosphorylases (Price and Stevens, 1999).

The active metabolites, associated receptor, and signaling pathways that mediate the increases in bTREK-1 gene expression have not been identified. It is possible that each of the five metabolites stimulates bTREK-1 expression. In this regard, it is interesting to note that both 8CPT-cAMP and the ESCA 8CPT-2'-OMe-cAMP can be converted to the same 8-(4-chlorophenylthio)-adenine derivative (Figure 5). It will be important to determine if 8CPT-Ade is the final common active metabolite.

Regardless of the identity of the active metabolite(s), it is clear that it does not function by activating a cAMP-dependent mechanism. First, although the hydrolyzable 8-(4-chlorophenylthio)-cAMP analogs and their metabolites at low concentrations all induced bTREK-1 expression, none of these down-regulated the Kv1.4 current whose expression is inhibited by a PKA-dependent mechanism (Enyeart et al., 2000). In contrast, 8-Br-cAMP and 6-Bnz-cAMP, as well as ACTH, at concentrations that induced bTREK-1, all inhibited the expression of Kv1.4 current (Figures 1,2,3,4 and 9). Further, we previously showed that neither 8CPT-2'-OMe-cAMP, 8CPT-2'-OMe-Ado, nor 8CPT-Ade activate PKA when applied to bovine AZF cells (Enyeart and Enyeart, 2009). Finally, in the current study we demonstrated that a non-hydrolyzable ESCA failed to induce bTREK-1 expression, even at concentrations that have been shown to activate Epac2.

The findings of this study are consistent with the possibility that cAMP metabolites induce bTREK-1 by activation of a cAMP-independent pathway that is also activated by ACTH. An *O*-nitrophenyl-derivative of ACTH, NPS-ACTH blocks the activity of bTREK-1 K<sup>+</sup> channels, induces

increases in  $[Ca^{2+}]_i$ , and stimulates large increases in cortisol synthesis at concentrations that produce little or no increase in cAMP synthesis (Liu et al., 2008;Moyle et al., 1973;Yamazaki et al., 1998). Although these effects are mediated through the MC2R receptor, the cAMP-independent signaling pathway has not been identified.

Previously, we showed that metabolites of 8CPT-2'-OMe-cAMP induced large, delayed increases in the expression of genes coding for steroidogenic proteins, resulting in corresponding increases in cortisol synthesis (Enyeart and Enyeart, 2009). With respect to kinetics and concentration-dependence, the effects of the metabolites on the expression of steroidogenic proteins resemble those on bTREK-1, suggesting a common mechanism.

The range of genes whose expression might be modulated by this novel signaling pathway is yet to be determined. It is clear that the presumptive metabolites did not produce a non-specific global increase in RNA synthesis. Total RNA synthesis was not increased in metabolite-treated cells. Expression of specific genes, including the transcriptional repressor Dax-1 and Kv1.4 were not increased by the metabolites (unpublished observations).

This and other studies demonstrate that ACTH and cAMP exert long-term control over the electrical properties of AZF cells by regulating the expression of genes that code for ion channels (Enyeart et al., 2000;Enyeart et al., 2003). ACTH may function through cAMP-dependent and – independent mechanisms. In this regard, it is possible that ACTH and metabolites of 8-(4-chlorophenylthio)-cAMP analogs may induce the expression of genes coding for bTREK-1  $K^+$  channels and steroidogenic proteins by activating a common, but yet to be identified, cAMP-independent pathway.

The extent to which metabolites of 8-(4-chlorophenylthio)-cAMP derivatives regulate gene expression and cell function in other tissues and organisms is unknown. Hydrolysis products of

selected cAMP analogs transform the protozoa *Trypanosoma brucei* from slender to stumpy-like forms (Laxman et al., 2006). It is possible that the effect of these metabolites in distantly related eukaryotes might be mediated through a common ancient signaling pathway.

Finally, 8CPT-cAMP analogs, including 8CPT-cAMP and 8CPT-2'-OMe-cAMP have been used in hundreds of studies to determine the roles of cAMP, PKA, and Epac proteins in cell signaling. The findings of our study indicate that the results of some of these previous studies may require re-evaluation.

## REFERENCES

- Ahlgren R, Simpson E R, Waterman M R and Lund J (1990) Characterization of the Promoter/Regulatory Region of the Bovine CYP11A (P-450<sub>scc</sub>) Gene. Basal and cAMP-Dependent Expression. *J Biol Chem* 265:3313-3319.
- Busca R, Abbe P, Mantoux R, Aberdam E, Peyssonnaud C, Eyche A, Ortonne J P and Ballotti R (2000) Ras Mediates the cAMP-Dependent Activation of Extracellular Signal-Regulated Kinases (ERKs) in Melanocytes. *EMBO J* 9:2900-2910.
- Chepurny OG, Leech C A, Kelley G G, Dzhura I, Dzura E, Xiangquan L, Rindler M J, Genieser H G and Holz G G (2009) Enhanced Rap1 Activation and Insulin Secretagogue Properties of an Acetoxymethyl Ester of an Epac-Selective Cyclic AMP Analogue in Rat INS-1 Cells. *J Biol Chem* 284:10728-10736.
- Christensen AE, Sleheim F, de Rooij J, Dremier S, Schwede F, Dao K, Martinez A, Maenhaut C, Bos J L, Genieser H G and Doskeland S O (2003) cAMP Analog Mapping of Epac1 and cAMP Kinase. Discriminating Analogs Demonstrate That Epac and cAMP Kinase Act Synergistically to Promote PC-12 Cell Neurite Extension. *J Biol Chem* 278:35394-35402.
- Davies SP, Reddy H, Caivano M and Cohen P (2000) Specificity and Mechanism of Action of Some Commonly Used Protein Kinase Inhibitors. *Biochem J* 351:95-105.



de Rooij J, Zwartkruis F J, Verheijen M H, Cool R H, Nijman S M, Wittinghofer A and Bos J L (1998) Epac Is a Rap1 Guanine-Nucleotide-Exchange Factor Directly Activated by Cyclic AMP. *Nature* 396:474-477.

Dostmann WR, Taylor S S, Genieser H G, Jastorff B, Døskeland S O and Ogried D (1990) Probing the Cyclic Nucleotide Binding Sites of cAMP-Dependent Protein Kinases I and II With Analogs of Adenosine 3',5'-Cyclic Phosphorothioates. *J Biol Chem* 265:10484-10491.

Enserink JM, Christensen A E, de Rooij J, van Triest M, Schwede F, Genieser H G, Døskeland S O, Blank J L and Bos J L (2002) A Novel Epac-Specific cAMP Analogue Demonstrates Independent Regulation of Rap1 and ERK. *Nature Cell Biology* 4:901-906.

Enyeart JA, Danthi S J and Enyeart J J (2003) Corticotropin Induces the Expression of TREK-1 mRNA and K<sup>+</sup> Current in Adrenocortical Cells. *Mol Pharmacol* 64:132-142.

Enyeart JA and Enyeart J J (2009) Metabolites of an Epac-Selective cAMP Analog Induce Cortisol Synthesis by Adrenocortical Cells Through a cAMP-Independent Pathway. *PLoS One* 4:e6088.

Enyeart JA, Xu L and Enyeart J J (2000) A Bovine Adrenocortical Kv1.4 K<sup>+</sup> Channel Whose Expression Is Potently Inhibited by ACTH. *J Biol Chem* 275:34640-34649.

Enyeart JJ, Gomora J C, Xu L and Enyeart J A (1997) Adenosine Triphosphate Activates a Noninactivating K<sup>+</sup> Current in Adrenal Cortical Cells Through Nonhydrolytic Binding. *J Gen Physiol* , 110:679-692.

**MOL #61861**

Enyeart JJ, Mlinar B and Enyeart J A (1993) T-Type  $\text{Ca}^{2+}$  Channels Are Required for ACTH-Stimulated Cortisol Synthesis by Bovine Adrenal Zona Fasciculata Cells. *Mol Endo* 7:1031-1040.

Enyeart JJ, Mlinar B and Enyeart J A (1996) Adrenocorticotrophic Hormone and cAMP Inhibit Noninactivating  $\text{K}^{+}$  Current in Adrenocortical Cells by an A-Kinase-Independent Mechanism Requiring ATP Hydrolysis. *J Gen Physiol* 108:251-264.

Enyeart JJ, Xu L, Danthi S and Enyeart J A (2002) An ACTH- and ATP-Regulated Background  $\text{K}^{+}$  Channel in Adrenocortical Cells Is TREK-1. *J Biol Chem* 277:49186-49199.

Fujita T, Meguro T, Fukuyama R, Nakamuta H and Koida M (2002) New Signaling Pathway for Parathyroid Hormone and Cyclic AMP Action on Extracellular-Regulated Kinase and Cell Proliferation in Bone Cells. Checkpoint of Modulation by Cyclic AMP. *J Biol Chem* 277:22191-22200.

Grahame-Smith DG, Butcher R W, Ney R L and Sutherland E W (1967) Adenosine 3',5'-Monophosphate As the Intracellular Mediator of the Action of Adrenocorticotrophic Hormone on the Adrenal Cortex. *J Biol Chem* 242:5535-5541.

Hamill OP, Marty A, Neher E, Sakmann B and Sigworth F J (1981) Improved Patch Clamp Techniques for High Resolution Current Recording From Cells and Cell-Free Membrane Patches. *Pflügers Arch* 391:85-100.

Haynes RC and Berthet L (1957) Studies on the Mechanism of Action of the Adrenocorticotrophic Hormone. *J Biol Chem* 225:115-124.

Holz GG, Chepurny O G and Schwede F (2008) Epac-Selective cAMP Analogs: New Tools With Which to Evaluate the Signal Transduction Properties of cAMP-Regulated Guanine Nucleotide Exchange Factors. *Cell Signalling* 20:10-20.

Holz GG, Kang G, Harbeck M, Roe M W and Chepurny O G (2006) Cell Physiology of cAMP Sensor Epac. *The Journal of Physiology* 577:5-15.

Iocovelli L, Capobianco L, Salvatore L, Sallese M, D'Ancona G M and De Blasi A (2001) Thyrotropin Activates Mitogen-Activated Protein Kinase Pathway in FRTL-5 by a cAMP-Dependent Protein Kinase A-Independent Mechanism. *Mol Pharmacol* 60:924-933.

Ivins JK, Parry M K and Long D A (2004) A Novel cAMP-Dependent Pathway Activates Neuronal Integrin Function in Retinal Neurons. *J Neurosci* 24:1212-1216.

John ME, John M C, Boggaram V, Simpson E R and Waterman M R (1986) Transcriptional Regulation of Steroid Hydroxylase Genes by Corticotropin. *Proc Nat Acad Sci USA* 83:4715-4719.

Kagawa N and Waterman M R (1990) cAMP-Dependent Transcription of the Human CYP21B (P-450C21) Gene Requires a Cis-Regulatory Element Distinct From the Consensus cAMP-Regulatory Element. *J Biol Chem* 265:11299-11305.

Kawasaki H, Springett G M, Mochizuki N, Toki S, Nakaya M, Matsuda M, Housman D E and Graybiel A M (1998) A Family of cAMP-Binding Proteins That Directly Activate Rap1. *Science* 282:2275-2279.

Laxman S, Riechers A, Sadilek M, Schwede F and Beavo J A (2006) Hydrolysis Products of cAMP Analogs Cause Transformation of Trypanosoma Brucei From Slender to Stumpy-Like Forms. *Proc Natl Acad Sci U S A* 103:19194-19199.

Liu H, Enyeart J A and Enyeart J J (2009) N6-Substituted cAMP Analogs Inhibit BTREK-1 K<sup>+</sup> Channels and Stimulate Cortisol Secretion by a PKA-Independent Mechanism. *Mol Pharmacol* 76(6):1290-301.

Liu H, Enyeart J A and Enyeart J J (2008) ACTH Inhibits BTREK-1 K<sup>+</sup> Channels Through Multiple cAMP-Dependent Signaling Pathways. *J Gen Physiol* 132:279-294.

Lund J, Ahlgren R, Wu D H, Kagimoto M, Simpson E R and Waterman M R (1990) Transcriptional Regulation of the Bovine CYP17 (P-450(17)Alpha) Gene. Identification of Two cAMP Regulatory Regions Lacking the Consensus cAMP-Responsive Element (CRE). *J Biol Chem* 265:3304-3312.

Mlinar B, Biagi B A and Enyeart J J (1993) A Novel K<sup>+</sup> Current Inhibited by ACTH and Angiotensin II in Adrenal Cortical Cells. *J Biol Chem* 268,No.12:8640-8644.

Mlinar B and Enyeart J J (1993) Voltage-Gated Transient Currents in Bovine Adrenal Fasciculata Cells II: A-Type K<sup>+</sup> Current. *J Gen Physiol* 102:239-255.

Moyle WR, Kong Y C and Ramachandran J (1973) Steroidogenesis and Cyclic Adenosine 3',5'-Monophosphate Accumulation in Rat Adrenal Cells. Divergent Effects of Adrenocorticotropin and Its O-Nitrophenyl Sulfenyl Derivative. *J Biol Chem* 248:2409-2417.

Parker KL and Schimmer B P (1995) Transcriptional Regulation of the Genes Encoding the Cytochrome P-450 Steroid Hydroxylases. *Vit and Horm* 51:339-370.

Payne AH and Hales D B (2004) Overview of Steroidogenic Enzymes in the Pathway From Cholesterol to Active Steroid Hormones. *Endocrine Reviews* 25:947-970.

Penhoat A, Jaillard C and Saez J M (1989) Corticotropin Positively Regulates Its Own Receptors and cAMP Response in Cultured Bovine Adrenal Cells. *Proc Natl Acad Sci U S A* 86:4978-4981.

Poppe H, Rybalkin S D, Rehmann H, Hinds T R, Tang X B, Christensen A E, Schwede F, Genieser H G, Bos J L, Doskeland S O, Beavo J A and Butt E (2008) Cyclic Nucleotide Analogs As Probes of Signaling Pathways. *Nature Methods* 5:277-278.

Price NC and Stevens L (1999) *Fundamentals of Enzymology: The Cell and Molecular Biology of Catalytic Proteins*. Oxford University Press, New York.

Raikhinstein M, Zohar M and Hanukoglu I (1994) CDNA Cloning and Sequence Analysis of the Bovine Adrenocorticotropic Hormone (ACTH) Receptor. *Biochim Biophys Acta* 1220:329-332.

Richardson MC and Schulster D (1973) The Role of Protein Kinase Activation in the Control of Steroidogenesis by Adrenocorticotrophic Hormone in the Adrenal Cortex. *Biochem J* 136:993-998.

Sala GB, Hayashi K, Catt K J and Dufau M L (1979) Adrenocorticotropin Action in Isolated Adrenal Cells. The Intermediate Role of Cyclic AMP in Stimulation of Corticosterone Synthesis. *J Biol Chem* 254:3861-3865.

Simpson ER and Waterman M R (1988) Regulation of the Synthesis of Steroidogenic Enzymes in Adrenal Cortical Cells by ACTH. *Ann Rev Physiol* 50:427-440.

Stork PJ and Schmitt J M (2002) Crosstalk Between cAMP and MAP Kinase Signaling in the Regulation of Cell Proliferation. *Trends Cell Biol* 12:258-266.

Waterman MR (1994) Biochemical Diversity of cAMP-Dependent Transcription of Steroid Hydroxylase Genes in the Adrenal Cortex. *J Biol Chem* 269:27783-27786.

Waterman MR and Simpson E R (1988) Regulation of Steroid Hydroxylase Gene Expression Is Multifactorial in Nature. *Recent Progress in Hormone Research* 45:532-566.

Yamazaki T, Kimoto T, Higuchi K, Ohta Y, Kawato S and Kominami S (1998) Calcium Ion As a Second Messenger for O-Nitrophenylsulfenyl-Adrenocorticotropin (NPS-ACTH) and ACTH in Bovine Adrenal Steroidogenesis. *Endocrinology* 139:4765-4771.

**MOL #61861**

Yamazaki T, Kawasaki H, Takamasa A, Yoshitomi T and Kominami S (2006)  $\text{Ca}^{2+}$  Signal Stimulates the Expression of Steroidogenic Acute Regulatory Protein and Steroidogenesis in Bovine Adrenal Fasciculata-Reticularis Cells. *Life Sci* 78:2923-2930.

**FOOTNOTE:**

This work was supported by the National Institutes of Health [Grant R01-DK47875]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Diabetes and Digestive and Kidney Diseases or the National Institutes of Health.



## FIGURE LEGENDS

### *Figure 1. Long-term effect of ACTH and 8-Br-cAMP on expression of bTREK-1 current.*

AZF cells were used for patch clamp experiments 1–48 h after plating. AZF cells were either plated in media containing no further addition (control - A,B), 2 nM ACTH (C), or 8-Br-cAMP (D). Whole cell  $K^+$  currents were recorded in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution (see Methods section 2.4). A-D) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1 amplitudes with (open circles) and without (closed circles) depolarizing pulses. Times indicated on traces correspond to those on graph at right. E) Summary of experiments as in A-D. Bars represent bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations at 1 and 48 h in control media and after 48 h exposure to ACTH (2nM) or 8-Br-cAMP (300  $\mu$ M), as indicated.

*Figure 2. 6-Bnz-cAMP induces expression of bTREK-1 current.* AZF cells were cultured in media containing no further addition (control) (A), 30  $\mu$ M 6-Bnz-cAMP (B) or 200  $\mu$ M 6-Bnz-cAMP (C). Whole cell  $K^+$  currents were recorded from AZF cells in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution (see Methods section 2.4). A-C) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1 amplitudes with (open circles) and without (closed circles) depolarizing pulses. Times indicated on traces correspond to those on graph at right. Cells were either untreated (A), or treated with 30  $\mu$ M (B) or 200  $\mu$ M (C) 6-Bnz-cAMP for 48 h before

recording. D) Summary of experiments as in A-C. Bars represent bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations after 48-96 h exposure to 6-Bnz-cAMP (30 or 300  $\mu$ M), as indicated.

**Figure 3. Effect of Epac2-selective cAMP analogs on bTREK-1 current expression.** Whole cell  $K^+$  currents were recorded from AZF cells in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution (see Methods section 2.4). A,B) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1 amplitudes with (open circles) and without (closed circles) depolarizing pulses. Time indicated on traces corresponds to those plotted on graph at right. AZF cells with treated with 30  $\mu$ M 8CPT-2'-OMe-cAMP (A) or 30  $\mu$ M Sp-8CPT-2'-OMe-cAMP (B) for 48 h before recording. C) Summary of experiments as in (A) and (B): Bars specify bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations after 24-96 h exposure to 8CPT-2'-OMe-cAMP (30  $\mu$ M) or Sp-8CPT-2'-OMe-cAMP (30  $\mu$ M) as indicated.

**Figure 4. Long-term effect of Sp-8CPT-2'-OMe-cAMP and 8CPT-2'-OMe-Ado on expression of bTREK-1 current.** AZF cells were cultured for 48 h in media containing no further addition (control), Sp-8CPT-2'-OMe-cAMP (100  $\mu$ M) (A) or 8CPT-2'-OMe-Ado (30  $\mu$ M) (B). Whole cell  $K^+$  currents were recorded from AZF cells in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution (see Methods section 2.4). A,B) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1

amplitudes with (open circles) and without (closed circles) depolarizing pulses. Times indicated on traces correspond to those plotted on graph at right. C) Summary of experiments as in (A) and (B): Bars represent bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations after 48 or 72 h exposure to Sp-8CPT-2'-OMe-cAMP (100  $\mu$ M) or 8CPT-2'-OMe-Ado (30  $\mu$ M) as indicated.

*Figure 5. Structures of 8CPT-2'-OMe-cAMP, 8CPT-cAMP and Metabolites.* Chemical structures of 8CPT-2'-OMe-cAMP, 8CPT-cAMP and their metabolites.

*Figure 6. Effects of ACTH, 8-Br-cAMP, and 6-Bnz-cAMP on bTREK-1 Gene Expression.*

AZF cells were incubated either without (control), or with ACTH, 8-Br-cAMP, 6-Bnz-cAMP, or 6-Bnz-cAMP + H-89, as indicated. Total RNA was isolated as described in the Methods. Membranes were hybridized with specific probe for bTREK-1. 18S rRNA bands from representative gels are shown as evidence of even loading. bTREK-1 mRNA levels are expressed as percent of the 3.6 KB control band value. A) Effect of time in culture on bTREK-1 mRNA. AZF cells were plated and total RNA isolated after 5, 24 or 48 h in culture, as indicated. B) Effect of ACTH and 8-Br-cAMP on bTREK-1 mRNA expression. AZF cells were cultured overnight before either no addition (control, white bar) or addition of ACTH (2 nM, black bar) or 8-Br-cAMP (30  $\mu$ M, light grey or 300  $\mu$ M, dark grey bar) for 48 hr before isolating total RNA. Asterisk designates statistically significant difference between control and treated cells (\* $P$  < 0.02). C) Effect of 6-Bnz-cAMP and H-89 on bTREK-1 mRNA. AZF cells were plated and cultured overnight before either no addition (control, white bar) or addition of H-89 (10  $\mu$ M, striped bar), 6-Bnz-cAMP (200  $\mu$ M, grey bar), or

6-Bnz-cAMP + H-89 (grey striped bar) for 24 h before isolating total RNA. Cells were pre-incubated with H-89 (10  $\mu$ M) for 1 h prior to 6-Bnz-cAMP (200  $\mu$ M) addition. (\* $P$  < 0.03).

**Figure 7. Effects of 8CPT-2'-OMe-cAMP and its metabolites on bTREK-1 Gene Expression.**

AZF cells were cultured overnight then incubated either without (control), or with 8CPT-2'-OMe-cAMP, Sp-8CPT-2'-OMe-cAMP, 8CPT-2'-OMe-5' AMP, or 8CPT-2'-OMe-Ado as indicated. Total RNA was isolated as described in the Methods. Membranes were hybridized with a specific probe for bTREK-1; bTREK-1 mRNA levels are expressed as percent of the 3.6 KB control band value. 18S rRNA bands from representative gels are shown as evidence of even loading. A) Concentration-dependent effect of 8CPT-2'-OMe-cAMP on bTREK-1 mRNA expression. AZF cells were untreated (control, white bar) or treated with 8CPT-2'-OMe-cAMP (1-50  $\mu$ M, grey bars) for 48 hr before isolating total RNA. B) Time-dependent effect of 8CPT-2'-OMe-cAMP on bTREK-1 mRNA expression. AZF cells were either untreated (control, white bars) or treated with 30  $\mu$ M 8CPT-2'-OMe-cAMP (grey bars) for 1-48 hr before isolating total RNA. C) Effect of 8CPT-2'-OMe-cAMP, Sp-8CPT-2'-OMe-cAMP, and 8-Br-cAMP on bTREK-1 mRNA expression. AZF cells were either untreated (control, white bar), or treated with 30  $\mu$ M 8CPT-2'-OMe-cAMP (grey bar, left panel), 30  $\mu$ M Sp-8CPT-2'-OMe-cAMP (grey striped bar, left panel), 100  $\mu$ M Sp-8CPT-2'-OMe-cAMP (grey striped bar, right panel), or 300  $\mu$ M 8-Br-cAMP (black bar, right panel) for 48 hr before isolating total RNA (\* $P$  < 0.005). D) Effect of metabolites of 8CPT-2'-OMe-cAMP on induction of bTREK-1 mRNA. AZF cells were treated with either 8CPT-2'-OMe-5' AMP (0.1-100  $\mu$ M, grey dotted bars), 8CPT-2'-OMe-Ado (0.1 to 50  $\mu$ M, dark grey bars) or adenosine (50  $\mu$ M, light grey bar) for 48 h before isolating total RNA.


**Figure 8. Long-term effect of 8CPT-cAMP and Sp-8CPT-cAMP on expression of bTREK-1 current.** AZF cells were cultured overnight then incubated either without (control), or with 8CPT-cAMP (30  $\mu$ M) or Sp-8CPT-cAMP (30 or 300  $\mu$ M), as indicated. Whole cell  $K^+$  currents were recorded from AZF cells in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution as described in the Methods section 2.4. A-C) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1 amplitudes with (open circles) and without (closed circles) depolarizing pulses. Time indicated on traces corresponds to those plotted on graph at right. Cells were treated with 30  $\mu$ M 8CPT-cAMP (A), 30  $\mu$ M Sp-8CPT-2'-OMe-cAMP (B) or 300  $\mu$ M Sp-8CPT-cAMP (C) for 48 h before recording. C) Summary of experiments as in A-C. Bars represent bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations after 48 h exposure to 8CPT-cAMP (30  $\mu$ M) or Sp-8CPT-2'-OMe-cAMP (30 or 300  $\mu$ M) as indicated.


**Figure 9. Metabolites of 8CPT-cAMP induce bTREK-1 current.** AZF cells were cultured overnight then incubated either without (control), or with 8CPT-Ado (30  $\mu$ M) or 8CPT-Ade (30  $\mu$ M) as indicated. Whole cell  $K^+$  currents were recorded from AZF cells in response to voltage steps to +20 mV applied from -80 mV at 30 s intervals with or without depolarizing prepulses to -20 mV. Pipettes contained standard solution as described in the Methods section 2.4. A, B) Representative  $K^+$  current traces recorded with (right traces) and without (left traces) depolarizing prepulses, and corresponding plot of bTREK-1 amplitudes with (open circles) and without (closed circles) depolarizing pulses. Time indicated on traces corresponds to those plotted on graph at right. Cells were treated with 30  $\mu$ M 8CPT-Ado (A) or 30  $\mu$ M 8CPT-Ade for 48 h before recording. C)

Summary of experiments as in A and B. Bars represent bTREK-1 current density expressed as mean  $\pm$  SEM of indicated number of determinations after 48 h exposure to 8CPT-Ado or 8CPT-Ade as indicated.


**Figure 10. Effects of 8CPT-cAMP, Sp-cAMP and 8CPT-Ade on expression of bTREK-1**


**mRNA.** AZF cells were cultured overnight then incubated either without (control), or with 8CPT-cAMP, Sp-cAMP, 8CPT-Ade, or ACTH as indicated. Total RNA was isolated as described in the Methods. Membranes were hybridized with a specific probe for bTREK-1; bTREK-1 mRNA levels are expressed as percent of the 3.6 KB control band value. 18S rRNA bands from representative gels are shown as evidence of even loading. A) 8CPT-cAMP but not Sp-8CPT-cAMP induces bTREK-1 mRNA. AZF cells were either untreated (control, white bar) or treated with 30  $\mu$ M 8CPT-cAMP (dark grey bar) or 30  $\mu$ M Sp-8CPT-cAMP (dark grey striped bar) for 48 hr before isolating total RNA (\* $P$  < 0.02). B) Concentration-dependent effect of 8CPT-Ade on bTREK-1 mRNA. AZF cells were either untreated (control, white bar) or treated with 8CPT-Ade (1-50  $\mu$ M, grey bars) for 48 hr before isolating total RNA (\* $P$  < 0.001). C) Time-dependent effect of 8CPT-Ade and ACTH on bTREK-1 mRNA expression. AZF cells were either untreated (control, white bar) or treated with 30  $\mu$ M 8CPT-cAMP (grey bars) or 2 nM ACTH (black bars) for 5, 24, and 48 hr before isolating total RNA (\* $P$  < 0.001).

















**8-(4-Chlorophenylthio)-2'-O-methyladenosine-3',5'-cyclic monophosphate (8CPT-2'-OMe-cAMP)**




**8-(4-Chlorophenylthio)adenosine-3',5'-cyclic monophosphate (8CPT-cAMP)**




**8-(4-Chlorophenylthio)-2'-O-methyladenosine-5'-O-monophosphate (8CPT-2'-OMe-5'AMP)**




**8-(4-Chlorophenylthio)adenosine-5'-O-monophosphate (8CPT-5'AMP)**




**8-(4-Chlorophenylthio)-2'-O-methyladenosine (8CPT-2'-OMe-Ado)**




**8-(4-Chlorophenylthio)adenosine (8CPT-Ado)**




**8-(4-Chlorophenylthio)adenine (8CPT-Ade)**







A




B



C



D



**FIGURE 9**





FIGURE 10





**Judith A. Enyeart, Haiyan Liu, and John J. Enyeart -  
cAMP Analogs and Their Metabolites Enhance TREK-1 mRNA and K<sup>+</sup>  
Current Expression in Adrenocortical Cells  
*MOLECULAR PHARMACOLOGY***

**SUPPLEMENTAL TABLE 1**


DRUG	PKA	Epac
ACTH	+	+
8-Br-cAMP	+	+
6-Bnz-cAMP	+	-
8CPT-2'-OMe-cAMP	-	+
Sp-8CPT-2'-OMe-cAMP	-	+
8CPT-2'-OMe-5'AMP	-	
8CPT-2'-OMe-adenosine	-	
8CPT-cAMP	+	+
Sp-8CPT-cAMP	+	+
8CPT-adenosine		
8CPT-adenine	-	

List of drugs used in the present study and indication of whether specific drug activates (+) or does not activate (-) PKA or Epac. Blanks indicate that PKA and/or Epac activity of drug has not been determined.

Judith A. Enyeart, Haiyan Liu, and John J. Enyeart

cAMP Analogs and Their Metabolites Enhance TREK-1 mRNA and K<sup>+</sup> Current Expression in Adrenocortical Cells  
MOLECULAR PHARMACOLOGY

### SUPPLEMENTAL FIGURE 1



**Supplemental Figure 1. Effects of 8-Br-cAMP and 6-Bnz-cAMP on cortisol secretion.** For cortisol determination, bovine AZF cells were plated as described in the Methods. After 24 h, medium was aspirated and replaced with the same medium either without (control) or with 8-Br-cAMP or 6-Bnz-cAMP as indicated. Cortisol was measured in duplicate from duplicate experimental media samples using a Cortisol EIA (DSL-10-200) from Diagnostic Systems Laboratories (Webster, TX), according to the manufacturer's directions. If necessary, media samples were diluted using DMEM/F12+. A) Cortisol concentration from media was determined after 24 h with no addition or after addition of 8-Br-cAMP (2-500 μM, grey bars). B) Cortisol concentration from AZF cell media was determined after 24 h with no addition or after addition of 6-Bnz-cAMP (30-300 μM, black bars).