

# MOLECULAR PHARMACOLOGY

May 2018

Volume 93

Number 5

molpharm.aspetjournals.org

ISSN 1521-0111

## ACCELERATED COMMUNICATION

High Constitutive Activity Accounts for the Combination of Enhanced Direct Activation and Reduced Potentiation in Mutated GABA<sub>A</sub> Receptors

*Allison L. Germann, Daniel J. Shin, Christina R. Kuhrau, Alexander D. Johnson, Alex S. Evers, and Gustav Akk*

468

## ARTICLES

Ethanol Reversal of Oxycodone Tolerance in Dorsal Root Ganglia Neurons

*Joanna C. Jacob, Kensuke Sakakibara, Ryan A. Mischel, Graeme Henderson, William L. Dewey, and Hamid I. Akbarali*

417

□ Hemodynamic Effects of Glutathione-Liganded Binuclear Dinitrosyl Iron Complex: Evidence for Nitroxyl Generation and Modulation by Plasma Albumin

*Taiming Liu, Meijuan Zhang, Michael H. Terry, Hobe Schroeder, Sean M. Wilson, Gordon G. Power, Qian Li, Trent E. Tipple, Dan Borchardt, and Arlin B. Blood*

427

□ Retinal-chitosan Conjugates Effectively Deliver Active Chromophores to Retinal Photoreceptor Cells in Blind Mice and Dogs

*Songqi Gao, Shirin Kahremany, Jianye Zhang, Beata Jastrzebska, Janice Querubin, Simon M. Petersen-Jones, and Krzysztof Palczewski*

438

□ Properties of Triheteromeric N-Methyl-D-Aspartate Receptors Containing Two Distinct GluN1 Isoforms

*Feng Yi, Linda G. Zachariassen, Katherine N. Dorsett, and Kasper B. Hansen*

453

□ Gedunin- and Khivorin-Derivatives Are Small-Molecule Partial Agonists for Adhesion G Protein-Coupled Receptors GPR56/ADGRG1 and GPR114/ADGRG5

*Hannah M. Stoveken, Scott D. Larsen, Alan V. Smrcka, and Gregory G. Tall*

477

□ CYP26C1 Is a Hydroxylase of Multiple Active Retinoids and Interacts with Cellular Retinoic Acid Binding Proteins

*Guo Zhong, David Ortiz, Alex Zelter, Abhinav Nath, and Nina Isoherranen*

489

□ “Selective” Class C G Protein-Coupled Receptor Modulators Are Neutral or Biased mGlu<sub>5</sub> Allosteric Ligands

*Shane D. Hellyer, Sabine Albold, Taide Wang, Amy N. Y. Chen, Lauren T. May, Katie Leach, and Karen J. Gregory*

504

The Novel C-terminal Truncated 90-kDa Isoform of Topoisomerase II $\alpha$  (TOP2 $\alpha$ /90) Is a Determinant of Etoposide Resistance in K562 Leukemia Cells via Heterodimerization with the TOP2 $\alpha$ /170 Isoform

*Ragu Kanagasabai, Soumendrakrishna Karmahapatra, Corey A. Kientz, Yang Yu, Victor A. Hernandez, Evan E. Kania, Jack C. Yalowich, and Terry S. Elton*

515

Differential Pharmacology and Binding of mGlu<sub>2</sub> Receptor Allosteric Modulators

*Daniel E. O'Brien, Douglas M. Shaw, Hyekyung P. Cho, Alan J. Cross, Steven S. Wesolowski, Andrew S. Felts, Jonas Bergare, Charles S. Elmore, Craig W. Lindsley, Colleen M. Niswender, and P. Jeffrey Conn*

526

- ❑ Cooperative Regulation of Intestinal UDP-Glucuronosyltransferases 1A8, -1A9, and 1A10 by CDX2 and HNF4 $\alpha$  Is Mediated by a Novel Composite Regulatory Element  
*Nurul Mubarakah, Julie-Ann Hulin, Peter I. Mackenzie, Ross A. McKinnon, Alex Z. Haines, Dong Gui Hu, and Robyn Meech* 541
- ❑ Mapping the Allosteric Action of Antagonists A740003 and A438079 Reveals a Role for the Left Flipper in Ligand Sensitivity at P2X7 Receptors  
*Rebecca C. Allsopp, Sudad Dayl, Anfal Bin Dayel, Ralf Schmid, and Richard J. Evans* 553
- ERRATUM**
- Correction to “Investigation into the Mechanism of Homo- and Heterodimerization of Angiotensin-Converting Enzyme” 416

❑ Supplemental material is available online at <http://molpharm.aspetjournals.org>.

*About the cover:* Effects of point mutants in the allosteric binding pocket on sensitivity to the antagonist A438079. See the article by Evans et al. ([dx.doi.org/10.1124/mol.117.111021](https://doi.org/10.1124/mol.117.111021)).