





















Figure S1. Comparison of locomotion activity of mice with same genotype between room temperature and 33°C. A-D, Locomotion distance (A) and activity time (C) of *Tph2* mice over 24 hrs. Mean locomotion distance was significantly increased at 33°C during both dark (p=0.0145) and light phase (p=0.0043) with wt mice and only during light phase (p=0.0002) with *Tph2*<sup>-/-</sup> mice (B). Mean activity time was significantly decreased at 33°C during dark phase (p=0.0384) with *Tph2*<sup>-/-</sup> mice (D). E-H, Locomotion distance (E) and activity time (G) of *Lmx1b* mice over 24 hours. Mean locomotion distance was increased at 33°C during both dark (p=0.0488) and light phase (p=0.0026) with *Lmx1b*<sup>ff</sup> mice (F). Mean activity time did not differ significantly between RT and 33°C with both *Lmx1b*<sup>ff</sup> and *Lmx1b*<sup>ff/p</sup> mice (H). Data are mean±sem. Unpaired t-test with Welch's correction for *Tph2* mice and Mann-Whitney test for *Lmx1b* mice (n=30 for *Tph2*<sup>+/+</sup>; n=29 for *Tph2*<sup>-/-</sup>; n=15 for *Lmx1b*<sup>ff</sup>; n=13 for *Lmx1b*<sup>ff/p</sup>).