MINIREVIEW

Small-Molecule Inhibitors Targeting Protein SUMOylation as Novel Anticancer Compounds

Yanfang Yang, Zijing Xia, Xixi Wang, Xinyu Zhao, Zenghua Sheng, Yang Ye, Gu He, Liangxue Zhou, Hongxia Zhu, Ningzhi Xu, and Shufang Liang

State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, and Collaborative Innovation Center for Biotherapy, Chengdu (Y.Ya., Z.X., X.W., X.Z., Z.S., Y.Ye., G.H., L.Z., N.X., S.L.); Departments of Nephrology (Z.X.) and Neurosurgery (L.Z.), West China Hospital, Sichuan University, Chengdu; and Laboratory of Cell and Molecular Biology, and State Key Laboratory of Molecular Oncology, Cancer Institute and Cancer Hospital, Chinese Academy of Medical Sciences, Beijing (H.Z., N.X.), People's Republic of China

Received February 26, 2018; accepted May 16, 2018

ABSTRACT

SUMOylation, one of post-translational modifications, is covalently modified on lysine residues of a target protein through an enzymatic cascade reaction similar to protein ubiquitination. Along with identification of many SUMOylated proteins, protein SUMOylation has been proven to regulate multiple biologic activities including transcription, cell cycle, DNA repair, and innate immunity. The dysregulation of protein SUMOylation and deSUMOylation modification is linked with carcinogenesis and tumor progression. The SUMOylation-associated enzymes are usually elevated in various cancers, which function as cancer biomarkers to relate to poor outcomes for patients. Considering the significance of protein SUMOylation in regulating diverse

biologic functions in cancer progression, numerous small-molecule inhibitors targeting protein SUMOylation pathway are developed as potentially clinical anticancer therapeutics. Here, we systematically summarize the latest progresses of associations of small ubiquitin-like modifier (SUMO) enzymes with cancers and small-molecular inhibitors against human cancers by targeting SUMOylation enzymes. We also compared the pros and cons of several special anticancer inhibitors targeting SUMO pathway. As more efforts are invested in this field, small-molecule inhibitors targeting the SUMOylation modification pathway are promising for development into novel anticancer drugs.

Introduction

The small ubiquitin-like modifier (SUMO) proteins, approximately with 11 kDa, covalently modify lysine (Lys) amino acid of a target substrate protein. Four major SUMO paralogues, including SUMO1, SUMO2, SUMO3, and SUMO4, are expressed in mammalian cells (Enserink, 2015). Most SUMOylation sites of a target protein reside in a consensus motif with the ψ KxE or ψ KxE/D sequence (ψ is an aliphatic residue, K is the target Lys, E/D is Asp or Glu, and x is any amino acid) (Yang et al., 2017). Except binding in consensus motif, SUMOs can noncovalently

interact with proteins through targeting specific SUMO-interacting motifs (Hecker et al., 2006).

SUMOylation regulates multiple biologic processes including transcription, cell cycle, DNA repair, and innate immunity (Seeler and Dejean, 2003; Bettermann et al., 2012). A multistep enzymatic cascade reaction is involved in the biochemical process of protein SUMOylation (Fig. 1). First, four SUMO paralogues are synthesized as preproteins that are cleaved by sentrin-specific proteases (SENPs) to expose a diglycine (GG) motif on the carboxy-terminus. After that, an ATP-requiring activation, mediated by a heterodimeric SUMO-activating enzyme E1 (SUMO E1), generates a SUMO-SUMO E1 thioester. Then SUMO is transferred to the SUMO-conjugating enzyme E2 (SUMO E2) to form a thioester again. Next, a SUMO ligase E3 (SUMO E3) stabilizes the interaction between SUMO E2 and the substrate, and brings about an isopeptide bond between the SUMO C-terminus and a Lys within the target

This work was supported by the National 863 High Tech Foundation [Grant 2014AA020608], the National Key Basic Research Program of China [Grants 2013CB911303 and 2011CB910703], the National Natural Science Foundation of China [Grants 31470810 and 31071235], the Science and Technology Department of Sichuan Province [Grant 2017JY0232], and the Health and Family Planning Commission of Sichuan Province [Grant 17ZD045].
<https://doi.org/10.1124/mol.118.112300>

ABBREVIATIONS: HCC, hepatocellular carcinoma; ML-792, [(1*R*,2*S*,4*R*)-4-[[5-[1-[(3-bromophenyl)methyl]pyrazole-3-carbonyl]pyrimidin-4-yl]amino]-2-hydroxycyclopentyl]methyl sulfamate; PTM, protein post-translational modification; SAE1/2, SUMO-activating enzyme subunit 1/2; SENP(s), sentrin-specific proteases; SUMO, small ubiquitin-like modifier; SUMO E1, SUMO-activating enzyme E1; SUMO E2, SUMO-conjugating enzyme E2; SUMO E3, SUMO ligase E3; Ubc9, ubiquitin-conjugating enzyme 9.

Fig. 1. Small-molecule inhibitors that target the protein SUMOylation process. Protein SUMOylation is a multistep enzymatic cascade reaction. First, all SUMO paralogues are synthesized as preproteins that are cleaved by SENPs to expose a carboxy-terminal diglycine (GG) motif. An ATP-requiring activation step by the heterodimeric SUMO E1 generates a SUMO-SUMO E1 thioester. SUMO is then transferred to the SUMO E2, again forming a thioester. This last step usually requires a SUMO E3 to form an isopeptide bond between the SUMO C terminus and a Lys within the target protein. SUMO E2 promotes SUMOs onto Lys residues of target substrate even in the absence of an SUMO E3. The small-molecule inhibitors targeting SUMOylation-associated enzymes can block the subsequent SUMO modification biochemical process. (A–C) Chemical structure of three inhibitors (ML-792, spectomycin B1, and triptolide).

protein (Yang et al., 2017). Generally, SENPs, SUMO E1, SUMO E2, and SUMO E3 are four key enzymes to in turn perform cleaving, activating, transferring, and ligating of SUMO to Lys residue of a substrate protein.

Protein SUMOylation is a dynamic regulation in physiologic conditions, and the imbalance of SUMOylation and deSUMOylation associates with the occurrence and progression of cancer and other diseases (Fig. 2) (Liang et al., 2017; Seeler and Dejean, 2017; Yang et al., 2017). To date, due to the importance of SUMOylation in regulating diverse biologic systems in carcinogenesis, several novel small-molecule inhibitors targeting SUMOylation have been developed as promising anticancer compounds. In this review, we summarize the latest research progresses of small-molecule inhibitors by targeting SUMOylation-associated enzymes as highly potent anticancer compounds.

Small-Molecule Inhibitors Targeting Protein SUMOylation-Associated Enzymes

The expression of SENPs, SUMO E1, SUMO E2, and SUMO E3 is usually increased in numerous cancers (Coppola et al.,

2009; Bellail et al., 2014; Seeler and Dejean, 2017). The SUMO pathway components are involved in various tumors (Coppola et al., 2009; Bellail et al., 2014; Liu et al., 2015; Seeler and Dejean, 2017). The SUMO pathway components are involved in various human cancers (Table 1), so small-molecule inhibitors targeting protein SUMOylation-associated enzymes exhibit promising antitumor activities.

SENP and Its Anticancer Inhibitors. SUMOylation is a reversible and dynamic process between protein SUMOylation and deSUMOylation for normal cellular activities (Yang et al., 2017). SENPs, as cysteine proteases, cleave the precursor or inactive form of SUMO at the C terminus via its hydrolase activity to expose two GG residues. Alternatively, SENPs can catalyze the deconjugation of SUMO molecules from a target protein. SENPs include six members, including SENP1, SENP2, SENP3, SENP5, SENP6, and SENP7. Among these SENPs, SENP1 is widely increased in various cancers (Table 1) (Zuo and Cheng, 2009; Xu et al., 2011). It plays an important role in cancer occurrence (Bawa-Khalfe et al., 2007), development (Bawa-Khalfe et al., 2010), and metastasis (Wang et al., 2013). For instance, SENP1 enhances vascular endothelial growth factor production by regulating the stability of hypoxia-inducible factor 1 (Xu et al., 2010), which is

In addition to SENP1, SENP2 has been confirmed to be a molecular target for chemical drugs that effectively enhance SUMO conjugation and suppress SENP2 activity. For example, compound 1, 2, 5-oxadiazole, has potential for development into a novel SENP2-targeting therapeutic agent against various diseases (Kumar et al., 2014). In addition, cytoprotective molecules are being developed as novel stroke therapies

TABLE 1
SUMOylation-associated enzymes in cancer cells

Enzyme	Cancer	Expression	Characteristics	Mechanisms	Study
SENPs	1) Prostate	Up-regulated	Involve in prostate cancer pathogenesis	Associate with androgen receptor signaling and hypoxia-HIF-1 α signaling	Zuo and Cheng (2009)
	2) Colorectal	Up-regulated	Play a role in cell cycle regulation	Up-regulate expression of CDK inhibitors (such as p16, p19, p21, and p27)	Xu et al. (2011)
	3) Pancreatic	Up-regulated	Associate with lymph node metastasis and TNM stage	Associate with MMP-9 which is pivotal for PDAC cell growth and migration	Ma et al. (2014)
	4) Neuroblastoma	Up-regulated	Promote cells migration and invasion	Regulate the expression of CDH1, MMP-9, and MMP-2	Xiang-Ming et al. (2016)
SEN2	5) Multiple myeloma	Up-regulated	Associate with poor prognosis	Associate with NF- κ B signaling	Xu et al. (2015)
	1) Bladder	Down-regulated	Suppress tumor metastasis	Inhibit the expression of MMP-13	Tan et al. (2013)
SEN3	2) Hepatocellular carcinoma	Down-regulated	Play a role in cell growth control	Modulate the stability of β -catenin	Shen et al. (2012)
	1) Gastric	Up-regulated	Induce specific mesenchymal gene expression in cancer metastasis	Potentiate the transcriptional activity of FOXO2	Ren et al. (2014)
SEN5	2) Oral squamous cell carcinoma	Up-regulated	Correlate with tumor differentiation	Associate with ROS	Sun et al. (2013)
	1) Breast	Up-regulated	Correlate with poor prognosis	Through SENP5-TGF β -MMP9 cascade process	Cashman et al. (2014)
SUMO E1	1) Lung adenocarcinoma	Up-regulated	Lower expression in lymph node metastases	—	Inamura et al. (2007)
	SAE1	Up-regulated	Correlate with tumorigenesis in vivo	Decrease chemotherapy sensitivity	Liu et al. (2015)
	SAE2	Up-regulated	Correlate with survival of the patients.	Associate with oncogenic transcription factor Myc	Kessler et al. (2012)
SUMO E2	1) Liver	Up-regulated	UBC9 directly phosphorylated by cell division cycle 2	Serine 71 of Ubc9 are required for phosphorylation	Tomasi et al. (2012)
	2) Colon	Up-regulated	Express at high levels in primary cancer	—	Moschos et al. (2010)
	3) Lung	Up-regulated	Express at high levels in primary cancer tissue and metastatic nodules	—	Li et al. (2013)
	4) Breast	Up-regulated	Correlate with poor response to chemotherapy and clinical prognosis	—	Chen et al. (2011)
	5) Melanoma	Up-regulated	Express at high levels in melanoma-positive lymph nodes	—	Moschos et al. (2007)
	6) Glioblastoma	Up-regulated	Promote DNA synthesis and cell growth	Block H2AX phosphorylation which indicates DNA double-strand damage, and G2/M cell cycle arrest	Yang et al. (2013)
	7) Lung adenocarcinoma	Up-regulated	Correlate positively with Dukes' stage	—	Moschos et al. (2010)

—, no relevant information found in the literature.

TABLE 2
Small-molecule inhibitors for SENP1/2
Inhibitors' chemical structure from corresponding literature or PubChem.

Target	Inhibitors	Chemical Structure	Source	Products	IC ₅₀ (μ M)	Solubility	Applicability (Cells)	Activity	Assay	Study
SENP1	Triptolide		Extracted from the Chinese herb <i>Tripterygium wilfordii</i> Hook F	Natural	0.0203	DMSO	LNCaP, PC-3	In vivo and in vitro	From the Chinese herb	Huang et al. (2012)
	Compound j5		2-(4-Chlorophenyl)-2-oxoethyl 4-benzamidobenzoate derivatives	Synthetic	2.385	DMSO	—	In vitro	Virtual screening and docking	Chen et al. (2012b)
	Compound 4		1-[4-(N-benzylamino) phenyl]-3-phenylurea derivatives	Synthetic	29.6	—	HeLa	In vitro	In silico screening	Uno et al. (2012)
	Compound 13m		4'-Methoxy-biphenyl-3-carboxylic acid 3(3-phenylpropionylamino)-benzylamide	Synthetic	3.5	DMSO	—	In vitro	In silico screening and rational drug design	Zhao et al. (2016)
	Momordin Ic (Mc)		A pentacyclic triterpenic compound	Natural	15.37	DMSO	PC-3	In vivo and in vitro	Cellular thermal shift assay	Wu et al. (2016)
	Compound 3		Based on hydrolysis of RanGAP-SUMO	Synthetic	3.3	—	—	In vitro	A quantitative assay based on readily available SDS-PAGE-Coomassie system	Xie et al. (2016)
SENP2	Compound 69		A class of 1,2,5-oxadiazoles	Synthetic	5.9	DMSO	—	In vitro	Combination of structure based virtual screening and quantitative FRET-based assay.	Kumar et al. (2014)
	Compound 117			Synthetic	3.7	DMSO	—	In vitro		
	Ebselen		2-phenyl-1,2-benzoselenazol-3-one	Synthetic	2	DMSO	B35	In vivo and In vitro	A quantitative high-throughput screen	Bernstock et al. (2018b)

— No relevant information found in the literature; CDH1, cadherin-1; CDK, cyclin-dependent kinase; H2AX, H2A histone family member X; HIF-1, hypoxia-inducible factor 1; MMP, matrix metalloproteinase; NF- κ B, nuclear factor κ B; PDAC, pancreatic ductal adenocarcinoma cells; ROS, reactive oxygen species; TGF β , transforming growth factor β ; TNM, tumor, node, and metastases staging system.

TABLE 3
Small-molecule inhibitors for SUMO-activating enzyme E1
Inhibitors' chemical structure from corresponding literature or PubChem.

Inhibitors	Chemical Structure	Source	Products	Substrate	IC ₅₀ (μM)	Solubility	Applicability (Cells)	Activity	Assay	Study
Ginkgolide acid Kerriamycin B Compound 21		An alkylphenol derivative Purified from actinomycetes Phenyl urea compound	Natural	RanGap1	3.0	DMSO	H1299	In vivo and in vitro	In situ cell-based SUMOylation assay	Fukuda et al. (2009a)
	Natural	RanGap1	11.7	MeOH	HEK293T	In vitro	In situ cell-based SUMOylation assay	Fukuda et al. (2009b)
	Synthetic	RanGap1	14.4	DMSO	—	In vitro	Structure-based virtual screening	Kumar et al. (2013)
Davidiin		Isolated from <i>Davidia involucrata</i>	Natural	RanGap1	0.15	MeOH	MKN-45, DU-145, NCI-H460	In vitro	In situ cell-based SUMOylation assay	Takemoto et al. (2014)
Tannic acid		A polyphenol	Natural	hLRH-1	12.8	DMSO	JEG3HepG2	In vivo and in vitro	Cell-based gene expression screen	Suzawa et al. (2015)
ML-792		Methyl sulfamate	Synthetic	RanGap1	0.003	DMSO	MDA-MB-468, HCT116, Colo- 205, MDA-MB-231, A375	In vitro	Pyrazole-carbonylpyrimidine-based scaffold and medicinal-chemistry efforts	He et al. (2017)

—, no relevant information found in the literature.

TABLE 4
Small-molecule inhibitors for SUMO-conjugating enzyme E2

Inhibitors	Chemical Structure	Source	Products	Substrate	IC ₅₀ (μM)	Solubility	Applicability (Cells)	Activity	Assay	Study
GSK145A		Diamino-pyrimidine compound	Synthetic	TRPS1 peptid	12.5	DMSO	—	In vitro	High-throughput fluorescence polarization assay	Brandt et al. (2013)
2-D08		A synthetic flavone	Synthetic	FL-AR peptide	6.0	DMSO	ZR-75-1, BT-474	In vitro	Microfluidic electrophoretic mobility shift system	Kim et al. (2013)
Spectomycin B1		<i>Streptomyces spectabilis</i> , originally identified as an antibiotic against gram-positive bacteria	Natural	RanGap1	4.4	DMSO	MCF-7	In vivo and in vitro	In situ cell-based screening system	Hirohama et al. (2013)
Compound 2		A degradation product from a commercial screening collection	Synthetic	RanGap1	75	DMSO	—	In vitro	Small-molecule microarray-based screening	Zlotkowski et al. (2017)

—, no relevant information found in the literature; DMSO, dimethylsulfoxide.

by quantitative high-throughput screening (Bernstock et al., 2018b).

SUMO-E1 and Its Anticancer Inhibitors. SUMO E1, an activating enzyme in SUMOylation pathway, is a heterodimer consisting of SUMO-activating enzyme subunit 1 (SAE1) and subunit 2 (SAE2), which has a particular significance due to its enhancing cell metastasis in tumorigenesis (Table 1). The increased SUMO enzymes are closely associated with pathogenesis of HCC; higher levels of SUMO E1 indicate lower survival rates of HCC patients (Lee and Thorgeirsson, 2004). Knock-down of SAE2 inhibits cancer malignancy and increases chemotherapy sensitivity of small cell lung cancer (Liu et al., 2015). Breast cancer patients with lower expression of SAE1/SAE2 have significantly fewer cases of metastatic cancer and increased survival compared with patients with higher SAE1/SAE2 levels (Kessler et al., 2012). SAE1/SAE2 is also up-regulated in lung adenocarcinoma (Inamura et al., 2007).

There has been growing interest in investigating selective SUMO E1-targeting inhibitors (Table 3). For example, several natural products—including ginkgolic acid (Fukuda et al., 2009a), kerriamycin B (Fukuda et al., 2009b), and davidiin (Takemoto et al., 2014)—are confirmed to inhibit SUMO E1 by blocking formation of SUMO E1–SUMO intermediate. Among these natural products, ginkgolic acid is the most widely used and commercially available chemical reagent that targets global SUMOylation, but its inhibitory effects on cells can vary greatly depending on the assay and a measured substrate. Tannic acid has been identified as targeting human liver receptor homolog-1 as a general nontoxic SUMOylation inhibitor via cell-based screening (Suzawa et al., 2015).

Most natural SUMO E1 inhibitors function in the micromolar range except for davidiin, which inhibits at submicromolar concentrations. To address this limitation, other synthetic SAE inhibitors—compound 21 (Kumar et al., 2013) and ML-792 ([[(1*R*,2*S*,4*R*)-4-[[5-[1-[(3-bromophenyl)methyl]pyrazole-3-carbonyl]pyrimidin-4-yl]amino]-2-hydroxycyclopentyl)methyl sulfamate] (He et al., 2017)—have been developed to inhibit cancer cell growth with good selective activity. ML-792 forms a covalent adduct with SUMO at its C terminus, which inhibits the enzyme activity of SAE2 at nanomolar concentrations (He et al., 2017). Therefore, ML-792 selectively blocks SAE activity in treating MYC-amplified tumors.

SUMO-E2 and Its Anticancer Inhibitors. Ubiquitin-conjugating enzyme 9 (Ubc9) is a sole SUMO E2 in the SUMOylation pathway, which promotes SUMO transferring into Lys residues of a target substrate even in the absence of SUMO E3 (Gareau and Lima, 2010). Ubc9 is increased in many cancers (Seeler and Dejean, 2017), including advanced melanomas, head and neck tumor, lung tumor, HCC, colon cancer, breast cancer, and glioblastoma (Ahn et al., 2001; Moschos et al., 2007, 2010; Wu et al., 2009; Chen et al., 2011; Tomasi et al., 2012; Li et al., 2013; Yang et al., 2013). Inhibition of Ubc9 expression sensitizes cytotoxic effects of chemotherapeutic drugs against melanoma cells (Ahn et al., 2001; Moschos et al., 2007). Ubc9 stability and its expression are increased by cell division cycle 2 (Cdc2)-mediated phosphorylation in HCC (Tomasi et al., 2012). Therefore, detection of Ubc9 expression level could be helpful to early diagnosis of adenocarcinoma, HCC, and other cancers.

Suppression of Ubc9 by small-molecule inhibitors represents a potential or emerging strategy. We have summarized

TABLE 5
The pros and cons of SUMO inhibitors

Protein SUMOylation Inhibitors	
Pros	Cons
Availability, compounds derived from natural products or synthetic chemicals	Unclear molecular mechanism of anticancer
Inhibitors target different SUMOylation-associated enzymes	Weak activity and low selectivity
Reduce the level of protein SUMOylation in different cancer	No high inhibition rate
Inhibit cell proliferation and migration of several specific cancer cell lines	No efficient antitumor activities for multiple cancers
A few inhibitors have been approved to enter clinical trials	No clinical applications

several Ubc9 inhibitors that suppress human malignant cells in vitro (Table 4). For instance, GSK145A was identified as an Ubc9 inhibitor using a high-throughput assay (Brandt et al., 2013). 2-D08 (2',3',4'-trihydroxyflavone) has been identified as a potent SUMOylation inhibitor by an electrophoretic mobility shift assay (Kim et al., 2013). In addition, spectomycin B1, an antibiotic against gram-positive bacteria (Staley and Rinehart, 1994) that has been identified as a novel SUMOylation inhibitor with direct binding to Ubc9, reduces estrogen receptor-dependent gene expression and suppresses proliferation of breast-cancer cells (Hirohama et al., 2013). Spectomycin B1 is a potential therapeutic agent against hormone-dependent breast cancers (Hirohama et al., 2013). The development of other Ubc9 chemical inhibitors is ongoing through a small-molecule microarray approach (Zlotkowski et al., 2017).

SUMO-E3 and Its Anticancer Inhibitors. SUMO E3 includes the protein inhibitor of activated STAT (PIAS), Ran binding protein 2 (RanBP2), the polycomb protein (Pc2), and other members (Martin et al., 2007). SUMO conjugation is always enhanced in the presence of SUMO E3, but in the absence of SUMO E3 Ubc9 can also promote SUMOs onto Lys residues of a target substrate. Most of current understanding on SUMO E3 functions mainly comes from studies of the PIAS family (Coppola et al., 2009; Chen et al., 2012a), which regulates protein stability and signaling transduction pathways, and controls inflammation (Rabellino et al., 2017). RanBP2, a nuclear pore complex protein that belongs to the second type of SUMO E3, has also been found to increase in multiple myeloma (Felix et al., 2009) and small cell lung cancer (Horio et al., 2010). Therefore, SUMO E3 ligase could be a potential biomarker for cancer and act as a drug target for cancer therapy. However, no small-molecule inhibitor has been designed to effectively inhibit SUMO-E3 enzymes up to now.

Pros and Cons of SUMOylation-Targeting Small-Molecule Inhibitors

SUMOylation-associated enzyme inhibitors have been derived from natural products or synthetic chemicals. Besides previous biologic activities, a few natural products have been found to have novel anticancer potentials via acting on the SUMO process. For example, kerriamycin B and spectomycin B1 are typical antibiotics against gram-positive bacteria, and ginkgolic acid and tannic acid have good anti-inflammatory or immunosuppressive activities (Staley and Rinehart, 1994; Antonoff et al., 2009). These reagents have been confirmed to have anticancer activities by inhibition of protein SUMOylation.

Although these compounds have yet to be approved for any clinical trials against tumors, they hold great promise for development into efficient therapeutics, thus extending new clinical applications to old drugs. As they are already used in clinical applications, these drugs would present few novel side effects if their disease treatment spectrum were enlarged.

The SUMOylation pathway is an ideal drug target to overcome oncogenic mechanisms that promote a proapoptotic state, but most SUMOylation inhibitors display weak anticancer activity and uncharacterized selectivity in their current stage (Table 5). Therefore, many more drug-like chemical scaffolds are needed to uncover the therapeutic potential of SUMOylation-associated enzyme inhibitors, along with their structural description.

Future Perspectives

Protein post-translational modification (PTM) is a step in protein biosynthesis, including phosphorylation, glycosylation, ubiquitination, SUMOylation, and other modifications (Kessler and Edelman, 2011). Among these PTMs, SUMOylation, a competitor of ubiquitination, has become a research hot spot in recent years (Yang et al., 2017). Mass spectrometry is the most sensitive method for identification and quantification of PTMs (Liang et al., 2012), including protein SUMOylation (McManus et al., 2016; Lamoliatte et al., 2017; Lumpkin et al., 2017). More and more SUMOylated proteins have been shown to be involved in the pathogenesis of multiple diseases, including brain ischemia and cancers (Liang et al., 2017; Yang et al., 2017; Bernstock et al., 2018a). Protein SUMOylation mediates in protein-protein interactions, is responsive to DNA damage, changes protein intracellular localization, or directs changes of protein activities (Dou et al., 2011; Hickey et al., 2012). Thus, the SUMOylation pathway has emerged as a promising therapeutic target for drug discovery because of its crucial role in various cancers.

Compared with conventional high-throughput screening, virtual filtering plays a noteworthy role in the identification and optimization of small-molecule inhibitors targeting the SUMO pathway. Through virtual screening in combination with biologic assay (Fukuda et al., 2009a,b; Kumar et al., 2013; Takemoto et al., 2014; Suzawa et al., 2015; He et al., 2017; Bernstock et al., 2018b). GSK145A, 2-D08, and spectomycin B1 inhibit Ubc9 (Brandt et al., 2013; Hirohama et al., 2013; Kim et al., 2013), and several compounds exhibit potential activity against tumors. However, the discovery phase of SUMOylation-associated enzyme inhibitors as a potential therapeutic remains in its infancy. Many of the available

inhibitors of SUMOylation-associated enzyme display weak activity and uncharacterized selectivity.

It is worth mentioning that a novel inhibitor, ML-792, specifically inhibits SAE at nanomolar concentrations without suppressing any other ubiquitin-activating enzymes or various other ATP-using enzymes (He et al., 2017). Therefore, the SUMO inhibitors with high selectivity facilitate further exploration of the novel cellular functions of protein SUMOylation, which opens a new avenue for specifically targeting SUMOylation as a potential clinical anticancer therapy.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Yang, Xia, Wang, Zhao, Sheng, Ye, He, Zhou, Zhu, Xu, Liang.

References

- Ahn JH, Xu Y, Jang WJ, Matunis MJ, and Hayward GS (2001) Evaluation of interactions of human cytomegalovirus immediate-early IE2 regulatory protein with small ubiquitin-like modifiers and their conjugation enzyme Ubc9. *J Virol* **75**: 3859–3872.
- Albrow VE, Ponder EL, Fasci D, Békés M, Deu E, Salvesen GS, and Bogoy M (2011) Development of small molecule inhibitors and probes of human SUMO deconjugating proteases. *Chem Biol* **18**:722–732.
- Antonoff MB, Chugh R, Borja-Cacho D, Dudeja V, Clawson KA, Skube SJ, Sorenson BS, Saltzman DA, Vickers SM, and Saluja AK (2009) Triptolide therapy for neuroblastoma decreases cell viability in vitro and inhibits tumor growth in vivo. *Surgery* **146**:282–290.
- Bawa-Khalife T, Cheng J, Lin SH, Ittmann MM, and Yeh ET (2010) SENP1 induces prostatic intraepithelial neoplasia through multiple mechanisms. *J Biol Chem* **285**: 25859–25866.
- Bawa-Khalife T, Cheng J, Wang Z, and Yeh ET (2007) Induction of the SUMO-specific protease 1 transcription by the androgen receptor in prostate cancer cells. *J Biol Chem* **282**:37341–37349.
- Bellail AC, Olson JJ, and Hao C (2014) SUMO1 modification stabilizes CDK6 protein and drives the cell cycle and glioblastoma progression. *Nat Commun* **5**:4234.
- Bernstock JD, Yang W, Ye DG, Shen Y, Pluchino S, Lee YJ, Hallenbeck JM, and Paschen W (2018a) SUMOylation in brain ischemia: patterns, targets, and translational implications. *J Cereb Blood Flow Metab* **38**:5–16.
- Bernstock JD, Ye D, Smith JA, Lee YJ, Gessler FA, Yasgar A, Kozunetsova J, Jadhav A, Wang Z, Pluchino S, et al. (2018b) Quantitative high-throughput screening identifies cytoprotective molecules that enhance SUMO conjugation via the inhibition of SUMO-specific protease (SENP)2. *FASEB J* **32**:1677–1691.
- Bettermann K, Benesch M, Weis S, and Haybaeck J (2012) SUMOylation in carcinogenesis. *Cancer Lett* **316**:113–125.
- Brandt M, Szewczuk LM, Zhang H, Hong X, McCormick PM, Lewis TS, Graham TI, Hung ST, Harper-Jones AD, Kerrigan JJ, et al. (2013) Development of a high-throughput screen to detect inhibitors of TRP51 sumoylation. *Assay Drug Dev Technol* **11**:308–325.
- Cashman R, Cohen H, Ben-Hamo R, Zilberberg A, and Efroni S (2014) SENP5 mediates breast cancer invasion via a TGF β RI SUMOylation cascade. *Oncotarget* **5**: 1071–1082.
- Chen P, Zhao D, Sun Y, Huang L, Zhang S, and Yuan Y (2012a) Protein inhibitor of activated STAT-1 is downregulated in gastric cancer tissue and involved in cell metastasis. *Oncol Rep* **28**:2149–2155.
- Chen SF, Gong C, Luo M, Yao HR, Zeng YJ, and Su FX (2011) Ubc9 expression predicts chemoresistance in breast cancer. *Chin J Cancer* **30**:638–644.
- Chen Y, Wen D, Huang Z, Huang M, Luo Y, Liu B, Lu H, Wu Y, Peng Y, and Zhang J (2012b) 2-(4-Chlorophenyl)-2-oxoethyl 4-benzamidobenzoate derivatives, a novel class of SENP1 inhibitors: virtual screening, synthesis and biological evaluation. *Bioorg Med Chem Lett* **22**:6867–6870.
- Coppola D, Parikh V, Boulware D, and Blanck G (2009) Substantially reduced expression of PIAS1 is associated with colon cancer development. *J Cancer Res Clin Oncol* **135**:1287–1291.
- Dou H, Huang C, Van Nguyen T, Lu LS, and Yeh ET (2011) SUMOylation and de-SUMOylation in response to DNA damage. *FEBS Lett* **585**:2891–2896.
- Enserink JM (2015) Sumo and the cellular stress response. *Cell Div* **10**:4.
- Felix RS, Colleoni GW, Caballero LO, Yamamoto M, Almeida MS, Andrade VC, Chaffaille MdeL, Silva WA, Jr, Begnami MD, Soares FA, et al. (2009) SAGE analysis highlights the importance of p53csv, ddx5, mapkapk2 and ranbp2 to multiple myeloma tumorigenesis. *Cancer Lett* **278**:41–48.
- Fukuda I, Ito A, Hirai G, Nishimura S, Kawasaki H, Saitoh H, Kimura K, Sodeoka M, and Yoshida M (2009a) Ginkgolic acid inhibits protein SUMOylation by blocking formation of the E1-SUMO intermediate. *Chem Biol* **16**:133–140.
- Fukuda I, Ito A, Uramoto M, Saitoh H, Kawasaki H, Osada H, and Yoshida M (2009b) Kerriamycin B inhibits protein SUMOylation. *J Antibiot (Tokyo)* **62**:221–224.
- Gareau JR and Lima CD (2010) The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. *Nat Rev Mol Cell Biol* **11**: 861–871.
- He X, Riceberg J, Soucy T, Koenig E, Minissale J, Gallery M, Bernard H, Yang X, Liao H, Rabino C, et al. (2017) Probing the roles of SUMOylation in cancer cell biology by using a selective SAE inhibitor. *Nat Chem Biol* **13**:1164–1171.
- Hecker CM, Rabiller M, Haglund K, Bayer P, and Dikic I (2006) Specification of SUMO1- and SUMO2-interacting motifs. *J Biol Chem* **281**:16117–16127.
- Hickey CM, Wilson NR, and Hochstrasser M (2012) Function and regulation of SUMO proteases. *Nat Rev Mol Cell Biol* **13**:755–766.
- Hirohama M, Kumar A, Fukuda I, Matsuoka S, Igarashi Y, Saitoh H, Takagi M, Shin-ya K, Honda K, Kondoh Y, et al. (2013) Spectomycin B1 as a novel SUMOylation inhibitor that directly binds to SUMO E2. *ACS Chem Biol* **8**: 2635–2642.
- Horio Y, Osada H, Shimizu J, Ogawa S, Hida T, and Sekido Y (2010) Relationship of mRNA expressions of RanBP2 and topoisomerase II isoforms to cytotoxicity of amrubicin in human lung cancer cell lines. *Cancer Chemother Pharmacol* **66**: 237–243.
- Huang W, He T, Chai C, Yang Y, Zheng Y, Zhou P, Qiao X, Zhang B, Liu Z, Wang J, et al. (2012) Triptolide inhibits the proliferation of prostate cancer cells and down-regulates SUMO-specific protease 1 expression. *PLoS One* **7**:e37693.
- Inamura K, Shimoi T, Ninomiya H, Hiramatsu M, Okui M, Satoh Y, Okumura S, Nakagawa K, Noda T, Fukayama M, et al. (2007) A metastatic signature in entire lung adenocarcinomas irrespective of morphological heterogeneity. *Hum Pathol* **38**: 702–709.
- Kessler BM and Edelman MJ (2011) PTMs in conversation: activity and function of deubiquitinating enzymes regulated via post-translational modifications. *Cell Biochem Biophys* **60**:21–38.
- Kessler JD, Kahle KT, Sun T, Meerbrey KL, Schlabach MR, Schmitt EM, Skinner SO, Xu Q, Li MZ, Hartman ZC, et al. (2012) A SUMOylation-dependent transcriptional subprogram is required for Myc-driven tumorigenesis. *Science* **335**: 348–353.
- Kim YS, Nagy K, Keyser S, and Schneekloth JS, Jr (2013) An electrophoretic mobility shift assay identifies a mechanistically unique inhibitor of protein sumoylation. *Chem Biol* **20**:604–613.
- Kumar A, Ito A, Hirohama M, Yoshida M, and Zhang KY (2013) Identification of sumoylation activating enzyme 1 inhibitors by structure-based virtual screening. *J Chem Inf Model* **53**:809–820.
- Kumar A, Ito A, Takemoto M, Yoshida M, and Zhang KY (2014) Identification of 1,2,5-oxadiazoles as a new class of SENP2 inhibitors using structure based virtual screening. *J Chem Inf Model* **54**:870–880.
- Lamoliatte F, McManus FP, Maarifi G, Chelbi-Alix MK, and Thibault P (2017) Uncovering the SUMOylation and ubiquitylation crosstalk in human cells using sequential peptide immunoprecipitation. *Nat Commun* **8**:14109.
- Lee JS and Thorgeirsson SS (2004) Genome-scale profiling of gene expression in hepatocellular carcinoma: classification, survival prediction, and identification of therapeutic targets. *Gastroenterology* **127**(5 Suppl 1):S51–S55.
- Li H, Niu H, Peng Y, Wang J, and He P (2013) Ubc9 promotes invasion and metastasis of lung cancer cells. *Oncol Rep* **29**:1588–1594.
- Liang S, Xu Z, Xu X, Zhao X, Huang C, and Wei Y (2012) Quantitative proteomics for cancer biomarker discovery. *Comb Chem High Throughput Screen* **15**:221–231.
- Liang Z, Yang Y, He Y, Yang P, Wang X, He G, Zhang P, Zhu H, Xu N, Zhao X, et al. (2017) SUMOylation of IQGAP1 promotes the development of colorectal cancer. *Cancer Lett* **411**:90–99.
- Liu X, Xu Y, Pang Z, Guo F, Qin Q, Yin T, Sang Y, Feng C, Li X, Jiang L, et al. (2015) Knockdown of SUMO-activating enzyme subunit 2 (SAE2) suppresses cancer malignancy and enhances chemotherapy sensitivity in small cell lung cancer. *J Hematol Oncol* **8**:67.
- Lumpkin RJ, Gu H, Zhu Y, Leonard M, Ahmad AS, Clauser KR, Meyer JG, Bennett EJ, and Komives EA (2017) Site-specific identification and quantitation of endogenous SUMO modifications under native conditions. *Nat Commun* **8**:1171.
- Ma C, Wu B, Huang X, Yuan Z, Nong K, Dong B, Bai Y, Zhu H, Wang W, and Ai K (2014) SUMO-specific protease 1 regulates pancreatic cancer cell proliferation and invasion by targeting MMP-9. *Tumour Biol* **35**:12729–12735.
- Martin S, Wilkinson KA, Nishimune A, and Henley JM (2007) Emerging extracellular roles of protein SUMOylation in neuronal function and dysfunction. *Nat Rev Neurosci* **8**:948–959.
- McManus FP, Altamirano CD, and Thibault P (2016) In vitro assay to determine SUMOylation sites on protein substrates. *Nat Protoc* **11**:387–397.
- Moschos SJ, Jukic DM, Athanassiou C, Bhargava R, Dacic S, Wang X, Kuan SF, Fayewicz SL, Galambos C, Acquafondata M, et al. (2010) Expression analysis of Ubc9, the small ubiquitin-like modifier (SUMO) E2 conjugating enzyme, in normal and malignant tissues. *Hum Pathol* **41**:1286–1298.
- Moschos SJ, Smith AP, Mandic M, Athanassiou C, Watson-Hurst K, Jukic DM, Edington HD, Kirkwood JM, and Becker D (2007) SAGE and antibody array analysis of melanoma-infiltrated lymph nodes: identification of Ubc9 as an important molecule in advanced-stage melanomas. *Oncogene* **26**:4216–4225.
- Qiao Z, Wang W, Wang L, Wen D, Zhao Y, Wang Q, Meng Q, Chen G, Wu Y, and Zhou H (2011) Design, synthesis, and biological evaluation of benzodiazepine-based SUMO-specific protease 1 inhibitors. *Bioorg Med Chem Lett* **21**:6389–6392.
- Rabellino A, Andreani C, and Scaglioni PP (2017) The role of PIAS SUMO E3-ligases in cancer. *Cancer Res* **77**:1542–1547.
- Ren YH, Liu KJ, Wang M, Yu YN, Yang K, Chen Q, Yu B, Wang W, Li QW, Wang J, et al. (2014) De-SUMOylation of FOXO2 by SENP3 promotes the epithelial-mesenchymal transition in gastric cancer cells. *Oncotarget* **5**:7093–7104.
- Seeler JS and Dejean A (2003) Nuclear and nuclear functions of SUMO. *Nat Rev Mol Cell Biol* **4**:690–699.
- Seeler JS and Dejean A (2017) SUMO and the robustness of cancer. *Nat Rev Cancer* **17**:184–197.
- Shen HJ, Zhu HY, Yang C, and Ji F (2012) SENP2 regulates hepatocellular carcinoma cell growth by modulating the stability of β -catenin. *Asian Pac J Cancer Prev* **13**:3583–3587.
- Staley AL and Rinehart KL (1994) Spectomycins, new antibacterial compounds produced by Streptomyces spectabilis: isolation, structures, and biosynthesis. *J Antibiot (Tokyo)* **47**:1425–1433.
- Sun Z, Hu S, Luo Q, Ye D, Hu D, and Chen F (2013) Overexpression of SENP3 in oral squamous cell carcinoma and its association with differentiation. *Oncol Rep* **29**: 1701–1706.

- Suzawa M, Miranda DA, Ramos KA, Ang KK, Faivre EJ, Wilson CG, Caboni L, Arkin MR, Kim YS, Fletterick RJ, et al. (2015) A gene-expression screen identifies a non-toxic sumoylation inhibitor that mimics SUMO-less human LRH-1 in liver. *eLife* **4**: e09003.
- Takemoto M, Kawamura Y, Hirohama M, Yamaguchi Y, Handa H, Saitoh H, Nakao Y, Kawada M, Khalid K, Koshino H, et al. (2014) Inhibition of protein SUMOylation by davidiin, an ellagitannin from *Davidia involucrata*. *J Antibiot (Tokyo)* **67**: 335–338.
- Tan MY, Mu XY, Liu B, Wang Y, Bao ED, Qiu JX, and Fan Y (2013) SUMO-specific protease 2 suppresses cell migration and invasion through inhibiting the expression of MMP13 in bladder cancer cells. *Cell Physiol Biochem* **32**:542–548.
- Tomasi ML, Tomasi I, Ramani K, Pascale RM, Xu J, Giordano P, Mato JM, and Lu SC (2012) S-adenosyl methionine regulates ubiquitin-conjugating enzyme 9 protein expression and sumoylation in murine liver and human cancers. *Hepatology* **56**:982–993.
- Uno M, Koma Y, Ban HS, and Nakamura H (2012) Discovery of 1-[4-(N-benzylamino)phenyl]-3-phenylurea derivatives as non-peptidic selective SUMO-sentrin specific protease (SEN)1 inhibitors. *Bioorg Med Chem Lett* **22**:5169–5173.
- Wang Q, Xia N, Li T, Xu Y, Zou Y, Fan Q, Bawa-Khalife T, Yeh ET, and Cheng J (2013) SUMO-specific protease 1 promotes prostate cancer progression and metastasis. *Oncogene* **32**:2493–2498.
- Wu F, Zhu S, Ding Y, Beck WT, and Mo YY (2009) MicroRNA-mediated regulation of Ubc9 expression in cancer cells. *Clin Cancer Res* **15**:1550–1557.
- Wu J, Lei H, Zhang J, Chen X, Tang C, Wang W, Xu H, Xiao W, Gu W, and Wu Y (2016) Momordin Ic, a new natural SENP1 inhibitor, inhibits prostate cancer cell proliferation. *Oncotarget* **7**:58995–59005.
- Xiang-Ming Y, Zhi-Qiang X, Ting Z, Jian W, Jian P, Li-Qun Y, Ming-Cui F, Hong-Liang X, Xu C, and Yun Z (2016) SENP1 regulates cell migration and invasion in neuroblastoma. *Biotechnol Appl Biochem* **63**:435–440.
- Xie W, Wang Z, Zhang J, Wang L, Zhao Y, and Zhou H (2016) Development and evaluation of a highly reliable assay for SUMO-specific protease inhibitors. *Bioorg Med Chem Lett* **26**:2124–2128.
- Xu J, Sun HY, Xiao FJ, Wang H, Yang Y, Wang L, Gao CJ, Guo ZK, Wu CT, and Wang LS (2015) SENP1 inhibition induces apoptosis and growth arrest of multiple myeloma cells through modulation of NF- κ B signaling. *Biochem Biophys Res Commun* **460**:409–415.
- Xu Y, Li J, Zuo Y, Deng J, Wang LS, and Chen GQ (2011) SUMO-specific protease 1 regulates the in vitro and in vivo growth of colon cancer cells with the upregulated expression of CDK inhibitors. *Cancer Lett* **309**:78–84.
- Xu Y, Zuo Y, Zhang H, Kang X, Yue F, Yi Z, Liu M, Yeh ET, Chen G, and Cheng J (2010) Induction of SENP1 in endothelial cells contributes to hypoxia-driven VEGF expression and angiogenesis. *J Biol Chem* **285**:36682–36688.
- Yang W, Wang L, Roehn G, Pearlstein RD, Ali-Osman F, Pan H, Goldbrunner R, Krantz M, Harms C, and Paschen W (2013) Small ubiquitin-like modifier 1-3 conjugation [corrected] is activated in human astrocytic brain tumors and is required for glioblastoma cell survival [published correction appears in Cancer Sci (2013) 104:274]. *Cancer Sci* **104**:70–77.
- Yang Y, He Y, Wang X, Liang Z, He G, Zhang P, Zhu H, Xu N, and Liang S (2017) Protein SUMOylation modification and its associations with disease. *Open Biol* **7**: 170167.
- Zhao Y, Wang Z, Zhang J, and Zhou H (2016) Identification of SENP1 inhibitors through in silico screening and rational drug design. *Eur J Med Chem* **122**: 178–184.
- Zlotkowski K, Hewitt WM, Sinniah RS, Tropea JE, Needle D, Lountos GT, Barchi JJ, Jr, Waugh DS, and Schneekloth JS, Jr (2017) A small-molecule microarray approach for the identification of E2 enzyme inhibitors in ubiquitin-like conjugation pathways. *SLAS Discov* **22**:760–766.
- Zuo Y and Cheng JK (2009) Small ubiquitin-like modifier protein-specific protease 1 and prostate cancer. *Asian J Androl* **11**:36–38.

Address correspondence to: Dr. Shufang Liang, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, and Collaborative Innovation Center for Biotherapy, No. 17, 3rd Section of People's South Road, Chengdu, 610041, People's Republic of China. E-mail: zizi2006@scu.edu.cn