

Special Section: From Insight to Modulation of CXCR4 and ACKR3 (CXCR7) Function – Minireview

CXCR4/ACKR3 Phosphorylation and Recruitment of Interacting Proteins: Key Mechanisms Regulating Their Functional Status

Amos Fumagalli, Aurélien Zarca,¹ Maria Neves,¹ Birgit Caspar, Stephen J. Hill, Federico Mayor, Jr., Martine J. Smit, and Philippe Marin

IGF, Université de Montpellier, CNRS, INSERM, Montpellier, France (A.F., P.M.); Division of Medicinal Chemistry, Faculty of Science, Amsterdam Institute for Molecules, Medicines and Systems (AIMMS), Vrije Universiteit Amsterdam, Amsterdam, The Netherlands (A.Z., M.J.S.); Departamento de Biología Molecular and Centro de Biología Molecular “Severo Ochoa” (UAM-CSIC), Madrid, Spain (M.N., F.M.); CIBERCV, Instituto de Salud Carlos III, Madrid, Spain (M.N., F.M.); and Division of Physiology, Pharmacology and Neuroscience, Medical School, University of Nottingham, Queen’s Medical Centre, Nottingham, United Kingdom (B.C., S.J.H.)

Received November 28, 2018; accepted February 21, 2019

ABSTRACT

The C-X-C motif chemokine receptor type 4 (CXCR4) and the atypical chemokine receptor 3 (ACKR3/CXCR7) are class A G protein-coupled receptors (GPCRs). Accumulating evidence indicates that GPCR subcellular localization, trafficking, transduction properties, and ultimately their pathophysiological functions are regulated by both interacting proteins and post-translational modifications. This has encouraged the development of novel techniques to characterize the GPCR interactome and to identify residues subjected to post-translational modifications, with a special focus on phosphorylation. This review first describes state-of-the-art methods for the identification of GPCR-interacting proteins

and GPCR phosphorylated sites. In addition, we provide an overview of the current knowledge of CXCR4 and ACKR3 post-translational modifications and an exhaustive list of previously identified CXCR4- or ACKR3-interacting proteins. We then describe studies highlighting the importance of the reciprocal influence of CXCR4/ACKR3 interactomes and phosphorylation states. We also discuss their impact on the functional status of each receptor. These studies suggest that deeper knowledge of the CXCR4/ACKR3 interactomes along with their phosphorylation and ubiquitination status would shed new light on their regulation and pathophysiological functions.

Introduction

The C-X-C chemokine receptor type 4 (CXCR4) and the atypical chemokine receptor 3 (ACKR3), earlier referred to as

This research was funded by a European Union’s Horizon2020 MSCA Program [Grant agreement 641833 (ONCORNET)]; A.F. and P.M. are also supported by CNRS, INSERM, Université de Montpellier and Fondation pour la Recherche Médicale (FRM); F.M. laboratory is also supported by grants from Ministerio de Economía, Industria y Competitividad (MINECO) of Spain [Grant SAF2017-84125-R]; CIBERCV-Instituto de Salud Carlos III, Spain [Grant CB16/11/00278] to F.M., cofunded with European FEDER contribution; Comunidad de Madrid-B2017/BMD-3671-INFLAMUNE; and Fundación Ramón Areces.

¹A.Z. and M.N. contributed equally to this work.

<https://doi.org/10.1124/mol.118.115360>.

CXCR7, are class A G protein-coupled receptors (GPCRs). Stromal cell-derived factor-1/C-X-C motif chemokine 12 (CXCL12) binds to both CXCR4 and ACKR3 receptors, whereas C-X-C motif chemokine 11 (CXCL11) binds only to the latter and to the C-X-C chemokine receptor type 3. CXCR4 and ACKR3 are coexpressed in various cell types [e.g., endothelial cells (Volin et al., 1998; Berahovich et al., 2014), neurons (Banisadr et al., 2002; Sánchez-Alcañiz et al., 2011); and glial cells (Banisadr et al., 2002, 2016; Odemis et al., 2010)], where they play a pivotal role in migration, proliferation, and differentiation. They are also overexpressed in various tumors and control invasion and metastasis (Sun et al., 2010; Zhao et al., 2015; Nazari et al., 2017).

ABBREVIATIONS: ACKR3 or CXCR7, atypical chemokine receptor 3; AIP4, E3 ubiquitin ligase atrophin-interacting protein 4; BRET, bioluminescence resonance energy transfer; CD164, endolyn; CD74, HLA class II histocompatibility antigen gamma chain; CHAPSO, 3-[(3-cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate; Co-IP, co-immunoprecipitation; CXCL12, C-X-C motif chemokine 12; CXCR4, C-X-C motif chemokine receptor 4; DDM, n-dodecyl- β -D-maltopyranoside; EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; ERK, extracellular signal-regulated kinases; FRET, fluorescence resonance energy transfer; GiP, G protein-coupled receptor-interacting protein; GPCR, G protein-coupled receptor; GRK, G protein-coupled receptor kinase; HEK, human embryonic kidney; IP, immunoprecipitation; LC-MS/MS, liquid chromatography coupled to tandem mass spectrometry; MIF, macrophage-migration inhibitory factor; PKC, protein kinase C; STAT, signal transducer and activator of transcription.

There is now considerable evidence indicating that GPCRs do not operate as isolated proteins within the plasma membrane. Instead, they physically interact with numerous proteins that influence their activity, trafficking, and signal transduction properties (Bockaert et al., 2004; Ritter and Hall, 2009; Magalhaes et al., 2012). These include proteins canonically associated with most GPCRs, such as G proteins, G protein-coupled receptor kinases (GRKs) and β -arrestins, specific partner proteins, and even GPCRs themselves. In fact, in comparison with monomers, GPCRs can form homomers and heteromers with specific pharmacological and signal transduction properties (Ferré et al., 2014).

Phosphorylation is another key mechanism contributing to the regulation of GPCR functional status and signal transduction (Tobin, 2008). Both canonical GRKs and other specific protein kinases are able to phosphorylate GPCRs at multiple sites (Luo et al., 2017), generating the so-called GPCR phosphorylation barcode that determines β -arrestin recruitment, receptor intracellular fate, and signaling outcomes (Reiter et al., 2012).

This review will describe recent data highlighting the influence of the CXCR4 and ACKR3 interactome on their functional activity and signal transduction properties. A special focus will be given to the reciprocal influence of the interactome on CXCR4/ACKR3 phosphorylation and its impact on the functional status and pathophysiological functions of each receptor.

Methods for the Identification of GPCR-Interacting Proteins

Considerable evidence suggests that GPCRs recruit GPCR-interacting proteins (GIPs) (Maurice et al., 2011). This prompted investigations aimed at identifying GIPs and at characterizing GPCR-GIP interactions, using either unbiased or targeted approaches. In unbiased methods, no knowledge of the GIPs is required beforehand and the GPCR of interest is used as bait to purify unknown GIPs. Meanwhile, targeted methods are devoted to the validation and characterization of previously identified GPCR-GIP interactions. Methods for identifying GIPs or characterizing GPCR-GIP interactions include genetic, biophysical, biochemical, or proteomic approaches and are summarized in Table 1.

Genetic Methods. The first method belonging to this class is the yeast two-hybrid assay (Fields and Song, 1989). In this method, the protein of interest (the bait protein) is expressed in yeast as a fusion to the DNA-binding domain of a transcription factor lacking the transcription activation domain. To identify partners of this bait, a plasmid library that expresses cDNA-encoded proteins fused to a transcription activation domain is introduced into the yeast strain. Interaction of a cDNA-encoded protein with the bait protein results in the activation of the transcription factor and expression of a reporter gene, enabling growth on specific media or a color reaction and the identification of the cDNA encoding the target proteins. A first disadvantage is the loss of spatial-temporal localization of the interaction; in fact, the yeast two-hybrid assay only captures a snapshot of potential interactions in an artificial biologic system. A second disadvantage is that it is not possible to investigate membrane-anchored proteins since the two proteins must cross the nuclear membrane to carry the reconstituted

transcription factor to the DNA. To overcome this issue, the membrane yeast two-hybrid assay (Stagljar et al., 1998) was developed. In this assay, the ubiquitin protein is split into two fragments that are fused to the two proteins of interest. The ubiquitin C-terminal fragment is then conjugated to a transcription factor that is released when the interaction occurs, and the ubiquitin protein is reformed. However, as in the yeast two-hybrid assay, the spatial-temporal localization of the interaction is lost. A second limitation is that the ubiquitin C-terminus carrying the transcription factor cannot be fused to soluble proteins because they could diffuse into the nucleus. Therefore, a mammalian version of the assay called mammalian membrane two-hybrid (Petschnigg et al., 2014) was developed. The kinase substrate sensor assay (Lievens et al., 2014), using the signal transducer and activator of transcription 3 (STAT3) as transcription factor, can also be used for investigating protein-protein interactions, including those involving cytosolic and membrane proteins in mammalian cells. However, the kinase substrate sensor assay cannot be used for studying GPCR interaction with proteins involved in the STAT3 cascade.

Biophysical Methods. Energy transfer-based methods, such as bioluminescence and fluorescent resonance energy transfer [BRET (Xu et al., 1999) and FRET (Clegg, 1995)] assays, are targeted methods that are generally used to investigate previously reported interactions. The basis of both is the transfer of energy from a donor to a nearby acceptor ($<100 \text{ \AA}$). Their high sensitivity allows the study of weak and transient interactions. The high spatial-temporal resolution permits accurate kinetic studies for investigating interaction dynamics.

Another biophysical method, with FRET as a basis and employed for the study of protein-protein interaction, is fluorescent lifetime imaging microscopy (Sun et al., 2012). The fluorescence lifetime is the average time that a molecule spends in the excited state before returning to the ground state. Since in FRET the energy transfer from the acceptor to the donor depopulates the excited state energy of the latter, it also shortens its lifetime. Fluorescent lifetime imaging microscopy can accurately measure the shorter donor lifetime that results from FRET; thus, it allows mapping of the spatial distribution of protein-protein interactions in living cells (Sun et al., 2011). Its main advantage over intensity-based FRET is the more accurate measurement of FRET, because only donor signals are measured, which eliminates the corrections for spectral bleed-through (Sun et al., 2011). Its main disadvantages are the necessity of a live specimen and the complexity of data recording and analysis.

Biochemical Methods. The proximity ligation assay (Fredriksson et al., 2002) is another powerful targeted method. In the direct version of the technique, two DNA oligonucleotide-conjugated antibodies are used against the proteins of interest. In the indirect version, secondary DNA-conjugated antibodies are used after the proteins of interest are targeted with an appropriate primary antibody. If the two conjugated antibodies are close enough (30–40 nm), they can bind together. The DNA connecting the two probes is then amplified and hybridized with fluorophores. This allows the visualization of the interaction in the place where it occurs, at a single-molecule resolution. The main disadvantages of the approach are the high costs and the necessity for specific antibodies, which are not always available.

In the bimolecular fluorescent complementation assay (Hu et al., 2002; Hu and Kerppola, 2003), a fluorescent protein

TABLE 1
Principal methods used to identify GPCR-interacting proteins and phosphorylated residues.
GPCR-interacting proteins (1–4) and phosphorylated residues (5).

Classification	Method	Screening	Advantages	Disadvantages
1. Genetic	Y2H	Highly suitable	Easy to perform. Inexpensive.	Loss of spatial-temporal information. Membrane-anchored proteins cannot be investigated. Performed in yeast.
	MYTH	Highly suitable	Easy to perform. Membrane-anchored proteins can be investigated.	Loss of spatial-temporal information. Soluble proteins cannot be investigated. Performed in yeast.
	MaMTH	Highly suitable	Easy to perform. Membrane anchored proteins can be investigated. Performed in mammalian cells.	Loss of spatial-temporal information. Soluble proteins cannot be investigated.
2. Biophysical	KISS	Possible	Sensitive enough for studying interaction dynamic. Both membrane and cytosolic proteins can be investigated.	Loss of spatial-temporal information. Proteins involved in the STAT3 cascade cannot be investigated.
	BRET/FRET	Not suitable	Precise spatial-temporal information. High sensitivity. Possibility to study interactions in living cells.	Generation of fusion proteins. Relies on the proximity and relative orientation between donor and acceptor.
3. Biochemical	Fluorescent lifetime microscopy	Suitable	More accurate than intensity-based FRET.	Data analysis more laborious than intensity-based FRET.
	PLA	Not suitable	Precise spatial information (single-molecule resolution). Possibility to perform in ex-vivo models.	Relies on antibodies. High cost. Not easy to scale up in large studies.
	BioID	Suitable	Precise spatial information. Several interactions in parallel. Possibility to perform in living cells.	Not well suited for studying interaction dynamic (fluorescent signal is delayed).

(continued)

TABLE 1—Continued

Classification	Method	Screening	Advantages	Disadvantages
	NanoBit	Suitable	Precise spatial information. Several interactions in parallel. Possibility to perform in living cells.	
4. Proteomic	Co-IP	Highly suitable	Purification of protein complexes in living cells and tissues.	Rely on antibodies. Loss of spatial-temporal information. Lysis conditions might influence results.
	Pull-down	Highly suitable	Can prove direct interaction.	Loss of spatial-temporal information. In vitro binding assays. Fusion of the receptor on the beads might alter receptor conformation.
	BioID	Highly suitable	Can detect weak and transient interactions in living cells.	Fusion of the biotin to the receptor might alter its targeting or functions.
5. Phosphorylation	[³² P]	Suitable	Very sensitive.	Radioactive method. Cannot give information on the number of phosphorylated residues nor their position.
	LC-MS	Highly suitable	Can pinpoint phosphorylated residues.	Can yield false negatives. Not quantitative unless combined with very expensive isotope tags.
	Mutagenesis	Suitable	Cheap and easy. Based on functional data in living cells. Can pinpoint phosphorylated residues.	Indirect method. Mutagenesis of the C-terminus can impair expression and/or localization of the receptor. Labor-intensive in case of multiple phosphosites. Not quantitative.
	Phospho-antibodies	Suitable	Direct and indirect. Can be used in any cell line. Semi-quantitative and qualitative.	Time consuming and expensive for the generation of the antibodies. Useless with low affinity antibodies. Cannot give information on contiguous phosphorylated residues.

BiFC, bimolecular fluorescent complementation assay; BioID, proximity-dependent biotin identification; KISS, kinase substrate sensor; MaMTH, mammalian membrane two-hybrid assay; MYTH, membrane yeast two-hybrid assay; PLA, proximity ligation assay; Y2H, yeast two-hybrid assay.

TABLE 2
CXCR4- and ACKR3-interacting proteins described in the literature.

Protein	Method of Identification	Cellular Context	Direct	Constitutive/Induced	Site of Interaction	Role	Reference
CXCR4-interacting proteins							
Filamin A	Pull-down Co-IP	HEK293 cells Recombinant protein	Yes	Constitutive and induced The ROCK inhibitor Y27632 reverses CXCL12-induced increased interaction	C-terminal tail and third loop of CXCR4	Stabilize CXCR4 at the surface	Gómez-Moutón et al. (2015)
AIP4	Pull Down Co-IP FRET	HEK293 cells	Yes	Constitutive and induced	CXCR4 C-tail serines and WW domains of AIP4. Serine 324 and 325 when phosphorylated increase interaction	Increase CXCR4 degradation	Bhandari et al. (2009)
RTN3	Y2H Co-IP	HEK293 cells	NA	Constitutive, induction not tested	Carboxyl terminal of RTN3	Increase cytoplasmic localization of CXCR4	Li et al. (2016)
CD74	Co-IP	HEK293 and MonoMac6 cells	NA	Constitutive, induction not tested	NA	Phosphorylation of AKT	Schwartz et al. (2009)
TLR2	Colocalization FRET Co-IP	Human monocyte and HEK293 cells	NA	Induced by Pg-fimbria	NA	CXCR4 inhibits TLR2-induced NF- κ B activation. In addition, CXCR4 found to be receptor of the pattern-recognition receptor complex	Hajjshengallis et al. (2008), Triantafyllou et al. (2008)
NMMHC	Pull-down Co-IP Colocalization	Jurkat T and Peer T cells lymphocytes	NA	Constitutive and not induced by CXCL12	CXCR4 C-terminus	Lymphocytes migration	Rey et al. (2002)
Drebrin	Pull Down Co-IP FRET	J77 T, HEK293T and HIV-infected T cells	YES	Constitutive and induced by superantigen E, which also relocalizes the edge of migrating lymphocytes	Drebrin N-terminus positively regulates interaction whereas the C-terminus seems to negatively regulate it	Drebrin affects key physiologic processes during antigen presentation in HIV entry	Pérez-Martínez et al. (2010), Gordo-Alonso et al. (2013)
CD164	Co-IP Colocalization	Jurkat and Ovarian surface epithelial cells	NA	Only induced when CXCL12 is presented on fibronectin	NA	CD164 participates to the CXCL12 mediated AKT and PKC- ζ phosphorylation	Forde et al. (2007), Huang et al. (2013)
mDIA2	Co-IP Colocalization	MDA-MB-231 cells	NA	Constitutive (very weak) and CXCL12 induced	NA	Cytoskeletal rearrangement necessary for nonapoptotic blebbing	Wyse et al. (2017)
ATP13A2 PI3K γ	MYTH Co-IP	Yeast Human myeloid cells	YES NA	Constitutive Only CXCL12 induced	NA NA	Integrin activation and chemotaxis	Usenovic et al. (2012) Schmid et al. (2011)
PECAM-1	PLA	Human coronary artery endothelial cells	NO	Constitutive Induction not studied	NA	CXCR4 part of a junctional mechano-sensitive complex	dela Paz et al. (2014)
MYBL2	2HY	Yeast	Yes	Induction not studied	NA	NA	Wang et al. (2011)
KCNK1	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
TMEM63B	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
GOLT1B	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
eIFB2	Co-IP Colocalization	Pre-B NALM6 cells	NA	Constitutive and negatively regulated by CXCL12	NA	NA	Palmesino et al. (2016)
CDC73	Co-IP	Pre-B NALM6 cells	NA	Constitutive	NA	NA	Palmesino et al. (2016)
CTR9	Co-IP	Pre-B NALM6 cells	NA	Induction not studied	NA	NA	Palmesino et al. (2016)
PAF1	Co-IP	Pre-B NALM6 cells	NA	Induction not studied	NA	NA	Palmesino et al. (2016)

(continued)

TABLE 2—Continued

Protein	Method of Identification	Cellular Context	Direct	Constitutive/Induced	Site of Interaction	Role	Reference
NPM	Co-IP	Pre-B NALM6 cells	NA	Constitutive Induction not studied	NA	NA	Palmesino et al. (2016)
CD11B	Co-IP	Pre-B NALM6 cells	NA	Constitutive Induction not studied	NA	NA	Palmesino et al. (2016)
PTPRS	Co-IP	Pre-B NALM6 cells	NA	Constitutive Induction not studied	NA	NA	Palmesino et al. (2016)
LGALS8	Co-IP	Pre-B NALM6 cells	NA	Constitutive Induction not studied	NA	NA	Palmesino et al. (2016)
ACKR3-interacting proteins							
EGFR	PLA Colocalization Co-IP	MCF7 cells, breast cancer tissues, CaP cells	Mediated by β -arrestin2	Interaction constitutive and induced by the epidermal growth factor.	NA	ACKR3 mediates phosphorylation of EGFR at tyrosine1110 after EGF-stimulation and phosphorylation of ERK1/2 with consequences on tumor proliferation. ACKR3 is involved in MIF-mediated ERK-1/2 and ζ -chain-associated protein kinase activation in addition to MIF-mediated chemotaxis.	Singh and Lokeshwar (2011), Salazar et al. (2014), Kallifatidis et al. (2016)
CD74	Co-IP Colocalization PLA	NIH/3T3 cells and human B cells	NA	Constitutive, induction not investigated.	NA	NA	Alampour-Rajabi et al. (2015)
PECAM-1	PLA	HCAECs cells	NA	Constitutive	NA	NA	dela Paz et al. (2014)
ATP13A2	MYTH	Yeast	Yes	NA	NA	NA	Uosenovic et al. (2012)
GJB2	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
MRPL4	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ATP5H	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ATP5B	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ATP5A1	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ATP5O	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ACKR3	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
ESPL1	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
HECTD2	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)
UBR7	Co-IP	HeLa cells	NA	NA	NA	NA	Hein et al. (2015)

ATP13A2, cation-transporting ATPase 13A2; ATP5H/ATP5B/ATP5A1/ATP5O, ATP synthase subunit *d*/beta/alpha/*O*, mitochondrial; CAP, human prostate cancer cells; CD11B, cyclin-dependent kinase 11B; CD164, endolysin; CDC73, parafibromin; CTR9, SH2 domain binding protein; eIF2B, eukaryotic translation initiation factor 2B; EGFR, epidermal growth factor receptor; ESPL1, HCAEC, Human Coronary Artery Endothelial Cells; HECTD2, probable E3 ubiquitin-protein ligase; GJB2, gap junction β -2 protein; GOL/T1B, vesicle transport protein GOT1B; KCNK1, potassium channel subfamily K member 1; LGALS8, galectin; mDIA2, diaphanous-related formin-2; MYBL2, myb-related protein B; MRPL4, 54S ribosomal protein L4; mitochondrial; MYTH, membrane yeast two-hybrid assay; NA, not applicable; NF, nuclear factor; NMMHC, motor protein nonmuscle myosin H chain; NPM, nucleophosmin; PAF1, hypothetical protein PD2; PECAM-1, platelet endothelial cell adhesion molecule; PI3K γ , PI3-kinase isoform p110 γ ; PLA, proximity ligation assay; PTPRS, receptor-type tyrosine-protein phosphatase S; RTN3, reticulon3; TLR2, Toll-like receptor 2; TMEM63B, CSC1-like protein 2; UBR7, putative E3 ubiquitin-protein ligase; Y2H, yeast two-hybrid assay.

is divided into two nonfluorescent fragments that are fused to the proteins of interest. Interaction reconstitutes the entire fluorescent protein. This method allows the direct visualization of the interaction and can be used for soluble or membrane-bound proteins. In addition, several interactions can be investigated in parallel through the use of different fluorescent proteins. Since there is a delay in fluorescence formation upon reconstitution of the fluorescent proteins, and the fluorophore formation is irreversible, these methods are usually not well suited for kinetic studies. To overcome these limitations, a novel assay called NanoBiT was developed. In this assay the nanoluciferase enzyme is divided in two subunits (LgBiT and SmBiT), with low affinity for each other, that can be brought together by the two interacting proteins. The low affinity makes the interaction reversible and therefore suitable for the investigation of kinetics (Duellman et al., 2017).

Proteomics Methods. Proteomic methods aim to identify GIPs of a receptor of interest via the use of affinity purification coupled with mass spectrometry (AP-MS). This approach is usually employed as an unbiased method for screening virtually all the GIPs of a GPCR of interest. Targeted versions of the method also exist and rely on GIP identification by Western blotting. However, the requirement for specific antibodies seriously limits its application. Several strategies can be used for the affinity purification step. In co-immunoprecipitation (Co-IP), specific antibodies against the protein of interest are used for precipitating the bait from a protein lysate. As specific GPCR antibodies providing high immunoprecipitation (IP) yields are rarely available, epitope-tagged versions of the receptor of interest are often expressed in the cell type or the organism of interest and precipitated using antibodies against the tag. The main advantages of Co-IP are the purification of proteins interacting with the entire receptor (whenever possible the native receptor) in living cells or tissues and its ability to purify the entire associated protein complex. The main limitations are the necessity for specific antibodies to precipitate GPCRs, the loss of spatial-temporal information, and the use of detergents for cell lysis that might denature the GPCR of interest and, accordingly, disrupt interactions with their protein partners. For this reason, special attention must be paid to lysis conditions that efficiently solubilize the receptor while conserving the receptor's native conformation and its interactions with GIPs. For instance, detergents such as Triton and NP-40 completely denature CXCR4 (Palmesino et al., 2016), whereas 3-([3-cholamidopropyl]-dimethylammonio)-2-hydroxy-1-propanesulfonate, also called CHAPSO (Babcock et al., 2001), and n-dodecyl- β -D-maltopyranoside, also called DDM (Palmesino et al., 2016), yield the highest proportion of receptor in the native conformation. Despite this limitation, several CXCR4-interacting proteins have been identified using Co-IP approaches (see Table 2).

Alternatively, pull-down assays can be performed to purify GPCR partners from a cell or tissue lysate. This approach uses the receptor (or one of its domains) fused with an affinity tag (e.g., glutathione-S-transferase) and immobilized on beads as bait. Such *in vitro* binding assays can also be used to prove direct physical interaction between two protein partners. In this case, the bait is incubated with a purified protein instead of a cell or tissue lysate. In all methods, affinity purified proteins are systematically identified by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS). A two-step version, named tandem affinity purification (Puig et al., 2001), has also

been reported (Daulat et al., 2007) and applies to both Co-IPs or pull-downs. Although tandem affinity purification methods drastically reduce the number of false-positive identifications, they require larger amounts of starting material.

In the proximity-dependent biotin identification method (Roux et al., 2013), the bait protein is fused to a prokaryotic biotin ligase molecule that biotinylates proteins in close proximity once expressed in cells. The method can detect weak and transient interactions occurring in living cells, and detergents do not affect the results. Though the fusion of biotin ligase to the bait might alter its targeting or functions, proximity-dependent approaches were recently applied to identify a GPCR-associated protein network with a high temporal resolution. Specifically, engineered ascorbic acid peroxidase was employed in combination with quantitative proteomics to decipher β -2 adrenergic (Lobingier et al., 2017; Paek et al., 2017) and angiotensin II type 1 (Paek et al., 2017) receptor-interacting proteins.

Methods for the Identification of GPCR Phosphorylated Sites

GPCR phosphorylation is a key regulatory mechanism of receptor function (Lefkowitz, 2004). Over the past years, numerous techniques have appeared with increasing resolution to pinpoint phosphorylated residues (summarized in Table 1), which consist of serines, threonines, or tyrosines.

Radioactive Labeling. The first method that was introduced for deciphering the phosphorylation status of GPCRs is a radioactivity-based technique, consisting of culturing cells in a medium in which phosphate is replaced with its radioactive counterpart, ^{32}P , resulting in radioactive phosphorylated residues (Meisenhelder et al., 2001). After culture, cells are lysed and receptors are immunoprecipitated using specific antibodies, and then resolved by SDS-PAGE. Receptors can then be digested using an enzyme such as trypsin, and the resulting peptides are separated by two-dimensional migration using electrophoresis and chromatography. Radioactive peptides are then detected in-gel by autoradiography or using a phosphorimager, yielding a phosphorylation map for a given receptor in a given cell line (Chen et al., 2013). This method is very sensitive but does not give precise information on the number of phosphorylated sites nor their position. Radioactive labeling was initially employed to characterize CXCR4 phosphorylation upon agonist stimulation (Haribabu et al., 1997). These studies characterized the C-terminal domain as the preferred site for phosphorylation (Haribabu et al., 1997) and identified a serine cluster present in the C-terminal domain and containing two residues (Ser³³⁸, Ser³³⁹) phosphorylated following CXCL12 challenge (Orsini et al., 1999).

LC-MS/MS. More recently, radioactive labeling-based methods have been progressively supplanted by the identification of phosphorylated residues by LC-MS/MS. In this method, the GPCR of interest is digested enzymatically, with one or several proteases, to generate peptides that cover a large part of the receptor sequence. The resulting peptides are then analyzed by LC-MS/MS (Dephoure et al., 2013). Although this approach can pinpoint any phosphorylated residue with high confidence, a few limitations complicate phosphorylated residue identification. First, phosphorylation can be lost during fragmentation. Second, since phosphorylation sites have a limited level of phosphorylation, only a small percentage of

peptide is actually phosphorylated (Wu et al., 2011). Third, the identification of the phosphorylated residues in peptides with multiple adjacent phosphorylated residues can be challenging. For each modified site, a phosphorylation index can be estimated by dividing the ion signal intensity corresponding to the phosphorylated peptide by the sum of the ion signals of the phosphorylated peptide and its nonphosphorylated counterpart. Absolute quantification, and thus the stoichiometry of phosphorylation, can also be determined for each modified residue by spiking the sample with a known concentration of high-purity, heavy isotope-labeled peptides (AQUA peptides) that correspond to the phosphorylated peptide and not phosphorylated one. The respective ion signals of unlabeled and labeled peptides are then compared (Gerber et al., 2003). This powerful technology allowed a first comprehensive phosphorylation map of CXCR4, stimulated or not with CXCL12 in human embryonic kidney (HEK)293 cells: LC-MS/MS analyses identified six phosphorylated residues: Ser³²¹, Ser³²⁴, Ser³²⁵, one between Ser^{338/339/341}, one between Ser^{346/347/348}, and either Ser³⁵¹ or Ser³⁵² (Busillo et al., 2010).

Mutagenesis. Another approach that can be used as a stand-alone technique or in complement with the previously described methods is to mutate potential or predicted phosphorylated residues (into alanine or aspartate to inhibit or mimic their phosphorylation, respectively) and assess functional differences between mutated and wild-type receptor (Okamoto and Shikano, 2017). Nevertheless, introducing those mutations can potentially alter expression of the receptor, its conformation, or its cellular localization. Despite these limitations, mutagenesis approaches have shown unequivocal efficiency in identifying or validating several phosphorylated residues on CXCR4 (Orsini et al., 1999; Mueller et al., 2013) in combination with a radioactive-labeling method or use of phospho-specific antibodies. Furthermore, mutating all the serine and threonine residues to alanine in the ACKR3 C-terminus abolished β -arrestin recruitment and receptor internalization, suggesting that receptor trafficking depends on the phosphorylation of some of these residues (Canals et al., 2012).

Phospho-Specific Antibody. To be able to detect and assess phosphorylation of residues in cells or tissues, antibodies that specifically target previously identified phosphorylated residues of GPCRs can be generated by immunizing animals with purified synthetic phosphorylated peptides encompassing the phosphorylated residues (Chen et al., 2013). After selection and functional validation, those antibodies can be used in Western blot or immunohistochemistry experiments. Phosphorylation can also be indirectly detected using antibodies specific to the unphosphorylated GPCR, showing decreased binding to the target when residues are phosphorylated, and recovery of the binding by using a protein phosphatase to dephosphorylate the receptor (Hoffmann et al., 2012). Antibodies that recognize several CXCR4 phosphorylated residues [Ser^{324/325}, Ser³³⁰, Ser³³⁹, Ser^{338/339}, and Ser^{346/347} (Woerner et al., 2005; Busillo et al., 2010; Mueller et al., 2013)] have been generated and used to investigate the receptor phosphorylation status in various conditions. To our knowledge, such phospho-specific antibodies are still lacking for ACKR3.

Association of CXCR4 and ACKR3 with Canonical GPCR-Interacting Proteins

G proteins, GRKs, and β -arrestins are the protein families considered canonical GPCR-interacting proteins controlling

receptor activity or being involved in signal transduction. GPCR activity is a result of a tightly regulated balance between activation, desensitization, and resensitization events. After receptor activation and interaction with G proteins, several mechanisms integrate to trigger GPCR desensitization and/or modulate additional signaling cascades, including phosphorylation by GRKs and recruitment of β -arrestins (Penela et al., 2010b; Nogués et al., 2018).

G Proteins. CXCR4 is known to couple to the *pertussis toxin*-sensitive $G\alpha_i$ protein family that mediates most of its signaling pathways (Busillo and Benovic, 2007). However, CXCR4 can also couple to other G proteins such as $G\alpha_{12/13}$ (Tan et al., 2006; Kumar et al., 2011) and $G\alpha_q$ (Soede et al., 2001). Tan and colleagues (2001) observed that both $G\alpha_i$ and $G\alpha_{13}$, as well as $G\beta\gamma$ subunits are involved in the CXCL12-dependent migration of Jurkat T cells. Specifically, $G\alpha_i$ proteins promote migration through the activation of Rac, whereas $G\alpha_{12/13}$ proteins activate Rho. Though CXCR4 is coupled to both $G\alpha_{12/13}$ and $G\alpha_i$ proteins in Jurkat cells, such a dual coupling has not been observed in other cell lines where the receptor specifically activates one or the other G protein family (Yagi et al., 2011). In fact, *pertussis toxin* inhibited the migration of nonmetastatic breast cancer cells (MCF-7), indicating that $G\alpha_i$ activation is required. However, it did not prevent the migration of metastatic breast cancer cells such as MDA-MB-231 and SUM-159. In those cell lines, $G\alpha_{12/13}$ activation mediates CXCL12-induced migration via the activation of the Rho signaling axis (Yagi et al., 2011). Therefore, CXCR4 coupling to one or the other G protein family might depend on the abundance of GPCRs, G proteins, and downstream targets.

As an atypical chemokine receptor, ACKR3 lacks the DRY-LAIV (Asp-Arg-Tyr-Leu-Ala-Ile-Val) motif necessary for interaction with G proteins. Nevertheless, using BRET, a study showed interaction of the receptor with G proteins in transfected HEK293 cells (Levoye et al., 2009). Yet, this interaction did not lead to the activation of G proteins (Levoye et al., 2009), reinforcing the common consensus that ACKR3 is unable to activate G proteins. Consistently, other studies showed that ACKR3 signals independently of G proteins through a mechanism requiring β -arrestins (Rajagopal et al., 2010; Canals et al., 2012).

Although these findings clearly indicate that ACKR3 cannot activate G proteins in most of the cell types, a report suggested that ACKR3 might activate G proteins in two specific cellular contexts, namely primary rodent astrocytes and human glioma cells (Ödemis et al., 2012). Using [³⁵S]GTP γ S-binding assay, calcium mobilization, and *pertussis toxin*-dependent activation of downstream signaling pathways [extracellular signal-regulated kinases (ERKs 1/2 and AKT phosphorylation), this group showed an ACKR3-dependent activation of $G\alpha_{i/o}$ proteins in primary astrocytes (Ödemis et al., 2012). They also reported *pertussis toxin*-dependent migration, proliferation, and activation of downstream signaling effectors in two human glioma cell lines (A764 and U343), further suggesting an ACKR3-dependent activation of $G\alpha_{i/o}$. So far, this is the only report suggesting a possible coupling of ACKR3 with G proteins. Though these data must be further confirmed, one possibility is that such a coupling is cell type-specific. Since ACKR3 was shown to form a heterodimer with CXCR4 in transfected cell lines (Levoye et al., 2009) and CXCR4 is well known for its coupling with G proteins (vide




Fig. 1. CXCR4 and ACKR3 residues potentially subjected to post-translational modifications. Schematic representation of the C-terminal tail of CXCR4 and ACKR3, in which serine/threonine (red), tyrosine (green), and lysine (blue) residues potentially subjected to post-translational modifications are highlighted.

supra), the authors also investigated the possibility that the ACKR3-dependent activation of $G_{\alpha_{i/o}}$ proteins was mediated by the ACKR3/CXCR4 complex. However, constitutive suppression of CXCR4 expression in primary astrocytes did not influence the ability of ACKR3 to activate G proteins in the [35 S]GTP γ S-binding assay (Ödemis et al., 2012). Consistently, transient suppression of CXCR4 expression did not influence the ACKR3-dependent calcium mobilization in primary cultures of astrocytes. This suggests that ACKR3 coupling to G proteins in astroglial cells, if any, occurs independently of the ACKR3/CXCR4 complex assembly.

Although in that specific case CXCR4 did not influence ACKR3 signaling, accumulating evidence supports the hypothesis that ACKR3 might conversely influence CXCR4 signaling. Specifically, the organization of CXCR4 and ACKR3 in heterodimers appears to inhibit CXCR4 interaction with G proteins in transfected HEK293T cells, as assessed by saturation BRET (Levoye et al., 2009). In accordance with a possible influence of ACKR3 on CXCR4-dependent G_{α_i} activation, Sierro and colleagues (2007) showed that the coexpression of ACKR3 with CXCR4 hindered the fast and G protein-dependent ERK activation triggered by CXCL12 exposure. In spite of these observations demonstrating that ACKR3 influences CXCR4-dependent G protein signaling, a direct proof of the role of the physical interaction between both receptors is still missing.

GRKs. Agonist-occupied GPCRs are specifically phosphorylated by different GRKs, a family of seven serine/threonine kinases (Ribas et al., 2007; Penela et al., 2010a). GRKs 2, 3, 5, and 6 phosphorylate CXCR4 in the C-terminus, which contains 15 serine and three threonine residues that are potential phosphorylation sites (Fig. 1). At least six of these residues were shown to be phosphorylated following receptor activation by CXCL12 (Busillo et al., 2010; Barker and Benovic, 2011; Mueller et al., 2013). In HEK293 cells, Ser³²¹, Ser³²⁴, Ser³²⁵, Ser³³⁰, Ser³³⁹, and two sites between Ser³⁴⁶ and Ser³⁵² were shown to be phosphorylated in response to CXCL12 in the CXCR4 C-terminus using LC-MS/MS and phosphosite-specific

antibodies (Busillo et al., 2010). GRK6 is able to phosphorylate Ser^{324/5}, Ser³³⁹ and Ser³³⁰, the latter with slower kinetics, whereas GRK2 and GRK3 phosphorylate residues between




Fig. 2. CXCR4 C-terminus phosphosites. Schematic representation of the C-terminal tail of CXCR4, in which serine residues known to be phosphorylated are highlighted in light blue. The kinases or the extracellular stimuli responsible for the phosphorylation are also specified. Hrg, heregulin.

Ser³⁴⁶ and Ser³⁵² (Fig. 2) (Busillo et al., 2010), and specifically Ser^{346/347} (Mueller et al., 2013). Interestingly, the latter study suggested a hierarchy in such phosphorylation events, since Ser^{346/347} phosphorylation is achieved faster and is needed for the subsequent phosphorylation of Ser^{338/339} and Ser^{324/325}. Notably, ligand washout resulted in rapid Ser^{324/325} and Ser^{338/339} dephosphorylation, whereas Ser^{346/347} residues did not exhibit major dephosphorylation during the 60-minute period studied (Mueller et al., 2013). Phosphorylation of CXCR4 by different GRKs can elicit several molecular responses, such as fluctuations in intracellular calcium concentration and phosphorylation of ERKs 1 and 2, leading to integrated cellular responses. In HEK293 cells, calcium mobilization is negatively regulated by GRK2, GRK6, and β -arrestin2. On the other hand, GRKs 3 and 6 together with β -arrestins act as positive regulators of ERK1/2 (Busillo et al., 2010). Overall, these studies show nonoverlapping roles for the different GRKs in the regulation of CXCR4 signaling. These differential roles may explain distinct cell type-dependent responses to CXCL12. However, what dictates the specific GRK subtype recruitment still needs to be investigated. Changes in the normal CXCR4 phosphorylation pattern as a result of receptor mutations or altered GRK activity can lead to abnormal receptor expression and/or responsiveness that promotes aberrant cell signaling and thus can contribute to several pathologies. Deletion of Ser^{346/347} leads to a gain of CXCR4-function and decreases receptor internalization and subsequent desensitization, indicating that mutations in the far C-terminus affect CXCR4-mediated signaling (Mueller et al., 2013). In this regard, a subpopulation of patients affected by WHIM (warts, hypogammaglobulinemia, infections, and myelokathexis) syndrome, a rare primary immunodeficiency disease, display C-terminally truncated CXCR4, leading to refractoriness to desensitization and enhanced signaling (Balabanian et al., 2008). On the contrary, increased CXCR4 phosphorylation at Ser³³⁹ is associated with poor survival in adults with B-cell acute lymphoblastic leukemia and correlates with poor prognosis in acute myeloid leukemia patients (Konoplev et al., 2011; Brault et al., 2014). Altered GRK expression/activity can also impair CXCR4 phosphorylation patterns. GRK3 suppression may contribute to abnormally sustained CXCR4 signaling in classic types of glioblastoma (Woerner et al., 2012), some WHIM patients (Balabanian et al., 2008), and in triple negative breast cancer, thus potentiating CXCR4-dependent migration, invasion, and metastasis (Billard et al., 2016; Nogués et al., 2018). It is interesting to note that, although GRKs 2 and 3 share a high homology and are able to phosphorylate the same residues in CXCR4 in model cells, their function is not redundant. Whereas both CXCR4 and GRK2 levels are increased in breast cancer patients, GRK3 is decreased, suggesting a differential role for both GRKs in a cancer context (Billard et al., 2016; Nogués et al., 2018). In fact, deregulation of GRK2 potentiates several malignant features of breast cancer cells, and its level positively correlates with tumor growth and increased metastasis occurrence (Nogués et al., 2016), but whether these roles involve changes in CXCR4 modulation is still under investigation. On the other hand, impaired chemotaxis of T and B cells toward CXCL12 is noted in the absence of GRK6, whereas GRK6 deficiency potentiates neutrophil chemotactic response to CXCL12 (Fong et al., 2002; Vroon et al., 2004), suggesting that the occurrence of highly cell type-specific mechanisms in the control of the CXCL12-CXCR4-GRK6 axis. Overall, these data

indicate the complexity of CXCR4 modulation by GRKs and support the need for a better characterization of cell type- or disease-specific CXCR4-GRKs interactions.

ACKR3 has lately been the focus of many studies, in particular because of its role in cancer progression and metastasis. However, the mechanisms underlying its regulation are still not well understood, although this receptor has been shown to interact with GRKs and arrestins and to associate with other partners. The C-terminus of ACKR3 contains five serine and four threonine residues that can potentially be phosphorylated (Fig. 1). Unlike for CXCR4, little is known about their actual phosphorylation status during the activation of the receptor, as no mass spectrometry data are available to date and only few mutational studies have been conducted (Canals et al., 2012; Hoffmann et al., 2012). In fact, only one study conducted in astrocytes showed that ACKR3 is phosphorylated by GRK2, but not other GRKs, and that this phosphorylation is essential for subsequent ACKR3-operated activation of ERK1/2 and AKT pathways (Lipfert et al., 2013). This study suggests that ACKR3 is indeed phosphorylated by GRKs, but the isoform(s) involved and subsequent responses are probably cell type-dependent and remain to be investigated in detail.

Arrestins. A study revealed that site-specific phosphorylation of CXCR4 by GRK isoforms has contrasting effects upon β -arrestin recruitment: Although receptor phosphorylation at its extreme C-terminus [two residues between Ser³⁴⁶ and Ser³⁵² (Busillo et al., 2010), and specifically Ser^{346/347} (Mueller et al., 2013)] by GRK2/3 is a necessary step in β -arrestin binding; its phosphorylation by GRK6 at upstream residues [Ser^{324/5}, Ser³³⁰, and Ser³³⁹ (Busillo et al., 2010)] appears to inhibit arrestin recruitment to CXCR4 or results in a receptor/arrestin complex that adopts a conformation that is distinct from that induced by phosphorylation of extreme C-terminal residues (Oakley et al., 2000; Busillo et al., 2010; Mueller et al., 2013). Further supporting the importance of Ser^{324/5} and Ser³³⁹ phosphorylation in β -arrestin recruitment, CXCR4 truncation mutants showing impaired phosphorylation at Ser^{324/325} and Ser^{338/339} also exhibit reduced CXCL12-induced receptor internalization (Mueller et al., 2013).

β -arrestins are also scaffold proteins for several signaling molecules, thus eliciting additional β -arrestin-dependent signaling pathways (Shenoy and Lefkowitz, 2011; Peterson and Luttrell, 2017). Following the recognition of β -arrestin-dependent signaling, the notion of biased ligands that preferentially induce G protein-dependent or independent signaling has emerged (Reiter et al., 2012). Biased signaling at chemokine receptors has been exhaustively reviewed elsewhere (Steen et al., 2014). For instance, a CXCR4-derived pepducin, ATI-2341, acts as a biased CXCR4 agonist that promotes G α_i signaling but not β -arrestin signaling, in contrast to CXCL12, which activates both G protein-dependent and independent pathways (Quoyer et al., 2013; Steen et al., 2014).

Upon activation by its cognate ligands, CXCL11 and CXCL12, ACKR3 recruits β -arrestin2 both in vitro (Rajagopal et al., 2010; Benredjem et al., 2016) and in vivo (Luker et al., 2009), a process leading to receptor internalization (Canals et al., 2012), transport to lysosomes, and degradation of the receptor-bound chemokine (Luker et al., 2010; Naumann et al., 2010; Hoffmann et al., 2012). The receptor is then mainly recycled back to the plasma membrane (Luker et al., 2010) even if a partial degradation of ACKR3 can be observed (Hoffmann et al., 2012). Interestingly,

the rate of receptor internalization is faster and recycling is lower in the presence of CXCL11, compared with CXCL12 (Montpas et al., 2018).

As previously mentioned, systematic mutation of C-terminal serine/threonine residues to alanine abolished ligand-induced β -arrestin2 recruitment to ACKR3, as monitored by BRET (Canals et al., 2012) and decreased ACKR3 internalization and subsequent degradation of radiolabeled CXCL12 in HEK293 cells (Hoffmann et al., 2012). Selective mutations of the two C-terminal serine/threonine clusters to alanine revealed differences in their functional properties. Mutating Ser³³⁵, Thr³³⁸, and Thr³⁴¹ (first cluster) or Ser³⁵⁰, Thr³⁵², and Ser³⁵⁵ (second cluster) to alanine decreased CXCL12 internalization only after a 5-minute challenge but not following longer agonist receptor stimulation. Yet, only mutation of the second cluster prevented CXCL12 degradation. Furthermore, ACKR3 appears to undergo ligand-independent internalization to a much greater extent than CXCR4 (Ray et al., 2012), and residues 339–362 (the two serine/threonine clusters) are essential for this peculiar cell fate in HEK293 cells. Although numerous studies have shown that ACKR3 internalization and the resulting chemokine degradation are dependent on β -arrestin, recent findings have been challenging this consensus (Montpas et al., 2018). Specifically, this study shows that β -arrestins are dispensable to chemokine degradation, suggesting that other scaffold proteins might be involved in this process.

Association of CXCR4 with Noncanonical GPCR-Interacting Proteins

Functional Interaction of CXCR4 with Second Messenger-Dependent Kinases and Receptor Tyrosine Kinases. Accumulating evidence indicates that phosphorylation of GPCRs by second messenger-dependent kinases such as protein kinase A and protein kinase C (PKC) (Lefkowitz, 1993; Ferguson et al., 1996; Krupnick and Benovic, 1998), as well as by members of the receptor tyrosine kinase family (Delcourt et al., 2007), participates in the regulation of GPCR signaling. CXCR4 is phosphorylated by PKC at Ser^{324/5} upon CXCL12 stimulation (Busillo et al., 2010), and this kinase has also been involved in Ser^{346/7} phosphorylation (Luo et al., 2017), even though these results are not entirely consistent with a previous study using different PKC inhibitors (Mueller et al., 2013). In some glioblastoma cell types, CXCR4 is phosphorylated at Ser³³⁹ in response to the PKC activator phorbol myristate acetate (Woerner et al., 2005). This suggests that Ser³³⁹ is also a PKC phosphorylation site, and that this phosphorylation event may serve as a crosstalk mechanism between CXCR4 and GPCRs that activate $G\alpha_q$ -PKC signaling. Sphingosine 1-phosphate receptors, neurokinin-1 and lysophosphatidic acid receptors may possibly be involved in glioblastoma progression via this means (Cherry and Stella, 2014). Nevertheless, the functional impact of Ser³³⁹ phosphorylation in glioblastoma remains to be established. Likewise, epidermal growth factor (EGF) through activation of its receptor can also promote CXCR4 phosphorylation at Ser³³⁹ in glioblastoma cells (Woerner et al., 2005), and both EGF and heregulin trigger Ser^{324/325} and Ser³³⁰ phosphorylation in the breast cancer T47D cell line (Sosa et al., 2010). Interestingly, in MCF7 breast cancer cells, heregulin also promotes CXCR4 phosphorylation on tyrosine residues via epidermal growth factor receptor (EGFR), leading to β -arrestin2 association with CXCR4 and downstream activation of the PRex1/Rac1 axis.

However, it is still unclear whether the EGFR-CXCR4 functional interaction is direct or depends on other kinases (Sosa et al., 2010). In another breast cancer line, BT-474, CXCR4 is phosphorylated on tyrosine residues in response to CXCL12 and through activation of ErbB2/ErbB3 and EGFR (Sosa et al., 2010). Although the specific Tyr residue(s) phosphorylated were not identified, it is worth noting that CXCR4 displays four intracellular Tyr residues (Ahr et al., 2005). Tyr¹⁵⁷ in the third intracellular loop has been involved in CXCR4-dependent STAT3 signaling (Ahr et al., 2005), whereas Tyr¹³⁵, within the conserved DRY motif, might be involved in receptor coupling to G proteins (Rovati et al., 2007). Consistent with this hypothesis, EGFR-mediated phosphorylation of the equivalent Tyr in another GPCR (the μ -opioid receptor) has been reported to reduce coupling to G proteins (Clayton et al., 2010). Therefore, identification of tyrosine residues phosphorylated in CXCR4 might add some insight into the mechanisms by which growth factor-receptor tyrosine kinases modulate CXCR4 activity. The crosstalk between CXCR4 and ErbB2/ErbB3 and EGFR remains an interesting avenue for future research, given the involvement of both receptors in cancer.

Physical Interaction with Noncanonical GPCR-Interacting Proteins. Beside canonical GIPs, CXCR4 has been shown to interact with additional proteins that modulate CXCR4 trafficking, subcellular localization and signaling, and proteins whose functions are still unknown. CXCR-interacting proteins, the methods used for the identification of these proteins, the site of their interaction in the receptor sequence, and their functional impact are summarized in Table 2.

Proteins controlling CXCR4 localization or trafficking. Filamin A directly interacts with CXCR4 and stabilizes the receptor at the plasma membrane by blocking its endocytosis (Gómez-Moutón et al., 2015). CXCR4 association with the E3 ubiquitin ligase atrophin-interacting protein 4 (AIP4) has opposite consequences: ubiquitination of CXCR4 by AIP4 targets the receptor to multivesicular bodies, which is followed by receptor degradation. In addition, agonist treatment increases CXCR4/AIP4 interaction, as assessed by Co-IP and FRET experiments (Bhandari et al., 2009), indicating that this interaction is dynamically regulated by a receptor conformational state. In addition, the authors identified Ser³²⁴ and Ser³²⁵ as critical sites for the formation of the CXCR4/AIP4 complex upon CXCL12 exposure. The mutation of both residues to alanine drastically reduces association of AIP4 with CXCR4, whereas their mutation to aspartic acid increases this interaction. Since Ser^{324/325} are phosphorylated by GRK6 (Busillo et al., 2010), these results suggest that CXCR4 activation by CXCL12 triggers recruitment of GRK6, which in turn phosphorylates the receptor at Ser^{324/325} to promote its interaction with AIP4. AIP4 then ubiquitinates CXCR4 and mediates its degradation (Bhandari et al., 2009). Reticulon-3 is another CXCR4-interacting protein that promotes its translocation to the cytoplasm (Li et al., 2016).

Proteins modulating CXCR4 signaling and functions. HLA class II histocompatibility antigen gamma chain (CD74), a single-pass type II membrane protein that shares with CXCR4 the ability to bind to the macrophage migration inhibitory factor (MIF), was also shown to interact with CXCR4. The CXCR4/CD74 complex is involved in AKT activation (Schwartz et al., 2009). In fact, blocking either CXCR4 or CD74 inhibits MIF-induced AKT activation. Using FRET, an interaction between CXCR4 and the Toll-like receptor 2 was observed in

human monocytes upon activation by Pg-fimbria (fimbriae produced by the major pathogen associated with periodontitis named *Porphyromonas gingivalis*). Analysis of a possible crosstalk between the two receptors showed that Pg-fimbria directly binds to CXCR4 and inhibits Toll-like receptor 2-induced nuclear factor- κ B activation by *P. gingivalis* (Hajishengallis et al., 2008; Triantafyllou et al., 2008). In Jurkat cells, endolyn (CD164) coprecipitates with CXCR4 in the presence of CXCL12 presented on fibronectin (Forde et al., 2007). CXCR4-CD164 interaction participates in CXCL12-induced activation of AKT and protein kinase C zeta (PKC ζ). In fact, the downregulation of CD164 reduces the activation of both kinases measured upon exposure of Jurkat cells to CXCL12. CXCR4/CD164 interaction has been detected in additional cell lines, such as primary human ovarian surface epithelial cells stably expressing CD164 (Huang et al., 2013).

The ability of CXCR4 to promote cell migration requires deep cytoskeletal rearrangements that can be modulated by CXCR4-interacting proteins. In Jurkat J77 cells, CXCR4 constitutively associates with drebrin (Pérez-Martínez et al., 2010), a protein known to bind to F-actin and stabilize actin filaments. Drebrin is also involved in CXCR4- and CD4-dependent HIV cellular penetration (Gordón-Alonso et al., 2013). CXCR4 interacts with diaphanous-related formin-2 (mDIA2). This interaction induces cytoskeletal rearrangements that lead to nonapoptotic blebbing. mDIA2-CXCR4 interaction is only detected during nonapoptotic amoeboid blebbing and is confined to nonapoptotic blebs upon CXCL12 stimulation (Wyse et al., 2017), suggesting a fine spatiotemporal regulation of the interaction. CXCR4 also constitutively associates with the motor protein nonmuscle myosin H chain (NMMHC) via its C-terminus (Rey et al., 2002). The authors showed that NMMHC and CXCR4 are colocalized in the leading edge of migrating lymphocytes, suggesting that this association might have a role in lymphocyte migration. The P13-kinase isoform p110 γ coprecipitates with CXCR4 in CXCL12-stimulated human myeloid cells. This interaction contributes to receptor-operated integrin activation and chemotaxis of myeloid cells (Schmid et al., 2011). Finally, CXCR4 was found to be part of a junctional mechano-sensitive complex through its interaction with the platelet endothelial cell adhesion molecule (PECAM-1) (dela Paz et al., 2014).

Proteins with unknown functions. Other potential CXCR4-interacting proteins have been identified using unbiased methods. These include the lysosomal protein cation-transporting ATPase ATP13A2 (Usenovic et al., 2012) and the nuclear protein Myb-related protein B that is involved in cell cycle progression (Wang et al., 2011). In a study aimed at characterizing the human interactome by Co-IP of 1125 green fluorescent protein-tagged proteins and LC-MS/MS analysis, CXCR4 was found to coprecipitate with the potassium channel subfamily K member 1, the CSC1-like protein 2, and the vesicle transport protein GOT1B (Hein et al., 2015). In another study, CXCR4 was found to interact with the eukaryotic translation initiation factor 2B complex in an acute lymphoblastic leukemia cell line (pre-B NALM-6 cells) but not in primary lymphocytes (Palmesino et al., 2016). The interaction was negatively regulated by CXCL12 exposure and confirmed by colocalization analysis. The same study showed that CXCR4 recruits parafibromin, SH2 domain binding protein, hypothetical protein PD2, nucleophosmin, cyclin-dependent kinase 11B, receptor-type tyrosine-protein phosphatase S and galectin (Palmesino et al., 2016).

Association of ACKR3 with Noncanonical GPCR-Interacting Proteins

Unlike for CXCR4, only few proteins are described as ACKR3-interacting proteins. Given the described role of ACKR3 in cancer, several studies have addressed ACKR3 crosstalk with well known pro-oncogenic growth factor receptors. ACKR3 colocalizes with and phosphorylates EGFR in breast and prostate cancer cells (Singh and Lokeshwar, 2011; Salazar et al., 2014; Kallifatidis et al., 2016), via cell type-specific mechanisms. However, a potential role for EGFR in ACKR3 crossactivation was not assessed in these studies. Some reports also suggest a possible functional interaction between ACKR3 and transforming growth factor beta (Rath et al., 2015) or vascular endothelial growth factor (Singh and Lokeshwar, 2011) receptors, but whether they involve physical interaction with ACKR3 and/or ACKR3 phosphorylation and activation was not assessed. ACKR3 weakly interacts with the MIF receptor CD74 (Alampour-Rajabi et al., 2015). Moreover, ACKR3 colocalizes with PECAM-1, the cell adhesion molecule required for leukocyte transendothelial migration in human coronary artery endothelial cells (dela Paz et al., 2014). Using a membrane yeast two-hybrid assay screen, cation-transporting ATPase ATP13A2 was identified as a putative ACKR3-interacting protein (Usenovic et al., 2012). In the study aimed at characterizing the human interactome of 1125 green fluorescent protein-tagged proteins, ACKR3 was found to interact with the gap junction β -2 protein, the 54S ribosomal protein L4, 54S ribosomal protein L4, mitochondrial (MRPL4), different ATP synthases (ATP5H, ATP5B, ATP5A1, ATP50), ACKR3 itself, the caspase separin ESPL1, the probable E3 ubiquitin-protein ligase HECTD2, and the putative E3 ubiquitin-protein ligase UBR7 (Hein et al., 2015). Ubiquitination is an essential mechanism of receptor regulation (Marchese and Benovic, 2001; Shenoy, 2007). ACKR3 can undergo ubiquitination in an agonist-dependent and -independent manner to regulate receptor trafficking. Ubiquitination is promoted by three enzymes, E1, E2, and E3, that ubiquitinate proteins on lysine residues (Dores and Trejo, 2012; Alonso and Friedman, 2013). Unexpectedly, ACKR3 is ubiquitinated by E3 ubiquitin ligase (E3) in the absence of an agonist and undergoes deubiquitination upon CXCL12 activation (Canals et al., 2012). Mutation of the five lysines in the receptor C-terminus to alanine, to prevent ubiquitination, impaired ACKR3 cell-trafficking and decreased ACKR3-mediated CXCL12 degradation (Hoffmann et al., 2012).

Conclusions

The identification of GPCR-interacting proteins and residues subjected to post-translational modification is of utmost importance. Several techniques are currently available to decipher the GPCR interactome and phosphorylation profile. These techniques have been successfully applied to CXCR4, revealing important interacting proteins as well as key residues involved in the regulation of receptor-mediated signal transduction. In contrast, ACKR3 interactome and phosphorylation sites have not been systematically investigated. Unbiased studies of the ACKR3 interactome and its phosphorylated residues and their control by ACKR3 ligands should open new avenues in the understanding of ACKR3 pathophysiological functions and the underlying signaling mechanisms.

Acknowledgments

This review is part of the minireview series “From insight to modulation of CXCR4 and ACKR3 (CXCR7) function.” We thank all our colleagues from the ONCORNET consortium for continuous scientific discussions and insights.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Fumagalli, Zarca, Neves, Caspar, Hill, Mayor, Smit, Marin.

References

- Ahr B, Denizot M, Robert-Hebmann V, Brelot A, and Biard-Piechaczyk M (2005) Identification of the cytoplasmic domains of CXCR4 involved in Jak2 and STAT3 phosphorylation. *J Biol Chem* **280**:6692–6700.
- Alampour-Rajabi S, El Bounkari O, Rot A, Müller-Newen G, Bachelier F, Gawaz M, Weber C, Schober A, and Bernhagen J (2015) MIF interacts with CXCR7 to promote receptor internalization, ERK1/2 and ZAP-70 signaling, and lymphocyte chemotaxis. *FASEB J* **29**:4497–4511.
- Alonso V and Friedman PA (2013) Minireview: ubiquitination-regulated G protein-coupled receptor signaling and trafficking. *Mol Endocrinol* **27**:558–572.
- Babcock GJ, Mirzabekov T, Wojtowicz W, and Sodroski J (2001) Ligand binding characteristics of CXCR4 incorporated into paramagnetic proteoliposomes. *J Biol Chem* **276**:38433–38440.
- Balabanian K, Levoe A, Klemm L, Lagane B, Hermine O, Harriague J, Baleux F, Arenzana-Seisdedos F, and Bachelier F (2008) Leukocyte analysis from WHIM syndrome patients reveals a pivotal role for GRK3 in CXCR4 signaling. *J Clin Invest* **118**:1074–1084.
- Banisadr G, Fontanges P, Haour F, Kitabgi P, Rostène W, and Mélik Parsadaniantz S (2002) Neuroanatomical distribution of CXCR4 in adult rat brain and its localization in cholinergic and dopaminergic neurons. *Eur J Neurosci* **16**:1661–1671.
- Banisadr G, Podojil JR, Miller SD, and Miller RJ (2016) Pattern of CXCR7 gene expression in mouse brain under normal and inflammatory conditions. *J Neuroimmune Pharmacol* **11**:26–35.
- Barker BL and Benovic JL (2011) G protein-coupled receptor kinase 5 phosphorylation of hip regulates internalization of the chemokine receptor CXCR4. *Biochemistry* **50**:6933–6941.
- Benredjem B, Girard M, Rhains D, St-Onge G, and Heveker N (2016) Mutational analysis of atypical chemokine receptor 3 (ACKR3/CXCR7) interaction with its chemokine ligands CXCL11 and CXCL12. *J Biol Chem* **292**:31–42.
- Berachovich RD, Zabel BA, Lewen S, Walters MJ, Ebsworth K, Wang Y, Jaen JC, and Schall TJ (2014) Endothelial expression of CXCR7 and the regulation of systemic CXCL12 levels. *Immunology* **141**:111–122.
- Bhandari D, Robia SL, and Marchese A (2009) The E3 ubiquitin ligase atrophin interacting protein 4 binds directly to the chemokine receptor CXCR4 via a novel WW domain-mediated interaction. *Mol Biol Cell* **20**:1324–1339.
- Billard MJ, Fitzhugh DJ, Parker JS, Brozowski JM, McGinnis MW, Timoshchenko RG, Serafini DS, Liningner R, Klaubner-Demore N, Sahagian G, et al. (2016) G protein-coupled receptor kinase 3 regulates breast cancer migration, invasion, and metastasis. *PLoS One* **11**:e0152856.
- Bockaert J, Dumuis A, Fagni L, and Marin P (2004) GPCR-GIP networks: a first step in the discovery of new therapeutic drugs? *Curr Opin Drug Discov Devel* **7**:649–657.
- Brault L, Rovó A, Decker S, Dierks C, Tzankov A, and Schwaller J (2014) CXCR4-SERINE339 regulates cellular adhesion, retention and mobilization, and is a marker for poor prognosis in acute myeloid leukemia. *Leukemia* **28**:566–576.
- Busillo JM, Armando S, Sengupta R, Meucci O, Bouvier M, and Benovic JL (2010) Site-specific phosphorylation of CXCR4 is dynamically regulated by multiple kinases and results in differential modulation of CXCR4 signaling. *J Biol Chem* **285**:7805–7817.
- Busillo JM and Benovic JL (2007) Regulation of CXCR4 signaling. *Biochim Biophys Acta* **1768**:952–963.
- Canals M, Scholten DJ, de Munnik S, Han MKL, Smit MJ, and Leurs R (2012) Ubiquitination of CXCR7 controls receptor trafficking. *PLoS One* **7**:e34192.
- Chen Y-J, Oldfield S, Butcher AJ, Tobin AB, Saxena K, Gurevich VV, Benovic JL, Henderson G, and Kelly E (2013) Identification of phosphorylation sites in the COOH-terminal tail of the μ -opioid receptor. *J Neurochem* **124**:189–199.
- Cherry AE and Stella N (2014) G protein-coupled receptors as oncogenic signals in glioma: emerging therapeutic avenues. *Neuroscience* **278**:222–236.
- Clayton CC, Bruchas MR, Lee ML, and Chavkin C (2010) Phosphorylation of the μ -opioid receptor at tyrosine 166 (Tyr3.51) in the DRY motif reduces agonist efficacy. *Mol Pharmacol* **77**:339–347.
- Clegg RM (1995) Fluorescence resonance energy transfer. *Curr Opin Biotechnol* **6**:103–110.
- Dault AM, Maurice P, Froment C, Guillaume J-L, Broussard C, Monsarrat M, Delagrèze P, and Jockers R (2007) Purification and identification of G protein-coupled receptor protein complexes under native conditions. *Mol Cell Proteomics* **6**:835–844.
- dela Paz NG, Melchior B, Shayo FY, and Frangos JA (2014) Heparan sulfates mediate the interaction between platelet endothelial cell adhesion molecule-1 (PECAM-1) and the Gαq/11 subunits of heterotrimeric G proteins. *J Biol Chem* **289**:7413–7424.
- Delcourt N, Bockaert J, and Marin P (2007) GPCR-jacking: from a new route in RTK signalling to a new concept in GPCR activation. *Trends Pharmacol Sci* **28**:602–607.
- Dephoure N, Gould KL, Gygi SP, and Kellogg DR (2013) Mapping and analysis of phosphorylation sites: a quick guide for cell biologists. *Mol Biol Cell* **24**:535–542.
- Dores MR and Trejo J (2012) Ubiquitination of G protein-coupled receptors: functional implications and drug discovery. *Mol Pharmacol* **82**:563–570.
- Duellman SJ, Machleidt T, Cali JJ, and Vidugiriene J (2017) Cell-based, bioluminescent assay for monitoring the interaction between PCSK9 and the LDL receptor. *J Lipid Res* **58**:1722–1729.
- Ferguson SSG, Downey WE III, Colapietro AM, Barak LS, Ménard L, and Caron MG (1996) Role of β -arrestin in mediating agonist-promoted G protein-coupled receptor internalization. *Science* **271**:363–366.
- Ferré S, Casadó V, Devi LA, Filizola M, Jockers R, Lohse MJ, Milligan G, Pin J-P, and Guitart X (2014) G protein-coupled receptor oligomerization revisited: functional and pharmacological perspectives. *Pharmacol Rev* **66**:413–434.
- Fields S and Song O (1989) A novel genetic system to detect protein-protein interactions. *Nature* **340**:245–246.
- Fong AM, Premont RT, Richardson RM, Yu Y-RA, Lefkowitz RJ, and Patel DD (2002) Defective lymphocyte chemotaxis in β -arrestin2- and GRK6-deficient mice. *Proc Natl Acad Sci USA* **99**:7478–7483.
- Forde S, Tye BJ, Newey SE, Roubelakis M, Smythe J, McGuckin CP, Pettengell R, and Watt SM (2007) Endolyn (CD114) modulates the CXCL12-mediated migration of umbilical cord blood CD133+ cells. *Blood* **109**:1825–1833.
- Fredriksson S, Gullberg M, Jarvius J, Olsson C, Pietras K, Gústafsdóttir SM, Östman A, and Landegren U (2002) Protein detection using proximity-dependent DNA ligation assays. *Nat Biotechnol* **20**:473–477.
- Gerber SA, Rush J, Stemman O, Kirschner MW, and Gygi SP (2003) Absolute quantification of proteins and phosphoproteins from cell lysates by tandem MS. *Proc Natl Acad Sci USA* **100**:6940–6945.
- Gómez-Moutón C, Fischer T, Peregil RM, Jiménez-Baranda S, Stossel TP, Nakamura F, and Mañes S (2015) Filamin A interaction with the CXCR4 third intracellular loop regulates endocytosis and signaling of WT and WHIM-like receptors. *Blood* **125**:1116–1125.
- Gordón-Alonso M, Rocha-Perugini V, Álvarez S, Ursa Á, Izquierdo-Useros N, Martínez-Picado J, Muñoz-Fernández MA, and Sánchez-Madrid F (2013) Actin-binding protein drebrin regulates HIV-1-triggered actin polymerization and viral infection. *J Biol Chem* **288**:28382–28397.
- Hajishengallis G, Wang M, Liang S, Triantafilou M, and Triantafilou K (2008) Pathogen induction of CXCR4/TLR2 cross-talk impairs host defense function. *Proc Natl Acad Sci USA* **105**:13532–13537.
- Haribabu B, Richardson RM, Fisher I, Sozzani S, Peiper SC, Horuk R, Ali H, and Snyderman R (1997) Regulation of human chemokine receptors CXCR4. Role of phosphorylation in desensitization and internalization. *J Biol Chem* **272**:28726–28731.
- Hein MY, Hubner NC, Poser I, Cox J, Nagaraj N, Toyoda Y, Gak IA, Weisswange I, Mansfeld J, Buchholz F, et al. (2015) A human interactome in three quantitative dimensions organized by stoichiometries and abundances. *Cell* **163**:712–723.
- Hoffmann F, Müller W, Schütz D, Penfold ME, Wong YH, Schulz S, and Stumm R (2012) Rapid uptake and degradation of CXCL12 depend on CXCR7 carboxyl-terminal serine/threonine residues. *J Biol Chem* **287**:28362–28377.
- Hu C-D, Chinenov Y, and Kerppola TK (2002) Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol Cell* **9**:789–798.
- Hu C-D and Kerppola TK (2003) Simultaneous visualization of multiple protein interactions in living cells using multicolor fluorescence complementation analysis. *Nat Biotechnol* **21**:539–545.
- Huang A-F, Chen M-W, Huang S-M, Kao C-L, Lai H-C, and Chan JY-H (2013) CD164 regulates the tumorigenesis of ovarian surface epithelial cells through the SDF-1 α /CXCR4 axis. *Mol Cancer* **12**:115.
- Kallifatidis G, Munoz D, Singh RK, Salazar N, Hoy JJ, and Lokeswar BL (2016) β -Arrestin-2 counters CXCR7-mediated EGFR transactivation and proliferation. *Mol Cancer Res* **14**:493–503.
- Konoplev S, Jorgensen JL, Thomas DA, Lin E, Burger J, Kantarjian HM, Andreeff M, Medeiros LJ, and Konopleva M (2011) Phosphorylated CXCR4 is associated with poor survival in adults with B-acute lymphoblastic leukemia. *Cancer* **117**:4689–4695.
- Krupnick JG and Benovic JL (1998) The role of receptor kinases and arrestins in G protein-coupled receptor regulation. *Annu Rev Pharmacol Toxicol* **38**:289–319.
- Kumar A, Kremer KN, Dominguez D, Tadi M, and Hedin KE (2011) G α 13 and Rho mediate endosomal trafficking of CXCR4 into Rab11+ vesicles upon stromal cell-derived factor-1 stimulation. *J Immunol* **186**:951–958.
- Lefkowitz RJ (1993) G protein-coupled receptor kinases. *Cell* **74**:409–412.
- Lefkowitz RJ (2004) Historical review: a brief history and personal retrospective of seven-transmembrane receptors. *Trends Pharmacol Sci* **25**:413–422.
- Levoe A, Balabanian K, Baleux F, Bachelier F, and Lagane B (2009) CXCR7 heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein signaling. *Blood* **113**:6085–6093.
- Li H, Liang R, Lu Y, Wang M, and Li Z (2016) RTN3 regulates the expression level of chemokine receptor CXCR4 and is required for migration of primordial germ cells. *Int J Mol Sci* **17**:382–395.
- Lievens S, Gerlo S, Lemmens I, De Clercq DJH, Risseuw MDP, Vanderroost N, De Smet A-S, Ruysssink E, Chevot E, Van Calenberg S, et al. (2014) Kinase Substrate Sensor (KISS), a mammalian in situ protein interaction sensor. *Mol Cell Proteomics* **13**:3332–3342.
- Lipfert J, Ödemis V, and Engle J (2013) Grk2 is an essential regulator of CXCR7 signalling in astrocytes. *Cell Mol Neurobiol* **33**:111–118.
- Lobingier BT, Hüttenhain R, Eichel K, Miller KB, Ting AY, von Zastrow M, and Krogan NJ (2017) An approach to spatiotemporally resolve protein interaction networks in living cells. *Cell* **169**:350–360.e12.
- Luker KE, Gupta M, Steele JM, Foerster BR, and Luker GD (2009) Imaging ligand-dependent activation of CXCR7. *Neoplasia* **11**:1022–1035.
- Luker KE, Steele JM, Mihalko LA, Ray P, and Luker GD (2010) Constitutive and chemokine-dependent internalization and recycling of CXCR7 in breast cancer cells to degrade chemokine ligands. *Oncogene* **29**:4599–4610.
- Luo J, Busillo JM, Stumm R, and Benovic JL (2017) G protein-coupled receptor kinase 3 and protein kinase C phosphorylate the distal C-terminal tail of the

- chemokine receptor CXCR4 and mediate recruitment of β -arrestin. *Mol Pharmacol* **91**:554–566.
- Magalhaes AC, Dunn H, and Ferguson SSG (2012) Regulation of GPCR activity, trafficking and localization by GPCR-interacting proteins. *Br J Pharmacol* **165**: 1717–1736.
- Marchese A and Benovic JL (2001) Agonist-promoted ubiquitination of the G protein-coupled receptor CXCR4 mediates lysosomal sorting. *J Biol Chem* **276**: 45509–45512.
- Maurice P, Guillaume J-L, Benleulmi-Chaachoua A, Daulat AM, Kamal M, and Jockers R (2011) GPCR-interacting proteins, major players of GPCR function. *Adv Pharmacol* **62**:349–380.
- Meisenhelder J, Hunter T, and van der Geer P (2001) Phosphopeptide mapping and identification of phosphorylation sites. *Curr Protoc Protein Sci* **Chapter 13**: Unit13.9.
- Montpas N, St-Onge G, Nama N, Rhainds D, Benredjem B, Girard M, Hickson G, Pons V, and Heveker N (2018) Ligand-specific conformational transitions and intracellular transport are required for atypical chemokine receptor 3-mediated chemokine scavenging. *J Biol Chem* **293**:893–905.
- Mueller W, Schütz D, Nagel F, Schulz S, and Stumm R (2013) Hierarchical organization of multi-site phosphorylation at the CXCR4 C terminus. *PLoS One* **8**: e64975.
- Naumann U, Cameroni E, Pruenster M, Mahabaleswar H, Raz E, Zerwes H-G, Rot A, and Thelen M (2010) CXCR7 functions as a scavenger for CXCL12 and CXCL11. *PLoS One* **5**:e9175.
- Nazari A, Khorramdelazad H, and Hassanshahi G (2017) Biological/pathological functions of the CXCL12/CXCR4/CXCR7 axes in the pathogenesis of bladder cancer. *Int J Clin Oncol* **22**:991–1000.
- Nogués L, Palacios-García J, Reglero C, Rivas V, Neves M, Ribas C, Penela P, and Mayor F Jr (2018) G protein-coupled receptor kinases (GRKs) in tumorigenesis and cancer progression: GPCR regulators and signaling hubs. *Semin Cancer Biol* **48**:78–90.
- Nogués L, Reglero C, Rivas V, Salcedo A, Lafarga V, Neves M, Ramos P, Mendiola M, Berjón A, Stamatakis K, et al. (2016) G protein-coupled receptor kinase 2 (GRK2) promotes breast tumorigenesis through a HDAC6-pin1 axis. *EBioMedicine* **13**: 132–145.
- Oakley RH, Laporte SA, Holt JA, Caron MG, and Barak LS (2000) Differential affinities of visual arrestin, β arrestin1, and β arrestin2 for G protein-coupled receptors delineate two major classes of receptors. *J Biol Chem* **275**:17201–17210.
- Odemis V, Boosmann K, Heinen A, Kürby P, and Engele J (2010) CXCR7 is an active component of SDF-1 signalling in astrocytes and Schwann cells. *J Cell Sci* **123**: 1081–1088.
- Ödemis V, Lipfert J, Kraft R, Hajek P, Abraham G, Hattermann K, Mentlein R, and Engele J (2012) The presumed atypical chemokine receptor CXCR7 signals through G(i/o) proteins in primary rodent astrocytes and human glioma cells. *Glia* **60**:372–381.
- Okamoto Y and Shikano S (2017) Differential phosphorylation signals control endocytosis of GPR15. *Mol Biol Cell* **28**:2267–2281.
- Orsini MJ, Parent JL, Mundell SJ, Marchese A, and Benovic JL (1999) Trafficking of the HIV coreceptor CXCR4. Role of arrestins and identification of residues in the c-terminal tail that mediate receptor internalization [published correction appears in *J Biol Chem* (2000) 275:25876]. *J Biol Chem* **274**:31076–31086.
- Paek J, Kalocsay M, Staus DP, Winkler L, Pascolutti R, Paulo JA, Gygi SP, and Kruse AC (2017) Multidimensional tracking of GPCR signaling via peroxidase-catalyzed proximity labeling. *Cell* **169**:338–349.e11.
- Palmesino E, Apuzzo T, Thelen S, Mueller B, Langen H, and Thelen M (2016) Association of eukaryotic translation initiation factor eIF2B with fully solubilized CXCR4. *J Leukoc Biol* **99**:971–978.
- Penela P, Murga C, Ribas C, Lafarga V, and Mayor F Jr (2010a) The complex G protein-coupled receptor kinase 2 (GRK2) interactome unveils new physiopathological targets. *Br J Pharmacol* **160**:821–832.
- Penela P, Rivas V, Salcedo A, and Mayor F Jr (2010b) G protein-coupled receptor kinase 2 (GRK2) modulation and cell cycle progression. *Proc Natl Acad Sci USA* **107**:1118–1123.
- Pérez-Martínez M, Gordón-Alonso M, Cabrero JR, Barrero-Villar M, Rey M, Mittelbrunn M, Lamana A, Morlino G, Calabia C, Yamazaki H, et al. (2010) F-actin-binding protein drebrin regulates CXCR4 recruitment to the immune synapse. *J Cell Sci* **123**:1160–1170.
- Peterson YK and Luttrell LM (2017) The diverse roles of arrestin scaffolds in G protein-coupled receptor signaling. *Pharmacol Rev* **69**:256–297.
- Petschnigg J, Groisman B, Kotlyar M, Taipale M, Zheng Y, Kurat CF, Sayad A, Sierra JR, Mattiazzi Usaj M, Snider J, et al. (2014) The mammalian-membrane two-hybrid assay (MaMTH) for probing membrane-protein interactions in human cells. *Nat Methods* **11**:585–592.
- Puig O, Caspary F, Rigaut G, Rutz B, Bouveret E, Bragado-Nilsson E, Wilm M, and Séraphin B (2001) The tandem affinity purification (TAP) method: a general procedure of protein complex purification. *Methods* **24**:218–229.
- Quoyer J, Janz JM, Luo J, Ren Y, Armando S, Lukashova V, Benovic JL, Carlson KE, Hunt SW III, and Bouvier M (2013) Pepducin targeting the C-X-C chemokine receptor type 4 acts as a biased agonist favoring activation of the inhibitory G protein. *Proc Natl Acad Sci USA* **110**:E5088–E5097.
- Rajagopal S, Kim J, Ahn S, Craig S, Lam CM, Gerard NP, Gerard C, and Lefkowitz RJ (2010) Beta-arrestin- but not G protein-mediated signaling by the “decoy” receptor CXCR7. *Proc Natl Acad Sci USA* **107**:628–632.
- Rath D, Chatterjee M, Borst O, Müller K, Langer H, Mack AF, Schwab M, Winter S, Gawaz M, and Geisler T (2015) Platelet surface expression of stromal cell-derived factor-1 receptors CXCR4 and CXCR7 is associated with clinical outcomes in patients with coronary artery disease. *J Thromb Haemost* **13**:719–728.
- Ray P, Mihalko LA, Coggins NL, Moudgil P, Ehrlich A, Luker KE, and Luker GD (2012) Carboxy-terminus of CXCR7 regulates receptor localization and function. *Int J Biochem Cell Biol* **44**:669–678.
- Reiter E, Ahn S, Shukla AK, and Lefkowitz RJ (2012) Molecular mechanism of β -arrestin-biased agonism at seven-transmembrane receptors. *Annu Rev Pharmacol Toxicol* **52**:179–197.
- Rey M, Vicente-Manzanares M, Viedma F, Yáñez-Mó M, Urzainqui A, Barreiro O, Vázquez J, and Sánchez-Madrid F (2002) Cutting edge: association of the motor protein nonmuscle myosin heavy chain-IIA with the C terminus of the chemokine receptor CXCR4 in T lymphocytes. *J Immunol* **169**:5410–5414.
- Ribas C, Penela P, Murga C, Salcedo A, García-Hoz C, Jurado-Pueyo M, Aymerich I, and Mayor F (2007) The G protein-coupled receptor kinase (GRK) interactome: Role of GRKs in GPCR regulation and signaling. *Biochim Biophys Acta* **1768**: 913–922.
- Ritter SL and Hall RA (2009) Fine-tuning of GPCR activity by receptor-interacting proteins. *Nat Rev Mol Cell Biol* **10**:819–830.
- Roux KJ, Kim DI, and Burke B (2013) BioID: a screen for protein-protein interactions. *Curr Protoc Protein Sci* **74**:19.23.1–19.23.14.
- Rovati GE, Capra V, and Neubig RR (2007) The highly conserved DRY motif of class A G protein-coupled receptors: beyond the ground state. *Mol Pharmacol* **71**: 959–964.
- Salazar N, Muñoz D, Kallifatidis G, Singh RK, Jordà M, and Lokeshwar BL (2014) The chemokine receptor CXCR7 interacts with EGFR to promote breast cancer cell proliferation. *Mol Cancer* **13**:198.
- Sánchez-Alcañiz JA, Haeghe S, Mueller W, Pla R, Mackay F, Schulz S, López-Bendito G, Stumm R, and Marin O (2011) Cxcr7 controls neuronal migration by regulating chemokine responsiveness. *Neuron* **69**:77–90.
- Schmid MC, Avraamides CJ, Dippold HC, Franco I, Foubert P, Ellies LG, Acevedo LM, Manglicmot JRE, Song X, Wrasidlo W, et al. (2011) Receptor tyrosine kinases and TLR/IL1Rs unexpectedly activate myeloid cell PI3ky, a single convergent point promoting tumor inflammation and progression. *Cancer Cell* **19**:715–727.
- Schwartz V, Lue H, Kraemer S, Korbel J, Krohn R, Ohl K, Bucala R, Weber C, and Bernhagen J (2009) A functional heteromeric MIF receptor formed by CD74 and CXCR4. *FEBS Lett* **583**:2749–2757.
- Shenoy SK (2007) Seven-transmembrane receptors and ubiquitination. *Circ Res* **100**: 1142–1154.
- Shenoy SK and Lefkowitz RJ (2011) β -Arrestin-mediated receptor trafficking and signal transduction. *Trends Pharmacol Sci* **32**:521–533.
- Sierro F, Biben C, Martínez-Muñoz L, Mellado M, Ransohoff RM, Li M, Woehl B, Leung H, Groom J, Batten M, et al. (2007) Disrupted cardiac development but normal hematopoiesis in mice deficient in the second CXCL12/SDF-1 receptor, CXCR7. *Proc Natl Acad Sci USA* **104**:14759–14764.
- Singh RK and Lokeshwar BL (2011) The IL-8-regulated chemokine receptor CXCR7 stimulates EGFR signaling to promote prostate cancer growth. *Cancer Res* **71**: 3268–3277.
- Soede RD, Zeelenberg IS, Wijnands YM, Kamp M, and Roos E (2001) Stromal cell-derived factor-1-induced LFA-1 activation during in vivo migration of T cell hybridoma cells requires Gq/11, RhoA, and myosin, as well as Gi and Cdc42. *J Immunol* **166**:4293–4301.
- Sosa MS, Lopez-Haber C, Yang C, Wang H, Lemmon MA, Busillo JM, Luo J, Benovic JL, Klein-Szanto A, Yagi H, et al. (2010) Identification of the Rac-GEF P-Rex1 as an essential mediator of ErbB signaling in breast cancer. *Mol Cell* **40**: 877–892.
- Staglar I, Korostensky C, Johnsson N, and te Heesen S (1998) A genetic system based on split-ubiquitin for the analysis of interactions between membrane proteins in vivo. *Proc Natl Acad Sci USA* **95**:5187–5192.
- Steen A, Larsen O, Thiele S, and Rosenkilde MM (2014) Biased and g protein-independent signaling of chemokine receptors. *Front Immunol* **5**:277.
- Sun X, Cheng G, Hao M, Zheng J, Zhou X, Zhang J, Taichman RS, Pienta KJ, and Wang J (2010) CXCL12 / CXCR4 / CXCR7 chemokine axis and cancer progression. *Cancer Metastasis Rev* **29**:709–722.
- Sun Y, Day RN, and Periasamy A (2011) Investigating protein-protein interactions in living cells using fluorescence lifetime imaging microscopy. *Nat Protoc* **6**: 1324–1340.
- Sun Y, Hays NM, Periasamy A, Davidson MW, and Day RN (2012) Monitoring protein interactions in living cells with fluorescence lifetime imaging microscopy. *Methods Enzymol* **504**:371–391.
- Tan W, Martin D, and Gutkind JS (2006) The G α 13-Rho signaling axis is required for SDF-1-induced migration through CXCR4. *J Biol Chem* **281**: 39542–39549.
- Tobin AB (2008) G-protein-coupled receptor phosphorylation: Where, when and by whom. *Br J Pharmacol* **153** (Suppl 1):S167–S176.
- Triantafilou M, Lepper PM, Briault CD, Ahmed MAE, Dmochowski JM, Schumann C, and Triantafilou K (2008) Chemokine receptor 4 (CXCR4) is part of the lipopolysaccharide “sensing apparatus”. *Eur J Immunol* **38**:192–203.
- Usenovic M, Knight AL, Ray A, Wong V, Brown KR, Caldwell GA, Caldwell KA, Staglar I, and Krainc D (2012) Identification of novel ATP13A2 interactors and their role in α -synuclein misfolding and toxicity. *Hum Mol Genet* **21**: 3785–3794.
- Volin MV, Joseph L, Shockley MS, and Davies PF (1998) Chemokine receptor CXCR4 expression in endothelium. *Biochem Biophys Res Commun* **242**:46–53.
- Vron A, Heijnen CJ, Raatgever R, Touw IP, Ploemacher RE, Premont RT, and Kavelaars A (2004) GRK6 deficiency is associated with enhanced CXCR4-mediated neutrophil chemotaxis in vitro and impaired responsiveness to G-CSF in vivo. *J Leukoc Biol* **75**:698–704.
- Wang J, Huo K, Ma L, Tang L, Li D, Huang X, Yuan Y, Li C, Wang W, Guan W, et al. (2011) Toward an understanding of the protein interaction network of the human liver [published correction appears in *Mol Syst Biol* (2017) 13:965]. *Mol Syst Biol* **7**: 536.
- Woerner BM, Luo J, Brown KR, Jackson E, Dahiya SM, Mischel P, Benovic JL, Pivnicka-Worms D, and Rubin JB (2012) Suppression of G-protein-coupled receptor kinase 3 expression is a feature of classical GBM that is required for maximal growth. *Mol Cancer Res* **10**:156–166.

- Woerner BM, Warrington NM, Kung AL, Perry A, and Rubin JB (2005) Widespread CXCR4 activation in astrocytomas revealed by phospho-CXCR4-specific antibodies. *Cancer Res* **65**:11392–11399.
- Wu R, Haas W, Dephore N, Huttlin EL, Zhai B, Sowa ME, and Gygi SP (2011) A large-scale method to measure absolute protein phosphorylation stoichiometries. *Nat Methods* **8**:677–683.
- Wyse MM, Goicoechea S, Garcia-Mata R, Nestor-Kalinoski AL, and Eisenmann KM (2017) mDia2 and CXCL12/CXCR4 chemokine signaling intersect to drive tumor cell amoeboid morphological transitions. *Biochem Biophys Res Commun* **484**:255–261.
- Xu Y, Piston DW, and Johnson CH (1999) A bioluminescence resonance energy transfer (BRET) system: application to interacting circadian clock proteins. *Proc Natl Acad Sci USA* **96**:151–156.
- Yagi H, Tan W, Dillenburg-Pilla P, Armando S, Amornphimoltham P, Simaan M, Weigert R, Molinolo AA, Bouvier M, and Gutkind JS (2011) A synthetic biology approach reveals a CXCR4-G13-Rho signaling axis driving transendothelial migration of metastatic breast cancer cells. *Sci Signal* **4**:ra60.
- Zhao H, Guo L, Zhao H, Zhao J, Weng H, and Zhao B (2015) CXCR4 over-expression and survival in cancer: a system review and meta-analysis. *Oncotarget* **6**: 5022–5040.

Address correspondence to: Dr. Amos Fumagalli, Institut de Génomique Fonctionnelle (CNRS), 141, rue de la Cardonille, 34094 Montpellier Cedex 5, France. E-mail: amos.fumagalli@igf.cnrs.fr; or Dr. Philippe Marin, Institut de Génomique Fonctionnelle (CNRS), 141, rue de la Cardonille, 34094 Montpellier Cedex 5, France. E-mail: philippe.marin@igf.cnrs.fr
