

Mammalian skeletal muscle voltage-gated sodium channels are affected by scorpion depressant  
'insect-selective' toxins when preconditioned

by

Lior Cohen, Yael Troub, Michael Turkov, Nicolas Gilles, Nitza Ilan, Morris Benveniste, Dalia  
Gordon and Michael Gurevitz

from the

Departments of Plant Sciences, George S. Wise Faculty of Life Sciences (LC, MT, NI, DG, MG), and  
Physiology and Pharmacology, Sackler School of Medicine (YT, MB), Tel-Aviv University, Ramat-Aviv  
69978, Tel-Aviv, Israel; CEA, Department d'Ingenierie et d'Etudes des Proteines, C.E. Saclay, F-91191  
Gif Sur Yvette Cedex, France (NG)

Running title: Scorpion depressant insect-toxins are active on rNa<sub>v</sub>1.4

Corresponding author: Michael Gurevitz, Department of Plant Sciences, George S. Wise Faculty of Life Sciences, Tel-Aviv University, Ramat-Aviv 69978, Tel-Aviv, Israel.

Phone: 972-3-6409844 Fax: 972-3-6406100 E-mail addresses: mamgur@post.tau.ac.il and

Number of text pages: 18

Number of tables: 2

Number of figures: 6

Number of references: 41

Number of words in Abstract: 250

Number of words in Introduction: 744

Number of words in Discussion: 1013

Abbreviations: C<sub>ss</sub>4, *Centruroides suffusus suffusus* toxin 4; LqhIT2 and Lqh-dprIT3, *Leiurus quinquestriatus hebraeus* anti-insect depressant toxins; Na<sub>v</sub>s, voltage gated sodium channels; PP, depolarizing prepulse

## ABSTRACT

Amongst scorpion  $\beta$ - and  $\alpha$ -toxins that modify the activation and inactivation of voltage-gated sodium channels ( $\text{Na}_v\text{s}$ ), depressant  $\beta$ -toxins have traditionally been classified as anti-insect selective on the basis of toxicity assays and lack of binding and effect on mammalian  $\text{Na}_v\text{s}$ . Here we show that the depressant  $\beta$ -toxins LqhIT2 and Lqh-dprIT3 from *Leiurus quinquestriatus hebraeus* (Lqh) bind in nM affinity to receptor site-4 on rat skeletal muscle  $\text{Na}_v\text{s}$ , but their effect on the gating properties can be viewed only after channel preconditioning, such as that rendered by a long depolarizing prepulse. This observation explains the lack of toxicity when depressant toxins are injected to mice. However, when the muscle channel  $\text{rNa}_v1.4$  expressed in *Xenopus* oocytes was modulated by the site-3  $\alpha$ -toxin Lqh $\alpha$ IT, LqhIT2 was capable of inducing a negative shift in the voltage-dependence of activation following a short prepulse, as was shown for other  $\beta$ -toxins. These unprecedented results suggest that depressant toxins may have a toxic impact on mammals in the context of the complete scorpion venom. To assess whether LqhIT2 and Lqh-dprIT3 interact with the insect and rat muscle channels in a similar manner we examined the role of Glu24, a conserved 'hot spot' at the bioactive surface of  $\beta$ -toxins. Whereas substitutions E24A/N abolished the activity of both LqhIT2 and Lqh-dprIT3 at insect  $\text{Na}_v\text{s}$ , they increased the affinity of the toxins for rat skeletal muscle channels. This result implies that depressant toxins interact differently with the two channel types and that substitution of Glu24 is key for converting toxin selectivity.

## INTRODUCTION

Voltage-gated sodium channels ( $\text{Na}_v\text{s}$ ) are critical in generation and propagation of action potentials in excitable cells, and are targeted by a large variety of chemically distinct compounds that bind at several receptor sites on the pore-forming  $\alpha$ -subunit (Catterall, 2000; Gordon, 1997). Most lipid-soluble  $\text{Na}_v$  activators, including pyrethroid insecticides, toxic alkaloids (e.g. veratridine and batrachotoxin) and marine cyclic polyether toxins (e.g. brevetoxins), affect  $\text{Na}_v\text{s}$  of both insects and mammals. Yet, despite the general conservation of  $\text{Na}_v$  structure, certain scorpion toxins show preference for  $\text{Na}_v$  subtypes in mammals or insects (Gordon et al., 1998, 2007; Cestele and Catterall, 2000; Gurevitz et al., 2007), which raised the idea of using some representatives for insect pest control (reviewed in Gurevitz et al., 2007).

Scorpion toxins that modulate  $\text{Na}_v$  gating are divided between the  $\alpha$  and  $\beta$  classes according to their mode of action and binding features to distinct receptor sites (Catterall, 2000).  $\alpha$ -Toxins prolong the action potential by inhibiting the fast inactivation of  $\text{Na}_v\text{s}$  upon binding to receptor site-3 (e.g. Lqh $\alpha$ IT from *Leiurus quinquestriatus hebraeus*; Eitan et al., 1990; Martin-Eauclaire and Couraud, 1995; Gordon et al., 1996) assigned mainly to extracellular loops in domains 1 and 4 (Catterall, 2000).  $\beta$ -Toxins shift the voltage dependence of channel activation to more hyperpolarized membrane potentials upon binding to receptor site-4 assigned mainly to external loops in domains 2 and 3 (Marcotte et al., 1997; Shichor et al., 2002; Cestele et al., 1998, 2006; Leipold et al., 2006). The  $\beta$ -toxins are further classified to: (i) Anti-mammalian  $\beta$ -toxins (e.g., C $\text{ss}2$  and C $\text{ss}4$  from *Centruroides suffusus suffusus*; Martin-Eauclaire and Couraud, 1995; Gordon et al., 1998; Gurevitz et al., 2007); (ii)  $\beta$ -Toxins that affect both insect and mammalian  $\text{Na}_v\text{s}$  (e.g. T $\text{s}1$  from *Tityus serrulatus* and Lqh $\beta$ 1; Possani et al., 1999; Gordon et al., 2003); (iii) Anti-insect selective excitatory  $\beta$ -toxins (e.g. AahIT from *Androctonus australis hector* and Bj-xtrIT from *Buthotus judaicus*) typified by the symptoms of contraction paralysis they produce in

blowfly larvae (Zlotkin et al., 1978; Froy et al., 1999), and by their unique structures (Oren et al., 1998; Li et al., 2005; Gurevitz et al., 2007); and (iv) Anti-insect depressant toxins, which upon injection to blowfly larvae induce flaccid paralysis due to sustained depolarization of the axonal membrane leading to block in the evoked action potentials and loss of muscle tonus (Lester et al., 1982; Zlotkin et al., 1991; Ben Kalipha et al., 1997; Strugatsky et al., 2005). These ~61 residue-long polypeptides were classified as  $\beta$ -toxins due to their ability to modulate  $\text{Na}_v$  activation and to compete with excitatory toxins on binding to receptor-site 4 in insect neuronal membranes (Gordon et al., 1984, 1992). Yet, depressant toxins did not compete with anti-mammalian  $\beta$ -toxins on binding to rat brain membranes, and were harmless when injected to mice (Lester et al., 1982; Herrmann et al., 1995; Strugatsky et al., 2005).

The bioactive surfaces of the anti-insect excitatory  $\beta$ -toxin, Bj-xtrIT, the anti-mammalian  $\beta$ -toxin, C<sub>ss</sub>4, and the anti-insect depressant  $\beta$ -toxin, LqhIT2, were described (Cohen et al., 2004, 2005; Karbat et al., 2006). These studies highlighted a conserved 'pharmacophore' composed of a key negatively-charged Glu in the  $\alpha$ -helix (Fig. 1), flanked by hydrophobic residues that may isolate the point of interaction with a counterpart channel residue from the bulk solvent. An additional hydrophobic cluster of bioactive residue, at the C-tail of Bj-xtrIT and on the loop connecting the second and third  $\beta$ -strands of C<sub>ss</sub>4, was suggested to determine toxin selectivity (Cohen et al., 2004, 2005). While the key residues involved in C<sub>ss</sub>4 and Bj-xtrIT activity form two topologically distinct domains, the bioactive surface of LqhIT2 is continuous, but involves the conserved pharmacophore (Glu24, Tyr26) and its vicinity (Karbat et al., 2006). The subset of residues common to the bioactive surfaces of C<sub>ss</sub>4, Bj-xtrIT, and LqhIT2 raised the possibility that these toxins might compete in binding for receptor site-4 on different  $\text{Na}_v$ s.

By analyzing the ability of depressant toxins to bind and affect various mammalian  $\text{Na}_v$ s, we found that despite the lack of effect on subcutaneously injected mice, depressant toxins bind to

receptor site-4 on rat muscle membranes in nM affinity. We show that LhqIT2 and Lqh-dprIT3 are effective on rNa<sub>v</sub>1.4 when the channel is excited prior to toxin application by either a long depolarizing prepulse or by modulation with an  $\alpha$ -toxin. These results suggest that the reported selectivity of depressant toxins to insect Na<sub>v</sub>s rests on the way they were tested, and that depressant toxins may have a toxic impact on mammals in the context of the complete venom.

## MATERIALS AND METHODS

**Toxins and their mutagenesis:** Production of Bj-xtrIT, LqhIT2, Css4, and Lqh-dprIT3 variant c in recombinant forms, PCR-driven mutagenesis, expression in *Escherichia coli*, *in-vitro* folding, and purification of toxin derivatives have been described in detail (Turkov et al., 1997; Froy et al., 1999; Strugatzky et al., 2005; Cohen et al., 2005)

**Binding experiments:** Neuronal membranes from cockroach were prepared from whole heads of adult *Periplaneta americana* according to a previously described method (Froy et al., 1999). Rat skeletal muscle membranes were prepared from adult albino Wistar strain (~300 g, laboratory bred) as was previously described (Gordon et al., 1988). Mammalian brain synaptosomes were prepared from the same rats as previously described (Gilles et al., 2001). Membrane protein concentration was determined by a Bio-Rad Protein Assay, using bovine serum albumin (BSA) as standard. Bj-xtrIT and Css4 (with a His-tag attached; His-Css4) were radioiodinated by lactoperoxidase (Sigma, Cat. No., L8257; 7 Units per 60  $\mu$ l reaction mix) using 10  $\mu$ g toxin and 0.5 mCi carrier-free Na<sup>125</sup>I (Amersham, UK) and the monoiodotoxin was purified as was previously described (Cohen et al., 2004, 2005). The media composition used in the binding assays and termination of the reactions were described elsewhere (Gilles et al., 2000, 2001). Non-specific toxin binding was determined in the presence of 1-10  $\mu$ M of the unlabeled toxin, and consisted typically of 10-30% of total binding. Equilibrium competition binding assays were

performed and analyzed as was previously described (Cohen et al., 2005). Each experiment was performed in duplicate and repeated at least three times as indicated ( $n$ ). Data are presented as mean  $\pm$  SD of the number of independent experiments.

**Expression of sodium channels in oocytes and two-electrode voltage clamp experiments:**

The genes encoding the *Drosophila melanogaster* sodium channel  $\alpha$ -subunit (DmNa<sub>v</sub>1) and the auxiliary TipE subunit were kindly provided by J. Warmke, Merck, New Jersey, USA, and M. S. Williamson, IACR-Rothamsted, UK, respectively. The gene encoding the rat skeletal muscle sodium channel, rNa<sub>v</sub>1.4, in the pAlter vector, was a gift from Dr. R.G. Kallen, University of Pennsylvania, Philadelphia, PA, USA. These genes and that for the auxiliary subunit h $\beta$ 1 were transcribed *in vitro* using T7 RNA-polymerase and the mMACHINE™ system (Ambion, Austin, TX) and were injected into *Xenopus laevis* oocytes as was previously described (Shichor et al., 2002).

**Two-electrode voltage-clamp recording:** Currents were measured 1-2 days after injection using a two-electrode voltage clamp and a Gene Clamp 500 amplifier (Axon Instruments, Union City, CA). Data were sampled at 10 kHz and filtered at 5 kHz. Data acquisition was controlled by a Macintosh PPC 7100/80 computer, equipped with ITC-16 analog/digital converter (Instrutech Corp., Port Washington, NY), utilizing Synapse (Synergistic Systems, Sweden). The bath solution contained (in mM): 96 NaCl, 2 KCl, 1 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 5 HEPES, pH 7.85. Oocytes were washed with bath solution flowing from a BPS-8 perfusion system (ALA Scientific Instruments, Westbury, NY) with a positive pressure of 4 psi. Toxins were diluted with bath solution and applied directly to the bath at their final desired concentration.

**Data analysis:** *Leak subtraction:* Capacitance transients and leak currents were removed by subtracting a scaled control trace utilizing a P/6 protocol (Armstrong and Bezanilla, 1974).

*GV analysis:* Mean conductance (G) was calculated from peak current/voltage relations using the

equation  $G = I/(V - V_{rev})$ , where  $I$  is the peak current elicited upon depolarization,  $V$  is the membrane potential, and  $V_{rev}$  is the reversal potential. Normalized conductance voltage relationship were fit with either a one or two component Boltzman distribution according to the following equation:

$$G/G_{max} = (1 - A)/(1 + \exp[(V_{1/2} - V)/k_1]) + A/(1 + \exp[(V_{2/2} - V)/k_2]) \quad [1]$$

where  $V_{1/2}$  and  $V_{2/2}$  are the respective membrane potentials for two populations of channels for which the mean conductance is half maximal;  $k_1$  and  $k_2$  are their respective slopes and  $A$  defines the proportion of the second population (amplitude) with respect to the total. For fits in which only one population of channels was apparent,  $A$  was set to zero.

*Steady state fast inactivation:* The voltage dependence of steady state fast inactivation is described by a single Boltzman distribution:

$$I/I_{max} = \alpha_0 + \alpha_1 / (1 + \exp[(V - V_{1/2})/k]) \quad [2]$$

where  $I$  is the peak current measured during the test depolarization step,  $I_{max}$  is the current obtained without a preceding conditioning step;  $V$  is the membrane potential of the conditioning step;  $V_{1/2}$  is the membrane potential at which half-maximal inactivation is achieved,  $k$  is the slope factor,  $\alpha_0$  is the remaining normalized peak current at very high depolarizing conditioning potentials, and  $\alpha_1$  is the normalized amplitude (Chen and Heinemann, 2001).

## RESULTS

### Binding of depressant toxins to rat skeletal muscle $Na_v$ s

Scorpion depressant toxins have traditionally been considered insect-selective on the basis of their exclusive toxicity to insects and high binding affinity for insect  $Na_v$ s (Lester et al., 1982; Zlotkin et al., 1991; Gordon et al., 1992; Strugatsky et al., 2005). Still, the elucidation of a pharmacophore common to the bioactive surface of scorpion  $\beta$ -toxins active on insects and



mammals (Cohen et al., 2005; Karbat et al., 2006) prompted us to analyze whether the toxins that show selectivity for insects would compete with the anti-mammalian  $\beta$ -toxin C<sub>ss4</sub> on binding to receptor site-4 on rat muscle and brain Na<sub>v</sub>s. While the excitatory toxin Bj-xtrIT did not displace <sup>125</sup>I-C<sub>ss4</sub> in concentrations up to 10  $\mu$ M, the depressant toxins LqhIT2 and Lqh-dprIT3 inhibited in a dose-dependent manner the binding of <sup>125</sup>I-C<sub>ss4</sub> to rat muscle membranes with K<sub>i</sub> values of  $45 \pm 7$  and  $30 \pm 3.2$  nM, respectively (Fig. 2, Table 1). In contrast, the excitatory and the depressant toxins did not displace <sup>125</sup>I-C<sub>ss4</sub> from rat brain synaptosomes at concentrations up to 10  $\mu$ M (inset in Fig. 2). The ability of depressant toxins to bind with an apparent high affinity to rat skeletal muscle Na<sub>v</sub>s raised the inevitable question as to why these toxins were inactive when injected subcutaneously into mice (Lester et al., 1982; Zlotkin et al., 1991, 1993). Therefore, we analyzed the effects of LqhIT2 and Lqh-dprIT3 on the rat muscle channel, rNa<sub>v</sub>1.4, and compared them to those obtained on the *Drosophila melanogaster* channel, DmNa<sub>v</sub>1, expressed in *Xenopus laevis* oocytes.

### **The effects of LqhIT2 and Lqh-dprIT3 on the activation of DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4**

The voltage-dependent activation of DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4 expressed in *Xenopus* oocytes was monitored by two-electrode voltage clamp in the absence and presence of LqhIT2 or Lqh-dprIT3 (Fig. 3, Table 2). Since the negative shift of voltage-dependent activation induced by  $\beta$ -toxins is better observed after a depolarizing prepulse (PP) (Cestele et al., 1998; Tsushima et al., 1999), we first examined the PP duration required to observe such a shift in the insect Na<sub>v</sub> in the presence of LqhIT2. While in its absence, DmNa<sub>v</sub>1 was not activated by a 50 ms test pulse to -50 mV independent of whether or not a depolarizing PP was provided, in the presence of 1  $\mu$ M LqhIT2, a hyperpolarizing shift in current-voltage relations at DmNa<sub>v</sub>1 was observed only after a preconditioning depolarizing PP (Fig. 3-A1). This shift was highly dependent on the PP length

(Fig. 3-A). A 100 ms PP to +60 mV provided the maximal effect measured by the developing currents elicited by a test pulse to -50 mV (Fig. 3-A3). Similar results were obtained with 200 nM Lqh-dprIT3 (Fig. 3-A2), a depressant toxin with higher toxicity to blowfly larvae (Strugatsky et al., 2005). Notably, Lqh-dprIT3 induced a maximal effect on DmNa<sub>v</sub>1 activation already after 10 ms PP (Fig. 3-A3). In comparison, a very short preconditioning depolarizing PP (2 ms) was sufficient to induce a hyperpolarizing shift in activation of rNa<sub>v</sub>1.2a by the anti mammalian  $\beta$ -toxin Csx4 (Cestele et al., 1998). These results have indicated that the effects of various  $\beta$ -toxins on the hyperpolarizing shift in channel activation clearly depend on the PP duration.

Based on the observation that LqhIT2 requires a relatively long PP to induce a hyperpolarizing shift in DmNa<sub>v</sub>1 activation, and that LqhIT2 and Lqh-dprIT3 bind with relatively high affinity to the rat muscle membranes (Fig. 2), we examined whether longer PP durations would facilitate the effect of these toxins on rNa<sub>v</sub>1.4 activation. We found that LqhIT2 was indeed capable of shifting the rNa<sub>v</sub>1.4 activation in the hyperpolarizing direction, but it required a longer PP to +60 mV, and even after a 2 s PP the effect was not maximal (Fig. 3-B). In the absence of toxin, a 500 ms PP had no effect on the peak current, and 1 or 2 s PP induced a 10% or 20% decrease in the peak current elicited at -10 mV (not shown), most likely due to the development of slow inactivation (Featherstone *et al.*, 1996; Mitrovitz *et al.*, 2000). Similar results were obtained with Lqh-dprIT3 (Fig. 3-B2 and B3), suggesting that depressant toxins not only bind with high affinity to rat muscle Na<sub>v</sub>s, but also have a clear effect under the appropriate conditions. In contrast, regardless of the PP length, the depressant toxins did not shift the activation curve of the rat brain channel rNa<sub>v</sub>1.2a expressed in oocytes (not shown), which was in concert with their inability to bind to rat brain synaptosomes (inset in Fig. 2). This data demonstrated that receptor site-4 varies in different Na<sub>v</sub>s.

### **Effect of substitutions at the 'hot spot' on selectivity of LqhIT2 and Lqh-dprIT3**

We have shown that substitution of a conserved Glu residue on the bioactive surfaces of the anti-mammalian  $\beta$ -toxin C<sub>ss</sub>4 (Glu28; Fig. 1) and the anti-insect selective excitatory  $\beta$ -toxin B<sub>j</sub>-xtrIT (Glu30) abolished their binding and activity at the brain and insect Na<sub>v</sub>s, respectively. Substitution of the adjacent Arg27 in C<sub>ss</sub>4 and His25 in B<sub>j</sub>-xtrIT also decreased the binding and activity of both toxins (Cohen et al., 2004, 2005). Therefore we analyzed whether the spatially equivalent residues in LqhIT2, Glu24 and Lys23 (Karbat et al., 2006), have a role in toxin binding and activity at rNa<sub>v</sub>1.4. Substitution K23A decreased LqhIT2 affinity for both cockroach neuronal and rat muscle membranes (Fig. 4, Table 1), as well as abolished the activity at DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4 (Fig. 5, Table 2). Although substitutions E24A/N decreased the toxin affinity for cockroach neuronal membranes by 450-fold (Fig. 4-B, Table 1) and no activity was observed at DmNa<sub>v</sub>1 with up to 5  $\mu$ M toxin (Fig. 5-A, Table 2), LqhIT2<sup>E24A/N</sup> affinity for the rat muscle *increased* 5-fold (Fig. 4-A, Table 1) and the effect at rNa<sub>v</sub>1.4 following a 500 ms PP was higher than that of LqhIT2 (Fig. 5-B, Table 2). Substitution E24N in Lqh-dprIT3 had a similar effect on toxin activity in that the effect at DmNa<sub>v</sub>1 declined, but increased at rNa<sub>v</sub>1.4 (Fig. 5-C and D, Table 2). These results not only indicate that depressant toxins have different requirements for modifying the activation of insect versus mammalian Na<sub>v</sub>s, they also show that a single amino acid substitution can invert the preference of depressant toxins and make them selective to skeletal muscle Na<sub>v</sub>s (Tables 1 and 2).

### **Lqh $\alpha$ IT modulates LqhIT2 activity at rNa<sub>v</sub>1.4**

We have previously shown that the  $\alpha$ -toxin Lqh $\alpha$ IT and the depressant  $\beta$ -toxin LqhIT2 allosterically increase the binding of one another at insect Na<sub>v</sub>s, which was manifested in a strong synergism in their toxicity to insects (Cohen et al., 2006). Considering that Lqh $\alpha$ IT was

shown to inhibit the fast inactivation of rNa<sub>v</sub>1.4 expressed in HEK cells with an EC<sub>50</sub> of 1.2 nM (Leipold et al., 2004), we examined whether Lqh $\alpha$ IT would modulate LqhIT2 activity at rNa<sub>v</sub>1.4. Lqh $\alpha$ IT in a concentration of 200 nM shifted the voltage-dependence of steady state fast inactivation of rNa<sub>v</sub>1.4 expressed in *Xenopus* oocytes by +15 mV ( $V_{1/2}$  = -48  $\pm$  0.3 mV in the control; -33.7  $\pm$  0.3 mV in the presence of Lqh $\alpha$ IT), thus increasing the percentage of channels available for activation at sub-threshold membrane potentials (under -40 mV) with no effect on the channels conductance-voltage relations (Fig. 6). These data suggested that Lqh $\alpha$ IT might have facilitated the activity of the depressant toxin. Upon co-application of Lqh $\alpha$ IT (200 nM) and LqhIT2 (5  $\mu$ M), a 50 ms PP to +60 mV induced a hyperpolarizing shift in rNa<sub>v</sub>1.4 conductance-voltage relations (Fig. 6-A). Such an effect on the channel activation was comparable to that obtained by a 500 ms PP to +60 mV when LqhIT2 was applied alone (Fig. 5-B, Table 2). This result demonstrates that Lqh $\alpha$ IT binding to rNa<sub>v</sub>1.4 and/or its influence on fast inactivation modulates the activity of LqhIT2 on this channel as indicated by the shorter PP required to observe the effect of the depressant toxin.

## DISCUSSION

The experiments described in this study reveal that the allegedly 'insect-selective' scorpion depressant toxins are capable of binding with high affinity and affecting mammalian skeletal muscle Na<sub>v</sub>s, a fact unnoticed for almost three decades. The classification of depressant toxins as insect-selective relied on the lack of toxicity when injected to mice and inability to bind rat brain synaptosomes (Zlotkin et al., 1993; Gordon et al., 1998; Gurevitz et al., 2007) or affect mammalian Na<sub>v</sub>s expressed in *Xenopus* oocytes (Gordon et al., 2003; Bosmans et al., 2005). However, a number of recent results have suggested that the issue of selectivity of depressant toxins toward insects deserves reexamination: (i) Scorpion  $\beta$ -toxins share a common

pharmacophore (Cohen et al., 2005; Karbat et al., 2006), which explains their ability to compete in binding; (ii) Receptor site-4 on rat skeletal muscle Na<sub>v</sub>s was suggested to differ from those of various mammalian neuronal and cardiac Na<sub>v</sub>s (Marcotte et al., 1997; Cestele et al., 1998; Leipold et al., 2006; Shciavon et al., 2006); and (iii) Scorpion  $\alpha$ - and  $\beta$ -toxins exert synergistic effects due to allosteric interactions between receptor sites 3 and 4 on insect Na<sub>v</sub>s (Cohen et al., 2006).

On the basis of these considerations we reanalyzed the activity of depressant toxins on insect and mammalian Na<sub>v</sub>s. Although depressant toxins were unable to bind rat brain Na<sub>v</sub>s, they exerted high affinity for the rat skeletal muscle Na<sub>v</sub>s (Fig. 2), which motivated us to analyze their effects on channel activation. The unexpected observation that the effect of the depressant toxin LqhIT2 on the insect Na<sub>v</sub> DmNa<sub>v</sub>1 necessitated a 10-fold longer preconditioning depolarizing prepulse (PP) than that required to observe a C<sub>ss</sub>4 effect on mammalian Na<sub>v</sub>s suggested that LqhIT2 should be analyzed on rNa<sub>v</sub>1.4 following a longer PP. The requirement for preconditioning depolarization (PP) has been attributed to a putative energetic barrier that needs to be overcome before the prebound  $\beta$ -toxin can trap the DII/S4 voltage sensor in its outward activated position, thus leading to enhanced channel activation upon subsequent depolarizations (Cestele et al., 1998). Thus far, a short PP (several ms) was ample for inducing a noticeable effect of most anti-mammalian  $\beta$ -toxins on mammalian Na<sub>v</sub>s (Tsushima et al., 1999; Cestele et al., 1998, 2006; Cohen et al., 2005). Here we show that a very long preconditioning PP (>500 ms) to +60 mV turned rNa<sub>v</sub>1.4 vulnerable to depressant toxins. It is likely that because depressant toxins did not modulate the gating properties of rNa<sub>v</sub>1.2 (Bosmans et al., 2005) and rNa<sub>v</sub>1.4 (Fig. 2; Gordon et al., 2003) following a depolarizing PP up to 50 ms, their activity on mammalian Na<sub>v</sub>s has not been noticed thus far.

The ability of depressant toxins to influence the activation of the mammalian  $\text{Na}_v$  following a long PP was surprising and raised the question as to their putative role *in vivo*. In light of the enhancement of LqhIT2 binding to insect  $\text{Na}_v$ s in the presence of a scorpion  $\alpha$ -toxin from the same venom, Lqh $\alpha$ IT (Cohen et al., 2006), and since Lqh $\alpha$ IT is highly potent on insect as well as a variety of mammalian  $\text{Na}_v$  subtypes, including r $\text{Na}_v$ 1.4 (Eitan et al., 1990; Chen et al., 2000; Leipold et al., 2004; Gordon et al., 2007), it was rational to analyze the joint effect of both toxins on r $\text{Na}_v$ 1.4. Indeed, the synergism between site-3 and site-4 toxins observed at r $\text{Na}_v$ 1.4 may be explained by the reduction of at least 10-fold in PP duration required for induction of LqhIT2 effect in the presence of Lqh $\alpha$ IT (Fig. 6). Such a mechanism may also apply to insect  $\text{Na}_v$ s where synergism between site-3 and site-4 toxins was reported (Cohen et al., 2006). Lqh $\alpha$ IT increases neuronal excitability and neuromuscular activity upon binding to receptor site-3 by induction of long plateau potentials in axons attributed to an increase in the probability of  $\text{Na}_v$ s to remain in open states due to inhibition of their fast inactivation (Eitan et al., 1990; Gilles et al., 2000; Lee et al., 2000; Benoit and Gordon, 2001). This inhibition of channel steady state fast inactivation expands the channel population available for activation at resting membrane potential, leading to increase in the frequency of action potentials, which may act as a prepulse to facilitate  $\beta$ -toxin activity. This, in turn, may facilitate  $\alpha$ -toxin action on the channels in their open states. Thus, the mutual enhancement in  $\alpha$ -toxin effect by  $\beta$ -toxin interaction with receptor site-4, and *vice versa*, results from an indirect modification of receptor sites-3 and 4, respectively and from alteration in the voltage-dependence of channel activation (Cohen et al., 2006). On the basis of these considerations we suggest that  $\text{Na}_v$ 1.4 in a stung mammal is preconditioned upon binding of the  $\alpha$ -toxin, thus enabling synergistic toxicity by the joint effects of  $\alpha$ - and depressant toxins.

The venom of *L. q. hebraeus* contains a number of  $\alpha$ -toxins active on mice (e.g., Lqh $\alpha$ IT, Lqh2, Lqh3, Lqh4, Lqh6, Lqh7; for references see Gordon et al., 2007), and  $\beta$ -toxins that exhibit preference for insects including various depressant toxins (e.g., LqhIT2, LqhIT5, Lqh-dprIT3, Lqh $\beta$ 1; for references see Gurevitz et al., 2007), which together increase the impact of stinging. In their isolated form, however, the depressant toxins may be considered selective to insects as their affinity for insect neuronal membranes is two orders of magnitude higher than the affinity for rat muscle membranes (Figs. 4 and 5, Table 2; Gordon et al., 1992; Strugatzky et al., 2005).

A surprising feature in the interaction of depressant toxins with the mammalian muscle Na<sub>v</sub> is the lack of function of Glu24, found to be conserved in other scorpion  $\beta$ -toxins and considered a hot spot on the bioactive surface of these toxins toward insect and rat brain Na<sub>v</sub>s (Cohen et al., 2004, 2005; Karbat et al., 2006; Fig. 1), suggesting that receptor site-4 on the muscle channels differs from that on DmNa<sub>v</sub>1. Apparently, further understanding of how scorpion  $\beta$ -toxins interact in a preferential manner with various receptor sites on Na<sub>v</sub> subtypes awaits elucidation of toxin-receptor interacting surfaces. Presently, our results indicate that a single substitution at this position (LqhIT2<sup>E24A/N</sup> and Lqh-dprIT3<sup>E24N</sup>; Figs. 4 and 5, Table 2) converts these depressant toxins from 'insect-selective' into 'mammal-selective'.

## ACKNOWLEDGEMENT

We thank Prof. W.A. Catterall, University of Washington, Seattle, for his critical comments on this manuscript.

## REFERENCES

Armstrong CM, and Bezanilla F (1974) Charge movement associated with the opening and closing of the activation gates of Na channels. *J Gen Physiol* 63:533-552.

- Ben Khalifa R, Stankiewicz M, Pelhate M, Serrano-Hernandez SE, Possani LD, Hinkel H, and Mebs D (1997) Action of babycurus-toxin 1 from the east African scorpion *Babycurus centrurimorphus* on the isolated cockroach giant axon. *Toxicon* 35:1069-1080.
- Benoit E, and Gordon D (2001) The scorpion  $\alpha$ -like toxin Lqh III specifically alters sodium channel inactivation in frog myelinated axons. *Neuroscience* 104:551-559.
- Catterall WA (2000) From ionic currents to molecular mechanisms: The structure and function of voltage-gated sodium channels. *Neuron* 26:13-25.
- Cestele S, and Catterall WA (2000) Molecular mechanisms of neurotoxin action on voltage-gated sodium channels. *Biochimie* 82:883-892.
- Cestele S, Qu Y, Rogers JC, Rochat H, and Catterall WA (1998) Voltage sensor-trapping: enhanced activation of sodium channels by  $\alpha$ -scorpion toxin bound to the S3-S4 loop in domain II. *Neuron* 21:919-931.
- Cestele S, Yarov-Yarovoy V, Qu Y, Sampieri F, Scheuer T, and Catterall WA (2006) Structure and function of the voltage sensor of sodium channels probed by a beta-scorpion toxin. *J Biol Chem* 281:21332-21344.
- Chen H, and Heinemann SH (2001) Interaction of scorpion alpha-toxins with cardiac sodium channels: binding properties and enhancement of slow inactivation. *J Gen Physiol* 117:505-518.
- Cohen L, Karbat I, Gilles N, Froy O, Angelovici R, Gordon D, and Gurevitz M (2004) Dissection of the functional surface of an anti-insect excitatory toxin illuminates a putative 'hot spot' common to all scorpion  $\beta$ -toxins affecting Na channels. *J Biol Chem* 279:8206-8211.



- Cohen L, Karbat I, Gilles N, Ilan N, Gordon D, and Gurevitz M (2005) Common features in the functional surface of scorpion  $\beta$ -toxins and elements that confer specificity for insect and mammalian voltage-gated Na-channels. *J Biol Chem* 280:5045-5053.
- Cohen L, Lipstein N, and Gordon D (2006) Allosteric interaction between scorpion toxin receptor sites on voltage-gated Na channels imply a novel role for weakly active components in arthropod venom. *FASEB J* 20:1933-1935.
- Eitan M, Fowler E, Herrmann R, Duval A, Pelhate M, and Zlotkin E (1990) A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: Purification, primary structure, and mode of action. *Biochemistry* 29:5941-5947.
- Featherstone DE, Richmond JE, and Ruben PC (1996) Interaction between fast and slow inactivation in Skm1 sodium channels. *Biophys J* 71:3098-3109.
- Froy O, Zilberberg N, Gordon D, Turkov D, Gilles N, Stankiewicz M, Pelhate M, Loret E, Oren D, Shaanan B, and Gurevitz M (1999) The putative bioactive surface of insect-selective scorpion excitatory neurotoxins. *J Biol Chem* 274:5769-5776.
- Gilles N, Chen H, Wilson H, Le Gall F, Montoya G, Molgo J, Schönherr R, Nicholson G, Heinemann SH, and Gordon D (2000) Scorpion  $\alpha$ - and  $\alpha$ -like toxins differentially interact with sodium channels in mammalian CNS and periphery. *Eur J Neurosci* 12:2823-2832.
- Gilles N, Leipold E, Chen H, Heinemann SH, and Gordon D (2001) Effect of depolarization on binding kinetics of scorpion  $\alpha$ -toxin highlights conformational changes of rat brain sodium channels. *Biochemistry* 40:14576-14584.
- Gordon D, Jover E, Couraud F, and Zlotkin E (1984) The binding of an insect selective neurotoxin (AaIT) from scorpion venom to locust synaptosomal membranes. *Biochim Biophys Acta* 778:349-358.

- Gordon D (1997) Sodium channels as targets for neurotoxins: mode of action and interaction of neurotoxins with receptor sites on sodium channels. in *Toxins and Signal Transduction* (Lazarowici P and Gutman Y eds) pp 119-149, Harwood Press, Amsterdam.
- Gordon D, Ilan N, Zilberberg N, Gilles N, Urbach D, Cohen L, Karbat I, Froy O, Gaathon A, Kallen RG, Benveniste M, and Gurevitz M (2003) An 'Old World' scorpion  $\beta$ -toxin that recognizes both insect and mammalian sodium channels: a possible link towards diversification of  $\beta$ -toxins. *Eur J Biochem* 270:2663-2670.
- Gordon D, Martin-Eauclaire MF, Cestele S, Kopeyan C, Carlier E, Ben Khalifa R, Pelhate M, and Rochat H (1996) Scorpion toxins affecting sodium current inactivation bind to distinct homologous receptor sites on rat brain and insect sodium channels. *J Biol Chem* 271:8034-8045.
- Gordon D, Merrick D, Wallner DA, and Catterall WA (1988) Biochemical properties of sodium channels in a wide range of excitable tissues studied with site-directed antibodies. *Biochemistry* 27:7032-7038.
- Gordon D, Moskowitz H, Eitan M, Warner C, Catterall WA, and Zlotkin E (1992) Localization of receptor sites for insect-selective toxins on sodium channels by site-directed antibodies. *Biochemistry* 31:7622-7628.
- Gordon D, Savarin P, Gurevitz M, and Zinn-Justin S (1998) Functional anatomy of scorpion toxins affecting sodium channels. *J Toxicol Toxin Rev* 17:131-158.
- Herrmann R, Moskowitz H, Zlotkin E, and Hammock BD (1995) Positive cooperativity among insecticidal scorpion neurotoxins. *Toxicon* 33:1099-1102.
- Karbat I, Turkov M, Cohen L, Kahn R, Gordon D, Gurevitz M, Frolov F (2006) X-ray structure and mutagenesis of the scorpion depressant toxin LqhIT2 reveals key determinants crucial for activity and anti-insect selectivity. *J Mol Biol* 366:586-601.

- Lee D, Gurevitz M, and Adams ME (2000) Modification of synaptic transmission and sodium channel inactivation by the insect-selective scorpion toxin Lqh $\alpha$ IT. *J Neurophysiol* 83:1181-1187.
- Lester D, Lazarovici P, Pelhate M, and Zlotkin E (1982) Purification, characterization and action of two insect toxins from the venom of the scorpion *Buthotus judaicus*. *Biochim Biophys Acta Prot Struct Mol Enzymol* 701:370-381.
- Li C, Guan RJ, Xiang Y, Zhang Y, Wang DC (2005) Structure of an excitatory insect-specific toxin with an analgesic effect on mammals from the scorpion *Buthus martensii* Karsch. *Acta Crystallogr D Biol Crystallogr* 61:14-21.
- Marcotte P, Chen LQ, Kallen RG, and Chahnin M (1997) Effects of *Tityus serrulatus* scorpion toxin  $\gamma$  on voltage-gated Na<sup>+</sup> channels. *Circ Res* 80:363-369.
- Martin-Eauclaire MF, and Couraud F (1995) Scorpion neurotoxins: Effects and mechanisms. in *Handk Neurotoxicology* (Chang LW and Dyer RS eds) pp 683-716, Marcel Dekker, NY, NY.
- Mitrovic N, George AL, and Horn R. (2000) Role of domain 4 in sodium channel slow inactivation. *J Gen Physiol* 115:707-717.
- Oren DA, Froy O, Amit E, Kleinberger-Doron N, Gurevitz M, Shaanan B (1998) An excitatory scorpion toxin with a distinctive feature: an additional alpha helix at the C terminus and its implications for interaction with insect sodium channels. *Structure* 6:1095-1103.
- Pintar A, Possani LD, Delepierre M (1999) Solution structure of toxin 2 from *Centruroides noxius* Hoffmann, a  $\beta$ -scorpion neurotoxin acting on sodium channels. *J Mol Biol* 287:359-367.
- Possani LD, Becerril B, Delepierre M, Tytgat J (1999) Scorpion toxins specific for Na<sup>+</sup>-channels. *Eur J Biochem* 264:287-300.

- Shichor I, Zlotkin E, Ilan N, Chikashvili D, Stuhmer W, Gordon D, Lotan I (2002) Domain 2 of *Drosophila* Para voltage-gated sodium channel confers insect properties to a rat brain channel. *J Neurosci* 22:4364-4371.
- Strugatsky D, Zilberberg N, Stankiewicz M, Ilan N, Turkov M, Cohen L, Pelhate M, Gilles N, Gordon D, and Gurevitz M (2005) Genetic polymorphism and expression of a highly potent scorpion depressant toxin enables refinement of the effects on insect Na-channels and illuminates the key role of Asn-58. *Biochemistry* 44:9179-9187.
- Tsushima RG, Borges A, and Backx PH (1999) Inactivated state dependence of sodium channel modulation by beta-scorpion toxin. *Pflugers Arch* 437:661-668.
- Turkov M, Rashi S, Zilberberg N, Gordon D, Ben Khalifa R, Stankiewicz M, Pelhate M, and Gurevitz M (1997) In vitro folding and functional analysis of an anti-insect selective scorpion depressant neurotoxin produced in *Escherichia coli*. *Prot Expr Purific* 9:123-131.
- Zlotkin E, Eitan M, Bindokas VP, Adams ME, Moyer M, Burkhart W, and Fowler E (1991) Functional duality and structural uniqueness of depressant insect-selective neurotoxins. *Biochemistry* 30:4814-4821.
- Zlotkin E, Fowler E, Gurevitz M, and Adams ME (1993) Depressant insect selective neurotoxins from scorpion venom: chemistry, action, and gene cloning, *Arch Insect Biochem Physiol* 22:55-73.
- Zlotkin E, Miranda F, and Rochat H (1978) Chemistry and pharmacology of Buthinae scorpion venoms. in *Arthropods Venoms* (Bettini S ed) pp 317-369, Springer Verlag, NY, NY.

### Footnotes

(1) This research was supported by the United States-Israel Binational Agricultural Research and Development grants IS-3928-06 (M.G. and D.G.) and IS-4066-07 (D.G. and M.G.); by the Israeli Science Foundation grants 909/04 (M.G.), 1008/05 (D.G.), 654/02 (M.B.); by a grant from the German-Israeli Foundation (G.I.F.) for Scientific Research and Development (No. G-770-242.1/2002 (D.G.); and by NIH 1 U01 NS058039-01 (M.G.).

(2) Current address of Morris Benveniste: Neuroscience Institute, Morehouse School of Medicine, 720 Westview Drive, Atlanta, GA 30310, U.S.A.

## Figure legends

**Fig. 1:** Sequence alignment and 3-D structure of  $\beta$ -toxin representatives. A, Sequences were aligned according to the conserved cysteine residues. The disulfide bonds formed between cysteine pairs in all 'long-chain' scorpion toxins are designated by lines. Dashes indicate gaps. Secondary structure motifs (B,  $\beta$ -strand; H,  $\alpha$ -helix) in C<sub>ss4</sub> follow the published structure of the  $\beta$ -toxin Cn2 (from *Centruroides noxius*; Pintar et al., 1999). B, A cartoon diagram of the C $\alpha$  structure of C<sub>ss4</sub> and LqhIT2 (in gray) covered by a semitransparent molecular surface of the toxins. LqhIT2 is derived from the Protein Data Bank accession 2I61. The C<sub>ss4</sub> model is from Cohen et al., 2005, and is spatially aligned with that of LqhIT2. The sulfur atoms in the disulfide bonds are highlighted in yellow, and the conserved glutamate is in red (see also in A). The figure was prepared using PyMOL (<http://www.pmol.org>).

**Fig. 2:** Competition of  $\beta$ -toxins with C<sub>ss4</sub> on binding to rat muscle membranes. Membranes were incubated 60 min at 22<sup>o</sup>C with 0.1 nM <sup>125</sup>I-C<sub>ss4</sub>, and increasing concentrations of the various  $\beta$ -toxins. Non-specific binding, determined in the presence of 1  $\mu$ M C<sub>ss4</sub>, was subtracted. The K<sub>i</sub> values are (in nM,  $n \geq 3$ ): C<sub>ss4</sub>,  $3.9 \pm 1.17$ ; Lqh-dprIT3,  $30 \pm 3.2$ ; LqhIT2,  $45 \pm 7$ ; Bj-xtrIT,  $\gg 10000$ . The binding of C<sub>ss4</sub>, LqhIT2, and Bj-xtrIT to rat brain synaptosomes under the same conditions is shown in the inset. The K<sub>i</sub> values are (in nM,  $n \geq 3$ ): C<sub>ss4</sub>,  $0.98 \pm 0.1$ ; LqhIT2 and Bj-xtrIT,  $\gg 10000$ . Representative experiments are shown.

**Fig. 3:** LqhIT2 and Lqh-dprIT3 effect on activation of DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4. A1,A2, Current-voltage (I-V) relations of DmNa<sub>v</sub>1 under control conditions and in the presence of 1  $\mu$ M LqhIT2 (A1), or 0.2  $\mu$ M Lqh-dprIT3 (A2) with or without a 50 ms preconditioning depolarizing prepulse (PP) to +60 mV from a -100 mV holding potential. B1,B2, Current-voltage relations of rNa<sub>v</sub>1.4

under control conditions and in the presence of 5  $\mu\text{M}$  LqhIT2 (B1), or 5  $\mu\text{M}$  Lqh-dprIT3 (B2) with a 50 ms or 500 ms preconditioning depolarizing PP to +60 mV from a -100 mV holding potential. A3 and B3, Analysis of the effect of depressant toxins on DmNa<sub>v</sub>1 (A3) and rNa<sub>v</sub>1.4 (B3) activation following various (0-1 s) PP durations to +60 mV and measuring at -50 mV. The current at each point was normalized to the maximal effect. Representative experiments are shown.

**Fig. 4:** Binding of LqhIT2 and its mutants to cockroach and rat muscle membrane. Competition of LqhIT2 and mutants with <sup>125</sup>I-Bj-xtrIT on binding to cockroach neuronal membranes (A), and with <sup>125</sup>I-Css4 to rat muscle membranes (B). Membranes were incubated 60 min at 22<sup>o</sup>C with 0.1 nM <sup>125</sup>I-Bj-xtrIT or <sup>125</sup>I-Css4, and increasing concentrations of the various mutants. Non-specific binding, determined in the presence of 1  $\mu\text{M}$  Bj-xtrIT or Css4, was subtracted. The K<sub>i</sub> values in nM,  $n \geq 3$  are given in Table 1. Representative experiments are shown.

**Fig. 5:** Alterations in conductance-voltage (G-V) relations induced by LqhIT2 and Lqh-dprIT3 mutants at DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4. The effects of LqhIT2 and its mutants K23A, E24A, and E24N on DmNa<sub>v</sub>1 (A) and on rNa<sub>v</sub>1.4 (B). The effects of Lqh-dprIT3 and its mutant E24N on DmNa<sub>v</sub>1 (C) and on rNa<sub>v</sub>1.4 (D); The labels in A cover B and those of C cover D. Toxin concentrations and the activation parameters ( $V_{0.5}$ ) are as described in Table 2. Conductance-voltage relations were determined as described in Fig. 1A with a 50 ms PP for DmNa<sub>v</sub>1 and a 500 ms PP for rNa<sub>v</sub>1.4. The data represent the mean  $\pm$  SEM of at least six independent experiments.

**Fig. 6:** Effects of combined application of LqhIT2 and Lqh $\alpha$ IT on rNa<sub>v</sub>1.4 activation. A, Conductance-voltage relations of rNa<sub>v</sub>1.4 in the absence of toxin ( $V_{0.5} = -33 \pm 0.3$  mV,  $n = 3$ ), in

the presence of 200 nM Lqh $\alpha$ IT ( $V_{0.5} = -33 \pm 0.4$  mV), and in the presence of 200 nM Lqh $\alpha$ IT and 5  $\mu$ M LqhIT2 ( $V_{0.5} = -38 \pm 0.1$  for all channels, and  $-52 \pm 1.5$  mV for 8% of the channel population; see Methods and Table 2). Conductance-voltage relations were determined as described in Fig. 2 with a 50 ms PP to +60 mV. Note that 5  $\mu$ M LqhIT2 have no effect on activation (see Fig. 3-B1). Inset, current traces obtained in the absence of toxins and upon co-application of 200 nM Lqh $\alpha$ IT and 5  $\mu$ M LqhIT2 at a test pulse to -45 mV following a 50 ms PP to +60 mV. B, Steady-state fast inactivation was determined from holding potential of -100 mV using a series of 50 ms PP from -80 to -20 mV in 5 mV increments prior to the test pulse of -20 mV. The steady-state inactivation of rNav1.4 fits a Boltzmann function with  $V_{0.5} = -48 \pm 0.3$  mV, and in the presence of 200 nM Lqh $\alpha$ IT,  $V_{0.5} = -33.7 \pm 0.3$  mV. Inset, effects of 200 nM Lqh $\alpha$ IT on the current fast inactivation elicited by a test pulse to -20 mV from a holding potential of -100 mV.



**Table 1.** Changes in apparent binding affinity of depressant  $\beta$ -toxins and selected mutants to rat muscle and cockroach neuronal membranes. The  $K_i$  values obtained from competition binding studies using  $^{125}\text{I}$ -Bj-xtrIT (cockroach neuronal membranes), and  $^{125}\text{I}$ -Css4 (rat muscle membrane preparation). See Fig. 3 for details.

Toxin	Ki, nM	Ki, nM	Ki ratio
	Muscle	Cockroach	Muscle/Cockroach
Lqh-dprIT3	30±3.2	0.2±0.04	150
LqhIT2	45±7	0.7±0.14	81
LqhIT2 <sup>K23A</sup>	900±122	11.1±1.2	86
LqhIT2 <sup>E24A</sup>	11±0.5	87±5	0.13
LqhIT2 <sup>E24N</sup>	8±1.2	91.5±1.5	0.09
Css4	3.9±1.17	>5000	
Bj-xtrIT	>10000	0.16±0.03	

**Table 2.** Alterations of the mid-voltage of activation ( $V_{0.5}$ ) derived from conductance-voltage (G-V) curves of DmNa<sub>v</sub>1 and rNa<sub>v</sub>1.4 induced by depressant toxins and their mutants. The  $V_{0.5}$  values are from the conductance-voltage (G-V) curves presented in Fig. 4. Depressant toxin effect on rNa<sub>v</sub>1.4 exhibit two components: a minor negative shift in the  $V_{0.5}$  of the entire channel population (upper number), and a stronger shift in the  $V_{0.5}$  of a fraction of the toxin-modified channel population (lower number), indicated by the numbers in parentheses. The data represent the mean  $\pm$  SEM of at least six independent experiments.

Toxin	DmNa <sub>v</sub> 1		rNa <sub>v</sub> 1.4	
	$\mu$ M	$V_{0.5}$ , mV	$\mu$ M	$V_{0.5}$ , mV
Control	-	-19.1 $\pm$ 0.5	-	-26 $\pm$ 0.2
<b>Lqh-dprIT3</b>	0.2	-35.4 $\pm$ 0.7	5	-31 $\pm$ 0.7 -50.7 $\pm$ 2.3 (23%)
Lqh-dprIT3 <sup>E24N</sup>	1	-20.2 $\pm$ 0.6	1	-38.5 $\pm$ 0.3 -60.4 $\pm$ 1.5 (11%)
<b>LqhIT2</b>	1	-33.1 $\pm$ 0.4	5	-30 $\pm$ 0.2 -52 $\pm$ 3.5 (8%)
LqhIT2 <sup>K23A</sup>	5	-20.8 $\pm$ 0.7	5	-27.2 $\pm$ 1.5
LqhIT2 <sup>E24A</sup>	5	-18.6 $\pm$ 0.6	5	-33.5 $\pm$ 1.5 -54.5 $\pm$ 9 (15%)
LqhIT2 <sup>E24N</sup>	5	-19 $\pm$ 0.5	5	-32.8 $\pm$ 1.4 -43.1 $\pm$ 4.1 (62%)

**A.**

	BB	HHHHHHH	BBBB	BBBB
Css4	KEGYLVNSYTGCK-FECFKLGDND--YCLR	<b>E</b> CRQQYGKGS	GGGYCYAF--GCWCTHLYEQAVVWPLPNKTCN	
LqhIT2	DGYIKRRD-GCK-VACLI----	GNEGCDK	<b>E</b> CKA-Y-GGSYGYCWTWGLACWCEGLPDDKT-WKSETNTCG	
Lqh-dprIT3	DGYIRGGD-GCK-VSCVI----	NHVFCDN	<b>E</b> CLAA--GGSYGYCWAWGLACWCEGLPADRE-WKYETNTC	
Bj-xtrIT	KKNGYPLDRN-G-KTTECSGVNAIAPHYCNS	<b>E</b> CTKVYVAESGYCCW--	GACYCFGLEDDKPIGPMKDI	TKKYCDVQIIP




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6