

MOL #57430

Docosahexaenoic acid inhibits superoxide dismutase 1 gene transcription in human cancer cells:
the involvement of PPAR α and HIF-2 α signaling

Erin R. Tuller, Charles T. Beavers, Jessica R. Lou, Michael A. Ihnat, Doris M. Benbrook, and
Wei-Qun Ding

From the Departments of Obstetrics and Gynecology (ERT, DMB), Biochemistry and Molecular
Biology (DMB), Pathology (CTB, JRL, WQD), and Cell Biology (MAI), University of
Oklahoma Health Sciences Center, Oklahoma City, OK 73104

MOL #57430

Running title: DHA inhibits SOD-1 gene transcription

Address correspondence to: Dr. Wei-Qun Ding, University of Oklahoma Health Sciences Center, 975 NE 10th Street, BRC 409, Oklahoma City, OK 73104. Telephone (405) 271-1605; fax (405) 271-3910; email weiqun-ding@ouhsc.edu

The number of text pages: 29

The number of tables: 0

The number of figures: 9

The number of references: 42

The number of words in Abstract: 246

The number of words in Introduction: 631

The number of words in Discussion: 1036

Abbreviations used: 4-HNE, 4-hydroxy Nonenal; DHA, docosahexaenoic acid; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HIF, hypoxia-inducible factor; HRE, hypoxia-response element; LPS, lipopolysaccharide; PPAR, peroxisome proliferator-activated Receptor; PPRE, peroxisome proliferation-responsive element; SOD-1, superoxide dismutase 1

MOL #57430

Abstract

Docosahexaenoic acid (DHA, n-3, 22:6) is known to have anticancer activity, but its mechanisms of action remain to be further elucidated. We recently demonstrated that DHA down-regulates superoxide dismutase 1 (SOD-1) gene expression, thereby weakening cellular antioxidant forces and enhancing cytotoxicity in various human cancer cells. The objective of this study was to investigate the mechanism of DHA's inhibitory effect on SOD-1 gene expression in human cancer cells. A reporter gene assay indicated that DHA suppresses SOD-1 gene transcription in a time- and concentration-dependent manner in human cancer cells. Pre-treatment with vitamin E did not block DHA's inhibitory effect, indicating that this suppression does not depend on lipid peroxidation. The suppressive effect of DHA on SOD-1 gene transcription could be mimicked by the peroxisome proliferator-activator receptor alpha (PPAR α) ligand clofibrate, but not the PPAR γ ligand troglitazone, suggesting the involvement of PPAR α signaling. Deletion analysis of the key DNA binding elements in the SOD-1 gene promoter identified the distal hypoxia response element (HRE), but not the the peroxisome proliferator response element (PPRE) or NF- κ B element, as essential for DHA's suppressive effects. Co-immunoprecipitation confirmed that PPAR α , but not PPAR γ , forms a complex with hypoxia-inducible factor (HIF)-2 α in cancer cells. Chromatin immunoprecipitation analysis indicated that both DHA and clofibrate reduce HIF-2 α binding to the HRE. Thus, we have identified the distal HRE in the SOD-1 gene promoter that mediates DHA's suppression on the transcription of this gene, and demonstrated the involvement of PPAR α and HIF-2 α signaling in this event.

MOL #57430

Introduction

DHA has been shown to induce anticancer activity in multiple experimental model systems. A number of cellular mechanisms have been proposed to play a role in DHA-induced cell death or growth inhibition. Lipid peroxidation is one well-established mechanism to explain the action of DHA on cancer cells. Accumulation of the lipid peroxidation products causes peroxidative damage, ultimately leading to the death of cells. The involvement of lipid peroxidation in DHA-induced growth inhibition and cytotoxicity of cancer cells was demonstrated by the measurement of the peroxidative products in DHA treated cells and by the abilities of antioxidants, such as vitamin E, to block DHA's cytotoxic effects (Gonzalez, 1995). Moreover, the killing of malignant cells by DHA can be accelerated by increasing cellular oxidative stress (Begin et al., 1988).

The balance between oxidants and antioxidants is critical for biological systems to maintain normal function. In eukaryotes, antioxidant defenses are provided by small molecules, such as glutathione, vitamin E, or vitamin C, and enzymes, such as superoxide dismutase (SOD), catalase, and glutathione peroxidase (GPx, (Pacifci and Davies, 1991)). SOD and catalase reduce superoxide anions and hydrogen peroxides, respectively. GPx uses glutathione to reduce hydrogen peroxides. The majority of the total cellular SOD is comprised of SOD-1, which is mainly located in the cytoplasm and nucleus and requires Cu and Zn for its activity (Crapo et al., 1992). SOD-2 is located primarily in the mitochondrial matrix and requires Mn for its activity. Both over-expression and under-expression of SOD-2 have been described in various types of human cancer (Kinnula and Crapo, 2004; Oberley and Oberley, 1997) and it has been suggested that SOD-2 is a tumor suppressor gene. Since all cells actively produce superoxide during

MOL #57430

routine metabolic processes, the altered expression of SOD-2 in many cancer cells (Fridovich, 1995) suggests that malignant cells may be more dependent for survival upon SOD-1 than normal cells. If so, targeted inactivation of SOD-1 is likely to result in preferential tumor cell killing. Indeed, a well-designed study demonstrated that drug therapy, believed to inhibit SOD-1, results in increased levels of superoxide, free radical-mediated damage to mitochondria, and induction of apoptosis in cancer cells (Huang et al., 2000). Furthermore, inhibition of SOD-1 attenuated angiogenesis and selectively induced apoptosis of tumor cells (Juarez et al., 2006).

The rat and human SOD-1 gene promoters contain PPREs (Rojo et al., 2004; Yoo et al., 1999). There is considerable evidence indicating that DHA is a natural ligand for PPARs, which bind to PPREs and regulate expression of a number of genes (Desvergne and Wahli, 1999). Additional transcription factors that might be implicated in DHA's regulation of SOD-1 expression are the HIF and NF- κ B, which have DNA binding elements present in the human SOD-1 gene promoter (Rojo et al., 2004). Our recent results indicate that DHA down-regulates expression of the SOD-1 gene, thereby enhancing oxidative stress and more effectively killing cancer cells (Ding et al., 2004). The down-regulation of SOD-1 gene expression by DHA was observed in a number of human cancer lines (Ding and Lind, 2007). This indicates that not only could DHA initiate lipid peroxidation due to its possessing multiple double bonds, which increases its oxidation potential, but DHA could also suppress the antioxidant enzyme system thereby enhancing oxidative stress. In order to understand how DHA regulates gene expression of antioxidant enzymes, the present study investigated the detailed cellular mechanisms of DHA-induced suppression of SOD-1 gene expression in human cancer cells. The A22780 line, a well-established ovarian cancer line, was chosen as our model system because it is one of the solid tumor lines showing a down-regulation

MOL #57430

of SOD-1 expression by DHA (Ding and Lind, 2007). We report here that DHA targets SOD-1 gene transcription through an interaction of HIF-2 α and PPAR α signaling that acts via the distal HRE in the SOD-1 gene promoter.

MOL #57430

Materials and Methods

Materials Dual-Luciferase[®] Reporter kit was purchased from Promega (Madison, WI). Antibodies were obtained from the following resources: PPAR α , PPAR γ , and HIF-1 α were from Santa Cruz Biotechnology Inc. (Santa Cruz, CA); glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was from ProMab Biotechnologies, Inc. (Albany, CA); and HIF-2 α from Novus Biologicals, Inc (Littleton, CO). The pGL-3-HRE-Lu Reporter construct was kindly provided by Dr. Konstantin Salnikow (Radiation Oncology Branch, NCI, Frederick, MD (Salnikow et al., 1999; Surazynski et al., 2008)). The QuickChange Site-Directed Mutagenesis Kit was from Stratagene (La Jolla, CA). 4-hydroxy Nonenal (4-HNE) was from Cayman (Ann Arbor, MI). DHA, clofibrate, troglitazone, lipopolysaccharide (LPS), and other reagents were analytic grade and purchased from Sigma Chemical Co. (St. Louis, MO).

Cell culture The ovarian carcinoma cell line A2780 was provided by Dr. Stephen Howell (University of California, San Diego). Cells were cultivated in RPMI 1640 medium supplemented with 10% FBS, 100 IU/ml penicillin, and 100 μ g/ml streptomycin, at 37°C in humidified air with 5% CO₂. The cells were sub-cultivated twice a week in 75 mm or 150 mm flasks following a protocol by American Type Culture Collection (ATCC, Manassas, VA). Other cell lines including Panc-1 (pancreatic cancer), MCF-7 (breast cancer), Raji (lymphoma), and CEM (leukemia) were from ATCC and cultured according to ATCC's instructions.

DNA deletion The SOD-1 promoter reporter constructs were kindly provided by Dr. Antonio Guadrado (Autonoma University of Madrid, Madrid, Spain (Rojo et al., 2004)). These include p1499-sod1, p750-sod1, p552-sod1, and p355-sod1. The QuickChange Site-Directed

MOL #57430

Mutagenesis Kit (Stratagene, Lo Jolla, CA) was used to delete DNA binding elements present in the p1499-sod1 promoter, including NF- κ B and HRE. The deletion was performed according to the manufacturer's protocol. The primers used for the deletions are as follows: NF- κ B-del forward primer: 5'-GAAAATTGCAGGGGAAGGATTGAGGTGTAGCGAC-3'; NF- κ B-del reverse primer: 5'-GTCGCTACACCTCAATCCTTCCCCTGCAATTTTC-3'; HRE-del forward primer: 5'-GCCAGACAAAAACGCTCTGTAGGGTTGTGG-3'; HRE-del reverse primer: CCACAACCCTACAGAGCGTTTTTGTCTGGC-3'. The sequence containing a putative NF- κ B (5'-AAAAGGTAAGTCCCGG-3', -480 through -464 of the human SOD-1 gene promoter (Kim et al., 1994)) and a putative HRE (5'-AGGTGATGCCTAGAAGCCAACTAGTTGCCGTTTGGTTA-3', -626 through -589 of the human SOD-1 gene promoter (Kim et al., 1994)) were deleted as confirmed by DNA sequencing.

Transient transfection and luciferase activity assay A2780 cells (2×10^6) were plated in 100 mm cell culture dishes with 8 ml RPMI medium. The cells proliferated to 60-80% confluence after 24 hours of culture. 3 μ g DNA of the reporter constructs were transfected into A2780 cells using the Lipofectamine 2000 Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. In some cases, an additional reporter construct encoding Renilla luciferase was co-transfected with the SOD-1 gene promoter constructs in order to determine the transfection efficiency in each well. 24 hours after the transfection, cells were lifted and randomly plated into 24-well plates at 250,000/well. This ensures a randomly even distribution of transfected cells. At 48 hours of transfection, cells were treated with DHA, clofibrate, troglitazone, or other reagents, at indicated concentrations and durations. Luciferase activity was assayed using the Dual-

MOL #57430

Luciferase[®] Reporter kit from Promega (Madison, WI). In short, cells were washed with cold PBS twice and harvested by adding 100 μ l lysis buffer. The lysate was quickly centrifuged at 13,000 g for 1 minute and the insoluble material was removed. The firefly luciferase reporter was measured first by adding Luciferase Assay Reagent II (50 μ l) to the cell extracts from each sample (50 μ l). The Renilla luciferase reaction was initiated sequentially by adding Stop & Glo[®] Reagent to the same sample. The firefly luciferase activity was normalized by the Renilla luciferase activity or by the amount of protein used for each determination. The data were expressed as percentages of luciferase activity detected in untreated cells.

Western blot analysis The protein samples (40 μ g each) were subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto polyvinylidene difluoride membranes. Following overnight blocking in 5% milk at 4°C, blots were incubated with primary antibodies (PPAR α , PPAR γ , HIF-1 α , HIF-2 α , and GAPDH) at appropriate dilution ratios for 2 hours at 22°C. After 3 \times 10 min washings with Tris-buffered saline with 0.1% Tween 20, the blots were incubated with horseradish peroxidase-conjugated secondary antibodies for 1 hour at 22°C. Chemiluminescent substrate (GE Healthcare) was used to detect binding of the primary antibodies to their cognate antigens and images were captured using the Konica SRX-101 x-ray developer.

Co-immunoprecipitation (Co-IP) Cells growing in 100-mm dishes were washed twice with cold PBS and harvested by adding 150 μ l of IP buffer containing 10 mM Tris-HCl (pH 7.4), 50 mM NaCl, 0.5 mM EDTA, 1 mM PMSF, and 1% Triton X-100. Cells were sonicated for 1-3 min on ice and centrifuged at 16,000 \times g for 30 min to remove insoluble material. Supernatants were

MOL #57430

collected and protein concentrations determined. In a typical IP reaction, the supernatant contained about 100 μ g protein from one sample. Supernatants were pre-cleared by adding 25 μ l of agarose-coupled protein A and gently rotated for 60 min at 4°C. The agarose beads were removed by centrifugation at 2,000 \times g for 5 min and the pre-cleared supernatants were collected. Appropriate primary antibodies or IgG were added to the supernatants (1:100 ratio) and the reaction was incubated at 4°C overnight with gentle rotation. The antibody-protein complexes were captured by adding 50 μ l of agarose-coupled protein A and rotating at 4°C for an additional 2 hours at 4°C. The supernatants were then removed by centrifuging at 2,000 \times g for 5 min. The pellets were washed with IP buffer 4 times and solubilized with 2 \times SDS-PAGE sample buffer at room temperature for 60 min. The samples were centrifuged and the supernatants were collected. The immunoprecipitated proteins were subjected to western blot analysis.

Chromatin immunoprecipitation assay (ChIP) The ChIP assay was done as previously described (Ding et al., 2006). In brief, A2780 cells were treated with 150 μ M DHA, 500 μ M clofibrate, or 20 μ M troglitazone for 16 hours. Formaldehyde was then added directly to the medium (1% final concentration) to fix the cells at 37°C for 10 min. Cells were washed, removed from the dish, pelleted, and lysed on ice in the SDS buffer. The lysate was sonicated to shear DNA into sizes ranging from 200-1,000 bp and the DNA was pre-cleared overnight at 4°C with salmon sperm DNA-saturated protein A agarose. The complex of protein and DNA was precipitated overnight at 4°C using HIF-2 α antibody and the precipitants were incubated at 65°C for 4 hours to reverse cross-links. The precipitants were then digested with protein kinase K and DNA was extracted with phenol/chloroform and precipitated with ethanol. DNA was analyzed by PCR amplification

MOL #57430

using primers specific to the promoter regions of interest and the PCR products were separated on a 1% agarose gel containing ethidium bromide and visualized under ultra violet light. The primers used for DNA amplification of the distal HRE fragment in the SOD-1 gene promoter were as follows: forward, 5'-GCCTTTAGGCCAGAC-3' (-649 to -635); reverse, 5'-CTGAGTTTGGCCACAGCGTC-3' (-333 through -352 (Kim et al., 1994)).

Statistical analysis One-way ANOVA analysis was performed to assess differences among groups of data using the Prism 4 program (GraphPad Software, San Diego, CA).

MOL #57430

Results

DHA down-regulates SOD-1 gene transcription. We recently demonstrated that DHA reduces SOD-1 gene expression at both mRNA and protein levels in human cancer cells (Ding and Lind, 2007; Ding et al., 2004). Because DHA can enter the nucleus and regulate gene transcription (Huang et al., 2002), we hypothesized that DHA down-regulates SOD-1 gene expression by targeting SOD-1 gene transcription regulation. To test this hypothesis, a SOD-1 promoter reporter construct (p1499-sod1) was transfected into A2780 cells and the effects of various DHA doses and treatment times on promoter activity were measured. DHA inhibited SOD-1 gene promoter activity in a time- and concentration-dependent manner (Figure 1). The inhibitory effects were most pronounced after 4 hours of treatment and lasted for at least 20 hours. These results, combined with our previous reports (Ding and Lind, 2007; Ding et al., 2004), clearly show that DHA targets SOD-1 gene transcription in human cancer cells. An additional cell line, Panc-1, was tested to determine whether DHA's inhibitory effects on SOD-1 gene transcription are detectable in cell lines other than A2780. Clearly DHA also inhibited SOD-1 gene transcription in this cell line (Figure 1d).

To understand if the inhibitory effect is DHA specific, cells were treated with 100 μ M linolenic acid (n-6, 18:2), a long-chain n-6 PUFA, for 4 hours. This long-chain n-6 PUFA was unable to reduce SOD-1 gene transcription (Figure 1c). Since lipid peroxidation is a well-established mechanism explaining DHA's cytotoxicity (Ding et al., 2004; Gonzalez, 1995), we tested whether it contributes to DHA's inhibitory effect on SOD-1 gene transcription. The antioxidant vitamin E blocked DHA's cytotoxicity (Figure 2c) but did not alter DHA-induced suppression of SOD-1 gene transcription (Figure 2a), suggesting that the lipid peroxidation products do not

MOL #57430

contribute to the mechanism by which DHA represses SOD-1 expression. This was further supported by the fact that 4-HNE, a lipid peroxide product, did not suppress SOD-1 gene transcription (Figure 2b).

PPAR α ligand mimics DHA's effects on SOD-1 gene transcription. DHA is an established ligand to PPAR and binding of DHA to PPAR regulates gene transcription in mammalian model systems (Diep et al., 2000). To understand whether PPAR α or PPAR γ is involved in DHA's inhibitory effect on SOD-1 gene expression, cells were treated with 20 μ M troglitazone, a PPAR γ agonist (Adams et al., 1997), and 500 μ M clofibrate, a PPAR α agonist (Gottlicher et al., 1992), for 20 hours. The concentrations of troglitazone and clofibrate were chosen based on previous studies (Canuto et al., 2003; Hashimoto et al., 2004; Yang et al., 2007). Luciferase activity assay indicated that clofibrate suppresses SOD-1 gene transcription, whereas troglitazone does not (Figure 3), suggesting that PPAR α , and not PPAR γ , is involved in transcriptional control of SOD-1 gene expression. On the other hand, lipopolysaccharide (LPS), an activator to the NF- κ B signaling pathway, was able to up-regulate SOD-1 gene transcription, a result consistent with a previous report (Marikovsky et al., 2003). There is a putative PPRE in the human SOD-1 gene promoter (Rojo et al., 2004), which is likely to mediate the suppressive effects of DHA and clofibrate on SOD-1 gene transcription through a PPAR α signaling pathway. To test this possibility, A2780 cells were transfected with a series of deletion mutants of the SOD-1 gene promoter reporter constructs, as indicated in Figure 4. Surprisingly, the inhibitory effects of both DHA and clofibrate were evident in cells transfected with the p750-sod1 reporter construct that does not retain the PPRE, but disappeared in cells transfected with the p552-sod1 that lacks the PPRE and distal HRE or the p355-sod1 reporter that lacks the PPRE, distal HRE

MOL #57430

and NF- κ B sites (Figure 5a). This indicates that the PPRE is not required in DHA-induced transcription suppression of the SOD-1 gene. Thus, the critical elements that mediate DHA suppression are located within the 200 bp fragment of the promoter between -755 bp and -552 bp, suggesting that the distal HRE is a likely candidate.

HRE mediates DHA and clofibrate's inhibitory effects on SOD-1 gene transcription. To locate the DNA binding elements that mediate DHA and clofibrate's inhibitory effects on SOD-1 gene transcription, deletions of the NF- κ B and distal HRE DNA binding elements in the intact SOD-1 gene promoter were constructed (Figure 5c). Although deletion of the NF- κ B element did not reverse DHA and clofibrate's inhibition, deletion of the distal HRE element completely reversed the inhibitory effects on SOD-1 gene transcription (Figure 5a,b), indicating the key role of this HRE in the transcription suppression of the SOD-1 gene by DHA and clofibrate. To further confirm this observation, the pGL-3-HRE-luc construct (Salnikow et al., 1999) containing only HRE binding elements in the promoter was transfected into A2780 cells, and the effects of DHA and clofibrate on the reporter activity were analyzed. Cobalt chloride was included as a positive control because it enhances HRE-mediated transcription (Bianchi et al., 1999). Consistent with the DHA inhibition of SOD-1 promoter activity in Figure 5, both DHA and clofibrate suppressed HRE-mediated luciferase activity (Figure 6).

PPAR α complexes with HIF-2 α to mediate DHA's effects on SOD-1 gene transcription. Western blot analysis with specific antibodies confirmed that PPAR γ and PPAR α are expressed in A2780 cells (Tuller et al., 2009). Furthermore, HIF-2 α is constitutively expressed in this cell line, whereas HIF-1 α is not detectable in the presence and absence of DHA. Treatment of the cells

MOL #57430

with cobalt induced HIF-1 α expression, consistent with a previous report (Bianchi et al., 1999), which serves as a positive control (Figure 7). To understand whether PPAR α might act with HIF-2 α to regulate HRE-responsive gene expression, we applied co-immunoprecipitation using cellular proteins extracted from DHA-treated cells. Interestingly, in the HIF-2 α precipitants, PPAR α but not PPAR γ was clearly detected (Figure 8). DHA decreased the levels of HIF2 α in ovarian cancer cells, as well as in several other human cancer cell lines including breast cancer (MCF-7), lymphoma (Raji), and leukemia (CEM) lines (data not shown). Consequently, the interaction of PPAR α and HIF-2 α was affected by DHA in these cells (Figure 8). We then performed ChIP analysis to determine whether HIF-2 α is capable of binding to the distal HRE of the SOD-1 promoter thereby regulating SOD-1 gene transcription. A2780 cells were treated with 150 μ M DHA and 500 μ M clofibrate for 20 hours. Specific primers covering the DNA fragment surrounding the distal HRE of the human SOD-1 gene promoter were used to amplify HIF-2 α antibody-precipitated DNA. This showed that HIF-2 α is able to bind the human SOD-1 promoter and this binding was indeed significantly attenuated in DHA- and clofibrate-treated cells, indicating that the HIF-2 α signaling mediates transcription suppression of the SOD-1 gene (Figure 9).

MOL #57430

Discussion

The present study demonstrates that DHA down-regulates SOD-1 gene transcription through a HRE-mediated mechanism, involving an interaction of PPAR α and HIF-2 α signaling. This provides a novel mechanism by which DHA weakens the antioxidant status of cancer cells, resulting in enhanced oxidative stress, growth inhibition, or apoptotic cell death (Ding et al., 2004). The concentration of DHA used in the present study has been frequently used in *in vitro* model systems (Collett et al., 2001; Leonardi et al., 2005; Merendino et al., 2005; Ng et al., 2005). A previous study in human subjects reported that the highest tolerated dose of EPA plus DHA is 188 mg/kg/day, orally (Burns et al., 1999) which is considerably higher than the usual (“physiological”) adult intake. There is no data published to indicate what molar concentration of DHA is achieved in plasma and cells when given at these pharmacological doses. However a recent study reported that daily consumption of 4.4g EPA plus DHA over 6 weeks resulted in a plasma DHA concentration at 51 μ g/mL (155 μ M) in humans, which is three times higher than the basal level (Garg et al., 2006). Based on these previous reports, we believe that the concentrations of DHA used in our study are physiologically relevant.

Targeting SOD-1 has been recognized as a potential strategy for cancer therapy (Huang et al., 2000). We recently demonstrated that DHA suppression of SOD-1 gene expression contributes to its cytotoxicity in human cancer cells (Ding et al., 2004). The cellular mechanisms of DHA’s action that lead to reduced expression of the SOD-1 gene remains unknown. In the present study, we approached this question by focusing on transcriptional regulation of the SOD-1 gene, because DHA is an established ligand to nuclear receptors and is known to regulate gene transcription in mammalian systems (Desvergne and Wahli, 1999). It was found that DHA

MOL #57430

suppresses SOD-1 gene transcription whereas linolenic acid (n-6, 18:2) does not affect SOD-1 promoter activity. Further, this suppression of SOD-1 gene transcription could not be reversed by pre-treatment of the cells with vitamin E, suggesting that lipid peroxidation products are not involved in this process. These findings support a new model for DHA to exert its cytotoxic effects on tumor cells: on the one hand, DHA is a long-chain n-3 PUFA which has high oxidative potential due to its possession of multiple double bonds (Hawkins et al., 1998). Indeed, lipid peroxidation is a well-accepted mechanism of DHA's action to kill cancer cells (Begin et al., 1988; Gonzalez, 1995). On the other hand, DHA is also a nuclear receptor ligand that can enter the nucleus by binding to fatty acid-binding proteins (Huang et al., 2002) and modulating gene transcription. Importantly, we have showed that DHA reduces expression of antioxidant enzymes, resulting in weakened antioxidant forces and more effective cancer cell killing (Ding et al., 2004). Thus, by initiating lipid peroxidation and inhibiting antioxidant power, DHA is able to more effectively kill cancer cells. As DHA is a natural product and holds the potential to be used for cancer prevention or treatment (Bougnoux, 1999; Hardman, 2004), understanding the detailed mechanisms of DHA's action is of importance in the development of novel strategies in using DHA as an anticancer agent.

Several DNA binding elements such as PPRE, NF- κ B, and HRE are previously described in the human SOD-1 gene promoter, which may contribute to SOD-1 gene transcription regulation (Rojo et al., 2004). Deletion analysis of the human SOD-1 gene promoter enabled us to locate the DNA binding element that is responsible for DHA-induced suppression of SOD-1 gene transcription. While we initially assumed that the PPRE and the NF- κ B DNA binding elements present in the SOD-1 gene promoter are most likely involved in DHA's inhibitory effect on

MOL #57430

SOD-1 gene expression, the experimental results disagreed with this assumption. In fact, several lines of evidence support the key role of the distal HRE of the SOD-1 gene promoter in mediating DHA's suppressive effect in our model system. First, while the PPAR α ligand clofibrate mimicked DHA's inhibitory effects on SOD-1 gene transcription, deletion of both the PPRE and the NF- κ B elements did not reverse or attenuate DHA and clofibrate's inhibitory effects. This suggests that these two DNA binding elements are not required for the transcription suppression of the SOD-1 gene by DHA. Additionally, deletion of the distal HRE element in the SOD-1 gene promoter completely reversed the inhibitory effects of DHA and clofibrate, indicating the essential role of the HRE in mediating this event. Second, application of the HRE-lu allowed us to further determine the involvement of this DNA binding element in DHA- and clofibrate-induced transcription suppression. It was found that DHA and clofibrate inhibited HRE-mediated promoter activity to an extent similar to their effects on the SOD-1 gene transcription, providing direct evidence of DHA targeting HRE-mediated transcription. Third, the essential role of HRE in mediating DHA's effects on the SOD-1 gene was further supported by the findings that HIF-2 α binds to the SOD-1 gene promoter as shown via ChIP analysis; and that both DHA and clofibrate can reduce this binding. A direct interaction of PPAR α , but not PPAR γ , with HIF-2 α was also evident in this model system. Our experimental results thus suggest that PPAR α and HIF-2 α signaling regulate DHA's inhibitory effect on SOD-1 gene transcription via the distal HRE element.

While the interaction of the PPAR α and NF- κ B signaling pathways has been well established in several model systems (Delerive et al., 1999; Delerive et al., 2000; Okamoto et al., 2005; Staels et al., 1998), the direct interaction of PPAR α and HIF proteins to regulate gene transcription has

MOL #57430

not been previously described. Our results indicate that PPAR α and HIF-2 α form a complex that might bind to the HREs and regulate gene expression. To the best of our knowledge, this is the first report of physical interaction between the HIF-2 α and PPAR α transcription factors. This novel observation of cross-talk between the PPAR and HIF signaling pathways provides insight into our understanding of the regulation of the SOD-1 gene. These observations are consistent with a recent study reporting that n-3 PUFAs, including EPA and DHA, down-regulate HIF signaling in colon cancer cells (Calviello et al., 2004), and a previous report that HRE mediates the induction of SOD-1 gene transcription (Yoo et al., 1999).

MOL #57430

Acknowledgement

MOL #57430

References

- Adams M, Montague CT, Prins JB, Holder JC, Smith SA, Sanders L, Digby JE, Sewter CP, Lazar MA, Chatterjee VK and O'Rahilly S (1997) Activators of peroxisome proliferator-activated receptor gamma have depot-specific effects on human preadipocyte differentiation. *J Clin Invest* **100**(12):3149-3153.
- Begin ME, Ells G and Horrobin DF (1988) Polyunsaturated fatty acid-induced cytotoxicity against tumor cells and its relationship to lipid peroxidation. *J Natl Cancer Inst* **80**(3):188-194.
- Bianchi L, Tacchini L and Cairo G (1999) HIF-1-mediated activation of transferrin receptor gene transcription by iron chelation. *Nucleic Acids Res* **27**(21):4223-4227.
- Bougnoux P (1999) n-3 polyunsaturated fatty acids and cancer. *Curr Opin Clin Nutr Metab Care* **2**(2):121-126.
- Burns CP, Halabi S, Clamon GH, Hars V, Wagner BA, Hohl RJ, Lester E, Kirshner JJ, Vinciguerra V and Paskett E (1999) Phase I clinical study of fish oil fatty acid capsules for patients with cancer cachexia: cancer and leukemia group B study 9473. *Clin Cancer Res* **5**(12):3942-3947.
- Calviello G, Di Nicuolo F, Gagnoli S, Piccioni E, Serini S, Maggiano N, Tringali G, Navarra P, Ranelletti FO and Palozza P (2004) n-3 PUFAs reduce VEGF expression in human colon cancer cells modulating the COX-2/PGE2 induced ERK-1 and -2 and HIF-1alpha induction pathway. *Carcinogenesis* **25**(12):2303-2310.
- Canuto RA, Muzio G, Maggiora M, Trombetta A, Martinasso G, Autelli R, Costelli P, Bonelli G and Baccino FM (2003) Apoptosis induced by clofibrate in Yoshida AH-130 hepatoma cells: role of HMG-CoA reductase. *J Lipid Res* **44**(1):56-64.

MOL #57430

Collett ED, Davidson LA, Fan YY, Lupton JR and Chapkin RS (2001) n-6 and n-3 polyunsaturated fatty acids differentially modulate oncogenic Ras activation in colonocytes. *Am J Physiol Cell Physiol* **280**(5):C1066-1075.

Crapo JD, Oury T, Rabouille C, Slot JW and Chang LY (1992) Copper,zinc superoxide dismutase is primarily a cytosolic protein in human cells. *Proc Natl Acad Sci U S A* **89**(21):10405-10409.

Delerive P, De Bosscher K, Besnard S, Vanden Berghe W, Peters JM, Gonzalez FJ, Fruchart JC, Tedgui A, Haegeman G and Staels B (1999) Peroxisome proliferator-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1. *J Biol Chem* **274**(45):32048-32054.

Delerive P, Gervois P, Fruchart JC and Staels B (2000) Induction of IkappaBalpha expression as a mechanism contributing to the anti-inflammatory activities of peroxisome proliferator-activated receptor-alpha activators. *J Biol Chem* **275**(47):36703-36707.

Desvergne B and Wahli W (1999) Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* **20**(5):649-688.

Diep QN, Touyz RM and Schiffrin EL (2000) Docosahexaenoic acid, a peroxisome proliferator-activated receptor-alpha ligand, induces apoptosis in vascular smooth muscle cells by stimulation of p38 mitogen-activated protein kinase. *Hypertension* **36**(5):851-855.

Ding WQ and Lind SE (2007) Phospholipid hydroperoxide glutathione peroxidase plays a role in protecting cancer cells from docosahexaenoic acid-induced cytotoxicity. *Mol Cancer Ther* **6**(4):1467-1474.

Ding WQ, Liu B, Vaught JL, Palmiter RD and Lind SE (2006) Clioquinol and docosahexaenoic acid act synergistically to kill tumor cells. *Mol Cancer Ther* **5**(7):1864-1872.

MOL #57430

Ding WQ, Vaught JL, Yamauchi H and Lind SE (2004) Differential sensitivity of cancer cells to docosahexaenoic acid-induced cytotoxicity: the potential importance of down-regulation of superoxide dismutase 1 expression. *Mol Cancer Ther* **3**(9):1109-1117.

Fridovich I (1995) Superoxide radical and superoxide dismutases. *Annu Rev Biochem* **64**:97-112.

Garg ML, Leitch J, Blake RJ and Garg R (2006) Long-chain n-3 polyunsaturated fatty acid incorporation into human atrium following fish oil supplementation. *Lipids* **41**(12):1127-1132.

Gonzalez MJ (1995) Fish oil, lipid peroxidation and mammary tumor growth. *J Am Coll Nutr* **14**(4):325-335.

Gottlicher M, Widmark E, Li Q and Gustafsson JA (1992) Fatty acids activate a chimera of the clofibrin acid-activated receptor and the glucocorticoid receptor. *Proc Natl Acad Sci U S A* **89**(10):4653-4657.

Hardman WE (2004) (n-3) fatty acids and cancer therapy. *J Nutr* **134**(12 Suppl):3427S-3430S.

Hashimoto F, Oguchi Y, Morita M, Matsuoka K, Takeda S, Kimura M and Hayashi H (2004) PPARalpha agonists clofibrate and gemfibrozil inhibit cell growth, down-regulate hCG and up-regulate progesterone secretions in immortalized human trophoblast cells. *Biochem Pharmacol* **68**(2):313-321.

Hawkins RA, Sangster K and Arends MJ (1998) Apoptotic death of pancreatic cancer cells induced by polyunsaturated fatty acids varies with double bond number and involves an oxidative mechanism. *J Pathol* **185**(1):61-70.

Huang H, Starodub O, McIntosh A, Kier AB and Schroeder F (2002) Liver fatty acid-binding protein targets fatty acids to the nucleus. Real time confocal and multiphoton fluorescence imaging in living cells. *J Biol Chem* **277**(32):29139-29151.

MOL #57430

Huang P, Feng L, Oldham EA, Keating MJ and Plunkett W (2000) Superoxide dismutase as a target for the selective killing of cancer cells. *Nature* **407**(6802):390-395.

Juarez JC, Betancourt O, Jr., Pirie-Shepherd SR, Guan X, Price ML, Shaw DE, Mazar AP and Donate F (2006) Copper binding by tetrathiomolybdate attenuates angiogenesis and tumor cell proliferation through the inhibition of superoxide dismutase 1. *Clin Cancer Res* **12**(16):4974-4982.

Kim HT, Kim YH, Nam JW, Lee HJ, Rho HM and Jung G (1994) Study of 5'-flanking region of human Cu/Zn superoxide dismutase. *Biochem Biophys Res Commun* **201**(3):1526-1533.

Kinnula VL and Crapo JD (2004) Superoxide dismutases in malignant cells and human tumors. *Free Radic Biol Med* **36**(6):718-744.

Leonardi F, Attorri L, Di Benedetto R, Di Biase A, Sanchez M, Nardini M and Salvati S (2005) Effect of arachidonic, eicosapentaenoic and docosahexaenoic acids on the oxidative status of C6 glioma cells. *Free Radic Res* **39**(8):865-874.

Marikovsky M, Ziv V, Nevo N, Harris-Cerruti C and Mahler O (2003) Cu/Zn superoxide dismutase plays important role in immune response. *J Immunol* **170**(6):2993-3001.

Merendino N, Loppi B, D'Aquino M, Molinari R, Pessina G, Romano C and Velotti F (2005) Docosahexaenoic acid induces apoptosis in the human PaCa-44 pancreatic cancer cell line by active reduced glutathione extrusion and lipid peroxidation. *Nutr Cancer* **52**(2):225-233.

Ng Y, Barhoumi R, Tjalkens RB, Fan YY, Kolar S, Wang N, Lupton JR and Chapkin RS (2005) The role of docosahexaenoic acid in mediating mitochondrial membrane lipid oxidation and apoptosis in colonocytes. *Carcinogenesis* **26**(11):1914-1921.

MOL #57430

Oberley TD and Oberley LW (1997) Antioxidant enzyme levels in cancer. *Histol Histopathol* **12**(2):525-535.

Okamoto H, Iwamoto T, Kotake S, Momohara S, Yamanaka H and Kamatani N (2005) Inhibition of NF-kappaB signaling by fenofibrate, a peroxisome proliferator-activated receptor-alpha ligand, presents a therapeutic strategy for rheumatoid arthritis. *Clin Exp Rheumatol* **23**(3):323-330.

Pacifici RE and Davies KJ (1991) Protein, lipid and DNA repair systems in oxidative stress: the free-radical theory of aging revisited. *Gerontology* **37**(1-3):166-180.

Rojo AI, Salinas M, Martin D, Perona R and Cuadrado A (2004) Regulation of Cu/Zn-superoxide dismutase expression via the phosphatidylinositol 3 kinase/Akt pathway and nuclear factor-kappaB. *J Neurosci* **24**(33):7324-7334.

Salnikow K, An WG, Melillo G, Blagosklonny MV and Costa M (1999) Nickel-induced transformation shifts the balance between HIF-1 and p53 transcription factors. *Carcinogenesis* **20**(9):1819-1823.

Staels B, Koenig W, Habib A, Merval R, Lebret M, Torra IP, Delerive P, Fadel A, Chinetti G, Fruchart JC, Najib J, Maclouf J and Tedgui A (1998) Activation of human aortic smooth-muscle cells is inhibited by PPARalpha but not by PPARgamma activators. *Nature* **393**(6687):790-793.

Surazynski A, Donald SP, Cooper SK, Whiteside MA, Salnikow K, Liu Y and Phang JM (2008) Extracellular matrix and HIF-1 signaling: the role of prolylase. *Int J Cancer* **122**(6):1435-1440.

MOL #57430

Tuller ER, Brook AL, Yu HJ, Lou JR, Benbrook DM and Ding WQ (2009) PPARalpha signaling mediates the synergistic cytotoxicity of clioquinol and docosahexaenoic acid in human cancer cells. *Biochem Pharmacol* 77(9):1480-1486.

Yang CC, Wang YC, Wei S, Lin LF, Chen CS, Lee CC, Lin CC and Chen CS (2007) Peroxisome proliferator-activated receptor gamma-independent suppression of androgen receptor expression by troglitazone mechanism and pharmacologic exploitation. *Cancer Res* 67(7):3229-3238.

Yoo HY, Chang MS and Rho HM (1999) The activation of the rat copper/zinc superoxide dismutase gene by hydrogen peroxide through the hydrogen peroxide-responsive element and by paraquat and heat shock through the same heat shock element. *J Biol Chem* 274(34):23887-23892.

MOL #57430

Footnotes

Financial support was obtained from the Oklahoma Center for the Advancement of Science and Technology [HR04-021], the American Cancer Society [IRG-05-066-01], and the College of Medicine, University of Oklahoma Health Sciences Center [COM 22134].

MOL #57430

Figure legend

Figure 1. DHA down-regulates human SOD-1 promoter activity. The p1499-sod1 reporter construct was transfected into A2780 cells and the cells were treated with 100 μ M DHA for 1, 4, and 20 hours (**A**), with various concentrations of DHA for 4 hours (**B**), or with 100 μ M linolenic acid (LA, n-6, 18:2) for 1 and 4 hours (**C**). Panc-1 cells were also transfected and treated with 150 μ M DHA for 4 hours (**D**). The cell lysates were prepared and luciferase activity assayed. Data are expressed as percentages of untreated cells (Bar, SEM, n=3). ★ $P < 0.01$, compared to untreated cells, using One-way ANOVA followed by Dunnett analysis.

Figure 2. Vitamin E blocks DHA's cytotoxicity but not inhibition of SOD-1 promoter activity. A2780 cells were pre-treated with 100 μ M vitamin E (Vit E) for 15 min before addition of 100 μ M DHA for 4 hours (**A**) or various concentrations of DHA for 72 hours (**C**). Cells were also treated with 100 μ M 4-HNE for 4 hours (**B**). Luciferase activity and cell viability were assayed as described in the Materials and Methods section. Data are expressed as percentages of untreated control cells. (Bar, SEM, n=3).

Figure 3. Clofibrate down-regulates human SOD-1 promoter activity. The p1499-sod1 reporter construct was transfected into A2780 cells and the cells were treated with 500 μ M clofibrate, 20 μ M troglitazone, or 1 μ g/ml LPS for 16 hours. The cell lysates were prepared and luciferase activity assayed. Data are expressed as percentages of untreated cells (Bar, SEM, n=3). ★ $P < 0.01$, compared to untreated cells, using One-way ANOVA followed by Dunnett analysis.

MOL #57430

Figure 4. Human SOD-1 gene promoter. Top panel, a diagram showing different DNA fragments of the human SOD-1 gene promoter that have been cloned into pGL-3 luciferase reporter construct. Major DNA binding elements in the promoter are indicated. **Bottom panel,** relative promoter activity detected in A2780 cells. Cells were transfected with different SOD-1 promoter reporter constructs along with the renilla luciferase construct. Luciferase activity was assayed and the firefly luciferase activity was normalized by the renilla luciferase activity. Data are expressed as percentages of the luciferase activity detected in p1499-sod1 transfected cells (Bar, SEM, n=3).

Figure 5. The distal HRE in the human SOD-1 gene promoter is essential for DHA-and clofibrate-induced suppression of SOD-1 gene transcription. A2780 cells were transfected with p750-sod1, p552-sod1, p355-sod1 (**A**), or p1499-sod1, HRE-del, NF- κ B del (**B**, **C**) reporter constructs. Cells were treated with 500 μ M clofibrate or 100 μ M DHA for 16 hours. The cell lysates were prepared and luciferase activity assayed. Data are expressed as percentages of untreated cells (Bar, SEM, n=3). ★ $P < 0.01$, compared to untreated cells, using One-way ANOVA followed by Dunnett analysis.

Figure 6. DHA and clofibrate suppress HRE-mediated gene transcription. A2780 cells were transfected with the pGL-3-HRE-Lu Reporter construct. Cells were treated with 500 μ M clofibrate or 150 μ M DHA for 16 hours. The cell lysates were prepared and luciferase activity assayed. Data are expressed as percentages of untreated cells (Bar, SEM, n=3). ★ $P < 0.01$, compared to untreated cells, using One-way ANOVA followed by Dunnett analysis.

MOL #57430

Figure 7. HIF-2 α is constitutively expressed in A2780 cells. A2780 cells were treated with 100 μ M DHA or 150 μ M cobalt for 16 hours. The cell lysates were prepared and blotted with antibodies against HIF-1 α , HIF-2 α , and GAPDH. Shown are representative gels of three separate experiments.

Figure 8. PPAR α complexes with HIF-2 α in A2780 cells. A2780 cells were treated with 100 μ M DHA for 16 hours and cell lysates were prepared. The cell lysates were precipitated with HIF-2 α (p-HIF-2 α) antibody or IgG (p-IgG) as described in the Materials and Methods section. The precipitants as well as the total cell lysates were separated onto SDS PAGE, and blotted with antibodies against PPAR γ , PPAR α , HIF-2 α and GAPDH. Shown are representative gels of three separate experiments.

Figure 9. DHA and clofibrate reduce HIF-2 α binding to the human SOD-1 gene promoter. A2780 cells were treated with 500 μ M clofibrate or 150 μ M DHA for 16 hours. ChIP assay was performed as described in the Materials and Methods section. The amplified DNA fragments were separated on 1% agarose gel and visualized under ultra violet light. Shown are representative gels of two separate experiments.

Figure 1






Figure 2

A



B



C




Figure 3




Figure 4





Figure 5






Figure 6

Figure 7




Figure 8




Figure 9

