

**MOL Manuscript # 80689**

## **A novel EPAC specific inhibitor suppresses pancreatic cancer cell migration and invasion**

**Muayad Almahariq, Tamara Tsalkova, Fang C. Mei, Haijun Chen, Jia Zhou, Sarita K. Sastry, Frank Schwede and Xiaodong Cheng**

Department of Pharmacology and Toxicology (T.T., F.M., M.A., H.C., J.Z., X.C.) and Sealy Center for Structural Biology and Molecular Biophysics (X.C.), Department of Biochemistry and Molecular Biology (S.S.), The University of Texas Medical Branch, Galveston, Texas 77555-0615; BIOLOG Life Science Institute, Forschungslabor und Biochemica-Vertrieb GmbH, Flughafendamm 9a, D-28199 Bremen, Germany (F.S.)

**MOL Manuscript # 80689**

**Running Title:** Characterization of an EPAC-specific inhibitor

**Address Correspondence to:** Xiaodong Cheng, Department of Pharmacology and Toxicology, The University of Texas Medical Branch, 301 University Boulevard, Galveston, Texas 77555-0615, Tel. 409-772-9656; Fax. 409-772-7050; Email: [xcheng@utmb.edu](mailto:xcheng@utmb.edu)

Number of text page: 16

Number of figures: 7

Number of references: 50

Number of words in abstract: 130

Number of words in introduction: 507

Number of words in discussion: 332

**List of nonstandard abbreviations used in the paper:** 007, 8-(4- Chlorophenylthio)- 2'- O-methyladenosine- 3', 5'- cyclic monophosphate; 007-AM, 007 acetoxymethyl ester; 8-NBD-cAMP, 8-[[2-[(7-nitro-4-benzofurazanyl)amino]ethyl]thio]adenosine-3',5'-cyclic monophosphate; cAMP, adenosine-3',5'-cyclic monophosphate. EPAC, exchange protein directly activated by cAMP; GEF, guanine nucleotide exchange factor; HTS, high throughput screening; Mant-GDP, 2'-/3'-O-(*N*'-methylantraniloyl)guanosine-5'-O- diphosphate; PKA, protein kinase A/cAMP-dependent protein kinase; Rap, Ras-proximate.

## **ABSTRACT**

Exchange protein directly activated by cAMP (EPAC) and cAMP-dependent protein kinase (PKA) are two intracellular receptors that mediate the effects of the prototypic second messenger cAMP. Identifying pharmacological probes for selectively modulating EPAC activity represents a significant unmet need within the research field. Herein, we report the identification and characterization of ESI-09, a novel non-cyclic nucleotide EPAC antagonist that is capable of specifically blocking intracellular EPAC-mediated Rap1 activation and Akt phosphorylation, as well as EPAC-mediated insulin secretion in pancreatic beta cells. Using this novel EPAC-specific inhibitor, we have probed the functional roles of over-expression of EPAC1 in pancreatic cancer cells. Our studies show that EPAC1 plays an important role in pancreatic cancer cell migration and invasion, and thus represents a potential target for developing novel therapeutic strategies for pancreatic cancer.

## **INTRODUCTION**

The major physiological effects of cAMP in mammalian cells are transduced by two ubiquitously expressed intracellular cAMP receptor families: the classic protein kinase A/cAMP-dependent protein kinase (PKA/cAPK), and the more recently discovered exchange protein directly activated by cAMP/cAMP regulated guanine nucleotide exchange factor (EPAC/cAMP-GEF) (Cheng et al., 2008). Like PKA, EPAC contains an evolutionarily conserved cAMP-binding domain that acts as a molecular switch for sensing intracellular levels of the second messenger cAMP, and activates the down-stream signaling molecules small GTPases Rap1 and Rap2 (de Rooij et al., 1998; Kawasaki et al., 1998). In addition, EPAC proteins exert their functions through interactions other cellular partners at specific cellular loci. For example, EPAC1 are known to associated with mitotic spindle, plasma membrane and nuclear membrane by interacting with tubulin (Qiao et al., 2002; Mei and Cheng, 2005), ezrin-radixin-moesin (ERM) proteins (Gloerich et al., 2010; Ross et al., 2011) and uncloporin RanBP2 (Liu et al., 2010; Gloerich et al., 2011), respectively. On the other hands, EPAC2 can interact with Rim (Rab3 interacting molecule) and Rim2 (Kashima et al., 2001; Ozaki et al., 2000), as well as a structurally related calcium sensor Piccolo (Fujimoto et al., 2002). In pancreatic beta cells, interactions among EPAC2, Rim2 and Piccolo are critical for cAMP-mediated insulin secretion (Ozaki et al., 2000; Kashima et al., 2001; Fujimoto et al., 2002).

More than a decade of extensive studies have now firmly established that many cAMP-related cellular processes, previously thought to be controlled by PKA alone, are also mediated by EPAC (Gloerich and Bos, 2010). For example, EPAC proteins have been implicated in regulating exocytosis and secretion (Ozaki et al., 2000; Seino and Shibasaki, 2005; Maillet et al., 2003; Li et al., 2007), cell adhesion (Enserink et al., 2004; Rangarajan et al., 2003), endothelial barrier junctions

## **MOL Manuscript # 80689**

(Cullere et al., 2005;Kooistra et al., 2005), leptin signaling (Fukuda et al., 2011), and cardiac functions (Metrich et al., 2010). Interestingly, although PKA and EPAC are activated by the same second messenger cAMP, they can act antagonistically in controlling various cellular functions such as Akt/PKB phosphorylation (Mei et al., 2002;Brennesvik et al., 2005) and hedgehog signaling (Ji et al., 2007). On the other hand, depending upon the specific cellular context, EPAC and PKA can exert synergistic effects on downstream signaling such as stimulation of neurotensin secretion (Li et al., 2007) and attenuation of cAMP signaling through phosphodiesterases (Dodge-Kafka et al., 2005). Despite these significant advances in our knowledge, the lack of EPAC specific antagonists has greatly limited our ability to discern the physiological functions of this family of proteins and understand their role in numerous diseases. For example, EPAC1 is over-expressed in human pancreatic ductal adenocarcinoma (PDA) specimens (Lorenz et al., 2008), but the implications of this over-expression are unclear. To address the lack of EPAC inhibitors, we have developed a robust high throughput screening (HTS) assay for identifying new chemical probes specifically targeting EPAC proteins (Tsalkova et al., 2012a). In this study, we describe the identification and characterization of ESI-09, a novel EPAC specific inhibitor, which enabled us to demonstrate that EPAC1 plays an important role in pancreatic cancer cell migration and invasion.

## **MATERIALS AND METHODS**

**Reagents.** 8-(4-Chlorophenylthio)-2'-O-methyladenosine-3', 5'-cyclic monophosphate, acetoxymethyl ester (007-AM), initially provided by and subsequently purchased from BioLog Life Science Institute (Bremen, Germany), is the highly membrane-permeant and potent acetoxymethyl ester prodrug (Vliem et al., 2008; Chepurny et al., 2009) of 8-(4-Chlorophenylthio)-2'-O-methyladenosine-3', 5'-cyclic monophosphate (007), the first selective EPAC agonist that is more than 100-fold less potent for PKA (Enserink et al., 2002; Christensen et al., 2003). After cell penetration of the uncharged prodrug, the active parental compound 007 is released by endogenous esterases and trapped inside the cell due to the negatively charged cyclic phosphate moiety. 8-[[2-[(7-nitro-4-benzofurazanyl)amino]ethyl]thio] adenosine-3',5'-cyclic monophosphate (8-NBD-cAMP) was also obtained from BioLog Life Science Institute. 2'-/3'-O-(*N'*-methylantraniloyl)guanosine-5'-O-diphosphate (MANT-GDP) was purchased from Molecular Probes, Invitrogen (Carlsbad, CA, USA). All other reagents were purchased through Sigma-Aldrich (St. Louis, MO, USA).

**Antibodies.** Antibodies specific against Akt (#9272), phosphor-Akt (T308) (#4056), phosphor-Akt (S473) (#9271), EPAC1 (#4155) and EPAC2 (#4156) were purchased from Cell Signaling Technology, while anti-Rap1 antibody (SC-65) was obtained from Santa Cruz Biotechnology, Inc.

**Protein expression and purification.** Recombinant EPAC1, EPAC2 and C-terminal truncated Rap1B(1-167) were purified as described previously (Mei and Cheng, 2005; Li et al., 2011). PKA RI $\alpha$ , RII $\beta$  and catalytic subunits were recombinantly expressed in *E. coli* and purified to homogeneity as reported (Cheng et al., 2001). Type I and II PKA holoenzymes were reconstituted from individually purified recombinant PKA R and C subunits (Yu et al., 2004). All proteins used in this study were at least 95% pure, as judged by SDS PAGE.

## **MOL Manuscript # 80689**

**High throughput screening (HTS).** Fluorescence intensity of 8-NBD-cAMP in complex with EPAC2 has been used as a primary signal in HTS (Tsalkova et al., 2012a). A primary screen of Maybridge HitFinder collection of compounds (Maybridge, Thermo Fisher Scientific, MA, USA) was performed in black 384-well low volume microplates (Corning Costar, Cambridge, MA, USA). Briefly, a protein solution containing 65 nM EPAC2 and 75 nM 8-NBD-cAMP was dispensed into 384-well plate (35µl/well) using BiomekFX Laboratory Automation Workstation equipped with 96-multichannel pipetting head (Beckman Coulter, CA, USA). Test compounds (1µl/well) were added from 10 mM stock solutions in DMSO. 10 mM cAMP or DMSO was used as a positive or negative control, respectively. Fluorescence intensity signal was recorded at room temperature before and after tested compounds were added using SpectaMaxM5 microplate reader (Molecular Devices, Silicon Valley, CA, USA) with excitation/emission wavelengths set at 470/540 nm.

**In vitro guanine nucleotide exchange factor (GEF) activity assay of EPACs proteins.** In vitro EPAC activity was measured using purified Rap1B(1-167) loaded with Mant-GDP as previously described (Tsalkova et al., 2009).

**In vitro PKA activity assay** – PKA kinase activities of the type I and II PKA holoenzymes were measured spectrophotometrically with a coupled enzyme assay using purified and reconstituted recombinant PKA holoenzymes (Cook et al., 1982).

**Cell culture.** AsPC-1 and PANC-1 cells with suppressed EPAC1 expression were generated using MISSION™ TRC lentiviral based shRNA according to the manufacturer's protocol. AsPC-1 and PANC-1 cells were maintained in RPMI 1640 medium with 10% fetal bovine serum (FBS). INS-1 cells (passage numbers 70–90) were maintained in RPMI 1640 medium containing 10 mM Hepes, 10% FBS, 2.0 mM L-glutamine, 1.0 mM sodium pyruvate, and 50 µM 2-mercaptoethanol. All cells were kept in a humidified incubator (95% air, 5% CO<sub>2</sub>) at 37 °C.

## MOL Manuscript # 80689

**Phosphorylation of Akt.** Cells were starved of serum for 24 hours before treatment with ESI-09 for 5 minutes. Cells were then treated with 10  $\mu$ M 007-AM for 15 minutes and cellular proteins were extracted in sodium dodecyl sulfate (SDS) lysis buffer (2% SDS, 10% glycerol, 60 mM Tris, pH 6.8). Non-treated cells, and cells treated with 10  $\mu$ M 007-AM only, were used as negative and positive controls respectively. Approximately 10  $\mu$ g of total protein extract were separated by SDS-PAGE and transferred to a polyvinylidene difluoride membrane. Akt phosphorylation was probed with anti-phospho-AKT T308 and S473 antibodies (1:1000). Phosphorylation levels were determined with densitometry and expressed as a percentage of basal Akt phosphorylation (negative control). At least three independent experiments were performed for each Western blot.

**Insulin secretion assay.** INS-1 cells were plated in 96-well plates pre-coated with poly-lysine at a density of  $1 \times 10^5$  cells/well. After overnight incubation, the medium was replaced with Krebs-Ringer buffer (KRB) containing 2.9 mM glucose. After an additional two-hour incubation, the cells were treated with ESI-09 or DMSO vehicle as a control in fresh KRB containing 11.8 mM glucose for 10 min, followed by a 30 min stimulation with 10  $\mu$ M 007-AM. The supernatant was collected and insulin was quantified using an Ultra-sensitive Rat Insulin ELISA kit from Crystal Chem Inc.

**Transwell migration/invasion assay.** The top chambers of 8 micron inserts (Costar Inc) were coated with BD Matrigel™ Basement Membrane Matrix (50  $\mu$ g/mL). Cells ( $2 \times 10^5$ ) pretreated with ESI-09 for 24 hours were added to the top chamber of the inserts in serum free RPMI medium containing 0.25% BSA. The bottom chamber was filled with 600.0  $\mu$ L of RPMI containing 10% FBS and ESI-09. The cells were then incubated at 37 °C in 5% CO<sub>2</sub> for 20 hours. Cells were removed from the top chamber and migrated cells were fixed in methanol and stained with crystal violet. The numbers of migrated cells were counted in four different fields. Three independent experiments were performed for each ESI-09 concentration.



## MOL Manuscript # 80689

**Wound healing assay.** Cells were grown to 95-100% confluency before a scratch wound was made. The medium was changed to RPMI 10% FBS containing 10  $\mu$ M ESI-09 or vehicle. The cells were then incubated at 37 °C in 5% CO<sub>2</sub>. The wound was imaged at 0 and 22 hrs after changing the medium. Healing rate was determined by calculating the percentage of wound closure according to the following equation: % wound closure = (initial wound width – wound width 22 hrs post-treatment with ESI-09 or vehicle)/initial wound width $\times$ 100. To make the results comparable across all assays, the widths of the initial wounds were all normalized to a 1 mm distance.

**Cell adhesion assay.** Cells were serum starved for 24 hours in RPMI, detached using 0.25% Trypsin-EDTA, and washed twice with serum-free RPMI before treatment with ESI-09 for 15 minutes. The cells were then treated with 10  $\mu$ M 007-AM and  $\sim 3\times 10^4$  in 100  $\mu$ L RPMI were immediately added to each well of a 96 well, non-tissue culture-treated EIA/RIA 96 well plate (Corning, NY). Non-treated cells and cells treated with 10  $\mu$ M 007-AM only were used for comparison. The plates were previously coated with collagen I for 2 hours (10  $\mu$ g/ml in PBS) and blocked with 5% BSA in PBS for 1 hour at 37 °C. The plates were incubated at 37 °C for 1 hour and unattached cells were removed by washing twice with PBS. The cells were fixed with 3.7% paraformaldehyde in PBS for 10 minutes, and stained with trypan blue. The attached cells were counted and the percentage of adherent cells in each treatment group was calculated using the following equation: % adhesion = number of adherent cells in treatment group/ number of adherent cells in non-treatment group  $\times$  100. The adhesion assays were repeated three times.

## RESULTS

**Identification and biochemical characterization of ESI-09.** We developed a sensitive and robust high throughput screening (HTS) assay (Tsalkova et al., 2012a) based on the principal that fluorescence intensity of a cAMP analog, 8-NBD-cAMP (8-(2-[7-Nitro-4-benzofurazanyl]

## MOL Manuscript # 80689

aminoethyl-thio) adenosine-3', 5'-cyclic monophosphate), is environmentally-sensitive. Binding of 8-NBD-cAMP to purified full-length EPAC2 led to a more than 100-fold increase in fluorescence signal. This increase was competitively reversed by the binding of cAMP (**Supplemental Figs. 1**). Through a HTS of the Maybridge HitFinder diversity library with 14,400 drug-like chemicals, we identified 3-[5-(tert-butyl)isoxazol-3-yl]-2-[2-(3-chlorophenyl)hydrazono]-3-oxopropanenitrile, referred to as ESI-09 (**Fig. 1**), which was capable of reducing both EPAC1 and EPAC2 GEF activity to basal levels at 25  $\mu$ M concentration in the presence of an equal concentration of cAMP (Tsalkova et al., 2012b). After the initial identification of the hit, individually vialled and highly purified ESI-09 was purchased from Maybridge, and its chemical structure and purity were validated by  $^1\text{H}$  and  $^{13}\text{C}$  NMR as well as mass spectrometry analysis. We then performed dose-dependent titrations to determine the relative binding affinity of ESI-09. When various amounts of cAMP or ESI-09 were added to the reaction mixture, with fixed concentrations of EPAC2 and 8-NBD-cAMP (8-[[2-[(7-nitro-4-benzofurazanyl)amino]ethyl]thio]adenosine-3',5'-cyclic monophosphate), a concentration-dependent decrease in 8-NBD-cAMP fluorescence was observed (**Fig. 2A**). While cAMP competed with 8-NBD-cAMP binding with an apparent  $\text{IC}_{50}$  of  $39 \pm 2.0$   $\mu$ M, ESI-09 showed an increased potency with an apparent  $\text{IC}_{50}$  of  $10 \pm 1.2$   $\mu$ M. To determine if this apparent higher affinity binding of EPAC2 antagonists can be translated to high potencies in suppressing the GEF activity of EPAC2, we determined the inhibition curves of EPAC-catalyzed Rap1-GDP exchange activity for ESI-09 in the presence of 25  $\mu$ M cAMP. As shown in **Figure 2B**, ESI-09 inhibited cAMP-mediated EPAC2 GEF activity with an apparent  $\text{IC}_{50}$  of  $1.4 \pm 0.1$   $\mu$ M. Similarly, ESI-09 was also effective in suppressing cAMP-mediated EPAC1 GEF activity with an apparent  $\text{IC}_{50}$  of  $3.2 \pm 0.4$   $\mu$ M. To determine the specificity of ESI-09, we performed counter-screening assays that measured type I and II PKA holoenzyme activities. An ESI-09 concentration of 25  $\mu$ M did not

## MOL Manuscript # 80689

significantly alter cAMP-induced type I $\alpha$  and II $\beta$  PKA holoenzymes activation, while H89, a selective PKA inhibitor, blocked type I and II PKA activities completely (**Supplemental Fig. 2**). When tested at 100  $\mu$ M concentration, ESI-09 inhibited type II PKA activity up to 20% in the presence of 100  $\mu$ M of cAMP. These data suggest that ESI-09 is at least 100 times more selective for EPAC proteins than PKA.

### **Molecular docking of ESI-09 into the cAMP binding domains (CBD) of EPAC1 and EPAC2.**

To gain insights into the structural basis for the binding of ESI-09 to the EPAC proteins, the intermolecular interactions of ESI-09 with EPAC1 and EPAC2 were analyzed and compared by molecular docking (Trott and Olson, 2010) using the cAMP-bound EPAC2 crystal structure (3CF6) (Rehmann et al., 2008) and a 3D homology model of EPAC1 (Arnold et al., 2006). AutoDock Vina docking results revealed that ESI-09 could be docked to the cAMP binding domain (CBD) of EPAC1 and EPAC2, with better energetic scores than those for cAMP. ESI-09 could fit well into the functional cAMP binding pocket of EPAC1; establishing favorable hydrophobic and hydrogen bonding interactions with the protein's active site residues. The *tert*-butyl-isoxazolyl moiety of ESI-09 could extend to a deep hydrophobic pocket around Phe268 and Met312, with its 3-chloro-phenyl fragment close to the hydrophobic pocket around Ala281 and Leu314. Moreover, ESI-09 could form five putative hydrogen bonds with the residues Gly269, Gln270, Leu271, Ala280 and Ala281 (**Fig. 3, A & B**). With EPAC2, the *tert*-butyl-isoxazolyl moiety could interact with a hydrophobic pocket formed by Phe367, Leu406, Ala407 and Ala415, while residues Val386, Val394, Leu397, Ala416 and Leu448 could form hydrophobic interactions with the 3-chloro-phenyl fragment of ESI-09. Meanwhile, the *tert*-butyl-isoxazolyl moiety could form a hydrogen bond with the residue Gly404 of EPAC2 (**Fig. 3, C & D**). These docking observations are in full agreement with the experimental EPAC inhibition results as shown in **Figure 2**, and provide useful information

## MOL Manuscript # 80689

to assist our ongoing efforts to design and synthesize more potent and specific inhibitors based on ESI-09 as a chemical lead.

**Cellular characterization of ESI-09.** To test if our newly identified EPAC antagonist ESI-09 is capable of modulating EPAC activation in living cells, we monitored its ability to suppress Akt phosphorylation as EPAC proteins are also known to activate Akt signaling while PKA inhibits it (Mei et al., 2002). To determine if ESI-09 is capable of blocking EPAC-mediated Akt activation, the phosphorylation status of T308 and S473 of Akt in the pancreatic cancer cell line AsPC-1, which over-express EPAC1, was followed using anti phospho-Akt antibodies. As shown in **Figure 4**, ESI-09 inhibited 007-AM stimulated Akt phosphorylation at T308 and S473 in a dose-dependent manner. Similar results were observed in the rat pancreatic  $\beta$  cell line, INS-1 (**Supplemental Fig. 3**). On the other hand, ESI-09 failed to suppress epidermal growth factor (EGF) induced phosphorylation of Akt in AsPC1 cells (**Fig. 4**).

After demonstrating that ESI-09 was capable of inhibiting EPAC-mediated Akt phosphorylation in cell culture, we proceeded to determine if ESI-09 was capable of suppressing EPAC-mediated physiological functions. EPAC proteins, particularly EPAC2, have been implicated in regulating cAMP-mediated exocytosis in a variety of secretory cells (Seino and Shibasaki, 2005). In pancreatic endocrine  $\beta$  cells, EPAC mediates the glucose-induced insulin secretion potentiated by glucagon-like peptide 1 (GLP-1) and gastric inhibitory polypeptide (GIP), which bind to receptors on the surface of  $\beta$  cells and promote intracellular cAMP production (Shibasaki et al., 2007; Leech et al., 2000; Kang et al., 2003; Kashima et al., 2001). When INS-1 was stimulated by 007-AM, a steady increase in insulin secretion was observed. This EPAC-mediated increase in the level of insulin secretion was inhibited in a dose-dependent manner by pre-incubation of the cells with ESI-09 (**Fig. 5**). The impact of ESI-09 on EPAC-mediated Akt phosphorylation and insulin

## MOL Manuscript # 80689

secretion is consistent with the biochemical Rap1 nucleotide exchange data shown in **Figure 2B** and demonstrates that this small molecule is a pan-EPAC inhibitor that is capable of suppressing EPAC cellular functions.

**ESI-09 Inhibits pancreatic cancer migration.** Increased levels of EPAC1 expression have been observed in human pancreatic ductal adenocarcinoma as compared with normal pancreas or surrounding tissue (Lorenz et al., 2008). However, the functional role of EPAC1 elevation in this neoplasm remains unclear. We sought to employ ESI-09 to determine the role of EPAC1 signaling in pancreatic cancer. Treatment of pancreatic cancer cells with ESI-09 did not significantly affect cell proliferation and viability (**Supplemental Fig. 4**). On the other hand, when pretreated with ESI-09, a significant decrease in cell migration was observed for the pancreatic cancer cell lines AsPC-1 and PANC-1, using both trans-well migration/invasion and wound healing assays (**Fig. 6, A & B**). To determine whether the observed impact on cell migration is EPAC1 specific, we examined the effect of suppressing EPAC1 expression on AsPC-1 and PANC-1 migration using RNA interference. Short hairpin construct shEPAC1 clone C28 led to a near complete knockdown of EPAC1 expression and significantly inhibited migration of both cell lines, while a partial reduction of EPAC1 expression by shEPAC1 clone C32 decreased AsPC-1 and PANC-1 cell migration slightly, but the changes were not significant (**Fig. 6C**). These results, combined with the fact that pancreatic cancer cells do not express detectable levels of EPAC2 (**Supplemental Fig. 5**), suggest that EPAC1 might play an important role in pancreatic cancer cell migration and invasion. To further determine how ESI-09 inhibits PDA cell migration and invasion, we performed a cell adhesion assay on a collagen I matrix. As shown in **Figure 7**, 007-AM led to an increase in cell adhesion for both AcPC-1 and PANC-1 cells, while pre-treatment with ESI-09 decreased 007-AM induced cell adhesion dose-dependently.

## **DISCUSSION**

Taken together, our results show that we have successfully identified ESI-09, a non-cyclic nucleotide small molecule that specifically inhibits EPAC1 and EPAC2. Consistent with its EPAC inhibition activity observed in a biochemical assay, ESI-09 also inhibited several well-known EPAC-mediated cellular functions including, Akt phosphorylation and insulin secretion. To the best of our knowledge, this is the first report of discovery and characterization of a small molecule inhibitor for the EPAC family of proteins since their initial discovery in 1998. McPhee et al. reported the discovery of compounds capable of competing with  $^3\text{H}$ -cAMP for binding with the isolated cAMP-binding domains of EPAC1 and EPAC2. However, it has not been shown that these compounds are capable of inhibiting EPAC functions biochemically or in cells (McPhee et al., 2005). The discovery of an EPAC specific antagonist may have immense implications for cAMP biology, as it will provide a means to pharmacologically dissect the roles of EPAC proteins in various cellular and diseases processes.

Pancreatic ductal adenocarcinoma is one of the most lethal human diseases, largely due to the fact that pancreatic cancer is resistant to treatments that are usually effective for other types of cancer. A better understanding of the molecular mechanism of PDA development and metastasis and novel, effective therapeutics are desperately needed. Recently, it has been shown that EPAC1 is markedly elevated in human PDA cells as compared with normal pancreas or surrounding tissue (Lorenz et al., 2008). While EPAC1 has been implicated in promoting cellular proliferation in prostate cancer (Misra and Pizzo, 2009; Misra and Pizzo, 2011) and migration and metastasis in melanoma (Baljinnyam et al., 2011; Baljinnyam et al., 2009; Baljinnyam et al., 2010), the role of EPAC1 in pancreatic cancer has never been investigated. Using our novel EPAC inhibitor ESI-09, we have demonstrated a functional role of EPAC1 overexpression in pancreatic cancer cell migration and

**MOL Manuscript # 80689**

invasion. These findings are consistent with similar results based on RNAi silencing techniques, suggesting that EPAC1 may represent a potential target for developing novel therapeutic strategies for PDA.

## MOL Manuscript # 80689

**Acknowledgement:** We are grateful to Dr. George Holz for providing the INS-1 cells and advice on insulin secretion analysis. We would also like to thank Professor Alfred Wittinghofer (Abteilung Strukturelle Biologie Max-Planck-Institut für molekulare Physiologie), Dr. Holger Rehman and Professor Johannes Bos (UMC Utrecht Universiteitsweg) for providing *E. coli* CK600K cells and pGEX-4T-EPAC2 constructs.

### **Author contributions:**

*Participated in research design:* Almahariq, Tsalkova, Mei, Chen, Zhou, Sastry, and Cheng.

*Conducted experiments:* Tsalkova, Mei, Almahariq, Chen, and Cheng.

*Contributed new reagents or analytic tools:* Schwede.

*Performed data analysis:* Tsalkova, Mei, Almahariq, Chen, Zhou, and Cheng.

*Wrote or contributed to the writing of the manuscript:* Tsalkova, Mei, Almahariq, Chen, Zhou, and Cheng.



## Reference List

- Arnold K, Bordoli L, Kopp J and Schwede T (2006) The SWISS-MODEL Workspace: a Web-Based Environment for Protein Structure Homology Modelling. *Bioinformatics* **22**:195-201.
- Baljinnyam E, De Lorenzo M S, Xie L H, Iwatsubo M, Chen S, Goydos J S, Nowycky M C and Iwatsubo K (2010) Exchange Protein Directly Activated by Cyclic AMP Increases Melanoma Cell Migration by a Ca<sup>2+</sup>-Dependent Mechanism. *Cancer Res* **70**:5607-5617.
- Baljinnyam E, Iwatsubo K, Kurotani R, Wang X, Ulucan C, Iwatsubo M, Lagunoff D and Ishikawa Y (2009) Epac Increases Melanoma Cell Migration by a Heparan Sulfate-Related Mechanism. *Am J Physiol Cell Physiol* **297**:C802-C813.
- Baljinnyam E, Umemura M, De Lorenzo M S, Iwatsubo M, Chen S, Goydos J S and Iwatsubo K (2011) Epac1 Promotes Melanoma Metastasis Via Modification of Heparan Sulfate. *Pigment Cell Melanoma Res* **24**:680-687.
- Brennesvik EO, Ktori C, Ruzzin J, Jebens E, Shepherd P R and Jensen J (2005) Adrenaline Potentiates Insulin-Stimulated PKB Activation Via CAMP and Epac: Implications for Cross Talk Between Insulin and Adrenaline. *Cell Signal* **17**:1551-1559.
- Cheng X, Ji Z, Tsalkova T and Mei F (2008) Epac and PKA: a Tale of Two Intracellular CAMP Receptors. *Acta Biochim Biophys Sin (Shanghai)* **40**:651-662.
- Cheng X, Phelps C and Taylor S S (2001) Differential Binding of CAMP-Dependent Protein Kinase Regulatory Subunit Isoforms Ialpha and Ibeta to the Catalytic Subunit. *J Biol Chem* **276**:4102-4108.
- Chepurny OG, Leech C A, Kelley G G, Dzhura I, Dzhura E, Li X, Rindler M J, Schwede F, Genieser H G and Holz G G (2009) Enhanced Rap1 Activation and Insulin Secretagogue Properties of an Acetoxymethyl Ester of an Epac-Selective Cyclic AMP Analog in Rat INS-1 Cells: Studies With 8-PCPT-2'-O-Me-CAMP-AM. *J Biol Chem* **284**:10728-10736.
- Christensen AE, Selheim F, de R J, Dremier S, Schwede F, Dao K K, Martinez A, Maenhaut C, Bos J L, Genieser H G and Doskeland S O (2003) CAMP Analog Mapping of Epac1 and CAMP Kinase. Discriminating Analogs Demonstrate That Epac and CAMP Kinase Act Synergistically to Promote PC-12 Cell Neurite Extension. *J Biol Chem* **278**:35394-35402.
- Cook PF, Neville M E, Jr., Vrana K E, Hartl F T and Roskoski R, Jr. (1982) Adenosine Cyclic 3',5'-Monophosphate Dependent Protein Kinase: Kinetic Mechanism for the Bovine Skeletal Muscle Catalytic Subunit. *Biochemistry* **21**:5794-5799.
- Cullere X, Shaw S K, Andersson L, Hirahashi J, Lusinskas F W and Mayadas T N (2005) Regulation of Vascular Endothelial Barrier Function by Epac, a CAMP-Activated Exchange Factor for Rap GTPase. *Blood* **105**:1950-1955.
- de Rooij J, Zwartkruis F J, Verheijen M H, Cool R H, Nijman S M, Wittinghofer A and Bos J L

**MOL Manuscript # 80689**

(1998) Epac Is a Rap1 Guanine-Nucleotide-Exchange Factor Directly Activated by Cyclic AMP. *Nature* **396**:474-477.

Dodge-Kafka KL, Soughayer J, Pare G C, Carlisle Michel J J, Langeberg L K, Kapiloff M S and Scott J D (2005) The Protein Kinase A Anchoring Protein MAKAP Coordinates Two Integrated CAMP Effector Pathways. *Nature* **437**:574-578.

Enserink JM, Christensen A E, de R J, van T M, Schwede F, Genieser H G, Doskeland S O, Blank J L and Bos J L (2002) A Novel Epac-Specific CAMP Analogue Demonstrates Independent Regulation of Rap1 and ERK. *Nat Cell Biol* **4**:901-906.

Enserink JM, Price L S, Methi T, Mahic M, Sonnenberg A, Bos J L and Tasken K (2004) The CAMP-Epac-Rap1 Pathway Regulates Cell Spreading and Cell Adhesion to Laminin-5 Through the Alpha3beta1 Integrin but Not the Alpha6beta4 Integrin. *J Biol Chem* **279**:44889-44896.

Fujimoto K, Shibasaki T, Yokoi N, Kashima Y, Matsumoto M, Sasaki T, Tajima N, Iwanaga T and Seino S (2002) Piccolo, a Ca<sup>2+</sup> Sensor in Pancreatic Beta-Cells. Involvement of CAMP-GEFII.Rim2.Piccolo Complex in CAMP-Dependent Exocytosis. *J Biol Chem* **277**:50497-50502.

Fukuda M, Williams K W, Gautron L and Elmquist J K (2011) Induction of Leptin Resistance by Activation of CAMP-Epac Signaling. *Cell Metab* **13**:331-339.

Gloerich M and Bos J L (2010) Epac: Defining a New Mechanism for CAMP Action. *Annu Rev Pharmacol Toxicol* **50**:355-375.

Gloerich M, Ponsioen B, Vliem M J, Zhang Z, Zhao J, Kooistra M R, Price L S, Ritsma L, Zwartkruis F J, Rehmann H, Jalink K and Bos J L (2010) Spatial Regulation of Cyclic AMP-Epac1 Signaling in Cell Adhesion by ERM Proteins. *Mol Cell Biol* **30**:5421-5431.

Gloerich M, Vliem M J, Prummel E, Meijer L A, Rensen M G, Rehmann H and Bos J L (2011) The Nucleoporin RanBP2 Tethers the CAMP Effector Epac1 and Inhibits Its Catalytic Activity. *J Cell Biol* **193**:1009-1020.

Ji Z, Mei F C, Johnson B H, Thompson E B and Cheng X (2007) Protein Kinase A, Not Epac, Suppresses Hedgehog Activity and Regulates Glucocorticoid Sensitivity in Acute Lymphoblastic Leukemia Cells. *J Biol Chem* **282**:37370-37377.

Kang G, Joseph J W, Chepurny O G, Monaco M, Wheeler M B, Bos J L, Schwede F, Genieser H G and Holz G G (2003) Epac-Selective CAMP Analog 8-PCPT-2'-O-Me-CAMP As a Stimulus for Ca<sup>2+</sup>-Induced Ca<sup>2+</sup> Release and Exocytosis in Pancreatic Beta-Cells. *J Biol Chem* **278**:8279-8285.

Kashima Y, Miki T, Shibasaki T, Ozaki N, Miyazaki M, Yano H and Seino S (2001) Critical Role of CAMP-GEFII--Rim2 Complex in Incretin-Potentiated Insulin Secretion. *J Biol Chem* **276**:46046-46053.

Kawasaki H, Springett G M, Mochizuki N, Toki S, Nakaya M, Matsuda M, Housman D E and

**MOL Manuscript # 80689**

- Graybiel A M (1998) A Family of CAMP-Binding Proteins That Directly Activate Rap1. *Science* **282**:2275-2279.
- Kooistra MR, Corada M, Dejana E and Bos J L (2005) Epac1 Regulates Integrity of Endothelial Cell Junctions Through VE-Cadherin. *FEBS Lett* **579**:4966-4972.
- Leech CA, Holz G G, Chepurny O and Habener J F (2000) Expression of CAMP-Regulated Guanine Nucleotide Exchange Factors in Pancreatic Beta-Cells. *Biochem Biophys Res Commun* **278**:44-47.
- Li J, O'Connor K L, Cheng X, Mei F C, Uchida T, Townsend C M, Jr. and Evers B M (2007) Cyclic Adenosine 5'-Monophosphate-Stimulated Neurotensin Secretion Is Mediated Through Rap1 Downstream of Both Epac and Protein Kinase A Signaling Pathways. *Mol Endocrinol* **21**:159-171.
- Li S, Tsalkova T, White M A, Mei F C, Liu T, Wang D, Woods V L, Jr. and Cheng X (2011) Mechanism of Intracellular CAMP Sensor Epac2 Activation: CAMP-INDUCED CONFORMATIONAL CHANGES IDENTIFIED BY AMIDE HYDROGEN/DEUTERIUM EXCHANGE MASS SPECTROMETRY (DXMS). *J Biol Chem* **286**:17889-17897.
- Liu C, Takahashi M, Li Y, Dillon T J, Kaech S and Stork P J (2010) The Interaction of Epac1 and Ran Promotes Rap1 Activation at the Nuclear Envelope. *Mol Cell Biol* **30**:3956-3969.
- Lorenz R, Aleksic T, Wagner M, Adler G and Weber C K (2008) The CAMP/Epac1/Rap1 Pathway in Pancreatic Carcinoma. *Pancreas* **37**:102-103.
- Maillet M, Robert S J, Cacquevel M, Gastineau M, Vivien D, Bertoglio J, Zugaza J L, Fischmeister R and Lezoualc'h F (2003) Crosstalk Between Rap1 and Rac Regulates Secretion of SAPPalpha. *Nat Cell Biol* **5**:633-639.
- McPhee I, Gibson L C, Kewney J, Darroch C, Stevens P A, Spinks D, Cooreman A and MacKenzie S J (2005) Cyclic Nucleotide Signalling: a Molecular Approach to Drug Discovery for Alzheimer's Disease. *Biochem Soc Trans* **33**:1330-1332.
- Mei FC and Cheng X D (2005) Interplay Between Exchange Protein Directly Activated by CAMP (Epac) and Microtubule Cytoskeleton. *Molecular Biosystems* **1**:325-331.
- Mei FC, Qiao J, Tsygankova O M, Meinkoth J L, Quilliam L A and Cheng X (2002) Differential Signaling of Cyclic AMP: Opposing Effects of Exchange Protein Directly Activated by Cyclic AMP and CAMP-Dependent Protein Kinase on Protein Kinase B Activation. *J Biol Chem* **277**:11497-11504.
- Metrich M, Berthouze M, Morel E, Crozatier B, Gomez A M and Lezoualc'h F (2010) Role of the CAMP-Binding Protein Epac in Cardiovascular Physiology and Pathophysiology. *Pflugers Arch* **459**:535-546.
- Misra UK and Pizzo S V (2009) Epac1-Induced Cellular Proliferation in Prostate Cancer Cells Is Mediated by B-Raf/ERK and MTOR Signaling Cascades. *J Cell Biochem* **108**:998-1011.

**MOL Manuscript # 80689**

Misra UK and Pizzo S V (2011) Upregulation of MTORC2 Activation by the Selective Agonist of EPAC, 8-CPT-2Me-CAMP, in Prostate Cancer Cells: Assembly of a Multi-Protein Signaling Complex. *J Cell Biochem*.

Ozaki N, Shibasaki T, Kashima Y, Miki T, Takahashi K, Ueno H, Sunaga Y, Yano H, Matsuura Y, Iwanaga T, Takai Y and Seino S (2000) CAMP-GEFII Is a Direct Target of CAMP in Regulated Exocytosis. *Nat Cell Biol* **2**:805-811.

Qiao J, Mei F C, Popov V L, Vergara L A and Cheng X (2002) Cell Cycle-Dependent Subcellular Localization of Exchange Factor Directly Activated by CAMP. *J Biol Chem* **277**:26581-26586.

Rangarajan S, Enserink J M, Kuiperij H B, de Rooij J, Price L S, Schwede F and Bos J L (2003) Cyclic AMP Induces Integrin-Mediated Cell Adhesion Through Epac and Rap1 Upon Stimulation of the Beta 2-Adrenergic Receptor. *J Cell Biol* **160**:487-493.

Rehmann H, Arias-Palomo E, Hadders M A, Schwede F, Llorca O and Bos J L (2008) Structure of Epac2 in Complex With a Cyclic AMP Analogue and RAP1B. *Nature* **455**:124-127.

Ross SH, Post A, Raaijmakers J H, Verlaan I, Gloerich M and Bos J L (2011) Ezrin Is Required for Efficient Rap1-Induced Cell Spreading. *J Cell Sci* **124**:1808-1818.

Seino S and Shibasaki T (2005) PKA-Dependent and PKA-Independent Pathways for CAMP-Regulated Exocytosis. *Physiol Rev* **85**:1303-1342.

Shibasaki T, Takahashi H, Miki T, Sunaga Y, Matsumura K, Yamanaka M, Zhang C, Tamamoto A, Satoh T, Miyazaki J and Seino S (2007) Essential Role of Epac2/Rap1 Signaling in Regulation of Insulin Granule Dynamics by CAMP. *Proc Natl Acad Sci U S A* **104**:19333-19338.

Trott O and Olson A J (2010) AutoDock Vina: Improving the Speed and Accuracy of Docking With a New Scoring Function, Efficient Optimization, and Multithreading. *J Comput Chem* **31**:455-461.

Tsalkova T, Blumenthal D K, Mei F C, White M A and Cheng X (2009) Mechanism of Epac Activation: Structural and Functional Analyses of Epac2 Hinge Mutants With Constitutive and Reduced Activities. *J Biol Chem* **284**:23644-23651.

Tsalkova T, Mei F C and Cheng X (2012a) A Fluorescence-Based High-Throughput Assay for the Discovery of Exchange Protein Directly Activated by Cyclic AMP (EPAC) Antagonists. *PLoS ONE* **7**:e30441.

Tsalkova T, Mei F C, Li S, Chepurny O G, Leech C A, Liu T, Woods V L, Jr. Holz G G and Cheng X (2012b) Isoform-specific Antagonists of Exchange Protein Directly Activated by cAMP. *Proc Natl Acad Sci U S A* **109**:in press.

Vliem MJ, Ponsioen B, Schwede F, Pannekoek W J, Riedl J, Kooistra M R, Jalink K, Genieser H G, Bos J L and Rehmann H (2008) 8-PCPT-2'-O-Me-CAMP-AM: an Improved Epac-Selective CAMP Analogue. *Chembiochem* **9**:2052-2054.

**MOL Manuscript # 80689**

Yu S, Mei F C, Lee J C and Cheng X (2004) Probing CAMP-Dependent Protein Kinase Holoenzyme Complexes I Alpha and II Beta by FT-IR and Chemical Protein Footprinting. *Biochemistry* **43**:1908-1920.

## **MOL Manuscript # 80689**

### **FOOTNOTE**

**Financial Support:** This work was supported by the National Institutes of Health National Institute of General Medicine (NIGMS) [Grant R01GM066170] and a training fellowship from the Keck Center for Interdisciplinary Bioscience Training of the Gulf Coast Consortia funded by NIGMS [Grant T32GM089657].

M. Almahariq, T. Tsalkova., and F. C. Mei contributed equally to the study.

## FIGURE LEGENDS

**Fig. 1. Chemical structure of 3-(5-*tert*-Butyl-isoxazol-3-yl)-2-[(3-chloro-phenyl)-hydrazono]-3-oxo-propionitrile (ESI-09).**

**Fig. 2. Relative potency of ESI-09.** (A) Dose-dependent competition of ESI-09 (open circles) and cAMP (closed squares) with 8-NBD-cAMP in binding to EPAC2. (B) Dose-dependent inhibition of EPAC1 (closed circles) or EPAC2 (open circles) GEF activity by ESI-09 in the presence of 25  $\mu$ M cAMP.


**Fig. 3. Predicted binding mode and molecular docking of ESI-09 into the cAMP binding domain (CBD) of EPAC1 protein (homology modeling) and EPAC2 protein (pdb code 3CF6).** (A) Binding mode of ESI-09 with EPAC1 protein (homology modeling); (B) Surface of the cAMP binding domain (CBD) of EPAC1 protein; (C) Binding mode of ESI-09 with EPAC2 protein (pdb code 3CF6); (D) Surface of the cAMP binding domain (CBD) of EPAC2 protein. Important residues are drawn in sticks. Hydrogen bonds are shown as dashed green lines. ESI-09 is shown in pink color and cAMP is shown in pale yellow color.

**Fig. 4. ESI-09 inhibits EPAC- but not EGF-mediated Akt phosphorylation in the pancreatic cancer cells AsPC-1.** Serum-starved AsPC-1 cells were stimulated with vehicle, 10  $\mu$ M 007-AM, or 0.1  $\mu$ M EGF after pretreatment with the indicated concentrations of ESI-09. Cell lysates were subjected to Western blot analyses using anti-phospho-Akt T308 and S473 antibodies (representative blot shown). Data points represent mean  $\pm$  s.d. (n=3). <sup>#</sup>Significantly higher than vehicle group (P < 0.02). \*Significantly lower than 007-AM stimulated group (P < 0.02).

**Fig. 5. ESI-09 inhibits EPAC2-mediated insulin secretion in INS-1 cells.** INS-1 cells were stimulated with vehicle or 10  $\mu$ M 007-AM after pretreatment with the indicated concentrations of ESI-09. Bars represent mean  $\pm$  s.d. (n=3). <sup>#</sup>Significantly higher than vehicle group (P < 0.05). \*Significantly lower than 007-AM stimulated group (P < 0.05).



**Fig. 6. EPAC1 inhibition decreases pancreatic cancer cells invasion and migration.** AsPC-1, and PANC-1 cells were pretreated with the indicated concentrations of ESI-09 for 24 hours before invasion/migration were measured by trans-well (A) and wound-healing assays (B). EPAC1 expression was suppressed by shEPAC1-C28 and shEPAC1-C32 and invasion/migration were measured by trans-well assays (C). Bars represent mean  $\pm$  s.d. (n=3). \*Significantly lower than vehicle group (P < 0.05). <sup>#</sup>Significantly lower than parental-cells group (P < 0.02).

**Fig. 7. ESI-09 inhibits EPAC1-mediated adhesion of PDA cells on collagen I.** AsPC-1 and PANC-1 cells were stimulated with vehicle or 10  $\mu$ M 007-AM after treatment with the indicated concentrations of ESI-09 for 5 minutes. Bars represent mean  $\pm$  s.d. (n=3). <sup>#</sup>Significantly higher than vehicle group (P < 0.03). \*Significantly lower than 007-AM stimulated group (P < 0.02).




**Figure 1**








**Figure 2**



**Figure 3**



**Figure 4**



**Figure 5**





Figure 6

- Vehicle
- 10  $\mu$ M 007-AM
- 10  $\mu$ M 007-AM + 1  $\mu$ M ESI-09
- 10  $\mu$ M 007-AM + 5  $\mu$ M ESI-09
- 10  $\mu$ M 007-AM + 10  $\mu$ M ESI-09



**Figure 7**