

Model Building Report

This document lists the results for the homology modelling project "EAAT1" submitted to SWISS-MODEL workspace on March 6, 2018, 7:19 a.m.. The submitted primary amino acid sequence is given in Table T1.

If you use any results in your research, please cite the relevant publications:


- Biasini, M., Bienert, S., Waterhouse, A., Arnold, K., Studer, G., Schmidt, T., Kiefer, F., Cassarino, T.G., Bertoni, M., Bordoli, L., Schwede, T. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res.* 42, W252-W258 (2014). [M doi>](#)
- Guex, N., Peitsch, M.C., Schwede, T. Automated comparative protein structure modeling with SWISS-MODEL and Swiss-PdbViewer: A historical perspective. *Electrophoresis* 30, S162-S173 (2009). [M doi>](#)
- Bienert, S., Waterhouse, A., de Beer, T.A., Tauriello, G., Studer, G., Bordoli, L., Schwede, T. The SWISS-MODEL Repository - new features and functionality. *Nucleic Acids Res.* 45, D313-D319 (2017). [M doi>](#)
- Benkert, P., Biasini, M., Schwede, T. Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics* 27, 343-350 (2011). [M doi>](#)
- Bertoni, M., Kiefer, F., Biasini, M., Bordoli, L., Schwede, T. Modeling protein quaternary structure of homo- and hetero-oligomers beyond binary interactions by homology. *Scientific Reports* 7 (2017). [M doi>](#)






Results



The SWISS-MODEL template library (SMTL version 2018-02-28, PDB release 2018-02-23) was searched with BLAST ([Camacho et al.](#)) and HHblits ([Remmert et al.](#)) for evolutionary related structures matching the target sequence in Table T1. For details on the template search, see Materials and Methods. Overall 85 templates were found (Table T2).

Models

The following model was built (see Materials and Methods "Model Building"):

Model #02	File	Built with	Oligo-State	Ligands	GMQE	QMEAN
	PDB	ProMod3 Version 1.1.0.	monomer	None	0.66	-5.58

QMEAN	-5.58	
C β	-1.55	
All Atom	-1.51	
Solvation	1.33	
Torsion	-5.67	



Template	Seq Identity	Oligo-state	Found by	Method	Resolution	Seq Similarity	Range	Coverage	Description
5llm.1.A	75.58	monomer	BLAST	X-ray	3.25Å	0.52	37 - 462	0.95	Excitatory amino acid transporter 1, Neutral amino acid transporter B(0), Excitatory amino acid transporter 1

Ligand	Added to Model	Description
6Z6	× - Binding site not conserved.	2-Amino-5,6,7,8-tetrahydro-4-(4-methoxyphenyl)-7-(naphthalen-1-yl)-5-oxo-4H-chromene-3-carbonitrile
ASP	× - Binding site not conserved.	ASPARTIC ACID
NA	× - Not biologically relevant.	SODIUM ION

Target MTKSNGEPRMGRMRERFQQGVRKRTLLAKKKVQNI TKEDVKS YLFRNAFVLLTVSAVIVGTILGFALRPYKMSYREVKY
5llm.1.A MTKSNGEPEPKMGRMRERFQQGVSRTLLAKKKVQNI TKEDVKS FLRRNALLLLTVLAVILGVVLGFLLRPYPLSPREVKY

Target FSPFGELLMRMLQMLVPLIISLVTGMAALDSKASGKMGMRVAVVYMTTIIAVVIGIIIVIIHPGKGTENMYREGK
5llm.1.A FAPFGELLMRMLKMLILPLIVSSLITGLASLDAKASGRLGMRVAVVYMTTIIAVVIGIILVLIHPGAASAAITASVGA

Target IVQVTAA-----DAFLDLIRNFPPNLVEACFKQFKTSYKRSFKVPIQANETLLGAVINNVSEAMETLTRIEMVVP
5llm.1.A AGSAENAPSKEVLDLFLARNIFPSNLVSAAFRSYSTTYEERTI-----TGTRVK---VPV

Target PGSVNGVNALGLVVFMCFGFVIGNMKEQQGALREFFDLSLNEAIMRLVAVIMWYAPLGILFLIAGKIVEMEDMGVIGGQL

```

511m. 1. A  GQEVEGMNIGLVVFSMVFGFALGKMGEGQQLLVDFNSLNEATMKLVAIIMWYAPLGILFLIAGKIVEMEDLEVLGGQL

Target      AMYTVTVIVGLLIHAVIVLPLLYFLVTRKNPWFVIGLLQALITALGTSSSSATLPITFKCLEENNGVDKRI TRFVLPVG
511m. 1. A  GMYMVTIVGLVIHGLIVLPLIYFLI TRKNPFVFIAGILQALITALGTSSSSATLPITFKCLEENNGVDKRI TRFVLPVG

Target      ATINMDGTALYEALAAIFIAQVNNFDLNFQGIITIRDLRITTT-----
511m. 1. A  ATINMDGTALYEAVAAIFIAQVNNYELDFGQIITISITATAASIGAAGIPQAGLVTMVIIVLTAVGLPTDDITLIIAVDWL

Target      -----NVLGDSL GAGIVEHL SRHELKNRD VEMGNSVIEENEMKKPYQLIAQDNEPEKPVADSETKM
511m. 1. A  LDRFRMTMNVLGDALGAGIVEHL SRKELEKQDAELGNSVIEENEMKKPYQLIAQDNETEKPI-DSETKM

```

Materials and Methods

Template Search

Template search with BLAST and HHblits has been performed against the SWISS-MODEL template library (SMTL, last update: 2018-02-28, last included PDB release: 2018-02-23).

The target sequence was searched with BLAST against the primary amino acid sequence contained in the SMTL. A total of 27 templates were found.

An initial HHblits profile has been built using the procedure outlined in (Remmert et al.), followed by 1 iteration of HHblits against NR20. The obtained profile has then be searched against all profiles of the SMTL. A total of 58 templates were found.

Template Selection

For each identified template, the template's quality has been predicted from features of the target-template alignment. The templates with the highest quality have then been selected for model building.

Model Building

Models are built based on the target-template alignment using ProMod3. Coordinates which are conserved between the target and the template are copied from the template to the model. Insertions and deletions are remodelled using a fragment library. Side chains are then rebuilt. Finally, the geometry of the resulting model is regularized by using a force field. In case loop modelling with ProMod3 fails, an alternative model is built with PROMOD-II (Guex et al.).

Model Quality Estimation

The global and per-residue model quality has been assessed using the QMEAN scoring function (Benkert et al.). For improved performance, weights of the individual QMEAN terms have been trained specifically for SWISS-MODEL.

Ligand Modelling

Ligands present in the template structure are transferred by homology to the model when the following criteria are met: (a) The ligands are annotated as biologically relevant in the template library, (b) the ligand is in contact with the model, (c) the ligand is not clashing with the protein, (d) the residues in contact with the ligand are conserved between the target and the template. If any of these four criteria is not satisfied, a certain ligand will not be included in the model. The model summary includes information on why and which ligand has not been included.

Oligomeric State Conservation

The quaternary structure annotation of the template is used to model the target sequence in its oligomeric form. The method (Bertoni et al.) is based on a supervised machine learning algorithm, Support Vector Machines (SVM), which combines interface conservation, structural clustering, and other template features to provide a quaternary structure quality estimate (QSQE). The QSQE score is a number between 0 and 1, reflecting the expected accuracy of the interchain contacts for a model built based a given alignment and template. Higher numbers indicate higher reliability. This complements the GMQE score which estimates the accuracy of the tertiary structure of the resulting model.

References

- **BLAST**
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L. BLAST+: architecture and applications. *BMC Bioinformatics* 10, 421-430 (2009). [doi>](#)
- **HHblits**
Remmert, M., Biegert, A., Hauser, A., Söding, J. HHblits: lightning-fast iterative protein sequence searching by HMM-HMM alignment. *Nat Methods* 9, 173-175 (2012). [doi>](#)

Table T1:

Primary amino acid sequence for which templates were searched and models were built.

```

MTKSNGEPRMGSRMERFQQGVRKRTLLAKKKVQNI TKEDVKS YLFRNAFVLLTVSAVIVGTILGFALRPYKMSYREVKYFSFPGELLMRMLQMLVLP LI
ISSLVGTMAALDSKASGKMGMRVYVYMTTII IAVVIGII IIVII IHPGKGTKENMYREGKIVQVTAADAFDLIRNMFPPNLVEACFKQFKTSY EKRSFK
VPIQANETLLGAVINNVSEAMETL TRIREEMVVPVGSVNGVNALGLVVFSMCFGVIGNMKEQGQALREFFDSLNEAIMRLVAVIMWYAPLGILFLIAGK

```

IVEMEDMGVIGGQLAMYTIVTVIVGLLIHAVIVLPLLYFLVTRKNPWFVFIGLLQALITALGTSSSSATLPITFKLEENNGVDKRI TRFVLPVIGATINMD
GTALYEALAAIFIAQVNNFDLNFQGIITIRDLRRTTTNVLGDSL GAGIVEHLRSRHELKNRDVMGNSVIEENEMKKPYQLIAQDNEPEKPVADSETKM

Table T2:

Template	Seq Identity	Oligo-state	QSQE	Found by	Method	Resolution	Seq Similarity	Coverage	Description
5mju.1.A	75.58	homo-trimer	0.64	BLAST	X-ray	3.71Å	0.52	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
5llm.1.A	75.58	monomer		BLAST	X-ray	3.25Å	0.52	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
5llu.1.A	75.16	homo-trimer	0.63	BLAST	X-ray	3.32Å	0.52	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
5mju.1.A	74.95	homo-trimer	0.65	HHblits	X-ray	3.71Å	0.52	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
5llm.1.A	74.95	monomer		HHblits	X-ray	3.25Å	0.52	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
5llu.1.A	74.31	homo-trimer	0.64	HHblits	X-ray	3.32Å	0.51	0.95	Excitatory amino acid transporter 1,Neutral amino acid transporter B(0),Excitatory amino acid transporter 1
3kbc.1.A	33.52	homo-trimer	0.47	HHblits	X-ray	3.51Å	0.36	0.72	425aa long hypothetical proton glutamate symport protein
4x2s.1.B	34.08	homo-trimer	0.48	HHblits	X-ray	4.21Å	0.36	0.71	425aa long hypothetical proton glutamate symport protein
4x2s.1.A	34.08	homo-trimer	0.48	HHblits	X-ray	4.21Å	0.36	0.71	425aa long hypothetical proton glutamate symport protein
2nwl.1.A	32.96	homo-trimer	0.51	HHblits	X-ray	2.96Å	0.36	0.71	glutamate symport protein
3v8f.1.A	33.24	homo-trimer	0.49	HHblits	X-ray	3.80Å	0.36	0.71	sodium-coupled L-aspartate transporter
2nwx.1.A	32.96	homo-trimer	0.49	HHblits	X-ray	3.29Å	0.36	0.71	425aa long hypothetical proton glutamate symport protein
2nww.1.A	32.96	homo-trimer	0.42	HHblits	X-ray	3.20Å	0.36	0.71	425aa long hypothetical proton glutamate symport protein
4oyf.1.C	32.96	homo-hexamer	0.20	HHblits	X-ray	3.41Å	0.36	0.71	GLUTAMATE SYMPORT PROTEIN
4x2s.1.B	36.94	homo-trimer	0.44	BLAST	X-ray	4.21Å	0.38	0.63	425aa long hypothetical proton glutamate symport protein
4x2s.1.A	36.94	homo-trimer	0.44	BLAST	X-ray	4.21Å	0.38	0.63	425aa long hypothetical proton glutamate symport protein
6bat.1.A	37.06	homo-trimer	0.52	BLAST	X-ray	3.40Å	0.38	0.63	Glutamate transporter homolog
6bau.1.A	37.06	homo-trimer	0.51	BLAST	X-ray	3.80Å	0.38	0.63	Glutamate transporter homolog
6bav.1.A	37.06	homo-trimer	0.50	BLAST	X-ray	3.70Å	0.38	0.63	Glutamate transporter homolog
3v8f.1.A	36.62	homo-trimer	0.46	BLAST	X-ray	3.80Å	0.38	0.63	sodium-coupled L-aspartate transporter
5cfy.1.A	36.31	homo-hexamer	0.15	BLAST	X-ray	3.50Å	0.38	0.63	425aa long hypothetical proton glutamate symport protein
4oye.1.A	36.31	homo-hexamer	0.14	BLAST	X-ray	4.00Å	0.38	0.63	425aa long hypothetical proton glutamate symport protein
6bat.1.A	17.07	homo-trimer	0.08	HHblits	X-ray	3.40Å	0.30	0.16	Glutamate transporter homolog
6bau.1.A	14.63	homo-trimer	0.08	HHblits	X-ray	3.80Å	0.30	0.16	Glutamate transporter homolog
6bav.1.A	14.63	homo-trimer	0.06	HHblits	X-ray	3.70Å	0.30	0.16	Glutamate transporter homolog
2nwx.1.A	14.81	homo-trimer	0.02	HHblits	X-ray	3.29Å	0.30	0.16	425aa long hypothetical proton glutamate symport protein
2nww.1.A	14.81	homo-trimer		HHblits	X-ray	3.20Å	0.30	0.16	425aa long hypothetical proton glutamate symport protein
4oyf.1.C	14.81	homo-trimer	0.09	HHblits	X-ray	3.41Å	0.30	0.16	GLUTAMATE SYMPORT PROTEIN

Template	Seq Identity	Oligo-state	QSQE	Found by	Method	Resolution	Seq Similarity	Coverage	Description
4p3j.1.A	12.35	homo-trimer	0.03	HHblits	X-ray	3.50Å	0.29	0.16	GltPh
4p6h.1.A	12.35	homo-trimer	0.03	HHblits	X-ray	4.08Å	0.29	0.16	GltPh

The table above shows the top 30 filtered templates. A further 29 templates were found which were considered to be less suitable for modelling than the filtered list.

4oye.1.A, 5llu.1.A, 4izm.1.A, 4p3j.1.A, 4ky0.1.A, 4ky0.1.C, 3v8f.1.A, 5e9s.1.B, 5e9s.1.A, 2nwl.1.A, 5dwy.1.C, 5dwy.1.A, 2nww.1.A, 4p6h.1.A, 5do7.2.B, 5do7.1.A, 6bau.1.A, 5cfy.1.A, 3v8g.1.A, 3v8g.1.C, 2nwx.1.A, 5llm.1.A, 4oyf.1.C, 3kbc.1.A, 5mju.1.A, 5t0o.1.A, 6bav.1.A, 5t0o.1.C, 6bat.1.A